

ABOUT THIS BOOK

This book introduces matrices through high school coordinate geometry. This approach makes it easier for beginners to learn Python for scientific computing. All problems in the book are from NCERT mathematics textbooks from Class 9-12. Exercises are from CBSE board exam papers.

The content is sufficient for industry jobs and covers nearly all matrix prerequisites for machine learning. There is no copyright, so readers are free to print and share.

License: <https://creativecommons.org/licenses/by-sa/3.0/>
and

<https://www.gnu.org/licenses/fdl-1.3.en.html>

First manual appeared in January 2018

First edition published on July 10, 2024

In this edition, some incorrect solutions were removed. All figures redrawn. More problems added.

CONTENTS

1	GATE CE-1 2026	3
1.1	Formulae	3
1.2	Problems	8

1. GATE CE-1 2026

A. Formulae

1.1.1 Intersection of Conics and Common Chord

- Conic in Vector Form: Every second-degree curve $ax^2 + 2bxy + cy^2 + 2dx + 2ey + f = 0$ can be written in matrix-vector form as:

$$\mathbf{x}^\top \mathbf{V} \mathbf{x} + 2\mathbf{u}^\top \mathbf{x} + f = 0, \quad \mathbf{V} = \begin{pmatrix} a & b \\ b & c \end{pmatrix} = \mathbf{V}^\top, \quad \mathbf{u} = \begin{pmatrix} d \\ e \end{pmatrix} \quad (1.1.1.1)$$

The conic is classified by the determinant of $\mathbf{V}$:

$$|\mathbf{V}| = \begin{vmatrix} a & b \\ b & c \end{vmatrix} = ac - b^2 \begin{cases} = 0, & \text{parabola} \\ > 0, & \text{ellipse} \\ < 0, & \text{hyperbola} \end{cases} \quad (1.1.1.2)$$

For a circle with centre $\mathbf{c}$ and radius r , $\|\mathbf{x} - \mathbf{c}\|^2 = r^2$ gives $\mathbf{V} = \mathbf{I}$, $\mathbf{u} = -\mathbf{c}$, and $f = \|\mathbf{c}\|^2 - r^2$.

- Line-Conic Intersection: The points of intersection of a line $L : \mathbf{x} = \mathbf{h} + \kappa \mathbf{m}$ ($\kappa \in \mathbb{R}$) with a conic section $\mathbf{x}^\top \mathbf{V} \mathbf{x} + 2\mathbf{u}^\top \mathbf{x} + f = 0$ are given by:

$$\mathbf{x}_i = \mathbf{h} + \kappa_i \mathbf{m} \quad (1.1.1.3)$$

where the parameter values κ_i are:

$$\kappa_i = \frac{-\mathbf{m}^\top (\mathbf{V} \mathbf{h} + \mathbf{u}) \pm \sqrt{[\mathbf{m}^\top (\mathbf{V} \mathbf{h} + \mathbf{u})]^2 - g(\mathbf{h}) (\mathbf{m}^\top \mathbf{V} \mathbf{m})}}{\mathbf{m}^\top \mathbf{V} \mathbf{m}} \quad (1.1.1.4)$$

with

$$g(\mathbf{h}) = \mathbf{h}^\top \mathbf{V} \mathbf{h} + 2\mathbf{u}^\top \mathbf{h} + f \quad (1.1.1.5)$$

The line meets the conic at two, one, or no real points according as the discriminant

$$\Delta = [\mathbf{m}^\top (\mathbf{V} \mathbf{h} + \mathbf{u})]^2 - g(\mathbf{h}) (\mathbf{m}^\top \mathbf{V} \mathbf{m}) \quad (1.1.1.6)$$

is positive, zero, or negative.

- Derivation of the Intersection Parameter: Substituting $\mathbf{x} = \mathbf{h} + \kappa \mathbf{m}$ into the conic equation and using symmetry $\mathbf{V} = \mathbf{V}^\top$:

$$(\mathbf{h} + \kappa \mathbf{m})^\top \mathbf{V} (\mathbf{h} + \kappa \mathbf{m}) + 2\mathbf{u}^\top (\mathbf{h} + \kappa \mathbf{m}) + f = 0 \quad (1.1.1.7)$$

$$\kappa^2 (\mathbf{m}^\top \mathbf{V} \mathbf{m}) + 2\kappa \mathbf{m}^\top (\mathbf{V} \mathbf{h} + \mathbf{u}) + g(\mathbf{h}) = 0 \quad (1.1.1.8)$$

When $\Delta = 0$, the two roots coincide and the line is tangent to the conic.

- Pencil of Conics: The intersection of two conics with parameters $(\mathbf{V}_1, \mathbf{u}_1, f_1)$ and $(\mathbf{V}_2, \mathbf{u}_2, f_2)$ lies on every member of the pencil:

$$\mathbf{x}^\top (\mathbf{V}_1 + \mu \mathbf{V}_2) \mathbf{x} + 2(\mathbf{u}_1 + \mu \mathbf{u}_2)^\top \mathbf{x} + (f_1 + \mu f_2) = 0 \quad (1.1.1.9)$$

- Degenerate Conic and Common Chord: The pencil represents a pair of straight

lines (a degenerate conic) if and only if:

$$\begin{vmatrix} \mathbf{V}_1 + \mu \mathbf{V}_2 & \mathbf{u}_1 + \mu \mathbf{u}_2 \\ (\mathbf{u}_1 + \mu \mathbf{u}_2)^\top & f_1 + \mu f_2 \end{vmatrix} = 0 \quad (1.1.1.10)$$

If $\mathbf{V}_1 = \mathbf{V}_2$ (e.g., two circles), choosing $\mu = -1$ removes the second-degree terms, leaving the common chord (radical axis):

$$2(\mathbf{u}_1 - \mathbf{u}_2)^\top \mathbf{x} + (f_1 - f_2) = 0 \quad (1.1.1.11)$$

1.1.2 Line Intersection and Angle Between Vectors

- **Forms of a Line:** A line through point $\mathbf{A}$ with direction vector $\mathbf{m}$ can be written in parametric form or normal form using normal vector $\mathbf{n}$ ($\mathbf{n}^\top \mathbf{m} = 0$):

$$\mathbf{x} = \mathbf{A} + \kappa \mathbf{m}, \quad \mathbf{n}^\top (\mathbf{x} - \mathbf{A}) = 0 \quad (1.1.2.1)$$

In 2D, if $\mathbf{m} = \begin{pmatrix} m_1 \\ m_2 \end{pmatrix}$, then

$$\mathbf{n} = \begin{pmatrix} -m_2 \\ m_1 \end{pmatrix} \quad (1.1.2.2)$$

- **Line Intersection and Solvability:** The lines $\mathbf{x} = \mathbf{A} + \kappa_1 \mathbf{m}_1$ and $\mathbf{x} = \mathbf{B} + \kappa_2 \mathbf{m}_2$ meet where:

$$(\mathbf{m}_1 \quad -\mathbf{m}_2) \begin{pmatrix} \kappa_1 \\ \kappa_2 \end{pmatrix} = \mathbf{B} - \mathbf{A} \quad (1.1.2.3)$$

This system has a unique solution if and only if $|\mathbf{m}_1 \quad \mathbf{m}_2| \neq 0$. If the determinant is zero, the lines are parallel or coincident.

- **Angle Between Vectors:** The angle θ between two vectors $\mathbf{a}$ and $\mathbf{b}$ is given by:

$$\cos \theta = \frac{\mathbf{a}^\top \mathbf{b}}{\|\mathbf{a}\| \|\mathbf{b}\|} \quad (1.1.2.4)$$

Two vectors are orthogonal if and only if $\mathbf{a}^\top \mathbf{b} = 0$.

- **Circle Representation and Chords:** A point on a circle of radius r centered at the origin is $\mathbf{x} = r \begin{pmatrix} \cos \alpha \\ \sin \alpha \end{pmatrix}$ with $\mathbf{e}_\alpha \begin{pmatrix} \cos \alpha \\ \sin \alpha \end{pmatrix}$. The chord length between $\mathbf{x}_\alpha = r \mathbf{e}_\alpha$ and $\mathbf{x}_\beta = r \mathbf{e}_\beta$ satisfies:

$$\mathbf{e}_\alpha^\top \mathbf{e}_\beta = \cos(\alpha - \beta) \quad (1.1.2.5)$$

$$\|\mathbf{x}_\alpha - \mathbf{x}_\beta\|^2 = 2r^2 (1 - \mathbf{e}_\alpha^\top \mathbf{e}_\beta) = 4r^2 \sin^2 \frac{\alpha - \beta}{2} \quad (1.1.2.6)$$

1.1.3 System of Linear Equations and Rouché–Capelli Theorem

- **Consistency and Solvability:** A system

$$\mathbf{A}\mathbf{x} = \mathbf{b}, \quad \mathbf{A} \in \mathbb{R}^{m \times n} \quad (1.1.3.1)$$

is consistent if and only if

$$\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A} \mid \mathbf{b}) = r \quad (1.1.3.2)$$

If $r = n$, the solution is unique; if $r < n$, there exist infinitely many solutions with $n - r$ free variables. If $\mathbf{A} \in \mathbb{R}^{n \times (n-1)}$, the system is consistent only if

$$|\mathbf{A} \mid \mathbf{b}| = 0 \quad (1.1.3.3)$$

- **Homogeneous System and Zero Norm Property:** For $\mathbf{A} \in \mathbb{R}^{2 \times 3}$ with rank 2, the solution of $\mathbf{Ax} = \mathbf{0}$ is given by the cross product of its rows:

$$\mathbf{x} = \kappa(\mathbf{a}_1 \times \mathbf{a}_2) \quad (1.1.3.4)$$

For any vector $\mathbf{v} \in \mathbb{R}^n$,

$$\|\mathbf{v}\|^2 = \mathbf{v}^\top \mathbf{v} = \sum_i v_i^2 = 0 \iff \mathbf{v} = \mathbf{0} \quad (1.1.3.5)$$

1.1.4 Determinants and Cofactor Expansion

- **Expansion Along a Row:** For $\mathbf{A} \in \mathbb{R}^{n \times n}$, expanding along row i with minor M_{ij} (the determinant after deleting row i and column j):

$$|\mathbf{A}| = \sum_{j=1}^n (-1)^{i+j} a_{ij} M_{ij} \quad (1.1.4.1)$$

For a 2×2 matrix:

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc \quad (1.1.4.2)$$

1.1.5 Relation Between Eigenvalues and Trace of a Matrix

- **Characteristic Polynomial and Trace Relation:** The characteristic polynomial of $\mathbf{A} \in \mathbb{R}^{n \times n}$ is

$$P(\lambda) = |\lambda \mathbf{I} - \mathbf{A}| \quad (1.1.5.1)$$

The sum of all eigenvalues equals the trace of the matrix:

$$\sum_{i=1}^n \lambda_i = \sum_{i=1}^n a_{ii} = \text{tr}(\mathbf{A}) \quad (1.1.5.2)$$

A matrix is symmetric if $\mathbf{A}^\top = \mathbf{A}$ and skew-symmetric if $\mathbf{A}^\top = -\mathbf{A}$.

1.1.6 Cholesky Decomposition

- **Factorization:** A symmetric positive definite matrix $\mathbf{A}$ can be factorized as $\mathbf{A} = \mathbf{LL}^\top$ where $\mathbf{L}$ is lower triangular.
- **2×2 Decomposition Form:**

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{12} & a_{22} \end{pmatrix} = \begin{pmatrix} l_{11} & 0 \\ l_{21} & l_{22} \end{pmatrix} \begin{pmatrix} l_{11} & l_{21} \\ 0 & l_{22} \end{pmatrix} = \begin{pmatrix} l_{11}^2 & l_{11}l_{21} \\ l_{11}l_{21} & l_{21}^2 + l_{22}^2 \end{pmatrix} \quad (1.1.6.1)$$

1.1.7 Bayes' Theorem in Vector Form

- Vector Formulation: Given prior vector $\pi = (\Pr(B_1) \dots \Pr(B_n))^\top$ and likelihood vector $\mathbf{l} = (\Pr(E | B_1) \dots \Pr(E | B_n))^\top$:

$$\Pr(E) = \pi^\top \mathbf{l} \quad (1.1.7.1)$$

$$\Pr(B_i | E) = \frac{\pi_i l_i}{\pi^\top \mathbf{l}} \quad (1.1.7.2)$$

1.1.8 Minimization via Completing the Square

- Inequality and Equality Condition: For real a and b :

$$ab - b^2 = \frac{a^2}{4} - \left(b - \frac{a}{2}\right)^2 \leq \frac{a^2}{4} \quad (1.1.8.1)$$

with equality holding if and only if $b = \frac{a}{2}$.

- Zero Integral Property: If $g(x) \geq 0$ is continuous on $[a, b]$ and $\int_a^b g(x) dx = 0$, then $g(x) = 0$ on $[a, b]$.

1.1.9 Laplace Transform and State-Space Form

- Transform Operational Properties:

$$y'(t) \xleftrightarrow{\mathcal{L}} sY(s) - y(0) \quad (1.1.9.1)$$

$$y''(t) \xleftrightarrow{\mathcal{L}} s^2 Y(s) - sy(0) - y'(0) \quad (1.1.9.2)$$

$$e^{at}u(t) \xleftrightarrow{\mathcal{L}} \frac{1}{s-a} \quad (1.1.9.3)$$

- State-Space Representation: Writing $\mathbf{z} = \begin{pmatrix} y \\ y' \end{pmatrix}^\top$, the ODE $y'' + py' + qy = 0$ becomes $\mathbf{z}' = \mathbf{A}\mathbf{z}$ where:

$$\mathbf{A} = \begin{pmatrix} 0 & 1 \\ -q & -p \end{pmatrix} \quad (1.1.9.4)$$

The exponents of the solution are the eigenvalues of $\mathbf{A}$.

1.1.10 Newton-Raphson Method

- First-Order Recurrence: For a differentiable function $f(x)$, the root is iteratively approximated by:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad (1.1.10.1)$$

1.1.11 Polynomial Interpolation (Vandermonde Matrix)

- Matrix Form: The polynomial $P_2(x) = c_0 + c_1x + c_2x^2$ through points (x_i, y_i) for $i = 1, 2, 3$ satisfies:

$$\begin{pmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \end{pmatrix} \begin{pmatrix} c_0 \\ c_1 \\ c_2 \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} \quad (1.1.11.1)$$

1.1.12 Equilibrium of Rigid Bodies in Vector Form

- Force and Moment Equilibrium: For a body in equilibrium under forces $\mathbf{F}_i$ acting

at points $\mathbf{r}_i$:

$$\sum_i \mathbf{F}_i = \mathbf{0}, \quad \sum_i (\mathbf{r}_i - \mathbf{O}) \times \mathbf{F}_i = \mathbf{0} \quad (1.1.12.1)$$

where in the 2D plane, $\mathbf{r} \times \mathbf{F} = r_x F_y - r_y F_x$.

- Bending Moment and Friction: The bending moment at section P is

$$M(P) = \sum_{\text{one side}} (\mathbf{r}_i - \mathbf{P}) \times \mathbf{F}_i \quad (1.1.12.2)$$

Limiting friction force is $F = \mu N$.

1.1.13 Direct Stiffness Method

- Member and Global Stiffness Formulation: For an axial member of length L , area A , modulus E , and direction vector $\mathbf{m}$, the nodal stiffness is

$$\mathbf{k} = \frac{AE}{L} \mathbf{m} \mathbf{m}^\top \quad (1.1.13.1)$$

and the member stiffness matrix is $\begin{pmatrix} \mathbf{k} & -\mathbf{k} \\ -\mathbf{k} & \mathbf{k} \end{pmatrix}$. The global system is

$$\mathbf{K} \mathbf{u} = \mathbf{F} \quad (1.1.13.2)$$

- Thermal Load Equivalence: A temperature rise ΔT with expansion coefficient α produces equivalent nodal loads of $AE\alpha\Delta T$ at member ends directed outward.

1.1.14 Cellular Update Rule in Matrix Form

- Matrix Formulation: For N bulbs in a line with state vector $\mathbf{x}_k \in \{0, 1\}^N$, the number of ON neighbours is given by $\mathbf{s}_k = \mathbf{A} \mathbf{x}_k$ with:

$$a_{ij} = \begin{cases} 1, & |i - j| = 1 \\ 0, & \text{otherwise} \end{cases} \quad (1.1.14.1)$$

1.1.15 Surveying and Traffic Relations

- Reduced Level: With height of instrument HI and staff reading h ,

$$RL = HI - h \quad (1.1.15.1)$$

- Space Mean Speed: For n vehicles covering distance d in times $\mathbf{t} = (t_1 \ \dots \ t_n)^\top$:

$$v_s = \frac{nd}{\mathbf{1}^\top \mathbf{t}} = \frac{n}{\sum_i \frac{1}{v_i}} \quad (1.1.15.2)$$

which is the harmonic mean of the spot speeds $v_i = d/t_i$.

B. Problems

1.1.1 At how many points will the curves $y = x^2$ and $y = -x^2 - 2x - 1$ intersect in the real (x, y) plane?

- a) 0
b) 1
c) 2
d) 3

Solution: Writing both curves in the form (1.1.1.1),

$$x^2 - y = 0 \implies \mathbf{V}_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \mathbf{u}_1 = \begin{pmatrix} 0 \\ -\frac{1}{2} \end{pmatrix}, f_1 = 0 \quad (1.1.1.1)$$

$$x^2 + 2x + y + 1 = 0 \implies \mathbf{V}_2 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \mathbf{u}_2 = \begin{pmatrix} 1 \\ \frac{1}{2} \end{pmatrix}, f_2 = 1 \quad (1.1.1.2)$$

Using (1.1.1.2),

$$|\mathbf{V}_1| = \begin{vmatrix} 1 & 0 \\ 0 & 0 \end{vmatrix} = 0, \quad |\mathbf{V}_2| = \begin{vmatrix} 1 & 0 \\ 0 & 0 \end{vmatrix} = 0 \quad (1.1.1.3)$$

so both curves are parabolas. Any common point lies on every member of the pencil (1.1.1.9):

$$\mathbf{x}^\top \begin{pmatrix} 1+\mu & 0 \\ 0 & 0 \end{pmatrix} \mathbf{x} + 2 \begin{pmatrix} \mu \\ \frac{\mu-1}{2} \end{pmatrix}^\top \mathbf{x} + \mu = 0 \quad (1.1.1.4)$$

The pencil is degenerate when (1.1.1.10) holds. Expanding along the second row using (1.1.4.1) and (1.1.4.2),

$$\begin{vmatrix} 1+\mu & 0 & \frac{\mu}{2} \\ 0 & 0 & \frac{\mu-1}{2} \\ \mu & \frac{\mu-1}{2} & \mu \end{vmatrix} = -\frac{\mu-1}{2} \begin{vmatrix} 1+\mu & 0 \\ \mu & \frac{\mu-1}{2} \end{vmatrix} \quad (1.1.1.5)$$

$$= -\frac{(\mu-1)^2(\mu+1)}{4} = 0 \implies \mu = -1, \mu = 1 \quad (1.1.1.6)$$

For $\mu = -1$, $\mathbf{V}_1 - \mathbf{V}_2 = \mathbf{0}$ and the second-degree terms vanish. From (1.1.1.11),

$$2(\mathbf{u}_1 - \mathbf{u}_2)^\top \mathbf{x} + (f_1 - f_2) = 0 \quad (1.1.1.7)$$

$$2 \begin{pmatrix} -1 & -1 \end{pmatrix} \mathbf{x} - 1 = 0 \implies \begin{pmatrix} 1 & 1 \end{pmatrix} \mathbf{x} = -\frac{1}{2} \quad (1.1.1.8)$$

The normal vector is $\mathbf{n} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$. From (1.1.2.2) and (1.1.2.1), a point on the line and its direction vector are

$$\mathbf{h} = \begin{pmatrix} 0 \\ -\frac{1}{2} \end{pmatrix}, \quad \mathbf{m} = \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \quad \mathbf{x} = \mathbf{h} + \kappa \mathbf{m} \quad (1.1.1.9)$$

Substituting this line into the first parabola using (1.1.1.8) and (1.1.1.5),

$$\mathbf{m}^\top \mathbf{V}_1 \mathbf{m} = \begin{pmatrix} 1 & -1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} = 1 \quad (1.1.1.10)$$

$$\mathbf{V}_1 \mathbf{h} + \mathbf{u}_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ -\frac{1}{2} \end{pmatrix} + \begin{pmatrix} 0 \\ -\frac{1}{2} \end{pmatrix} = \begin{pmatrix} 0 \\ -\frac{1}{2} \end{pmatrix} \quad (1.1.1.11)$$

$$\mathbf{m}^\top (\mathbf{V}_1 \mathbf{h} + \mathbf{u}_1) = \begin{pmatrix} 1 & -1 \end{pmatrix} \begin{pmatrix} 0 \\ -\frac{1}{2} \end{pmatrix} = \frac{1}{2} \quad (1.1.1.12)$$

$$g(\mathbf{h}) = \mathbf{h}^\top \mathbf{V}_1 \mathbf{h} + 2\mathbf{u}_1^\top \mathbf{h} + f_1 = 0 + \frac{1}{2} + 0 = \frac{1}{2} \quad (1.1.1.13)$$

which gives

$$\kappa^2 + \kappa + \frac{1}{2} = 0 \quad (1.1.1.14)$$

The discriminant from (1.1.1.6) is

$$\Delta = [\mathbf{m}^\top (\mathbf{V}_1 \mathbf{h} + \mathbf{u}_1)]^2 - g(\mathbf{h}) (\mathbf{m}^\top \mathbf{V}_1 \mathbf{m}) = \frac{1}{4} - \frac{1}{2} = -\frac{1}{4} < 0 \quad (1.1.1.15)$$

From (1.1.1.4),

$$\kappa = -\frac{1}{2} \pm \frac{j}{2} \quad (1.1.1.16)$$

which is not real (here j is i). As a check, the other degenerate member $\mu = 1$ gives

$$(x^2 - y) + (x^2 + 2x + y + 1) = 2x^2 + 2x + 1 = 0 \implies x = \frac{-1 \pm j}{2} \quad (1.1.1.17)$$

which is again not real. Hence the curves do not meet in the real (x, y) plane and the number of intersection points is 0.

Since PQ is a diameter, $\mathbf{P}$ and $\mathbf{Q}$ are opposite points on the x -axis. Let $\mathbf{R} = r\mathbf{w}_\alpha$ and $\mathbf{S} = r\mathbf{w}_\beta$ with $0 < \beta < \alpha < 180^\circ$. The points are listed in Table 1.1.3.

Point	Vector	Description
O	$\mathbf{0}$	centre
P	$-r\mathbf{i}$	end of diameter
Q	$r\mathbf{i}$	end of diameter
R	$r\mathbf{w}_\alpha$	on the circle
S	$r\mathbf{w}_\beta$	on the circle

TABLE 1.1.3

CONSTRUCTION MATRIX OF THE POINTS

The construction matrix is

$$(\mathbf{P} \quad \mathbf{Q} \quad \mathbf{R} \quad \mathbf{S}) = r \begin{pmatrix} -\mathbf{i} & \mathbf{i} & \mathbf{w}_\alpha & \mathbf{w}_\beta \end{pmatrix} \quad (1.1.3.2)$$

From (1.1.2.5),

$$\cos \angle ROS = \mathbf{w}_\alpha^\top \mathbf{w}_\beta = \cos(\alpha - \beta) \implies \alpha - \beta = 80^\circ \quad (1.1.3.3)$$

The direction vectors of the lines PR and QS are

$$\mathbf{R} - \mathbf{P} = r \begin{pmatrix} 1 + \cos \alpha \\ \sin \alpha \end{pmatrix} = 2r \cos \frac{\alpha}{2} \begin{pmatrix} \cos \frac{\alpha}{2} \\ \sin \frac{\alpha}{2} \end{pmatrix} \quad (1.1.3.4)$$

$$\mathbf{S} - \mathbf{Q} = r \begin{pmatrix} \cos \beta - 1 \\ \sin \beta \end{pmatrix} = 2r \sin \frac{\beta}{2} \begin{pmatrix} -\sin \frac{\beta}{2} \\ \cos \frac{\beta}{2} \end{pmatrix} \quad (1.1.3.5)$$

The point T lies beyond R on PR and beyond S on QS . Hence $\overrightarrow{TR}$ is along $\mathbf{P} - \mathbf{R}$ and $\overrightarrow{TS}$ is along $\mathbf{Q} - \mathbf{S}$. Using (1.1.2.4),

$$\cos \angle RTS = \begin{pmatrix} \cos \frac{\alpha}{2} & \sin \frac{\alpha}{2} \end{pmatrix} \begin{pmatrix} -\sin \frac{\beta}{2} \\ \cos \frac{\beta}{2} \end{pmatrix} \quad (1.1.3.6)$$

$$= \sin \frac{\alpha}{2} \cos \frac{\beta}{2} - \cos \frac{\alpha}{2} \sin \frac{\beta}{2} = \sin \frac{\alpha - \beta}{2} \quad (1.1.3.7)$$

$$= \sin 40^\circ = \cos 50^\circ \quad (1.1.3.8)$$

$$\implies \angle RTS = 50^\circ \quad (1.1.3.9)$$

For a numerical check, take $r = 1$, $\alpha = 130^\circ$, $\beta = 50^\circ$. From (1.1.2.3),

$$(\mathbf{R} - \mathbf{P} \quad -(\mathbf{S} - \mathbf{Q})) \begin{pmatrix} \kappa_1 \\ \kappa_2 \end{pmatrix} = \mathbf{Q} - \mathbf{P} \implies \mathbf{T} = \begin{pmatrix} 0 \\ 2.1445 \end{pmatrix} \quad (1.1.3.10)$$

and (1.1.2.4) gives $\angle RTS = 50^\circ$.




Fig. 1.1.3.1. Circle with diameter PQ , $\angle ROS = 80^\circ$ and $\angle RTS = 50^\circ$

`codes/ce1q7/q07.py`


`codes/ce1q7/q07plot.py`

1.1.4 Consider a linear arrangement of seven bulbs, each of which can be in the ON or OFF states. The initial configuration of the bulbs is shown in the figure. In every Step, the states of the bulbs are changed based on the following rules: Any OFF bulb with exactly one ON neighbor at the end of the previous Step is turned ON. Any ON bulb with both neighbors ON at the end of the previous Step is turned OFF. The state of any bulb not meeting the conditions above is left unchanged. The state of bulbs at the end of Step 1 and Step 2 are also shown in the figure. The number of bulbs which are ON at the end of Step 8 is (GATE CE 2026)


- | | |
|------|------|
| a) 5 | c) 3 |
| b) 4 | d) 0 |

[border=10pt]standalone tikz amsmath

Initial



Step 1 ↓



 ON

Step 2 ↓

 OFF

