

# Preliminary gamified ultrasound visual biofeedback on rhotic syllable tongue movement for speech therapy

Sarah R. Li,<sup>1</sup> Kathryn J. Eary,<sup>1</sup> Sarah Dugan, PhD CCC-SLP,<sup>2,3</sup> Maurice Lamb, PhD,<sup>4</sup> McKenna Becker,<sup>5</sup> Reneé Seward, MGD,<sup>5</sup> Suzanne Boyce, PhD CCC-SLP,<sup>2</sup> Michael A. Riley, PhD,<sup>3</sup> & T. Douglas Mast, PhD<sup>1</sup>

<sup>1</sup>Biomedical Engineering; <sup>2</sup>Communication Sciences and Disorders; <sup>3</sup>Rehabilitation, Exercise, and Nutrition Sciences; <sup>5</sup>Design | University of Cincinnati  
<sup>4</sup>University of Skövde, Sweden

## Ultrasound biofeedback

- Ultrasound imaging provides visual biofeedback of the tongue (Fig. 1) in real time [1], which is particularly useful for refining tongue movement towards a pattern (i.e., a desired speech sound production).
- This refinement is helpful for:
  - Speech sound disorder (SSD) remediation [2]: changing “wabbit” to be the desired “rabbit.”
  - Accent expansion in second language learning [3].
- Understanding tongue movement from ultrasound images can be difficult due to the rapidly moving appearance of the tongue surface, possibly in different directions for different parts of the tongue [4].




Figure 1: Ultrasound imaging of the tongue.

## Tongue movement measurement

- Tongue movement from ultrasound images (Fig. 2) can be automatically measured as tongue part displacement trajectories [5, 6].
- These represent changes in vocal tract constriction due to the root, dorsum, and blade. (Positive displacement = narrowing of the vocal tract.)




Figure 2: Example ultrasound images (left and center) from accurate (top) and misarticulated (bottom) /r/ (“are”) productions, with measured displacement trajectories (right).

## Simple biofeedback parameter

- A linear combination of blade and dorsum displacement can be used as a parameter  $\delta$  (Fig. 3) to represent the accuracy of a production, demonstrated previously for the syllables “ear,” “are,” “booeer,” “reap,” “raw,” and “rue” [5, 6].




Figure 3: Averages (solid lines) and standard deviations (shading) of  $\delta$  in “are” productions, with a discrimination threshold from a receiver operating characteristic curve (ROC).

## Design considerations

- A **gamified** presentation of this biofeedback may help elicit an external focus of attention, demonstrated in non-speech motor control studies to improve motor learning [7].
- **Difficulty level** affects motivation and the amount of information that an individual speaker can learn (see the Challenge Point framework) [8, 9].
- Difficulty level can be changed by adjusting **target width**. Fig. 4 shows a representation of a **target** with a width (between ..... lines) and amplitude (between 0 and .....).
- Fitts’ Law relates index of difficulty (from target width and amplitude) to movement time (i.e., slower movement at harder difficulties retains accuracy) [10].
- **Feedback conditions** (e.g., delayed timing) impact motor learning [8].




Figure 4: Representation of target; accurate /r/ is achieved by reaching the target area. The green arrow (left) shows the target amplitude and the red (right) the target width.

## References

- [1] J. L. Preston et al., “Ultrasound images of the tongue: a tutorial for assessment and remediation of speech sound errors,” *JoVE*, no. 119, pp. e55123, Jan. 2017, doi: 10.3791/55123.
- [2] E. Sugden, S. Lloyd, J. Lam, and J. Cleland, “Systematic review of ultrasound visual biofeedback in intervention for speech sound disorders,” *Int J Lang Commun*, vol. 54, no. 5, pp. 705–728, Sep. 2019, doi: 10.1111/1460-6984.12478.
- [3] T. L. Face, “Ultimate attainment of Spanish rhotics by native English-speaking immigrants to Spain,” *Lengua y Migracion*, vol. 10, pp. 57–80, Jan. 2018.
- [4] S. Boyce, “The articulatory phonetics of /r/ for residual speech errors,” *Semin Speech Lang*, vol. 36, no. 04, pp. 257–270, Oct. 2015, doi: 10.1055/s-0035-1562909.
- [5] S. R. Li et al., “An automatic, simple ultrasound biofeedback parameter for distinguishing accurate and misarticulated rhotic syllables,” in *Proc Interspeech 2021*, 636–640, doi: 10.21437/Interspeech.2021-1749.
- [6] S. R. Li et al., “Classification of accurate and misarticulated /ar/ for ultrasound biofeedback using tongue part displacement trajectories,” *Clin Linguist Phon*, pp. 1–27, Mar. 2022, doi: 10.1080/02699206.2022.2039777.
- [7] G. Wulf, C. Shea, and R. Lewthwaite, “Motor skill learning and performance: a review of influential factors: Motor skill learning and performance,” *Med Educ*, vol. 44, no. 1, pp. 75–84, Jan. 2010, doi: 10.1111/j.1365-2923.2009.03421.x.
- [8] E. Maas et al., “Principles of Motor Learning in Treatment of Motor Speech Disorders,” *Am J Speech Lang Pathol*, vol. 17, no. 3, pp. 277–298, Aug. 2008, doi: 10.1044/1058-0360(2008/025).
- [9] McAllister Tara, Hitchcock Elaine R., and Ortiz José A., “Computer-Assisted Challenge Point Intervention for Residual Speech Errors,” *Perspect ASHA Spec Interest Groups*, vol. 6, no. 1, pp. 214–229, Feb. 2021, doi: 10.1044/2020\_PERSP-20-00191.
- [10] S. R. Kuberski and A. I. Gafos, “Fitts’ Law in Tongue Movements of Repetitive Speech,” *Phonetica*, 2019, doi: 10.1159/000501644.

## STAR and GO, GO Goat! games




Figure 5: TonguePART Graphical user interface (GUI) for measuring tongue part displacement trajectories and STAR game (center). The STAR game shows the  $\delta$  parameter during all the frames of a production as a moving star, with the representation of the target (here, the red and green dashed lines above the star) directly compared to the  $\delta$  parameter.

- Both games: are audio activated, use the simple parameter  $\delta$



Figure 6: GO, GO, Goat! game. The green arrow (top) shows the mapping of target amplitude and the red (bottom) that of target width. After a production finishes, the goat jumps; where this goat lands is the  $\delta$  parameter at one frame (estimated midpoint; i.e., calculated from the displacements between the initial and midpoint frames) of a production linearly scaled to the representation of the target as platforms in the game.

Table: Differences in feedback conditions between the STAR and GO, GO, Goat! games.

| STAR                           | GO, GO Goat!                   |
|--------------------------------|--------------------------------|
| • Concurrent                   | • Delayed                      |
| • Greater amount (more frames) | • Lesser amount (fewer frames) |
| • Direct target                | • Scaled target                |

## Pilot data (two participants)

- One participant with typical speech showed a change in articulation (Fig. 7).
- Participants found both games likeable, with the STAR game liked slightly better.

**Conclusion:** These gamified biofeedback interfaces allow for testing of motor learning principles and have potential use for speech therapy.

Figure 7: Comparison of tongue surface contours at end (/r/) of productions from an adult participant with typical speech, with the task of reaching a specified target for /r/ (“are”). Averages (solid lines) and standard deviations (shading) are shown for a selection of productions. Contours were rotated so that mandible shadows on the images matched, and interpolation was used to standardize the horizontal sampling.

