

# Contents

<b>I</b>	<b>Multivariable Calculus</b>	<b>1</b>
<b>11</b>	<b>Parametric Equations and Polar Coordinates</b>	<b>2</b>
11.1	Curves Defined by Parametric Equations . . . . .	2
11.2	Calculus with Parametric Curves . . . . .	3
11.3	Polar Coordinates . . . . .	9
11.4	Areas and Lengths in Polar Coordinates . . . . .	9
11.5	Conic Sections . . . . .	9
11.6	Conic Sections in Polar Coordinates . . . . .	9
<b>12</b>	<b>Infinite Sequences and Series</b>	<b>10</b>
12.1	Sequences . . . . .	11
12.2	Series . . . . .	11
12.3	The Integral Test and Estimates of Sums . . . . .	11
12.4	The Comparison Tests . . . . .	11
12.5	Alternating Series . . . . .	11
12.6	Absolute Convergence and the Ratio and Root Tests . . . . .	11
12.7	Strategy for Testing Series . . . . .	11
12.8	Power Series . . . . .	11
12.9	Representation of Functions as Power Series . . . . .	11
12.10	Taylor and Maclaurin Series . . . . .	11
12.11	The Binomial Series . . . . .	11
12.12	Applications of Taylor Polynomials . . . . .	11
<b>13</b>	<b>Vectors and the Geometry of Space</b>	<b>12</b>
13.1	Three-Dimensional Coordinate Systems . . . . .	12
13.2	Vectors . . . . .	12
13.3	The Dot Product . . . . .	12
13.4	The Cross Product . . . . .	12
13.5	Equations of Lines and Planes . . . . .	12
13.6	Cylinders and Quadric Surfaces . . . . .	12
13.7	Cylindrical and Spherical Coordinates . . . . .	12
<b>14</b>	<b>Vector Functions</b>	<b>13</b>
14.1	Vector Functions and Space Curves . . . . .	13
14.2	Derivatives and Integrals of Vector Functions . . . . .	13

14.3	Arc Length and Curvature . . . . .	13
14.4	Motion in Space: Velocity and Acceleration . . . . .	13
<b>15</b>	<b>Partial Derivatives</b>	<b>14</b>
15.1	Functions of Several Variables . . . . .	14
15.2	Limits and Continuity . . . . .	14
15.3	Partial Derivatives . . . . .	14
15.4	Tangent Planes and Linear Approximations . . . . .	14
15.5	The Chain Rule . . . . .	14
15.6	Directional Derivatives and the Gradient Vector . . . . .	14
15.7	Maximum and Minimum Values . . . . .	14
15.8	Lagrange Multipliers . . . . .	14
<b>16</b>	<b>Multiple Integrals</b>	<b>15</b>
16.1	Double Integrals over Rectangles . . . . .	15
16.2	Iterated Integrals . . . . .	15
16.3	Double Integrals over General Regions . . . . .	15
16.4	Double Integrals in Polar Coordinates . . . . .	15
16.5	Applications of Double Integrals . . . . .	15
16.6	Surface Area . . . . .	15
16.7	Triple Integrals . . . . .	15
16.8	Triple Integrals in Cylindrical and Spherical Coordinates . . . . .	15
16.9	Change of Variables in Multiple Integrals . . . . .	15
<b>17</b>	<b>Vector Calculus</b>	<b>16</b>
17.1	Vector Fields . . . . .	16
17.2	Line Integrals . . . . .	16
17.3	The Fundamental Theorem for Line Integrals . . . . .	16
17.4	Green's Theorem . . . . .	16
17.5	Curl and Divergence . . . . .	16
17.6	Parametric Surfaces and Their Areas . . . . .	16
17.7	Surface Integrals . . . . .	16
17.8	Stokes' Theorem . . . . .	16
17.9	The Divergence Theorem . . . . .	16
17.10	Summary . . . . .	16
<b>18</b>	<b>Second-Order Differential Equations</b>	<b>17</b>
18.1	Second-Order Linear Equations . . . . .	17
18.2	Nonhomogenous Linear Equations . . . . .	17
18.3	Applications of Second-Order Differential Equations . . . . .	17
18.4	Series Solutions . . . . .	17

**Part I**

**Multivariable Calculus**

## Chapter 11

# Parametric Equations and Polar Coordinates

### 11.1 Curves Defined by Parametric Equations

Suppose that  $x$  and  $y$  are both given as functions of a third variable  $t$  (called a **parameter** by the equations)

$$x = f(t) \quad y = g(t)$$


(called **parametric equations**). Each value of  $t$  determines a point  $(x,y)$ . As  $t$  changes,  $(x,y) = (f(t), g(t))$  changes and traces out a curve  $C$ , which is called a **parametric curve**. The direction of the arrows on curve  $C$  show the change in the position of the equation as  $t$  increases.

We can also restrict  $t$  to a finite interval. In general, the curve with parametric equations

$$x = f(t) \quad y = g(t) \quad a \leq t \leq b$$


has **initial point**  $(f(a), g(a))$  and **terminal point**  $(f(b), g(b))$ .

#### The Cycloid



**Example 11.1.1.** A circle with radius  $r$  rolls along the  $x$ -axis. The curve traced out by a point  $P$  on the circumference of the circle is called a **cycloid**. Find parametric equations for the cycloid.

*Solution.* We will use the angle of rotation  $\theta$  as the parameter ( $\theta = 0$  when  $P$  is at the origin).



Suppose the circle has rotated  $\theta$  radians. Using the figure, the distance it has rolled from the origin is

$$|OT| = \text{arc } PT = r\theta$$

because  $P$  starts at the origin. Therefore, the center of the circle is  $C(r\theta, r)$ . Let the coordinates of  $P$  be  $(x, y)$ . Then from the figure,

$$x = |OT| - |PQ| = r\theta - r\sin\theta = r(\theta - \sin\theta)$$

$$y = |TC| - |QC| = r - r\cos\theta = r(1 - \cos\theta)$$

**Definition 11.1.1.** Parametric equations of the cycloid are

$$x = r(\theta - \sin\theta) \quad y = r(1 - \cos\theta)$$

## 11.2 Calculus with Parametric Curves

We will mainly solve problems involving tangents, area, arc length, and surface area.

### Tangents

In the previous section, we saw that some curves defined by parametric equations  $x = f(t)$  and  $y = g(t)$  can also be expressed, by eliminating the parameter, in the form  $y = F(x)$ . If we substitute  $x = f(t)$  and  $y = g(t)$  in the equation  $y = F(x)$ , we get

$$g(t) = F(f(t))$$

If  $g$ ,  $f$ , and  $F$  are differentiable, the Chain Rule gives

$$g'(t) = F'(f(t))f'(t) = F'(x)f'(t)$$

If  $f'(t) \neq 0$ , we can solve for  $F'(x)$ :

**Definition 11.2.1.** The slope of the tangent to the parametric curve  $y = F(x)$  is  $F'(x)$ .

$$F'(x) = \frac{g'(t)}{f'(t)}$$

This enables us to find tangents to parametric curves without having to eliminate the parameter. We can rewrite the previous equation in an easily remembered form.

**Definition 11.2.2.** We can use this to find tangents to parametric curves without having to eliminate the parameter.

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} \quad \text{if} \quad \frac{dx}{dt} \neq 0$$

The curve has a

- horizontal tangent when  $\frac{dy}{dt} = 0$  (provided that  $\frac{dx}{dt} \neq 0$ )
- vertical tangent when  $\frac{dx}{dt} = 0$  (provided that  $\frac{dy}{dt} \neq 0$ )

This is useful when sketching parametric curves.

**Definition 11.2.3.** We can also find  $\frac{d^2y}{dx^2}$  by replacing  $y$  with  $\frac{dy}{dx}$

$$\frac{d^2y}{dx^2} = \frac{d}{dx} \left( \frac{dy}{dx} \right) = \frac{\frac{d}{dt} \left( \frac{dy}{dx} \right)}{\frac{dx}{dt}}$$

*Proof.* Find  $\frac{dy}{dx}$  and  $\frac{d^2y}{dx^2}$  considering  $y(t)$  and  $g(t)$ .

1.

$$\text{Chain rule: } \frac{dy}{dt} = \frac{dy}{dx} \frac{dx}{dt} \implies \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} \quad (\implies \text{ means "implies" })$$

2.

$$\text{Chain rule: } \frac{d}{dt} \left( \frac{dy}{dx} \right) = \left( \frac{d}{dx} \frac{dy}{dx} \right) \frac{dx}{dt} = \frac{d^2y}{dx^2} \frac{dx}{dt}$$

$$\text{Substitute: } \frac{d}{dt} \left( \frac{dy}{dx} \right) = \frac{d}{dt} \left( \frac{\frac{dy}{dt}}{\frac{dx}{dt}} \right)$$

$$\text{Quotient rule: } = \frac{\frac{d^2y}{dt^2} \frac{dx}{dt} - \frac{dy}{dt} \frac{d^2x}{dt^2}}{\left( \frac{dx}{dt} \right)^2}$$

Set equation from line 1 and line 3 equal and divide both sides by  $\frac{dx}{dt}$

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{\frac{d^2y}{dt^2} \frac{dx}{dt} - \frac{dy}{dt} \frac{d^2x}{dt^2}}{\left( \frac{dx}{dt} \right)^2 \left( \frac{dx}{dt} \right)} \\ &= \frac{\frac{d^2y}{dt^2} \frac{dx}{dt} - \frac{dy}{dt} \frac{d^2x}{dt^2}}{\left( \frac{dx}{dt} \right)^3} \end{aligned}$$

□

**Example 11.2.1.** A curve C is defined by the parametric equations  $x = t^2$ ,  $y = t^3 - 3t$ .

1. Show that C has two tangents at the point (3,0) and find their equations.
2. Find the points on C where the tangent is horizontal or vertical.
3. Determine where the curve is concave upward or downward.

*Solution.* A curve C is defined by the parametric equations  $x = t^2$ ,  $y = t^3 - 3t$ .

1. Rewrite  $y = t^3 - 3t = t(t^2 - 3) = 0$  when  $t = 0$  or  $t = \pm\sqrt{3}$ . This indicates that C intersects itself at (3,0).

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{3t^2 - 3}{2t} = \frac{3}{2} \left( t - \frac{1}{t} \right)$$

$$t = \pm\sqrt{3} \rightarrow dy/dx = \pm 6/(2\sqrt{3})$$

so the equations of the tangents at (3,0) are

$$y = \sqrt{3}(x - 3) \quad \text{and} \quad y = -\sqrt{3}(x - 3)$$

2.  $C$  has a horizontal tangent when  $dy/dx = 0$ . In other words, when  $dy/dt = 0$  and  $dx/dt \neq 0$ .  $dy/dt = 3t^2 - 3 = 0$  when  $t^2 = 1$  so  $t = \pm 1$ . This means there are horizontal tangents on  $C$  at  $(1, -2)$  and  $(-1, 2)$ .  $C$  has a vertical tangent when  $dx/dt = 2t = 0$ , so  $t = 0$ . This means  $C$  has a vertical tangent at  $(0, 0)$ .
3. To determine concavity we calculate the second derivative:

$$\frac{d^2y}{dx^2} = \frac{\frac{d}{dt}\left(\frac{dy}{dx}\right)}{\frac{dx}{dt}} = \frac{\frac{3}{2}\left(1 + \frac{1}{t^2}\right)}{2t} = \frac{3(t^2 + 1)}{4t^3}$$

The curve is concave upward when  $t > 0$  and concave downward when  $t < 0$ .

## Area

We already know that area under a curve  $y = F(x)$  from  $a$  to  $b$  is  $A = \int_a^b F(x)dx$ . We can apply this to parametric equations using the Substitution Rule for Definite Integrals.

**Definition 11.2.4.** If the curve  $C$  is given by parametric equations  $x = f(t)$  and  $y = g(t)$  and  $t$  increases from  $\alpha$  to  $\beta$ ,

$$A = \int_a^b ydx = \int_{\alpha}^{\beta} g(t)f'(t)dt$$

(Switch  $\alpha$  to  $\beta$  if the point on  $C$  at  $\beta$  is more left than  $\alpha$ .)

**Example 11.2.2.** Find the area under one arch of the cycloid  $x = r(\theta - \sin \theta)$ ,  $y = r(1 - \cos \theta)$ .

*Solution.* One arch of the cycloid is given by  $0 \leq \theta \leq 2\pi$ . Using the Substitution Rule with  $y = r(1 - \cos \theta)$  and  $dx = r(1 - \cos \theta)d\theta$ , we have

$$\begin{aligned} A &= \int_0^{2\pi} ydx = \int_0^{2\pi} r(1 - \cos \theta)r(1 - \cos \theta)d\theta \\ &= r^2 \int_0^{2\pi} (1 - \cos \theta)^2 d\theta = r^2 \int_0^{2\pi} (1 - 2\cos \theta + \cos^2 \theta) d\theta \\ &= r^2 \int_0^{2\pi} \left[ 1 - 2\cos \theta + \frac{1}{2}(1 + \cos 2\theta) \right] d\theta \\ &= r^2 \left[ \frac{3}{2}\theta - 2\sin \theta + \frac{1}{4}\sin 2\theta \right]_0^{2\pi} \\ &= r^2 \left( \frac{3}{2} \cdot 2\pi \right) = 3\pi r^2 \end{aligned}$$



## Arc Length

We already know how to find length  $L$  of a curve  $C$  given in the form  $y = F(x)$ ,  $a \leq x \leq b$ .

**Definition 11.2.5.** If  $F'$  is continuous, then

$$L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

If  $C$  can describe the parametric equations  $x = f(t)$  and  $y = g(t)$ ,  $\alpha \leq t \leq \beta$ , where  $dx/dt = f'(t) > 0$ . Using the substitution rule, we obtain

$$L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_\alpha^\beta \sqrt{1 + \left(\frac{dy/dt}{dx/dt}\right)^2} \frac{dx}{dt} dt$$

Since  $dx/dt > 0$ , we have

**Theorem 11.1.** If a curve  $C$  is described by the parametric equations  $x = f(t)$ ,  $y = g(t)$ ,  $\alpha \leq t \leq \beta$ , where  $f'$  and  $g'$  are continuous on  $[\alpha, \beta]$  and  $C$  is traversed exactly once as  $t$  increases from  $\alpha$  to  $\beta$ , then the length of  $C$  is

$$L = \int_\alpha^\beta \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

This is consistent with the general formula  $L = \int ds$  and  $(ds^2) = (dx^2) + (dy^2)$ .

*Proof.* Prove the length formula of a parametric curve

$$\vec{ds} = \vec{i} dx + \vec{j} dy$$

$$ds^2 = \vec{ds} \cdot \vec{ds} = \left(\vec{i} dx + \vec{j} dy\right) \cdot \left(\vec{i} dx + \vec{j} dy\right) = dx^2 + dy^2$$

$$ds = \sqrt{dx^2 + dy^2} = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$L = \int_\alpha^\beta ds = \int_\alpha^\beta \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

**Example 11.2.3.** Find the length of the unit circle as  $(x, y)$  moves both once and twice around the circle.

*Solution.* For one traversal around the unit circle,

$$x = \cos t \quad y = \sin t \quad 0 \leq t \leq 2\pi$$

so  $dx/dt = -\sin t$  and  $dy/dt = \cos t$

$$\begin{aligned} L &= \int_0^{2\pi} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \int_0^{2\pi} \sqrt{\sin^2 t + \cos^2 t} dt \\ &= \int_0^{2\pi} dt = 2\pi \end{aligned}$$

For two traversals around the unit circle,

$$x = \sin 2t \quad y = \cos 2t \quad 0 \leq t \leq 2\pi$$

so  $dx/dt = 2 \cos 2t$  and  $dy/dt = -2 \sin 2t$

$$L = \int_0^{2\pi} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \int_0^{2\pi} \sqrt{4 \cos^2 2t + 4 \sin^2 2t} dt = \int_0^{2\pi} 2 dt = 4\pi$$

## Surface Area

We can also adapt the surface area formula to a parametric curve.

**Definition 11.2.6.** If a curve  $C$  is described by the parametric equations  $x = f(t)$ ,  $y = g(t)$ ,  $\alpha \leq t \leq \beta$ , is rotated about the **x-axis**, where  $f', g'$  are continuous and  $g(t) \geq 0$ , the surface area is

$$S = \int_{\alpha}^{\beta} 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

If the curve  $C$  is rotated about the **y-axis**, the surface area is

$$S = \int_{\alpha}^{\beta} 2\pi x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

The generic formulas  $S = \int 2\pi y ds$  for rotation about the x-axis and  $S = \int 2\pi x ds$  for rotation about the y-axis are still valid, but for parametric curves we use

$$ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

**Example 11.2.4.** Show that the surface area of a sphere of radius  $r$  is  $4\pi r^2$

*Solution.* The sphere is obtained by rotating the semicircle

$$x = r \cos t \quad y = r \sin t \quad 0 \leq t \leq \pi$$

about the x-axis.

$$\begin{aligned} S &= \int_0^{\pi} 2\pi r \sin t \sqrt{(-r \sin t)^2 + (r \cos t)^2} dt \\ &= 2\pi \int_0^{\pi} r \sin t \sqrt{r^2(\sin^2 t + \cos^2 t)} dt \\ &= 2\pi \int_0^{\pi} r \sin t \cdot r dt = 2\pi r^2 \int_0^{\pi} \sin t dt \\ &= 2\pi r^2 (-\cos t) \Big|_0^{\pi} = 4\pi r^2 \end{aligned}$$

**11.3 Polar Coordinates**

**11.4 Areas and Lengths in Polar Coordinates**

**11.5 Conic Sections**

**11.6 Conic Sections in Polar Coordinates**



## Chapter 12

# Infinite Sequences and Series

12.1 Sequences

12.2 Series

12.3 The Integral Test and Estimates of Sums

12.4 The Comparison Tests

12.5 Alternating Series

12.6 Absolute Convergence and the Ratio and Root Tests

12.7 Strategy for Testing Series

12.8 Power Series

12.9 Representation of Functions as Power Series

12.10 Taylor and Maclaurin Series

12.11 The Binomial Series

12.12 Applications of Taylor Polynomials

## Chapter 13

# Vectors and the Geometry of Space

13.1 Three-Dimensional Coordinate Systems

13.2 Vectors

13.3 The Dot Product

13.4 The Cross Product

13.5 Equations of Lines and Planes

13.6 Cylinders and Quadric Surfaces

13.7 Cylindrical and Spherical Coordinates

## Chapter 14

# Vector Functions

14.1 Vector Functions and Space Curves

14.2 Derivatives and Integrals of Vector Functions

14.3 Arc Length and Curvature

14.4 Motion in Space: Velocity and Acceleration

## Chapter 15

# Partial Derivatives

15.1 Functions of Several Variables

15.2 Limits and Continuity

15.3 Partial Derivatives

15.4 Tangent Planes and Linear Approximations

15.5 The Chain Rule

15.6 Directional Derivatives and the Gradient  
Vector

15.7 Maximum and Minimum Values

15.8 Lagrange Multipliers



## Chapter 16

# Multiple Integrals

16.1 Double Integrals over Rectangles

16.2 Iterated Integrals

16.3 Double Integrals over General Regions

16.4 Double Integrals in Polar Coordinates

16.5 Applications of Double Integrals

16.6 Surface Area

16.7 Triple Integrals

16.8 Triple Integrals in Cylindrical and Spherical Coordinates

16.9 Change of Variables in Multiple Integrals

## Chapter 17

# Vector Calculus

17.1 Vector Fields

17.2 Line Integrals

17.3 THE Fundamental Theorem for Line Integrals

17.4 Green's Theorem

17.5 Curl and Divergence

17.6 Parametric Surfaces and Their Areas

17.7 Surface Integrals

17.8 Stokes' Theorem

17.9 The Divergence Theorem

17.10 Summary

## Chapter 18

# Second-Order Differential Equations

18.1 Second-Order Linear Equations

18.2 Nonhomogenous Linear Equations

18.3 Applications of Second-Order Differential Equations

18.4 Series Solutions