

Patient Name : T GEETHA SAGARI

Age / Gender : 24 years / Female

Patient ID : 21496

Source : DIRECT

Referral : Dr V. MADHAVA RAO

Collection Time : Dec 04, 2025, 06:13 a.m.

Receiving Time : Dec 04, 2025, 06:22 a.m.

Reporting Time : Dec 04, 2025, 08:49 a.m.

Sample ID :



1253380004

Test Description	Result(s)	Biological Reference Intervals	Unit(s)
<u>COMPLETE URINE EXAMINATION</u>			
Colour	Pale yellow	Pale yellow	
(Sample Type:Urine)			
pH	6.0	4.5 – 8.5	
Specific Gravity	1.025	1.015 – 1.025	
Glucose	Negative	Negative	
Protein	Negative	Negative	
Ketones	Negative	Negative	
Bilirubins	Negative	Negative	
Nitrite	Negative	Negative	
Blood	Negative	Negative	
Urobilinogen	Negative	Negative	
Leukocytes	Negative	Negative	
<u>MICROSCOPIC EXAMINATION</u>			
Pus cells (WBCs)	2 - 3	0 - 2/hpf	/hpf
Epithelial cells	5 - 6	0 - 4/hpf	/hpf
Red blood cells	Nil	0 - 2/hpf	/hpf
Cast	Nil	Nil	
Crystals	Nil	Nil	
Other	Nil	Nil	
Note			
<u>Electrolytes</u>			
Sodium	136.1	136 - 145	mmol/L
Method: ISE Direct(Sample Type:Serum)			
Potassium	4.34	3.3 - 5.1	mmol/L
Method: ISE Direct(Sample Type:Serum)			
Ionised calcium	4.96	4.5 - 5.3	mg/dL
Method: ISE Direct(Sample Type:Serum)			
Chloride	100.1	90 - 110	mmol/L
Method: ISE Direct(Sample Type:Serum)			
Instrument:Test Done On Fully Automated Biochemistry Analyser, Sensa Core ST-200			
Note			
<u>CREATININE</u>			
CREATININE	0.58	0.5 - 0.9	mg/dL
Method:JAFTE COMPENSATED (Sample Type:Serum)			

Patient Name : T GEETHA SAGARI

Age / Gender : 24 years / Female

Patient ID : 21496

Source : DIRECT

Referral : Dr V. MADHAVA RAO

Collection Time : Dec 04, 2025, 06:13 a.m.

Receiving Time : Dec 04, 2025, 06:22 a.m.

Reporting Time : Dec 04, 2025, 08:49 a.m.

Sample ID :



1253380004

Test Description	Result(s)	Biological Reference Intervals	Unit(s)
------------------	-----------	--------------------------------	---------

Instrument: Test Done On Fully Automated Biochemistry Analyser, BIOSYSTEMS BA 400

Note

URINE ALBUMIN/CREATININE RATIO

*URINE ALBUMIN Latex Method(Sample Type:Urine)	1.63	0 - 15	mg/L
*URINE CREATININE Jaffe's Compensated(Sample Type:Urine)	51.33	37 - 250	mg/dL
*URINE ALBUMIN/CREATININE RATIO calculated	3.18	0 - 30	mg/gram creatinine

Instrument: Tests performed on Biosystems BA 400, A Fully Automatic Biochemistry analyser

Normal: 0 - 30

Microalbuminuria: 30 - 300 mg/gram creatinine

Macroalbuminuria : > 300 mg/gram creatinine

Note

* Marked parameter are not under NABL Accreditation.

POST PRANDIAL PLASMA GLUCOSE

Post Prandial Plasma Glucose Method : Glucose oxidase/Peroxidase	148.53	< 180	mg/dL
---	--------	-------	-------

Instrument: Done on BIOSYSTEMS BA 400, a Fully Automated Biochemical Analyser

Note

ESR

*ESR Method	40	0 - 20	mm/Hour
----------------	----	--------	---------

Interpretation:

- It indicates presence and intensity of an inflammatory process. It does not diagnose a specific disease. Changes in the ESR are more significant than the abnormal results of a single test.
- It is a prognostic test and used to monitor the course or response to treatment of diseases like tuberculosis, bacterial endocarditis, acute rheumatic fever, rheumatoid arthritis, SLE, Hodgkins disease, temporal arteritis and polymyalgia rheumatica.
- It is also increased in pregnancy, multiple myeloma, menstruation, and hypothyroidism.

Note

* Marked parameter are not under NABL Accreditation.

Patient Name : T GEETHA SAGARI

Age / Gender : 24 years / Female

Patient ID : 21496

Source : DIRECT

Referral : Dr V. MADHAVA RAO

Collection Time : Dec 04, 2025, 06:13 a.m.

Receiving Time : Dec 04, 2025, 06:22 a.m.

Reporting Time : Dec 04, 2025, 08:49 a.m.

Sample ID :



1253380004

Test Description	Result(s)	Biological Reference Intervals	Unit(s)
VITAMIN B12			
*Vitamin B12	201	222-1439	pg/mL
CLIA			
Method : CLIA ChemiLuminescence Immunoassay			

Method: ChemiLuminescence Immunoassay(Serum)

Interpretation:

1. Vit B12 levels are decreased in megaloblastic anemia, partial/total gastrectomy, pernicious anemia, peripheral neuropathies, chronic alcoholism, senile dementia, and treated epilepsy.
2. An associated increase in homocysteine levels is an independent risk marker for cardiovascular disease and deep vein thrombosis.
3. HoloTranscobalamin II levels are a more accurate marker of active VitB12 component.

Note: Patients on Biotin supplement may have interference in some immunoassays. With individuals taking high dose Biotin (more than 5 mg per day) supplements, at least 8-hour wait time before blood draw is recommended.

Ref: Arch Pathol Lab Med—Vol 141, November 2017

Note

* marked parameter are not under NABL Accreditation.

Thyroid Function Test			
*T3	106.28	87 - 178	ng/dL
*T4	16.50	4.82 - 15.65	µg/dL
*TSH	0.760	0.38 - 5.33	µIU/mL

Method: Chemi Luminescence Immunoassay(Sample Type:Serum)

Interpretation

- Measurement of TSH in conjunction with T3,T4 or FT3,FT4 is used for assessment of thyroid gland functional status and also for monitoring patients on treatment for hyperthyroidism & hypothyroidism. Suppressed TSH (<0.01 µIU/mL) suggests a diagnosis of hyperthyroidism and elevated concentration (>7 µIU/mL) suggest hypothyroidism. TSH levels may be affected by acute illness and several medications including dopamine and glucocorticoids. TSH concentration are decreased (low or undetectable) in Graves disease. Increased in TSH secreting pituitary adenoma (secondary hyperthyroidism); PPTH. TSH concentration are primary elevated in hypothyroidism (along with decreased T4) except for pituitary & hypothalamic disease.
- Mild to modest TSH elevations in patient with normal T3 & T4 levels indicates impaired thyroid hormone reserves & incipient hypothyroidism (subclinical hypothyroidism).
- Mild to modest decrease in TSH with normal T3 & T4 indicates subclinical hyperthyroidism.
- Degree of TSH suppression does not reflect the severity of hyperthyroidism, therefore, measurement of free thyroid hormone levels is required in patient with a suppressed TSH level.
- If free T4 is normal, free T3 should be checked as it is the first hormone to increase in early hyperthyroidism. Though TSH levels can also be used to effectively monitor patients being treated with thyroid hormones but it may be misleading. Therefore total or free T4 generally serve as front line assay during this period. The free T3 & T4 (FT3 & FT4) measures concentrations of free hormones, which are not affected by changing in concentrations of binding proteins, therefore more reliable indicator of true thyroid status.

Patient Name : T GEETHA SAGARI

Age / Gender : 24 years / Female

Patient ID : 21496

Source : DIRECT

Referral : Dr V. MADHAVA RAO

Collection Time : Dec 04, 2025, 06:13 a.m.

Receiving Time : Dec 04, 2025, 06:22 a.m.

Reporting Time : Dec 04, 2025, 08:49 a.m.

Sample ID :



1253380004

Test Description	Result(s)	Biological Reference Intervals	Unit(s)
------------------	-----------	--------------------------------	---------

Note:* marked parameter are not under NABL Accreditation.

HbA1c

HbA1c Method:HPLC (BIORAD D 10)(Sample Type:Blood)	6.6	Non Diabetic : 4.0 - 5.6 % Border line : 5.7 - 6.4 % Good control in Diabetes : < 6 Years : < 8.5 % 6 Years - 12 Years : < 8.0 % 13 Years - 18 Years : < 7.5 % > 18 Years : < 7 %	%
Estimated Average Glucose (eAG) Calculated	142.72		mg/dL

Interpretation :

- HbA1c is used for monitoring diabetic control. It reflects the estimated average glucose (eAG).
- HbA1c has been endorsed by clinical groups & ADA (American Diabetes Association) guidelines 2012, for diagnosis of diabetes using a cut-off point of 6.5%. ADA defined biological reference range for HbA1c is 4% to 5.7%. Patients with HbA1c value between 5.7% to 6.5% are considered Pre-diabetic.
- Trends in HbA1c are a better indicator of diabetic control than a solitary test.
- Low glycated haemoglobin(below 4%) in a non-diabetic individual are often associated with systemic inflammatory diseases,chronic anaemia(especially severe iron deficiency &haemolytic), chronic renal failure and liver diseases. Clinical correlation suggested.
- To estimate the eAG from the HbA1C value, the following equation is used: $eAG(mg/dl) = 28.7 \times A1c - 46.7$
- Interference of Haemoglobinopathies in HbA1c estimation.

A. For HbF > 25%, an alternate platform (Fructosamine) is recommended for testing of HbA1c.

B. Homozygous hemoglobinopathy is detected, fructosamine is recommended for monitoring diabetic status

C. Heterozygous state detected is corrected for HbS and HbC trait.

Hemoglobin electrophoresis (HPLC method) is recommended for detecting hemoglobinopathy.

Note

LIPID PROFILE

TOTAL CHOLESTEROL Method:Cholestrol Oxidase/Peroxidase(Sample Type:Serum)	164.35	Desirable : Up to 200 mg/dL Borderline High: 200-239 mg/dL High : > 240 mg/dL	mg/dL
TRIGLYCERIDES Method:Glycerol phosphate oxidase/Peroxidase(Sample Type:Serum)	117.54	Normal : Up to 150 mg/dL Borderline High : 150-199 mg/dL High : 200-499 mg/dL Very high : > 500 mg/dL	mg/dL

Patient Name : T GEETHA SAGARI

Age / Gender : 24 years / Female

Patient ID : 21496

Source : DIRECT

Referral : Dr V. MADHAVA RAO

Collection Time : Dec 04, 2025, 06:13 a.m.

Receiving Time : Dec 04, 2025, 06:22 a.m.

Reporting Time : Dec 04, 2025, 08:49 a.m.

Sample ID :



1253380004

Test Description	Result(s)	Biological Reference Intervals	Unit(s)
HDL CHOLESTEROL Method:Direct(Sample Type:Serum)	39.17	> 45 mg/dL High Risk : < 35 mg/dL Low Risk : > 60 mg/dL	mg/dL
*LDL CHOLESTEROL Calculated	101.67	Optimal : Up to 100 mg/dL Near optimal/above optimal : 100-129 mg/dL Borderline High : 130-159 mg/dL High : 160-189 mg/dL Very High : > 190 mg/dL	mg/dL
*VLDL CHOLESTEROL Calculated	23.51	5 - 51 mg/dL	mg/dL
*CHOL / HDL RATIO Calculated	4.20	Low risk : < 3.5 Average risk : 3.5 - 5.0 High risk : > 5.0	
*LDL/HDL RATIO Calculated	2.60	Desirable : < 0.5-3.0 Borderline : 3.1- 6.0 Elevated : > 6.0	

INSTRUMENT: DONE ON BA 400 BIOSYSTEMS FULLY AUTOMATED BIOCHEMISTRY ANALYZER

Note

* marked parameter are not under NABL Accreditation.

URIC ACID

URIC ACID Method:URICASE/PEROXIDASE(Sample Type:Serum)	3.86	Male:3.5-7.2 Female:2.6 - 6.0	mg/dL
---	------	----------------------------------	-------

Note

Liver Function Tests

BILIRUBIN (TOTAL) DICHLOROPHENYL DIAZONIUM (Sample Type:Serum)	0.53	<24Hrs: 1.0 - 8.0 mg/dL <48Hrs: 6.0 - 12.0 mg/dL 3 - 5 Days: 10.0 - 14.0 mg/dL Adults : <1.0 mg/dL	mg/dL
*BILIRUBIN (DIRECT) DICHLOROPHENYL DIAZONIUM (Sample Type:Serum)	0.32	< 0.2 mg/dL	mg/dL
BILIRUBIN (INDIRECT)(Sample Type:Serum) calculated Method : Calculated	0.21	0.2 -0.8 mg/dL	mg/dL
SGOT IFCC without pyridoxal 5-P(Sample Type:Serum)	35.59	0 - 40	U/L
SGPT IFCC without pyridoxal 5-P(Sample Type:Serum)	61.98	0 - 41	U/L

Patient Name : T GEETHA SAGARI

Age / Gender : 24 years / Female

Patient ID : 21496

Source : DIRECT

Referral : Dr V. MADHAVA RAO

Collection Time : Dec 04, 2025, 06:13 a.m.

Receiving Time : Dec 04, 2025, 06:22 a.m.

Reporting Time : Dec 04, 2025, 08:49 a.m.

Sample ID :



1253380004

Test Description	Result(s)	Biological Reference Intervals	Unit(s)
TOTAL PROTEIN BIURET (Sample Type:Serum)	7.78	Pre mature : 0.6 -6.0 g/dL New born : 4.6 - 7.0 g/dL One week : 4.4 - 7.6 g/dL 7 months - 1 year : 5.1-7.3 g/dL 1 Year - 2 years : 5.6 - 7.5 g/dL > 3 years : 6.5 - 8.0 g/dL	g/dL
ALBUMIN BROMOCRESOL GREEN(Sample Type:Serum)	4.45	3.5 - 5.0 g/dL	g/dL
*GLOBULIN Calculated	3.33	2.5 - 4.5	g/dL
*A:G RATIO	1.34	1.5 - 3.5	
ALP (ALKALINE PHOSPHATASE) 2Amino 2Methyl - 1Propanol Buffer IFCC Method(Sample Type:Seru	93.62	<105 U/L	U/L
*gGT Method : IFCC	34.21	< 38 Note: * marked parameter are not under NABL Accreditation.	U/L

S.PHOSPHORUS

PHOSPHORUS Method: PHOSPHOMOLYBDATE/UV (Sample Type:Serum)	4.31	2.5 - 4.5	mg/dL
---	------	-----------	-------

Method:TEST DONE ON BIOSYSTEMS BA 400. A FULLY AUTOMATED BIOCHEMISTRY ANALYSER

S.IRON

IRON Method:FERROZINE(Sample Type:Serum)	26.30	Men : 65 - 175 µg/dL Women : 50 - 170 µg/dL	µg/dL
---	--------------	--	-------

Instrument:Test Done On Fully Automated Biochemistry Analyser, BIOSYSTEMS BA 400

Note

UREA

UREA Method:UREASE / GLUTAMATE DEHYDROGENASE(Sample Type:Serum)	15.26	Cord: 21 - 40 mg/dL Pre - Mature : 3 - 20 mg/dL New Born : 4 - 12 mg/dL Child : 5 - 18 mg/dL Adult : 15 - 39 mg/dL <60 Years : 8 - 23 mg/dL	mg/dL
--	-------	--	-------

Note

HEMOGRAM (CBP)

*Haemoglobin	9.4	12.0 - 18.0	g/dL
--------------	------------	-------------	------

Patient Name : T GEETHA SAGARI

Age / Gender : 24 years / Female

Patient ID : 21496

Source : DIRECT

Referral : Dr V. MADHAVA RAO

Collection Time : Dec 04, 2025, 06:13 a.m.

Receiving Time : Dec 04, 2025, 06:22 a.m.

Reporting Time : Dec 04, 2025, 08:49 a.m.

Sample ID :



1253380004

Test Description	Result(s)	Biological Reference Intervals	Unit(s)
*Leucocyte Count (WBC Count)	8700	4000- 11000	μL
*Platelet Count	540000	120000 - 380000	μL
*R.B.C count	5.12	3.80 - 5.30	10 ⁶ μL
*MCV	60.0	80.0 - 100.0	fL
*PCV	30.7	36.0 - 56.0	%
*MCH	18.4	27-32	pg
*MCHC	30.6	32.0 - 36.0	g/dL
*RDW-CV	17.7	10.0 - 16.5	%
RDW-SD	42.5		fl
RDWI	212		
*Mean Platelet Volume (MPV)	9.2	5.0 - 10.0	fL
*PDW	17.1	12.0 - 18.0	%
*P-LCR	33.0		%
*P-LCC	17.82		
*PCT	0.50		%
MentzerIndex	12		
NLR	1.89		
*DIFFERENTIAL COUNT			
*Granulocyte	64.1	42 - 85	%
*Lymphocytes	33.9	11.0 - 49.0	%
*Monocytes	2.0	0.0 - 9.0	%
Comments:	Anemia Hypochromia Microcytosis Thrombocytosis		

Instrument: TEST PERFORMED ON NIHON KOHDEN HEMATOLOGY ANALYZER

Note

* Marked parameter are not under NABL Accreditation.

FASTING PLASMA GLUCOSE

Fasting Plasma Glucose (F)	108.80	Desired Fasting Plasma Glucose:	mg/dL
Method : Glucose oxidase/Peroxidase			
		For nondiabetic: 70 - 100 mg/dL	
		For Diabetic: 80 - 130 mg/dL	

****END OF REPORT****


Patient Name : T GEETHA SAGARI
Age / Gender : 24 years / Female
Patient ID : 21496
Source : DIRECT


Referral : Dr V. MADHAVA RAO
Collection Time : Dec 04, 2025, 06:13 a.m.
Receiving Time : Dec 04, 2025, 06:22 a.m.
Reporting Time : Dec 04, 2025, 08:49 a.m.
Sample ID :



1253380004

Test Description	Result(s)	Biological Reference Intervals	Unit(s)
------------------	-----------	--------------------------------	---------


D. Rajitha
Quality Manager

Approved By

Dr. V. Madhava Rao
MD (PGI-Chandigarh)
DM (Endocrinology) (AIIMS-New Delhi)