PARISHRAM 2.0 2026 (Class 12th)

BIOLOGY (BOTANY)

DPP: 3

Sexual Reproduction in Flowering Plants

Q1 Which of the following device is not used by plants to prevent autogamy-
(A) Self - incompatibility
(B) Production of unisexual flowers
(C) Heterostyly
(D) Production of cleistogamous flowers

Q2 Chasmogamy is pollination in -
(A) Bud condition
(B) Closed flowers
(C) Unrelated flowers
(D) Opened flowers

Q3 Pollination in water hyacinth and water lily is brought about by the agency of:
(A) Birds
(B) Bats
(C) Insects or wind
(D) Water

Q4 Pollen grain of water pollinated plants are coated by covering to prevent it from wetting
(A) mucilage (B) cuticle
(C) exine (D) intine

Q5 The process whereby a perfect flower is pollinated by its pollen is called -
(A) Allogamy (B) Autogamy
(C) Xenogamy (D) Hydrogamy

Q6 Which type of pollination occurs when pollen grains are transferred from the anther to the stigma of a different plant?
(A) Autogamy (B) Xenogamy
(C) Geitonogamy (D) Cleistogamy

Q7 What is the main advantage of cleistogamous flowers in terms of pollination?
(A) They ensure assured seed-set without the need for pollinators.
(B) They promote cross-pollination.
(C) They attract more pollinators.
(D) They produce fewer pollen grains.

Q8 Which of the following is not an agent of pollination?
(A) Wind
(B) Water
(C) Insect and birds
(D) None of these

Q9 Which of the following condition is helpful for wind pollination of flowers?

(A) sticky pollens.
(B) flowers with nectar.
(C) well exposed stamens.
(D) attractive colored petals.

Q10 Which of the following best describes geitonogamy in pollination?
(A) Transfer of pollen grains between flowers of different plants
(B) Transfer of pollen grains from the anther to the stigma of another flower of the same plant
(C) Transfer of pollen grains from the anther to the stigma of a different species
(D) Transfer of pollen grains within the same flower

Answer Key

Q1	D
Q2	D
Q3	C
Q4	A
Q5	B
Q6	B
Q7	A
Q8	D
Q9	C
Q10	B

Hints & Solutions

Note: scan the QR code to watch video solution

Q1 Text Solution:

Cleistogamous flowers are those that do not open and are self-pollinating. These flowers actually encourage autogamy (self-fertilization), which is the opposite of what the plant aims to prevent.

Video Solution:

Q2 Text Solution:

Chasmogamous flowers which are similar to flowers of other species with exposed anthers and stigma.

Video Solution:

Q3 Text Solution:

In a majority of aquatic plants such as water hyacinth and water lily, the flowers emerge above the level of water and are pollinated by insects or wind as in most of the land plants.

Video Solution:

Q4 Text Solution:

In most of the water-pollinated species, pollen grains are protected from wetting by a mucilaginous covering.

Video Solution:

Q5 Text Solution:

Autogamy: Transfer of pollen grains from the anther to the stigma of the same flower.

Video Solution:

Q6 Text Solution:

Xenogamy – Transfer of pollen grains from anther to the stigma of a different plant.

Video Solution:

Q7 Text Solution:

Cleistogamous flowers do not open and self-pollinate, ensuring seed production even in the absence of pollinators. This is advantageous because it guarantees the plant's reproduction without relying on external agents for pollination.

Video Solution:



Q8 Text Solution:

Plants use two abiotic (wind and water) and one biotic (animals) agents to achieve pollination.

Video Solution:



Q9 Text Solution:

(3)

Wind pollinated flowers have well exposed stamens.

NCERT Page No. 29

Video Solution:



Q10 Text Solution:

Geitonogamy – Transfer of pollen grains from the anther to the stigma of another flower of the same plant.

Video Solution:



Android App | iOS App | PW Website