PARISHRAM 2.0 2026 (Class 12th)
Biology (Botany)
Sexual Reproduction in Flowering Plants

DPP: 6

Q1 Which of the following helps during seed germination by allowing water and oxygen to enter?
(A) Pericarp (B) Micropyle
(C) Cotyledons (D) Endosperm

Q2 Which part of the seed forms from the integuments of the ovule?
(A) Endosperm (B) Pericarp
(C) Seed coat (D) Cotyledon

Q3 In which type of seeds is the endosperm completely consumed during embryo development?
(A) Albuminous seeds
(B) Non-albuminous (ex-albuminous) seeds
(C) Seeds with perisperm
(D) Seeds with both cotyledons and endosperm

Q4 Persistant nucellus present in some seeds is called
(A) Perisperm (B) Pericarp
(C) Periderm (D) Phelloderm

Q5 Which of the following is correctly matched?
(A) Pea - Albuminous seed
(B) Beet - Perisperm present
(C) Maize - Non-Albuminous seed
(D) Castor - Non-albuminous seed

Q6 The micropyle in the seed coat facilitates the entry of:
(A) Pollen grains
(B) Nutrients
(C) Oxygen and water
(D) Hormones

Q7 The scutellum observed in a grain of wheat or maize is comparable to which part of the seed in other monocotyledons?
(A) Cotyledon (B) Endosperm
(C) Aleurone layer (D) Plumule

Q8 In which part of the embryo maximum growth takes place in epigeal germination: -
(A) Radicle
(B) Plumule
(C) Epicotyl
(D) Hypocotyl

Q9 Which of the following statements about seed moisture is true?
(A) Mature seeds contain 20-30% of water by mass.
(B) Seed coat remain soft in mature seeds
(C) Mature seeds become relatively dry with 10-15% moisture
(D) Germination reduces the seed moisture content further

Q10 In albuminous seed, the food is stored in-
(A) Testa (B) Plumule
(C) Cotyledon (D) Endosperm

Answer Key

Q1	B
Q2	C
Q3	B
Q4	A
Q5	B
Q6	C
Q7	A
Q8	D
Q9	C
Q10	D

Hints & Solutions

Note: scan the QR code to watch video solution

Q1 **Text Solution:**
Micropyle

Video Solution:

Q2 **Text Solution:**
Seed coat

Video Solution:

Q3 **Text Solution:**
Non-albuminous (ex-albuminous) seeds

Video Solution:

Q4 **Text Solution:**
Occasionally, in some seeds such as black pepper and beet, remnants of nucellus are also persistent. This residual, persistent nucellus is the perisperm.

Video Solution:

Q5 **Text Solution:**
Occasionally, in some seeds such as black pepper and beet, remnants of nucellus are also persistent. This residual, persistent nucellus is the perisperm.

Video Solution:

Q6 **Text Solution:**
Oxygen and water

Video Solution:

Q7 **Text Solution:**
The scutellum is the single cotyledon present in the seeds of monocotyledons like wheat or maize.

Video Solution:

Q8 **Text Solution:**
In epigeal germination, the hypocotyl elongates and pushes the cotyledons above the ground.

Video Solution:

Q9 Text Solution:

As the seed matures, its water content is reduced and seeds become relatively dry (10-15 per cent moisture by mass).

Video Solution:

Q10 Text Solution:

Albuminous seed retain a portion of their endosperm at maturity , which acts as a food reserve for the developing embryo.

Video Solution:

Android App | iOS App | PW Website