Parishram 2.0 2026 (Class 12th)


Biology (Botany)

DPP:07

Sexual reproduction in flowering plants

Q1 Which of the following is an example of false fruit?
(A) Mango (B) Guava
(C) Apple (D) Mustard

Q2 Observe the image and identify A,B,C,D



(A) A - Thalamus B - Mesocarp C - Endocarp D - Seed
(B) A -Seed B - Thalamus C - Endcarp D - Mesocarp
(C) A - Thalamus B - seed C - Mesocarp D - Endocarp
(D) A - Thalamus B - Seed C - Endocarp D - Mesocarp

Q3 A true fruit develops from
(A) Ovary (B) Thalamus
(C) Petals (D) Receptacle

Q4 Viability of Seeds
(A) Maybe lost within a few months
(B) Is 1000 years in Lupinus arcticus
(C) Recorded in Phoenix is 10000 years
(D) All of these

Q5 The period of viability is maximum in the seed of
(A) Lupine (B) Date palm
(C) Lotus (D) Orobanche

Q6 The world's oldest viable seed, excavated from Arctic Tundra is of -
(A) Phoenix dactylifera
(B) Calotropis
(C) Lupine (Lupinus)
(D) Victoria

Q7 Fruits which develop without fertilization are called as
(A) True fruits
(B) Parthenocarpic fruits
(C) Dry fruits
(D) None of these

Q8 Seed dormancy allows the plants to-
(A) Overcome unfavourable climate conditions
(B) Develop healthy seeds
(C) Reduce viability
(D) Prevent deterioration of seeds

Q9 In which of the following plants does the endosperm persist in the mature seed?
(A) Pea and groundnut
(B) Bean and gram
(C) Castor and coconut
(D) Gram and sunflower

Q10 Which of the following will lose its economic value if fruits are produced through parthenocarpy?
(A) Citrus (B) Banana
(C) Grape (D) Pomegranate

Answer Key

Q1	C
Q2	D
Q3	A
Q4	A
Q5	A
Q6	C
Q7	B
Q8	A
Q9	C
Q10	D

Hints & Solutions

Note: scan the QR code to watch video solution

Q1 **Text Solution:**

In a few species such as apple, strawberry, cashew, etc., the thalamus also contributes to fruit formation.Such fruits are called false fruits.

Video Solution:

Q2 **Text Solution:**

A - Thalamus B - Seed C - Endocarp D - Mesocarp

Video Solution:

Q3 **Text Solution:**

Most fruits however develop only from the ovary and are called true fruits

Video Solution:

Q4 **Text Solution:**

In a few species the seeds lose viability within a few months.

Video Solution:

Q5 **Text Solution:**

The oldest is that of a lupine, Lupinus arcticus excavated from Arctic Tundra. The seed germinated and flowered after an estimated record of 10,000 years of dormancy.

Video Solution:

Q6 **Text Solution:**

Lupine (*Lupinus*)

There are several records of very old yet viable seeds. The oldest is that of a lupine, *Lupinus arcticus* excavated from Arctic Tundra. The seed germinated and flowered after an estimated record of 10,000 years of dormancy.

Video Solution:

Q7 **Text Solution:**

Although in most of the species, fruits are the results of fertilisation, there are a few species in which fruits develop without fertilisation. Such fruits are called parthenocarpic fruits.

Video Solution:

Q8 **Text Solution:**

Overcome unfavourable climate conditions

Video Solution:

Q9 **Text Solution:**

Castor and coconut

Video Solution:

Q10 **Text Solution:**

1. Pomegranate seeds with fleshy testa are the edible parts.
2. Parthenocarpy renders the fruit seedless and thus ineffective i.e. In the case of induced parthenocarpy, the seeds are not produced, rendering the fruit seedless.
3. As a result, they become obsolete and lose economic significance.
4. The fleshy part of pomegranate is the outer layer of the seed coat called the testa, which is eaten.
5. When parthenocarpic fruits are produced, the ovaries ripen without the ovule being fertilized.
6. Seeds are not produced in the absence of fertilization.
7. Because the seeds are the only edible part of this fruit, its economic value will decline.
8. Because the seeds of oranges, bananas, and grapes are not consumed, parthenocarpy is beneficial.

Video Solution: