
PARISHRAM 2.0 2026 (Class 12th)
Biology (Zoology)
DPP 4
Human Reproduction

Q1 Before ovulation, the secondary oocyte forms a membrane called as _____.
(A) chorion
(B) zona pellucida
(C) corona radiata
(D) vitelline membrane.

Q2 Identify P, Q and R in the given diagram and choose the correct option.



(A) P-Primary follicle, Q-Tertiary follicle, R-Graafian follicle
(B) P-Graafian follicle, Q-Secondary oocyte, R-Corpus luteum
(C) P- Secondary oocyte, Q-Tertiary follicle, R-Graafian follicle
(D) P- Primary follicle, Q-Tertiary follicle, R-Corpus luteum

Q3 The rupture of the mature (Graafian) follicle and the release of the secondary oocyte (ovum) from the ovary is termed:
(A) Oogenesis (B) Ovulation
(C) Fertilization (D) Implantation

Q4 Which of the following correctly describes the changes that occur when a secondary follicle transforms into a tertiary follicle?
(A) The follicle ruptures to release the ovum into the fallopian tube.
(B) The tertiary follicle develops a fluid-filled cavity called the antrum, and the theca layer differentiates into theca interna and theca externa.
(C) The follicle degenerates and forms the corpus luteum.
(D) The secondary oocyte completes meiosis II within the follicle.

Q5 At puberty, how many primary follicles are approximately left in each ovary?
(A) 1,000–2,000
(B) 10,000–20,000
(C) 60,000–80,000
(D) 100,000–200,000

Q6 What do primary follicles become when they are surrounded by more layers of granulosa cells and a new theca?
(A) Graafian follicles
(B) Secondary follicles
(C) Tertiary follicles
(D) Corpus luteum

Q7 During embryonic development, oogonia begin meiosis and get temporarily arrested in which phase, forming primary oocytes?
(A) Metaphase-I (B) Anaphase-I
(C) Prophase-I (D) Telophase-I

Q8 _____ is the characteristic feature of a tertiary follicle.
(A) Presence of a zygote
(B) Formation of the corpus luteum
(C) Fluid-filled cavity called antrum
(D) Release of the secondary oocyte

Q9 Identify the incorrectly matched pair related to oogenesis and follicle development.

(A) Primary oocyte Undergoes first meiotic division in tertiary follicle.
(B) First meiotic division Produces secondary oocyte and first polar body.
(C) Secondary oocyte Retains bulk of the nutrient rich cytoplasm of the primary oocyte.
(D) Graafian follicle Directly formed from primary follicle.

Q10 **Assertion:** One primary oocyte produces two functional female gametes.

Reason: The divisions during oogenesis are unequal, resulting in one large ovum and smaller polar bodies.
(A) Both Assertion and Reason are true, and Reason is the correct explanation of Assertion.
(B) Both Assertion and Reason are true, but Reason is not the correct explanation of Assertion.
(C) Assertion is true, and Reason is false.
(D) Assertion is false, and Reason is true.

Answer Key

Q1	B
Q2	B
Q3	B
Q4	B
Q5	C
Q6	B
Q7	C
Q8	C
Q9	D
Q10	D

Hints & Solutions

Note: scan the QR code to watch video solution


Q1 Text Solution:

The secondary oocyte forms a new membrane called zona pellucida. The Graafian follicle ruptures to release the secondary oocyte (ovum) from the ovary by the process called ovulation.

Video Solution:

Q2 Text Solution:

P-Graafian follicle, Q-Secondary oocyte, R-Corpus luteum



Video Solution:

Q3 Text Solution:

Ovulation is the specific event when the graafian follicle ruptures and releases the secondary oocyte into the fallopian tube.

Video Solution:

Q4 Text Solution:

- Option A describes ovulation, which occurs later.
- Option C refers to events post-ovulation.
- Option D is incorrect, as the secondary oocyte completes meiosis II only after fertilization.

Video Solution:

Q5 Text Solution:

At puberty only 60,000-80,000 primary follicles are left in each ovary.

Video Solution:

Q6 Text Solution:

When a primary follicle is surrounded by multiple layers of granulosa cells and a newly formed theca layer, it matures into a secondary follicle.

Video Solution:

Q7 **Text Solution:**

Oogonia enter meiosis-I during foetal development but pause at prophase-I, becoming primary oocytes. They stay in this arrested stage until puberty.

Video Solution:

Q8 **Text Solution:**

As the secondary follicle matures into a tertiary follicle, it develops a fluid-filled cavity called the antrum, which is a key distinguishing feature.

Video Solution:

Q9 **Text Solution:**

The Graafian follicle is not formed directly from the primary follicle. It develops step-by-step:

Primary follicle Secondary follicle Tertiary follicle Graafian follicle.

Video Solution:

Q10 **Text Solution:**

Correct Assertion: Only one functional female gamete (the ovum) is produced from a single primary oocyte.

Video Solution:

Android App | iOS App | PW Website