

Assignment 3

AI24BTECH11008- Sarvajith

Question 16: A natural number has prime factorization given by $n = 2^x 3^y 5^z$, where y and z are such that $y + z = 5$ and $y^{-1} + z^{-1} = 5/6, y > z$. Then the number of odd divisors of n , including 1, is:

- a. 11
- b. $6x$
- c. 12
- d. 6

Question 17: Let $f(x) = \sin^{-1}(x)$ and $g(x) = [x^2 - x - 2]/[2x^2 - x - 6]$. If $g(2) = \lim_{x \rightarrow 2} g(x)$, then the domain of the function $f \circ g$ is:

- a. $(-\infty, -2] \cup [-4/3, \infty]$
- b. $(-\infty, -1] \cup [2, \infty]$
- c. $(-\infty, -2] \cup [-1, \infty]$
- d. $(-\infty, -2] \cup [-3/2, \infty]$

Question 18: If the mirror image of the point $(1, 3, 5)$ with respect to the plane $4x - 5y + 2z = 8$ is (α, β, γ) , then $5(\alpha + \beta + \gamma)$:

- a. 47
- b. 39
- c. 43
- d. 41

Question 19: Let $f(x) = \int_0^x e^t f(t) dt + e^x$ be a differentiable function for all $x \in \mathbf{R}$. Then $f(x)$ equals:

- a. $2e^{e^x-1} - 1$
- b. e^{e^x-1}
- c. $2e^{e^x} - 1$
- d. $e^{e^x} - 1$

Question 20: The triangle of the maximum area that can be inscribed in a given circle of radius 'r' is:

- a. A right-angle triangle having two of its sides of length $2r$ and r .
- b. An equilateral triangle of height $2r/3$.
- c. Isosceles triangle with base equal to $2r$.
- d. An equilateral triangle having each of length $\sqrt{3}r$

1 SECTION-B

Question 1: The total number of 4-digit numbers whose greatest common divisor with 18 is 3, is

Question 2: Let α and β be two real numbers such that $\alpha + \beta = 1$ and $\alpha\beta = -1$. Let $P_n = \alpha^n + \beta^n$, $P_{n-1} = 1$ and $P_{n+1} = 29$ for some integer $n = 1$. Then the value of P_n^2 is

Question 3: Let X_1, X_2, \dots, X_{18} be eighteen observations such that $\sum_{i=1}^{18} (X_i - \alpha) = 36$ and $\sum_{i=1}^{18} (X_i - \beta)^2 = 90$, where α and β are distinct real numbers. If the standard deviation of these observations is 1, then the value of $|\alpha - \beta|$ is

Question 4: In $I_{m,n} = \int_0^1 x^{m-1} (1-x)^{n-1} dx$, for $m, n \geq 1$ and $\int_0^1 [x^{m-1} + x^{n-1}] / [(1+x)^{m+n}] dx = \alpha I_{m,n}$, $\alpha \in R$, then α is

Question 5: Let L be a common tangent line to the curves $4x^2 + 9y^2 = 36$ and $(2x)^2 + (2y)^2 = 31$. Then the square of the slope of the line L is

Question 6: If the matrix $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 3 & 0 & -1 \end{pmatrix}$ satisfies the equation $A^{20} + \alpha A^{19} + \beta A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{pmatrix}$

for some real numbers α and β , then $\beta - \alpha$ is equal to?

Question 7: If the arithmetic mean and the geometric mean of the p^{th} and q^{th} terms of the sequence $-16, 8, -4, 2, \dots$ satisfy the equation $4x^2 - 9x + 5 = 0$, then $p+q$ is equal to?

Question 8: Let the normals at all the points on a given curve pass through a fixed point (a, b) . If the curve passes through $(3, -3)$ and $(4, -2\sqrt{2})$, and given that $a - 2\sqrt{2}b = 3$, then $(a^2 + b^2 + ab)$ is equal to?

Question 9: Let z be those complex number which satisfies $|z + 5| \leq 4$ and $z(i + 1) + \bar{z}(1 - i) \geq -10$, $i = \sqrt{-1}$. If the maximum value of $|z + 1|^2$ is $\alpha + \beta\sqrt{2}$, then the value of $\alpha + \beta$ is?

Question 10: Let a be an integer such that all the real roots of the polynomial $2x^5 + 5x^4 + 10x^3 + 10x^2 + 10x + 10$ lie in the interval $(a, a + 1)$, then $|a|$ is equal to?