## 1

## Question 1.5.2

Find the intersection I of the angle bisectors of B and C

## **Solution**

From ?? the bisectors of B and C are obtained as

$$\left(\frac{11}{\sqrt{122}} + \frac{7}{\sqrt{74}} \quad \frac{1}{\sqrt{122}} + \frac{5}{\sqrt{74}}\right) \begin{pmatrix} \mathbf{x} \\ \mathbf{y} \end{pmatrix} = \frac{2}{\sqrt{74}} - \frac{38}{\sqrt{122}}$$
 (1)

and

$$\left(\frac{11}{\sqrt{122}} + \frac{1}{\sqrt{2}} \quad \frac{1}{\sqrt{122}} - \frac{1}{\sqrt{2}}\right) \begin{pmatrix} \mathbf{x} \\ \mathbf{y} \end{pmatrix} = \frac{2}{\sqrt{2}} - \frac{38}{\sqrt{122}}$$
 (2)

respectively.

The pair of linear equations can be solved using the Augmented matrix  $\begin{bmatrix} P|Q \end{bmatrix}$  Here,

$$P = \begin{bmatrix} \frac{11}{\sqrt{122}} + \frac{7}{\sqrt{74}} & \frac{1}{\sqrt{122}} + \frac{5}{\sqrt{74}} \\ \frac{11}{\sqrt{122}} + \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{122}} - \frac{1}{\sqrt{2}} \end{bmatrix}$$
(3)

$$Q = \begin{bmatrix} \frac{2}{\sqrt{74}} - \frac{38}{\sqrt{122}} \\ \frac{2}{\sqrt{2}} - \frac{38}{\sqrt{122}} \end{bmatrix} \tag{4}$$

$$\left[P|Q\right] = \begin{bmatrix}
\frac{11}{\sqrt{122}} + \frac{7}{\sqrt{74}} & \frac{1}{\sqrt{122}} + \frac{5}{\sqrt{74}} & \frac{2}{\sqrt{74}} - \frac{38}{\sqrt{122}} \\
\frac{11}{\sqrt{122}} + \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{122}} - \frac{1}{\sqrt{2}} & \frac{2}{\sqrt{2}} - \frac{38}{\sqrt{122}}
\end{bmatrix}$$
(5)

The augmented matrix is converted into decimal notations for easier calculations and then can be solved using row reduction as follows

$$\begin{bmatrix}
1.81 & 0.67 & | & -3.21 \\
1.7 & -0.62 & | & -2.03
\end{bmatrix}
\xrightarrow{R_2 \leftarrow R_1 - (\frac{1.81}{1.7})R_2}
\begin{bmatrix}
1.81 & 0.67 & | & -3.21 \\
0 & 1.33 & | & -1.05
\end{bmatrix}$$

$$\xrightarrow{R_1 \leftarrow R_1 - \frac{0.67}{1.33}R_2}
\begin{bmatrix}
1.81 & 0 & | & -2.68 \\
0 & 1.33 & | & -1.05
\end{bmatrix}$$

$$\xrightarrow{R_1 \leftarrow \frac{R_1}{1.81}}
\begin{bmatrix}
1 & 0 & | & -1.48 \\
0 & 1.33 & | & -1.05
\end{bmatrix}$$

$$\xrightarrow{R_1 \leftarrow \frac{R_1}{1.81}}
\begin{bmatrix}
1 & 0 & | & -1.48 \\
0 & 1.33 & | & -1.05
\end{bmatrix}$$

$$\xrightarrow{R_1 \leftarrow \frac{R_1}{1.81}}
\begin{bmatrix}
1 & 0 & | & -1.48 \\
0 & 1 & -0.79
\end{bmatrix}$$

$$\xrightarrow{R_1 \leftarrow \frac{R_1}{1.81}}$$

We obtain

$$\mathbf{I} = \begin{pmatrix} -1.48 \\ -0.79 \end{pmatrix} \tag{10}$$