GSEA functional enrichment

GO_MF; stk: TSB vs SN
All study−significant terms (FDR < 0.1); observed source models only; no Top−N limit

Oxidoreductase activity, acting on the aldehyde or oxo group of donors, NAD or NADP as acceptor [GO:0016620]
Oxidoreductase activity, acting on the aldehyde or oxo group of donors [GO:0016903]
Ligase activity, forming carbon−nitrogen bonds [GO:0016879]
Hydrolase activity, acting on carbon−nitrogen (but not peptide) bonds [GO:0016810]
Transferase activity [GO:0016740]
Transferase activity, transferring phosphorus−containing groups [GO:0016772]
Nucleic acid binding [GO:0003676]
Phosphotransferase activity, alcohol group as acceptor [GO:0016773]
Isomerase activity [GO:0016853]
Catalytic activity, acting on a nucleic acid [GO:0140640]
Channel activity [GO:0015267]
Passive transmembrane transporter activity [GO:0022803]
Ion channel activity [GO:0005216]
Proton transmembrane transporter activity [GO:0015078]
Catalytic activity, acting on RNA [GO:0140098]
Carbohydrate transmembrane transporter activity [GO:0015144]
protein−N(PI)−phosphohistidine−sugar phosphotransferase activity [GO:0008982]
RNA binding [GO:0003723]
aminoacyl−tRNA ligase activity [GO:0004812]
Ligase activity, forming carbon−oxygen bonds [GO:0016875]
Catalytic activity, acting on a tRNA [GO:0140101]
tRNA binding [GO:0000049]
rRNA binding [GO:0019843]
Structural molecule activity [GO:0005198]
Structural constituent of ribosome [GO:0003735]

−2
−1
0
1
Normalized enrichment score (signed NES)

−log10(study FDR)
2
3
4
Eligible pathway members
100
200
300

Rows: largest positive NES to most negative NES. stk: TSB vs SN: NES > 0 = Higher in TSB; NES < 0 = Higher in SN. Source lod_imputed models excluded before enrichment; no minimum−replicate filter added. Enrichment is not a direct measure of pathway activity.