GSEA functional enrichment

KEGG; stk: TSB vs SN
All study−significant terms (FDR < 0.1); observed source models only; no Top−N limit

Purine metabolism [map00230]
Citrate cycle (TCA cycle) [map00020]
beta−Lactam resistance [map01501]
Lysine biosynthesis [map00300]
Glyoxylate and dicarboxylate metabolism [map00630]
One carbon pool by folate [map00670]
Lysine degradation [map00310]
Arginine and proline metabolism [map00330]
Carbon metabolism [map01200]
Quorum sensing [map02024]
Biosynthesis of secondary metabolites [map01110]
Biosynthesis of amino acids [map01230]
Porphyrin and chlorophyll metabolism [map00860]
Phosphotransferase system (PTS) [map02060]
Glycolysis / Gluconeogenesis [map00010]
Pyrimidine metabolism [map00240]
Fructose and mannose metabolism [map00051]
Aminoacyl−tRNA biosynthesis [map00970]
Ribosome [map03010]

−2
−1
0
1
2

Normalized enrichment score (signed NES)

Eligible pathway members
100
200
300

−log10(study FDR)
2
3
4

Rows: largest positive NES to most negative NES. stk: TSB vs SN: NES > 0 = Higher in TSB; NES < 0 = Higher in SN. Source lod_imputed models excluded before enrichment; no minimum−replicate filter added. Enrichment is not a direct measure of pathway activity.