

AI1110 Assignment-9

Gollapudi Sasank CS21BTECH11019

May 25, 2022

Outline

1 Question

2 Solution

Question

Players X and Y roll dice alternately starting with X . The player that rolls 11 wins. Show that the probability p that X wins equals $18/35$.

Solution

Let the Random Variables A, B denote the following :

$A = 0$: The person who starts the game wins

$A = 1$: The person who starts the game loses

$B = 0$: 11 occurs at the first throw

$B = 1$: 11 does not occur at the first throw

Given,

$$\Pr(A = 0) = p \quad (1)$$

$$\Rightarrow \Pr(A = 1) = 1 - p \quad (2)$$

$$\Pr(B = 0) = \frac{2}{36} = \frac{1}{18} \quad (3)$$

$$\Rightarrow \Pr(B = 1) = \frac{34}{36} = \frac{17}{18} \quad (4)$$

The Events $B = 0$ and $B = 1$ form a partition to the Sample Space.

$$\Rightarrow \Pr(A = 0) = \Pr((A = 0)((B = 0) + (B = 1))) \quad (5)$$

$$\Rightarrow \Pr(A = 0) = \Pr((A = 0)(B = 0) + (A = 0)(B = 1)) \quad (6)$$

The Events $(A = 0)(B = 0)$ and $(A = 0)(B = 1)$ are mutually exclusive

$$\Rightarrow \Pr(A = 0) = \Pr((A = 0)(B = 0)) + \Pr((A = 0)(B = 1)) \quad (7)$$

$$\Rightarrow \Pr(A = 0) = \Pr(A = 0|B = 0) \Pr(B = 0) + \Pr(A = 0|B = 1) \Pr(B = 1) \quad (8)$$

$\Pr(A = 0|B = 0) = 1$ because if 11 occurs at first throw X wins.

Now the event $(A = 0|B = 1)$ is the case where X wins when 11 does not occur at first throw. So in this case if we consider the game from the second throw then Y throws first. But here we need the probability of the case where the person who starts the game loses i.e $\Pr(A = 1) = 1 - p$.

$$\therefore \Pr(A = 0|B = 1) = 1 - p$$

$$\Rightarrow p = 1 \times \frac{1}{18} + (1 - p) \times \frac{17}{18} \quad (9)$$

$$\Rightarrow p = \frac{1 + 17 - 17p}{18} \quad (10)$$

$$\Rightarrow 18p = 18 - 17p \quad (11)$$

$$\Rightarrow 35p = 18 \quad (12)$$

$$\Rightarrow p = \frac{18}{35} \quad (13)$$