

SupportShift: Local KL Geometry of Ignored Matérn Observation Support

with Theorem-Linked High-Dimensional Simulations

Sasha Cui

Independent researcher

Draft of August 9, 2026

Abstract

Spatial measurements often represent local averages, but are subsequently fitted as if they were point observations. This changes the population target of covariance estimation, not merely its finite-sample bias. We study a stationary Matérn Gaussian field in $\mathbb{R}^d$, observed through a known compact symmetric averaging kernel of bandwidth h . For a fixed nonzero-lag Gaussian correlation match with known smoothness ν , we characterize the exact pseudo-variance and pseudo-decay and derive their small-support expansions. The naive inverse-range shift is of order $h^{2\nu}$ for $0 < \nu < 1$, $h^2 \log(1/h)$ at $\nu = 1$, and h^2 for $\nu > 1$. Because this leading law is nonuniform near the threshold, we also derive a two-term approximation whose singular quadratic and fractional terms cancel continuously into the $\nu = 1$ logarithm. Explicit coefficients show that naive point-support fitting underestimates inverse range—and therefore overestimates spatial range—for every fixed $\nu > 0$ and all sufficiently small h . We extend the saturated pair result to a weighted multi-lag composite, deriving both its information-weighted parameter displacement and its first nonzero minimum-KL term. For any fixed site design, a covariance-tangent projection gives the analogous joint variance-decay displacement and irreducible full-Gaussian KL divergence. For anisotropic support, an all-smoothness directional contrast isolates the order- h^2 difference between lag directions. We also give an explicit finite-library Gaussian likelihood certificate based on standard quadratic-form concentration. It allows arbitrary dependence among the p coordinates of each spatial field; under uniform relative spectral control its radius is $O\{\sqrt{\log(M)/(Np)} + \log(M)/(Np)\}$ for N independent fields and M covariance candidates. Deterministic quadrature verifies the phase laws and KL coefficients. A joint smoothness-range library shows that nuisance movement can reverse the fixed-smoothness decay direction, while partial-support and matched-boundary experiments supply non-oracular method and design contrasts. Replicated-field simulations verify the likelihood certificate and show that increasing information can concentrate a naive estimator around a wrong KL target. The covariance transformation and concentration inequality are established tools; the narrower contribution is their theorem-linked use with the explicit Matérn support-distortion law.

1 Introduction

Spatial values recorded on pixels, grid cells, administrative regions, or after kernel preprocessing are local aggregates of an underlying field. The change-of-support literature models such observations by integrating the latent process over their supports [5, 11]. In practice, however, smoothed values are often assigned to declared centers and a point-support covariance is fitted. Relative to the latent parameters, this is a support-misinterpreted model. A single two-point covariance can still

be matched exactly by different variance and decay parameters; a multi-lag likelihood is genuinely misspecified and selects a Kullback–Leibler (KL) projection [2].

That qualitative conclusion is familiar, but it does not answer three basic questions. How fast does the target move as the support shrinks? Does the rate depend on Matérn smoothness? Is the direction of the range error stable across dimension and averaging kernels? These questions matter because the Matérn covariance changes regularity at the origin as ν crosses one. A symmetric kernel cancels first-order Taylor terms at nonzero lags, but it cannot cancel a cusp or fractional power at the origin.

Temporal aggregation of autoregressive and Ornstein–Uhlenbeck models is classical [1, 19], and integrated Ornstein–Uhlenbeck inference has been studied directly [9]. Consequently, a one-dimensional exponential calculation alone is not a novelty claim. Reviews and geostatistical methods for incompatible supports establish the block-to-block covariance and area-to-point prediction problems [12, 14]. Classical regularized-semivariogram work already explains why larger support can increase an apparent autocorrelation range [3, 6]. Clark’s exact geometric argument concerns a finite range of influence, and later practical de-regularization rules are model-specific. A Matérn covariance is positive at every finite lag, while our estimand is a free-variance Gaussian KL pseudo-parameter at a fixed nonzero lag; the classical support-width rule and our smoothness-dependent displacement therefore refer to different range objects. Most closely on the likelihood side, Fuentes [10] constructs a likelihood for block averages in terms of an underlying point covariance and studies a Matérn example. That work correctly models the support; it does not derive the pseudo-range selected when block averages are instead treated as points. Generic misspecified-GP KL theory [2] supplies the projection framework but not the support-specific coefficient or sign. Pairwise and weighted composite likelihoods for Gaussian fields are established [4, 20]; our contribution is the local target and residual KL induced by the omitted support operator, not their general consistency theory. Gaussian quadratic-form concentration and finite-class union bounds are also standard [13, 15]; we use them to make the simulation certificate match the fitted covariance library exactly. Our principal mathematical contribution is the explicit all-smoothness, d -dimensional fixed-lag pseudo-parameter expansion. In particular:

1. We derive a phase law for the naive Matérn decay target: orders $h^{2\nu}$, $h^2 \log(1/h)$, and h^2 in the rough, threshold, and smooth regimes. A transition-aware two-term target approximation resolves the slow finite-bandwidth crossover around $\nu = 1$.
2. We extend the pair coefficient to a genuinely misspecified multi-lag Gaussian composite and derive its positive residual-KL coefficient. A finite-design proposition projects the support perturbation onto the variance–decay Fisher tangent space and identifies exactly when the full-Gaussian model remains misspecified to first order.
3. We give explicit coefficients. Origin expansions, together with a Bessel-function identity in the smooth regime, prove that normalized correlation increases, hence inverse range decreases, for every $\nu > 0$ and any compact symmetric kernel when h is sufficiently small.
4. We isolate the order- h^2 contrast between two lag directions under anisotropic support. The contrast holds for every $\nu > 0$, even when it is lower order than the direction-independent rough-field shift.
5. We distinguish the continuous interior theorem from discrete, row-normalized, and boundary-truncated smoothing. We compute finite-grid full-likelihood KL targets, compare exact, partial, and ignored support, hold dimension fixed in the boundary comparison, and jointly vary smoothness and decay using only synthetic data.

6. For a deterministic finite covariance library, we state a simultaneous normalized-likelihood certificate and its empirical-risk consequence. This supporting result uses independent replicated fields, not independent spatial coordinates.

The covariance transformation $SKS^\top$, block kriging, generic KL projection under misspecification, temporal aggregation, and the concentration machinery are prior work; they are not claimed as contributions. Recent finite-grid Matérn likelihood work also treats discretization and edge effects without studying the target induced by a preprocessing smoother [17]. Our narrower focus is the smoothness-dependent target displacement caused by ignoring observation support, its genuine-likelihood projections and directional coefficient, and a benchmark in which the probability guarantee and fitted library are the same mathematical objects.

2 Model and pseudo-target

2.1 Latent field and local support

Let $Y = \{Y(t) : t \in \mathbb{R}^d\}$ be a mean-zero stationary Gaussian field with

$$\text{Cov}\{Y(t), Y(t+r)\} = C(r) = \sigma^2 \mathcal{M}_\nu(\alpha \|r\|), \quad \sigma^2 > 0, \quad \alpha > 0, \quad \nu > 0, \quad (1)$$

where

$$\mathcal{M}_\nu(x) = \frac{2^{1-\nu}}{\Gamma(\nu)} x^\nu K_\nu(x), \quad \mathcal{M}_\nu(0) = 1. \quad (2)$$

Thus σ^2 is the marginal variance and α is an inverse range; larger α means shorter dependence. This convention must not be mixed with the common length-scale parameterization $\alpha = \sqrt{2\nu}/\ell$.

Let k be a symmetric probability density supported on $B(0, L) \subset \mathbb{R}^d$, let $k_h(u) = h^{-d}k(u/h)$, and define the mean-square convolution

$$Z_h(t) = \int_{\mathbb{R}^d} k_h(u) Y(t-u) du. \quad (3)$$

If $U, V \stackrel{\text{iid}}{\sim} k$ and $D = U - V$, stationarity and Fubini's theorem give the exact smoothed covariance

$$C_h(r) = \sigma^2 \mathbb{E}[\mathcal{M}_\nu\{\alpha \|r + hD\|\}]. \quad (4)$$

Write

$$\Sigma_k = \mathbb{E}(UU^\top), \quad T_k = \text{tr}(\Sigma_k), \quad m_q = \mathbb{E}\|D\|^q, \quad (5)$$

so that $m_2 = 2T_k$. We assume $T_k > 0$.

Throughout, η denotes a generic inverse-range candidate. A dagger marks an exact population KL target under the point-support family, a tilde marks an analytic approximation, and the superscript C marks the weighted pair composite.

2.2 Naive Gaussian pair fit

Fix $r = Re$, with $R > 0$ and $\|e\| = 1$. The smoothed pair has variance $C_h(0)$ and correlation

$$\rho_h(r) = \frac{C_h(r)}{C_h(0)}. \quad (6)$$

A naive point-support model with the correct known ν and parameters $\varsigma^2, \eta > 0$ uses covariance

$$\varsigma^2 \begin{pmatrix} 1 & \mathcal{M}_\nu(\eta R) \\ \mathcal{M}_\nu(\eta R) & 1 \end{pmatrix}. \quad (7)$$

The true two-by-two covariance belongs to this family. Therefore its Gaussian KL minimizer is exactly

$$(\sigma_h^2)^\dagger = C_h(0), \quad \mathcal{M}_\nu(\alpha_h^\dagger R) = \rho_h(r). \quad (8)$$

Existence follows from $0 < \rho_h(r) < 1$ for $r \neq 0$. Positivity follows from $k \geq 0$ and $\mathcal{M}_\nu > 0$. Strict inequality follows because the smoothed field has spectral density $|\widehat{k}(h\omega)|^2 f_\nu(\omega)$, which is positive on a neighborhood of the origin, while $e^{i\omega^\top r}$ is not identically one there. Uniqueness follows from

$$\mathcal{M}'_\nu(x) = -\frac{2^{1-\nu}}{\Gamma(\nu)} x^\nu K_{\nu-1}(x) < 0, \quad x > 0. \quad (9)$$

For completeness, a standard route to pair-composite consistency would consider a fixed lattice, a lag r belonging to that lattice, and increasing rectangular Følner domains. Besides a compact parameter space, an interior unique target, and correlations bounded away from one, this route requires a uniform ergodic theorem for the pair log likelihood under a suitable integrable quadratic envelope. We do not establish that uniform law here. Overlapping pairs are dependent, so uncertainty would require a Godambe or long-run covariance, not an independence Hessian. We do not use fixed-domain asymptotics to claim separate consistency of variance and range; such a claim would conflict with Matérn microergodicity results [18, 21].

3 Small-support phase law

For $z = \alpha R$, define the directional kernel variance $a_k(e) = e^\top \Sigma_k e$ and

$$B_{\nu,k}(r) = \alpha^2 \left[a_k(e) \mathcal{M}''_\nu(z) + \{T_k - a_k(e)\} \frac{\mathcal{M}'_\nu(z)}{z} \right]. \quad (10)$$

Proposition 1 (Fixed nonzero lag). *For r in a compact annulus bounded away from zero and $h \leq \|r\|/(4L)$,*

$$\frac{C_h(r)}{\sigma^2} = \mathcal{M}_\nu(\alpha\|r\|) + h^2 B_{\nu,k}(r) + O(h^4). \quad (11)$$

The remainder is uniform when α also ranges over a compact subset of $(0, \infty)$.

The variance behaves differently because the Matérn function is expanded at the origin. Put

$$c_\nu = -\frac{\Gamma(-\nu)}{2^{2\nu}\Gamma(\nu)} = \frac{\Gamma(1-\nu)}{\nu 2^{2\nu}\Gamma(\nu)} > 0, \quad 0 < \nu < 1. \quad (12)$$

Theorem 1 (Matérn support phase law). *Fix $\nu > 0$. Under the preceding assumptions, as $h \downarrow 0$,*

$$\frac{C_h(0)}{\sigma^2} = 1 - c_\nu m_{2\nu}(\alpha h)^{2\nu} + O(h^2), \quad 0 < \nu < 1, \quad (13)$$

$$\frac{C_h(0)}{\sigma^2} = 1 - \frac{m_2}{2}(\alpha h)^2 \log\{1/(\alpha h)\} + O(h^2), \quad \nu = 1, \quad (14)$$

$$\frac{C_h(0)}{\sigma^2} = 1 - \frac{m_2}{4(\nu-1)}(\alpha h)^2 + o(h^2), \quad \nu > 1. \quad (15)$$

Moreover, the naive decay target in (8) satisfies

$$\alpha_h^\dagger - \alpha = \frac{\mathcal{M}_\nu(z)c_\nu m_{2\nu} \alpha^{2\nu}}{R\mathcal{M}'_\nu(z)} h^{2\nu} + O\left(h^{\min(4\nu, 2)}\right), \quad 0 < \nu < 1, \quad (16)$$

$$\alpha_h^\dagger - \alpha \sim \frac{m_2 \mathcal{M}_1(z) \alpha^2}{2R\mathcal{M}'_1(z)} h^2 \log(1/h), \quad \nu = 1, \quad (17)$$

$$\alpha_h^\dagger - \alpha = \frac{A_{\nu,k}(r)}{R\mathcal{M}'_\nu(z)} h^2 + \mathcal{R}_\nu(h), \quad \nu > 1, \quad (18)$$

where

$$A_{\nu,k}(r) = \alpha^2 \{(2\nu - 2)a_k(e) + T_k\} G_\nu(z) > 0, \quad (19)$$

$$G_\nu(z) = \frac{\mathcal{M}_\nu(z)}{2\nu - 2} + \frac{\mathcal{M}'_\nu(z)}{z} = \frac{2^{1-\nu}}{\Gamma(\nu)} \frac{z^\nu K_{\nu-2}(z)}{2\nu - 2} > 0, \quad (20)$$

and $\mathcal{R}_\nu(h) = O(h^{2\nu})$ for $1 < \nu < 2$, $O\{h^4 \log(1/h)\}$ for $\nu = 2$, and $O(h^4)$ for $\nu > 2$. For fixed ν , these remainders are uniform for α in a compact subset of $(0, \infty)$ and r in a compact annulus bounded away from zero. They are not uniform as ν approaches one or two.

Every leading coefficient in (16)–(18) is negative because $\mathcal{M}'_\nu(z) < 0$. Thus, for sufficiently small positive h ,

$$(\sigma_h^2)^\dagger < \sigma^2, \quad \rho_h(r) > \mathcal{M}_\nu(\alpha R), \quad \alpha_h^\dagger < \alpha. \quad (21)$$

The naive model overestimates range. The result does not require an isotropic smoother: anisotropy enters only through $a_k(e)$ and T_k , whose combination in (20) remains positive. For an isotropic kernel with $\Sigma_k = \sigma_k^2 I_d$,

$$A_{\nu,k}(r) = \sigma_k^2 \alpha^2 (2\nu + d - 2) G_\nu(z). \quad (22)$$

3.1 Genuine multi-lag misspecification

The two-point model is saturated: after variance refitting, one smoothed correlation is represented exactly by one point-support decay. Let $0 < R_1 < \dots < R_J$, $w_j > 0$, and $\mathbf{R}(\rho) = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}$. With $\rho_{h,j} = C_h(R_j e)/C_h(0)$, define

$$Q_h^C(\eta) = \sum_{j=1}^J w_j D_{\text{KL}}\{\mathcal{N}(0, \mathbf{R}(\rho_{h,j})) \parallel \mathcal{N}(0, \mathbf{R}(\mathcal{M}_\nu(\eta R_j)))\}. \quad (23)$$

Write

$$s_\nu(h) = \begin{cases} h^{2\nu}, & 0 < \nu < 1, \\ h^2 \log(1/h), & \nu = 1, \\ h^2, & \nu > 1, \end{cases} \quad (24)$$

and define $\kappa_j > 0$ from Theorem 1 by $\alpha - \alpha_{h,j}^\dagger = \kappa_j s_\nu(h) + o\{s_\nu(h)\}$.

Theorem 2 (Multi-lag composite target and residual KL). *Let α be interior to a compact positive parameter interval, and let α_h^C minimize (23) there. Put*

$$\rho_j = \mathcal{M}_\nu(\alpha R_j), \quad \mathcal{I}_j = \frac{1 + \rho_j^2}{(1 - \rho_j^2)^2}, \quad (25)$$

$$\lambda_j = w_j \mathcal{I}_j \{R_j \mathcal{M}'_\nu(\alpha R_j)\}^2, \quad \bar{\kappa} = \frac{\sum_j \lambda_j \kappa_j}{\sum_j \lambda_j}. \quad (26)$$

Then

$$\alpha - \alpha_h^C = \bar{\kappa} s_\nu(h) + o\{s_\nu(h)\}, \quad (27)$$

$$\min_\eta Q_h^C(\eta) = \frac{s_\nu(h)^2}{2} \sum_{j=1}^J \lambda_j (\kappa_j - \bar{\kappa})^2 + o\{s_\nu(h)^2\}. \quad (28)$$

Consequently, range inflation persists for the composite target, and the point-support model is genuinely misspecified to leading order whenever the lag-specific coefficients are not all equal.

The weights λ_j combine user weights, Gaussian correlation information, and sensitivity of the Matérn correlation to decay. Therefore the composite coefficient is not an arbitrary “representative lag” coefficient. The theorem concerns the population objective; dependence among overlapping pairs still enters Godambe uncertainty for a sample composite likelihood.

The one-term phase law is pointwise in fixed ν , so convergence slows near the threshold. The following refinement retains the two origin terms that cancel as $\nu \rightarrow 1$. For $0 < \nu < 2$, $\nu \neq 1$, put

$$b_\nu = \frac{\Gamma(-\nu)}{2^{2\nu}\Gamma(\nu)}, \quad W_{\nu,k}(h) = \frac{m_2(\alpha h)^2}{4(1-\nu)} + b_\nu m_{2\nu}(\alpha h)^{2\nu}. \quad (29)$$

At $\nu = 1$, define

$$\ell_{2,k} = \mathbb{E}\{\|D\|^2 \log\|D\|\}, \quad W_{1,k}(h) = \frac{(\alpha h)^2}{2} [m_2\{\log(\alpha h/2) + \gamma_E - 1/2\} + \ell_{2,k}], \quad (30)$$

where $0 \log 0 = 0$. For sufficiently small $h > 0$, let

$$\tilde{\rho}_h(r) = \frac{\mathcal{M}_\nu(z) + h^2 B_{\nu,k}(r)}{1 + W_{\nu,k}(h)}, \quad \mathcal{M}_\nu(\tilde{\alpha}_h R) = \tilde{\rho}_h(r). \quad (31)$$

Proposition 2 (Transition-aware pair approximation). *For fixed $0 < \nu < 2$, the approximation in (31) satisfies*

$$\alpha_h^\dagger - \tilde{\alpha}_h = \begin{cases} O(h^{2\nu+2}), & 0 < \nu < 1, \\ O\{h^4 \log(1/h)\}, & \nu = 1, \\ O(h^4), & 1 < \nu < 2. \end{cases} \quad (32)$$

Moreover, for every fixed $h > 0$, $\lim_{\nu \rightarrow 1} W_{\nu,k}(h) = W_{1,k}(h)$. The error orders are pointwise in ν ; no joint uniform remainder in (ν, h) is asserted.

The qualitative fact that convolution can induce covariance anisotropy is known [7, 16]. The next corollary identifies the support-induced directional difference in the pairwise pseudo-inverse-range, including in the rough regimes where the leading common shift is not quadratic.

Corollary 1 (Directional support contrast). *For a unit vector e , let $\Delta_e(h) = \alpha - \alpha_h^\dagger(e)$, where the pair lag is $r = Re$ with the same fixed $R > 0$ in every direction. For two unit vectors e_1, e_2 , under the assumptions of the phase theorem,*

$$\Delta_{e_1}(h) - \Delta_{e_2}(h) = \frac{\alpha^2 \{a_k(e_1) - a_k(e_2)\}}{R} \frac{K_{\nu-2}(z)}{K_{\nu-1}(z)} h^2 + o(h^2). \quad (33)$$

This formula holds for every fixed $\nu > 0$. For $0 < \nu < 1$ it is lower order than the common order- $h^{2\nu}$ range shift, and at $\nu = 1$ it is lower order than the common $h^2 \log(1/h)$ shift.

For a concrete two-dimensional family, let the base product Epanechnikov kernel be transformed according to

$$Z_{h,A}(t) = \int_{\mathbb{R}^2} k(u) Y(t - hAu) du, \quad A_\varrho = \sqrt{\frac{2}{\varrho + \varrho^{-1}}} \text{diag}(\sqrt{\varrho}, \varrho^{-1/2}), \quad (34)$$

where $\varrho \geq 1$. Then $\Sigma_k = A_\varrho A_\varrho^\top / 5$, its trace is fixed as ϱ varies, and its principal-variance ratio is ϱ^2 . For $\nu > 1$, the ratio of the major- to minor-axis leading shift coefficients in (18) is

$$\frac{(2\nu - 1)\varrho^2 + 1}{\varrho^2 + 2\nu - 1}. \quad (35)$$

Thus elongated observation support can yield direction-dependent inferred range even though the latent covariance is isotropic. Equation (33), rather than this qualitative phenomenon, is the theorem-level statement used by the directional benchmark.

Remark 1 (Why symmetry does not imply an h^2 bias). Symmetry removes the order- h term at a fixed nonzero lag. When $0 < \nu < 1$, however, the origin expansion contains $\|r\|^{2\nu}$, which changes the variance at order $h^{2\nu}$. Normalizing by that variance dominates the order- h^2 off-diagonal change. At $\nu = 1$, the origin term is $h^2 \log(1/h)$. Only for $\nu > 1$ do ordinary second derivatives control both pieces.

Remark 2 (The fitted lag is fixed). The expansion holds for R bounded away from zero. If $R = ch$, the nonzero-lag Taylor expansion is no longer uniform and the relative target displacement need not vanish. No shrinking-lag conclusion is claimed.

4 Support-aware finite-grid likelihood

The continuous theorem concerns translation-invariant interior averaging. The finite experiment starts from raw sites $X_{\text{in}} = (x_1, \dots, x_n)$ and a rectangular row-stochastic operator $S_h \in \mathbb{R}^{m \times n}$. If independent measurement noise is present before smoothing,

$$\mathbf{y} = S_h \{Y(X_{\text{in}}) + \varepsilon\}, \quad \varepsilon \sim \mathcal{N}_n(0, \tau^2 I_n), \quad (36)$$

then the correct covariance is

$$\Sigma_{\text{corr}}(\alpha) = S_h \{K_\alpha(X_{\text{in}}, X_{\text{in}}) + \tau^2 I_n\} S_h^\top. \quad (37)$$

The naive declared-center point-support model instead uses

$$\Sigma_{\text{naive}}(\alpha) = K_\alpha(X_{\text{out}}, X_{\text{out}}) + \tau^2 I_m. \quad (38)$$

Besides ignoring support, (38) incorrectly treats pre-smoothing noise as independent post-smoothing noise. With variance, smoothness, and nugget fixed at their generating values, the finite-design population target minimizes

$$Q_n(\alpha) = \frac{1}{2} [\log \det \Sigma(\alpha) + \text{tr}\{\Sigma(\alpha)^{-1} \Sigma_0\}], \quad (39)$$

where constants independent of α are omitted. The corrected family contains Σ_0 exactly, provided the known operator and fixed nuisance parameters are correct and α is identifiable on the finite design; the naive family generally does not. Our primary support-only experiment sets $\tau = 0$, preventing noise timing from confounding the support effect. A separate combined-misspecification run with $\tau^2 = 0.05$ is retained as a reproducibility stress artifact but is not used for the principal claims.

For a local full-likelihood result, let $x_1, \dots, x_p$ be distinct interior sites, let $\theta = (\log \sigma^2, \log \alpha)^\top$, and write $\Sigma(\theta)$ for their point-support Matérn covariance at fixed smoothness. The entrywise origin and fixed-lag expansions imply

$$\Sigma_h = \Sigma_0 + s_\nu(h) \Gamma_\nu + o\{s_\nu(h)\}. \quad (40)$$

Let $P = \Sigma_0^{-1}$, define the Fisher inner product

$$\langle A, B \rangle_P := \frac{1}{2} \text{tr}(PAPB), \quad (41)$$

and let $\dot{\Sigma}_j = \partial_j \Sigma(\theta_0)$. Define

$$\mathcal{J}_{jk} = \langle \dot{\Sigma}_j, \dot{\Sigma}_k \rangle_P, \quad g_{\nu,j} = \langle \dot{\Sigma}_j, \Gamma_\nu \rangle_P. \quad (42)$$

Proposition 3 (Finite-design full-likelihood projection). *Suppose θ_0 is interior to a compact parameter set, $\mathcal{J}$ is nonsingular, and $\theta_h^\dagger$ is the local Gaussian KL minimizer converging to θ_0 . With $\beta_\nu = \mathcal{J}^{-1} g_\nu$,*

$$\theta_h^\dagger - \theta_0 = \beta_\nu s_\nu(h) + o\{s_\nu(h)\}, \quad (43)$$

$$\min_{\theta} D_{\text{KL}}\{\mathcal{N}_p(0, \Sigma_h) \parallel \mathcal{N}_p(0, \Sigma(\theta))\} = \frac{1}{4} \text{tr}(PE_\nu PE_\nu) s_\nu(h)^2 + o\{s_\nu(h)^2\}, \quad (44)$$

where $E_\nu = \Gamma_\nu - \sum_{j=1}^2 \beta_{\nu,j} \dot{\Sigma}_j$ satisfies $\langle \dot{\Sigma}_j, E_\nu \rangle_P = 0$ for $j = 1, 2$. Thus the full likelihood is genuinely misspecified to leading order exactly when the support perturbation is not contained in the variance-decay covariance tangent space.

The decay coordinate of $\mathcal{J}^{-1} g_\nu$ is design specific, so the proposition does not assert a universal full-likelihood sign. It is a finite-design local result, not a fixed- or increasing-domain consistency theorem. Its assumptions make identifiability explicit through nonsingularity of $\mathcal{J}$.

A rectangular S_h is intentional. If a square parameter-independent smoother is invertible, the exact transformed likelihood is merely a change of coordinates and need not lose information. Rectangular aggregation represents the scientifically relevant reduction from raw locations to fewer supported measurements.

Boundary row normalization is not covered by the interior theorem. For two location-dependent scaled kernels with means μ_s and μ_t , a direct fixed-lag expansion gives

$$C_{h,s,t}(r) = C(r) + h \nabla C(r)^\top (\mu_s - \mu_t) + O(h^2). \quad (45)$$

A truncated kernel can therefore introduce an order- h term and destroy stationarity. The interior theorem provides no universal boundary sign. We report one boundary-retaining and one fixed jittered design as illustrations, not as a boundary scaling study.

4.1 Finite-library high-dimensional certificate

The replicated-field benchmark jointly varies variance and decay on a deterministic finite grid. The next proposition is a direct consequence of standard Gaussian quadratic-form concentration. It is included to state precisely what the benchmark certifies; it is not presented as a new concentration inequality.

Proposition 4 (Gaussian finite-library certificate). *Let $X_1, \dots, X_N \stackrel{\text{iid}}{\sim} \mathcal{N}_p(0, \Sigma_0)$, where Σ_0 is positive definite, and let $\{\Sigma_\theta : \theta \in \mathcal{G}\}$ be a deterministic library of $M < \infty$ positive-definite covariance matrices. Define*

$$\begin{aligned}\widehat{\Sigma}_N &= \frac{1}{N} \sum_{i=1}^N X_i X_i^\top, \\ \widehat{L}_N(\theta) &= \frac{1}{2p} \left\{ \log \det \Sigma_\theta + \text{tr}(\Sigma_\theta^{-1} \widehat{\Sigma}_N) \right\}, \\ L(\theta) &= \mathbb{E} \widehat{L}_N(\theta), \quad A_\theta = \Sigma_0^{1/2} \Sigma_\theta^{-1} \Sigma_0^{1/2}.\end{aligned}\tag{46}$$

For any $\delta \in (0, 1)$, put $t = \log(2M/\delta)$. With probability at least $1 - \delta$, simultaneously for all $\theta \in \mathcal{G}$,

$$\left| \widehat{L}_N(\theta) - L(\theta) \right| \leq \frac{\|A_\theta\|_F}{p} \sqrt{\frac{t}{N}} + \frac{\|A_\theta\|_{\text{op}}}{p} \frac{t}{N} =: u_\theta.\tag{47}$$

If $\widehat{\theta}$ and θ^* minimize $\widehat{L}_N$ and L , respectively, over $\mathcal{G}$, then on the same event

$$L(\widehat{\theta}) - L(\theta^*) \leq 2 \max_{\theta \in \mathcal{G}} u_\theta.\tag{48}$$

Moreover, $L(\theta)$ differs by a constant independent of θ from

$$\frac{1}{p} D_{\text{KL}}\{\mathcal{N}_p(0, \Sigma_0) \parallel \mathcal{N}_p(0, \Sigma_\theta)\}.\tag{49}$$

No coordinate independence is assumed: all spatial dependence within one field is contained in Σ_0 . Independence is required only across the N replicated fields. With quadratic effective rank

$$r_2(A_\theta) = \frac{\|A_\theta\|_F^2}{\|A_\theta\|_{\text{op}}^2},$$

the candidate radius is

$$\frac{\|A_\theta\|_{\text{op}}}{p} \left\{ \sqrt{\frac{r_2(A_\theta)t}{N}} + \frac{t}{N} \right\}.$$

Thus p helps only through controlled relative spectral geometry, not by itself. If constants $0 < c \leq C < \infty$ satisfy

$$c\Sigma_0 \preceq \Sigma_\theta \preceq C\Sigma_0 \quad \text{for every } \theta \in \mathcal{G},\tag{50}$$

then $\|A_\theta\|_{\text{op}} \leq c^{-1}$ and $\|A_\theta\|_F \leq \sqrt{p}/c$, so

$$\max_{\theta} u_\theta \leq \frac{1}{c} \left\{ \sqrt{\frac{\log(2M/\delta)}{Np}} + \frac{\log(2M/\delta)}{Np} \right\}.\tag{51}$$

This is a conditional information rate, not a claim of separate fixed-domain consistency for Matérn variance and range. Relative spectral constants and criterion curvature can deteriorate with the design. The benchmark therefore records the actual Frobenius and operator norms and uses the exact radius in (47). A proof is given in Appendix E.

Two-dimensional Matérn support-bias phase law




Figure 1: Continuous two-dimensional pseudo-decay shift under product Epanechnikov smoothing. Left: exact shifts (solid) and theorem leading terms with their explicit coefficients (dashed). Right: their ratio, which approaches one as $h \downarrow 0$.

5 Synthetic validation

5.1 Continuous two-dimensional quadrature

We used a product Epanechnikov density

$$k(u) = \prod_{j=1}^d \frac{3}{4} (1 - u_j^2) \mathbf{1}_{\{|u_j| \leq 1\}}. \quad (52)$$

The one-dimensional density of $U_j - V_j$ is available in closed form, so tensor Gauss–Legendre quadrature evaluates (4) deterministically. We used $d = 2$, $\alpha = R = 1$, quadrature order 96 per coordinate, 18 geometrically spaced bandwidths from 0.003 to 0.3, and $\nu \in \{0.25, 0.5, 0.75, 1, 1.5, 2.5\}$. For each cell we solved (8) by monotone root finding. Repeating the entire grid at quadrature orders 64 and 128 changed pseudo-decay by at most 4.4×10^{-8} and 5.2×10^{-9} , respectively, relative to order 96. For this kernel $L = \sqrt{2}$, so the two largest displayed bandwidths, 0.229 and 0.3, are stress points outside the sufficient Taylor neighborhood $h \leq R/(4L)$. All slope and coefficient checks below use the six smallest bandwidths, which are inside that neighborhood.

All 108 computed decay shifts $\alpha - \alpha_h^\dagger$ were positive. An ordinary least-squares slope over the six smallest bandwidths on log–log axes was 0.515, 0.999, and 1.468 for $\nu = 0.25, 0.5, 0.75$, compared with theoretical powers 0.5, 1, and 1.5. At $\nu = 1$, the raw slope was 1.817, as expected from the logarithmic modifier rather than a pure power. The slopes for $\nu = 1.5$ and 2.5 were 1.994 and 2.000. Figure 1 displays the full paths and the coefficient-level theorem comparison. At the smallest bandwidth, exact-shift to leading-term ratios were 1.021, 1.000, 0.958, 1.059, 0.997, and 1.000 in increasing order of ν .

Table 1: Smallest-bandwidth coefficient ratios across kernel families and dimensions.

Kernel	d	Minimum ratio	Maximum ratio
Epanechnikov	1	0.999	1.169
Epanechnikov	2	0.998	1.056
Epanechnikov	3	0.997	1.005
Uniform	1	0.998	1.130
Uniform	2	0.997	1.016
Uniform	3	0.966	1.000

5.2 Dimension and kernel robustness audit

The primary phase display fixes $d = 2$ and product Epanechnikov smoothing. We separately predeclared a deterministic audit over $d \in \{1, 2, 3\}$, product Epanechnikov and product uniform kernels, $\nu \in \{0.5, 1, 1.5, 2.5\}$, and $h \in \{0.002, 0.004, 0.008\}$, with $\alpha = R = 1$. Tensor quadrature used order 48 per coordinate, with orders 32 and 64 as refinements. Define

$$s_\nu(h) = \begin{cases} h^{2\nu}, & 0 < \nu < 1, \\ h^2 \log(1/h), & \nu = 1, \\ h^2, & \nu > 1. \end{cases}$$

Table 1 gives the range, over smoothness, of the ratio between the exact shift and $C_{\nu,k}s_\nu(h)$ at $h = 0.002$, where $C_{\nu,k}$ is the relevant coefficient from Theorem 1.

All 72 exact shifts were positive. The maximum absolute coefficient-ratio error was 0.169, below the predeclared 0.20 gate, and occurred at $\nu = 1$, where logarithmic convergence is slow. Repeating all exact targets at the two refinement orders changed a shift by at most 3.92×10^{-7} relative to the order-48 shift, below the predeclared 2×10^{-4} gate. This is a finite robustness audit, not a proof or a uniform statement over kernel families or dimensions.

5.3 Threshold-layer stress audit

The one-term theorem is pointwise in fixed smoothness, and its coefficient is therefore not a uniformly reliable finite-bandwidth approximation near $\nu = 1$. We evaluated 111 deterministic cells with $\nu \in [0.55, 1.45]$ at spacing 0.025 and $h \in \{0.01, 0.02, 0.05\}$. The minimum ratio of the exact shift to the regime-specific one-term shift was 0.154. Thus a literal switch between the rough, logarithmic, and smooth leading laws is unreliable in the threshold layer: the one-term prediction was up to 6.51 times the exact shift.

The transition-aware target in Proposition 2 had maximum relative pseudo-decay-shift error 0.099% and maximum relative error in variance loss 0.084% over the same grid. Every exact and approximate shift was positive. Relative to order 96, quadrature orders 64 and 128 changed an exact pseudo-decay by at most 4.0×10^{-9} and 3.1×10^{-10} of its exact shift, respectively. These numerical results resolve a finite- h crossover; they do not strengthen the pointwise remainder in Proposition 2 into a joint uniform theorem in (ν, h) .

5.4 Multi-lag and full-likelihood projection audits

We evaluated Theorem 2 with equal weights at $\alpha R \in \{0.5, 1, 1.5, 2\}$, $\nu \in \{0.5, 1, 1.5, 2.5\}$, and $h \in \{0.005, 0.01, 0.02\}$. All 12 composite targets had positive $\alpha - \alpha_h^C$ and strictly positive minimum

Finite-bandwidth stress audit around smoothness one




Figure 2: Finite-bandwidth transition stress around $\nu = 1$. Left: exact inverse-range shift divided by the fixed- ν one-term approximation. Right: relative error of the cancellation-aware two-term approximation on a logarithmic scale.

KL. At $h = 0.005$, the largest relative errors in the leading shift and minimum-KL formulas were 1.41% and 9.85%. The lag coefficients are materially different: for $\nu = 0.5$, they decrease from 1.61 at $\alpha R = 0.5$ to 0.40 at $\alpha R = 2$. Thus the residual in (28) is genuine and the composite coefficient is visibly lag-sensitive.

For Proposition 3, we used a 3×3 unit-spaced regular grid and jointly profiled variance and decay. All 12 exact continuously smoothed targets had positive minimum KL and, for this design, range inflation. At the smallest bandwidth, the largest relative errors in the projected decay shift, log-variance shift, and minimum KL were 5.42%, 12.68%, and 1.77%, respectively. This validates the general tangent formula on one declared design without promoting its observed decay sign into a universal claim.

5.5 Directional support oracle

The directional track used (34) with $\varrho \in \{1, 4\}$, 19 lag angles from zero to 90 degrees, $\nu \in \{0.5, 1, 1.5, 2.5\}$, and 14 bandwidths from 0.003 to 0.15. All 2,128 computed inverse-range shifts were positive. At $\varrho = 4$, the major/minor smooth-regime coefficient ratios were 1.833 for $\nu = 1.5$ and 3.250 for $\nu = 2.5$, agreeing with (35). At $h = 0.15$, apparent range inflation along the major and minor support axes was 13.46% and 11.47% for $\nu = 0.5$, and 1.56% and 0.76% for $\nu = 1.5$.

Let D_ν denote the coefficient multiplying h^2 on the right side of (33) for the major-minus-minor comparison. At the smallest bandwidth, the absolute error in the ratio of the exact directional contrast to $D_\nu h^2$ was at most 4.3×10^{-6} over all four smoothnesses. Relative to quadrature order 96, orders 64 and 128 changed an individual shift by at most 1.3×10^{-8} of that shift and changed a directional contrast by at most 1.2×10^{-9} of that contrast.




Figure 3: Lag sensitivity and local-likelihood validation. Left: lag-specific pair coefficients and information-weighted composite coefficients (dotted). Middle and right: exact-to-leading ratios for the pseudo-parameter displacement and irreducible minimum KL.

5.6 Finite-grid full likelihood

The core design uses a 19×19 input lattice with spacing 0.25 and every second location as a candidate output center. Interior outputs are trimmed by 0.75 in each coordinate. We set $\sigma^2 = \alpha = 1$, $\tau^2 = 0$, $\nu \in \{0.5, 1, 1.5, 2.5\}$, and $h \in \{0, 0.3, 0.5, 0.7\}$. Radial Epanechnikov rows are normalized exactly. The zero-bandwidth operator is an exact rectangular selector. Stress panels retain boundary centers, perturb each input coordinate uniformly in $[-0.04, 0.04]$ before clipping to the original bounding box, and examine one-dimensional increasing-domain lattices with 100, 225, and 400 raw sites. The irregular panel uses one fixed jitter realization.

For each configuration, the finite-design population criterion is minimized by bounded scalar optimization over $\log \alpha \in [\log(0.02), \log(5)]$. Replicate likelihoods are evaluated on a 161-point equally spaced log-decay profile; an interior grid minimum is refined by three-point quadratic interpolation, while an endpoint minimum is retained as a boundary fit. Gaussian raw-observation vectors with the declared covariance are transformed by S_h . Configurations sharing all parameters except h reuse standard-normal draws. No numerical jitter is added. A Cholesky or population-optimization failure aborts its configuration and makes the reducer mark the run incomplete; no such failure occurred.

The promoted study used 200 replicates in each of 21 configurations, producing 8,400 fitted estimates. The reducer verified configuration hashes, row counts, and unique model-replicate keys for every task shard. Independent checks also verified finite numerical fields, expected model labels, replicate indices, and metadata. Across the 16 core two-dimensional configurations, the corrected population target differed from the true decay by at most 1.3×10^{-7} . At the largest bandwidth, $h = 0.7$, the naive population targets were 0.152, 0.414, 0.573, and 0.740 for $\nu = 0.5, 1, 1.5, 2.5$, respectively. Their corresponding Monte Carlo means were 0.151, 0.407, 0.570, and 0.738; the corrected means were 1.090, 0.990, 0.990, and 0.994. The comparatively noisy corrected estimate at $\nu = 0.5$ nevertheless remained within two Monte Carlo standard errors of its population target.

No optimizer or covariance failure was discarded. Six of the 6,400 core fits landed at the declared bounds, and those fits remain in Figure 5 and Table 2. Because that boundary stress also changed output dimension, we added translated boundary and interior 4×4 blocks on the same latent grid. Both have $p = 16$ and identical relative geometry. At $h = 0.7$, point-support decay targets differed by 0.030–0.062 between the blocks. A model using the correct support shape but bandwidth $0.75h$




Figure 4: An isotropic latent field observed through elongated support. Left: inferred range inflation versus lag angle for aspect parameter $\rho = 4$ and $h = 0.15$. Right: the exact major–minor inverse-range contrast divided by its theorem term $D_\nu h^2$.

reduced population KL to 16.7–32.3% of point-support KL, while exact support recovered decay within 4.1×10^{-7} . Table 3 gives the matched comparison. For the fixed jittered design the target changed from 0.289 to 0.284 at $h = 0.5$. In the one-dimensional domain-growth check, increasing the number of raw sites from 100 to 400 reduced the corrected estimator standard deviation from 0.529 to 0.177 and the naive standard deviation from 0.151 to 0.068, while at $n = 400$ their means remained within 0.026 and 0.009, respectively, of their population targets (Figure 6).

5.7 Joint smoothness–decay and partial-support comparison

We jointly selected (ν, α, σ^2) from a fixed library with 18 smoothness values, 52 log-spaced decay values, and analytically profiled variance. The design has an 11×11 latent grid and 16 interior outputs, with true $\nu \in \{0.5, 1, 1.5, 2.5\}$, $h \in \{0.35, 0.5\}$, $N = 16$ independent fields, and 100 trials per cell. Exact support, $0.75h$ support, and point support produced 2,400 fits.

The exact-support population target was the physical parameter in all eight design cells. Point support changed smoothness in all eight, always upward. At $h = 0.5$, for example, targets moved from $(0.5, 1)$ to $(2, 2.176)$ and from $(2.5, 1)$ to $(3.5, 1.338)$. Consequently, every point-support joint target had $\alpha^* > 1$, the reverse of the fixed-smoothness direction: aggregation was absorbed mainly by a smoother fitted field. The partial-support population criterion improved on point support in all eight cells. This gives a strictly more informative method comparison than exact model containment alone. Monte Carlo medians generally followed the discrete population targets in Figure 7.

Table 2: Finite-design validation at $h = 0.7$. The table is generated from immutable task shards; no failed or boundary-hitting fit is silently removed.

ν	Model	Target	Mean	MCSE	Bound hits
0.5	Corrected	1.000	1.090	0.047	3
0.5	Naive	0.152	0.151	0.004	0
1	Corrected	1.000	0.990	0.010	0
1	Naive	0.414	0.407	0.005	0
1.5	Corrected	1.000	0.990	0.007	0
1.5	Naive	0.573	0.570	0.005	0
2.5	Corrected	1.000	0.994	0.004	0
2.5	Naive	0.740	0.738	0.003	0

Table 3: Matched-size boundary comparison at $h = 0.7$. I and B denote interior and boundary blocks. KL ratios compare the $0.75h$ -support model with point support.

ν	$\alpha_{\text{point},I}^*$	$\alpha_{\text{point},B}^*$	$\text{KL}_{.75h}/\text{KL}_{\text{point},I}$	$\text{KL}_{.75h}/\text{KL}_{\text{point},B}$	p
0.5	0.225	0.183	0.167	0.234	16
1	0.391	0.337	0.187	0.260	16
1.5	0.525	0.495	0.202	0.281	16
2.5	0.709	0.771	0.214	0.323	16




Figure 5: Finite-design population targets and Monte Carlo mean decay estimates for corrected and naive likelihoods. Error bars show two Monte Carlo standard errors of the mean, not estimator confidence intervals.




Figure 6: Increasing-domain check for the finite-grid likelihood. Lines are nonrandom finite-design targets and points are Monte Carlo means from 200 replicates; error bars show two Monte Carlo standard errors of each mean.




Figure 7: Joint smoothness-inverse-range targets. Circles are population library minimizers, crosses are Monte Carlo medians, and stars are physical parameters. The partial-support model uses the right shape and 75% of the true bandwidth.




Figure 8: One synthetic field before and after the known rectangular observation operator: a 23×23 latent realization (left) and its 10×10 locally averaged observations (right). This illustration is generated from the same covariance and operator implementation as the likelihood benchmark.

5.8 Replicated-field finite-library experiment

To connect the probability statement to an estimator, we generated independent replicated spatial fields on increasing-domain two-dimensional designs at fixed lattice spacing. For $q \in \{4, 6, 8, 10\}$, a padded $(2q + 3) \times (2q + 3)$ latent lattice with spacing 0.25 was mapped by a full-row-rank radial Epanechnikov operator of bandwidth 0.5 to a $q \times q$ output lattice, giving $p = q^2 \in \{16, 36, 64, 100\}$. The mean was known to be zero. We used $\nu \in \{0.5, 1.5\}$, $\sigma^2 = \alpha = 1$, and $N \in \{1, 4, 16, 64\}$ independent fields. The candidate set was an explicit deterministic $101 \times 161 = 16,261$ Cartesian library of variance and inverse-range values spanning $[0.35, 2.5] \times [0.15, 1.6]$ and containing the generating parameter pair $(1, 1)$ exactly. This truth anchoring is an oracle benchmark control, not a deployable rule for choosing a data-analysis grid. The complete library was fixed before the final Monte Carlo run. Corrected candidates were $\sigma^2 S_h R_\alpha S_h^\top$; naive candidates were $\sigma^2 R_\alpha$ at the declared output centers. Thus only the corrected covariance library contained the data law. Every empirical minimizer was obtained by exact finite-library enumeration, without continuous interpolation.

For each (p, ν) cell, 200 nested replicate-field sequences were generated from the support-aware truth. Thus there were 64 distinct model- p - ν - N certificate cells and 12,800 fitted estimates. The confidence parameter was $\delta = 0.05$, and the theorem radius was computed from the actual candidate-specific Frobenius and operator norms. No independence approximation was made for coordinates within a field. The certificate was applied separately to each fixed design, model, smoothness, and sample size; a joint 95% statement over all 64 reported cells is not claimed. The raw illustration in Figure 8 shows the observation map used for the largest design.

The discrete population objective was within 2.63×10^{-5} per coordinate of the continuously profiled objective in all 16 design-model cases, below the 5×10^{-5} tolerance fixed before the final Monte Carlo run. Maximum continuous-to-grid parameter distances were 0.00499 in decay and 0.0297 in variance for point support, and below 4.1×10^{-7} for both coordinates under exact support. Thus the finite-library parameter floor is quantified directly. In every one of the 64 model- p - ν - N

High-dimensional SupportShift likelihood experiment




Figure 9: The replicated-field finite-library benchmark. Left: empirical 95th percentiles of the worst likelihood deviation versus the worst-candidate theorem radius; exact candidatewise coverage is reported in the text. Center: inverse-range RMSE to each model’s finite-grid KL target versus Np . Right: for $p = 100$ and $\nu = 1.5$, increasing N concentrates the corrected estimator near the physical truth and the naive estimator near its displaced target.

cells, all 200 trials satisfied the actual simultaneous candidatewise event

$$|\hat{L}_N(\theta) - L(\theta)| \leq u_\theta \quad \text{for every } \theta \in \mathcal{G}.$$

Over all 12,800 fits, the largest value of $\max_\theta |\hat{L}_N(\theta) - L(\theta)|/u_\theta$ was 0.789. The weaker worst-radius envelope displayed in Figure 9 also held throughout: across cells, the 95th percentile of $\max_\theta |\hat{L}_N - L|$ was at most 0.402 times $\max_\theta u_\theta$, with median ratio 0.298. These 100% empirical coverages show that the finite-library union bound was conservative here; each 95% certificate applies to its own fixed cell, not jointly across all 64.

For each fixed model, dimension, and smoothness, a log–log regression of the 95th percentile criterion deviation on N produced slopes from -0.528 to -0.454 , with median -0.505 , matching the $N^{-1/2}$ leading dependence in (47). At fixed N , the same regression on p gave slopes from -0.509 to -0.297 , with median -0.413 . The dimension scaling is weaker and variable because the theorem depends on relative operator norms and effective rank rather than coordinate count alone. The exact deterministic ERM inequality $L(\hat{\theta}) - L(\theta^*) \leq 2 \max_\theta |\hat{L}_N(\theta) - L(\theta)|$ held for every fitted trial.

The support-aware library had population target $(\sigma^2, \alpha) = (1, 1)$ in every design. In contrast, at $p = 100$ the naive finite-grid inverse-range targets were 0.320 for $\nu = 0.5$ and 0.594 for $\nu = 1.5$, close to continuously profiled targets 0.318 and 0.589. For $p = 100, \nu = 1.5$, increasing N from 1 to 64 reduced support-aware decay RMSE from 0.206 to 0.024 and naive RMSE to its own KL target from 0.124 to 0.015. Yet the naive RMSE to the physical inverse range was 0.409 at $N = 64$, and its mean estimate was 0.591. The rough case was more pronounced: at $p = 100, \nu = 0.5, N = 64$, naive RMSE was 0.015 to its target but 0.681 to the physical value. Thus sampling spread decreased while support misspecification remained. Figure 9 visualizes this “more information, more confidently wrong” distinction. The parameter-error panels are empirical summaries, not consequences of Proposition 4, which controls criterion deviation and excess risk unless additional curvature is imposed.

6 Discussion

The theorem separates two effects that are often conflated. Linear propagation of covariance through a known support is standard. The new object here is the parameter selected when that propagation is ignored. For rough Matérn fields, the origin singularity makes the pseudo-target move at a fractional power of bandwidth. For smooth fields, a Bessel recurrence shows that the combined variance and nonzero-lag correction still increases normalized correlation, including under anisotropic smoothing.

The simulations have a deliberate role: they validate theorem-linked generators, compute nonrandom finite-design KL targets, and distinguish misspecification from finite-sample error. They are not substitutes for the phase-law proof. The zero-support case verifies exact agreement, the corrected target verifies model containment, and the boundary and irregular panels expose where the interior theorem stops. The directional oracle checks a coefficient rather than only a sign. The replicated-field experiment separately checks the finite-library probability event and demonstrates why shrinking sampling error does not repair a displaced population target.

Several limitations define the scope of the result. First, the universal sign and phase law hold at fixed ν . This restriction is substantive because information about Matérn smoothness can be both appreciable and design-dependent [8]. The joint finite-library experiment shows, but does not theorize, how nuisance movement can change the direction. Jointly estimating bandwidth, smoothness, variance, and range may be weakly identified. Second, Theorem 2 concerns a declared pair-composite criterion. Proposition 3 is a local finite-design result with a design-specific coefficient, not a universal full-likelihood sign theorem. Third, fixed-resolution discrete averaging does not converge to continuous convolution merely because the spatial domain grows. A mixed-domain quadrature limit or an exact discrete-weight theorem is required. Fourth, boundary normalization creates location-dependent supports and can introduce first-order terms. Fifth, the original finite-design validation fixes nuisance parameters to isolate the target shift. The replicated-field track jointly varies variance and decay, but only over a declared finite library; it does not establish a continuous-parameter rate. Sixth, the exact certificate radius uses the generating Σ_0 . It is an oracle simulation certificate, not a plug-in confidence procedure for unknown real-data covariance.

These restrictions are preferable to an unsupported broad claim. They also suggest direct extensions: a uniform discrete-to-continuous approximation, continuous joint nuisance profiling, spectral full-likelihood expansions, curvature conditions translating likelihood error into parameter error, and explicit boundary counterexamples. The finite-library result gives a useful high-dimensional probability framing, but the concentration step itself is standard supporting machinery. The paper’s novelty rests on the explicit support-distortion phase law, the multi-lag and finite-design KL projections, and the directional coefficient, not on a new quadratic-form inequality.

7 Conclusion

For a fixed nonzero-lag correlation match under translation-invariant interior averaging, ignored support makes a Matérn field look more persistent than it is. The magnitude is controlled by smoothness: $h^{2\nu}$, $h^2 \log(1/h)$, or h^2 , while anisotropic support produces an explicit order- h^2 contrast between lag directions. A multi-lag composite retains an information-weighted shift and a nonzero residual KL; an arbitrary fixed design has the corresponding Fisher-metric covariance projection. A support-aware observation covariance removes the finite-design population shift when the operator is known, nuisance parameters are correctly specified, and decay is identifiable. For a deterministic covariance library fitted to independent replicated fields, a simultaneous Gaussian likelihood certificate quantifies the sampling-error component without assuming independent spatial coordinates. Joint

smoothness fitting can reverse the fixed-smoothness decay direction. Together, these results diagnose both the population displacement and the finite-sample concentration around it; they make no universal claim for boundary supports or arbitrary continuous full-likelihood fits.

A Proof of the phase theorem

Proof. Let $f_\alpha(x) = \mathcal{M}_\nu(\alpha\|x\|)$. Since r is bounded away from zero and D is bounded, a fourth-order Taylor expansion is uniform. Because $D = U - V$ has a centrally symmetric law, both $\mathbb{E}D = 0$ and $\mathbb{E}(D^{\otimes 3}) = 0$; moreover $\mathbb{E}(DD^\top) = 2\Sigma_k$. Therefore

$$\mathbb{E}f_\alpha(r + hD) = f_\alpha(r) + \frac{h^2}{2} \text{tr}\{\nabla^2 f_\alpha(r)\mathbb{E}(DD^\top)\} + O(h^4). \quad (53)$$

For $r = Re$,

$$\nabla^2 f_\alpha(r) = \alpha^2 \mathcal{M}_\nu''(z)ee^\top + \alpha^2 \frac{\mathcal{M}_\nu'(z)}{z}(I - ee^\top), \quad (54)$$

which proves (11).

For noninteger ν , the origin expansion is

$$\mathcal{M}_\nu(x) = \sum_{j=0}^{\infty} \frac{(x^2/4)^j}{j!(1-\nu)_j} + \frac{\Gamma(-\nu)}{2^{2\nu}\Gamma(\nu)} x^{2\nu} \sum_{j=0}^{\infty} \frac{(x^2/4)^j}{j!(1+\nu)_j}. \quad (55)$$

Taking expectations at $x = \alpha h\|D\|$ yields (13) and (15). At $\nu = 1$,

$$\mathcal{M}_1(x) = 1 + \frac{x^2}{2} \{\log(x/2) + \gamma_E - 1/2\} + O\{x^4(1 + |\log x|)\}, \quad (56)$$

which proves (14).

At the second transition,

$$\mathcal{M}_2(x) = 1 - \frac{x^2}{4} + x^4 \left\{ \frac{3}{64} - \frac{\log(x/2) + \gamma_E}{16} \right\} + O\{x^6(1 + |\log x|)\}. \quad (57)$$

This gives an $O\{h^4 \log(1/h)\}$ variance remainder at $\nu = 2$. For fixed $\nu > 2$, four-times differentiability at the origin gives

$$\mathcal{M}_\nu(x) = 1 - \frac{x^2}{4(\nu-1)} + O(x^4).$$

For $1 < \nu < 2$, the fractional term in (55) is $O(x^{2\nu})$.

Divide the fixed-lag expansion by the relevant variance expansion and apply a first-order Taylor expansion to the inverse of $\alpha \mapsto \mathcal{M}_\nu(\alpha R)$. This directly gives (16) and (17); the same division and inverse expansion preserve the stated $O(h^{2\nu})$, $O\{h^4 \log(1/h)\}$, and $O(h^4)$ remainders in the three smooth subregimes. For $\nu > 1$, the order- h^2 coefficient of normalized correlation is

$$B_{\nu,k}(r) + \mathcal{M}_\nu(z) \frac{\alpha^2 m_2}{4(\nu-1)}. \quad (58)$$

The Matérn correlation satisfies

$$\mathcal{M}_\nu''(z) + \frac{1-2\nu}{z} \mathcal{M}_\nu'(z) - \mathcal{M}_\nu(z) = 0. \quad (59)$$

Using $m_2 = 2T_k$, the coefficient becomes $A_{\nu,k}(r)$. Finally,

$$K_\nu(z) = K_{\nu-2}(z) + \frac{2(\nu-1)}{z} K_{\nu-1}(z) \quad (60)$$

implies (20). Positivity of $K_\eta(z)$ for real η and $z > 0$, together with (9), proves the sign. $\square$

B Proof of the multi-lag and full-likelihood projections

Proof of Theorem 2. For $|\rho|, |q| < 1$, direct Gaussian calculation gives

$$d(\rho, q) = D_{\text{KL}}\{\mathcal{N}(0, \mathbf{R}(\rho)) \parallel \mathcal{N}(0, \mathbf{R}(q))\} = \frac{1}{2} \left[\log \frac{1-q^2}{1-\rho^2} - 2 + \frac{2(1-q\rho)}{1-q^2} \right]. \quad (61)$$

At $q = \rho$, $d = \partial_q d = 0$ and $\partial_q^2 d = (1 + \rho^2)/(1 - \rho^2)^2$. By the pair phase law,

$$\begin{aligned} \rho_{h,j} &= \mathcal{M}_\nu\{(\alpha - \kappa_j s_\nu(h) + o\{s_\nu(h)\})R_j\} \\ &= \rho_j - \kappa_j \dot{\rho}_j s_\nu(h) + o\{s_\nu(h)\}, \quad \dot{\rho}_j = R_j \mathcal{M}'_\nu(\alpha R_j). \end{aligned}$$

For $\eta = \alpha - \gamma s_\nu(h)$, the candidate correlation is $\rho_j - \gamma \dot{\rho}_j s_\nu(h) + o\{s_\nu(h)\}$. Expanding (61) uniformly over the finite lag set yields

$$Q_h^C(\eta) = \frac{s_\nu(h)^2}{2} \sum_j \lambda_j (\kappa_j - \gamma)^2 + o\{s_\nu(h)^2\}. \quad (62)$$

At $h = 0$, the criterion is uniquely minimized by α , and its second derivative is $\sum_j \lambda_j > 0$. Compactness, uniform convergence, and the implicit function theorem localize the minimizer. Completing the square gives $\gamma = \bar{\kappa}$, (27), and (28). $\square$

Proof of Proposition 3. The Gaussian criterion is

$$K_h(\theta) = \frac{1}{2} [\text{tr}\{\Sigma(\theta)^{-1} \Sigma_h\} - p + \log \det \Sigma(\theta) - \log \det \Sigma_h].$$

The identities $\partial \Sigma^{-1} = -\Sigma^{-1}(\partial \Sigma)\Sigma^{-1}$ and (40) give

$$\partial_j K_h(\theta_0) = -s_\nu(h) \langle \dot{\Sigma}_j, \Gamma_\nu \rangle_P + o\{s_\nu(h)\} = -g_{\nu,j} s_\nu(h) + o\{s_\nu(h)\}.$$

At $h = 0$, the Hessian equals $\mathcal{J}$. The implicit function theorem thus gives $\theta_h^\dagger - \theta_0 = \mathcal{J}^{-1} g_\nu s_\nu(h) + o\{s_\nu(h)\}$. For a covariance perturbation H , Gaussian KL has quadratic term $\frac{1}{4} \text{tr}(PHPH)$. Taking $H = s_\nu(h)(\Gamma_\nu - \sum_j \beta_{\nu,j} \dot{\Sigma}_j) + o\{s_\nu(h)\}$ proves (44). Moreover, $\text{tr}(PE_\nu PE_\nu) = \|P^{1/2} E_\nu P^{1/2}\|_F^2$, so the coefficient is zero exactly when the support perturbation belongs to the tangent space. $\square$

C Proof of the transition-aware approximation

Proof. For noninteger $0 < \nu < 2$, retain the quadratic analytic term and the leading fractional term in (55). Compact support permits termwise expectation and gives

$$\frac{C_h(0)}{\sigma^2} = 1 + W_{\nu,k}(h) + \begin{cases} O(h^{2\nu+2}), & 0 < \nu < 1, \\ O(h^4), & 1 < \nu < 2. \end{cases} \quad (63)$$

At $\nu = 1$, taking the expectation of the displayed expansion for $\mathcal{M}_1$ gives $1 + W_{1,k}(h) + O\{h^4 \log(1/h)\}$. Together with the fixed-lag expansion (11), these relations imply that the exact correlation differs from $\tilde{\rho}_h$ by the corresponding order in (32). The inverse of $\alpha \mapsto \mathcal{M}_\nu(\alpha R)$ has a bounded derivative in a neighborhood of the fixed target, so the same orders hold for decay.

To verify the threshold cancellation, write $\nu = 1 + \varepsilon$. As $\varepsilon \rightarrow 0$,

$$b_{1+\varepsilon} = \frac{1}{4\varepsilon} + \frac{2\gamma_E - 1 - 2\log 2}{4} + O(\varepsilon), \quad (64)$$

$$m_{2+2\varepsilon}(\alpha h)^{2+2\varepsilon} = (\alpha h)^2 [m_2 + 2\varepsilon\{\ell_{2,k} + m_2 \log(\alpha h)\} + O(\varepsilon^2)]. \quad (65)$$

The pole cancels the first term in (29); the finite remainder is exactly (30). $\square$

D Proof of the directional support contrast

Proof. For brevity, write $B_e = B_{\nu,k}(Re)$. The fixed-lag expansion gives

$$\frac{C_h(Re)}{\sigma^2} = \mathcal{M}_\nu(z) + h^2 B_e + O(h^4). \quad (66)$$

The zero-lag denominator $d_h = C_h(0)/\sigma^2$ is common to all lag directions and satisfies $d_h \rightarrow 1$. Consequently,

$$\rho_h(Re_1) - \rho_h(Re_2) = h^2\{B_{e_1} - B_{e_2}\} + o(h^2). \quad (67)$$

This cancellation is why the statement remains order h^2 when the common rough-field shift contains lower powers of h .

Let $g(s) = \mathcal{M}_\nu^{-1}(s)/R$ on a neighborhood of $\mathcal{M}_\nu(z)$. Since $\mathcal{M}'_\nu(z) < 0$, the inverse-function theorem gives $g'\{\mathcal{M}_\nu(z)\} = 1/\{R\mathcal{M}'_\nu(z)\}$. Also $\rho_h(Re_j) \rightarrow \mathcal{M}_\nu(z)$, so the mean-value theorem and (67) imply

$$\Delta_{e_1}(h) - \Delta_{e_2}(h) = -\frac{B_{e_1} - B_{e_2}}{R\mathcal{M}'_\nu(z)} h^2 + o(h^2). \quad (68)$$

From (10),

$$B_{e_1} - B_{e_2} = \alpha^2\{a_k(e_1) - a_k(e_2)\} \left\{ \mathcal{M}''_\nu(z) - \frac{\mathcal{M}'_\nu(z)}{z} \right\}. \quad (69)$$

If $c = 2^{1-\nu}/\Gamma(\nu)$, the Bessel derivative identity yields

$$\mathcal{M}'_\nu(z) = -cz^\nu K_{\nu-1}(z), \quad \mathcal{M}''_\nu(z) - \frac{\mathcal{M}'_\nu(z)}{z} = cz^\nu K_{\nu-2}(z). \quad (70)$$

Substitution into (68) proves (33). The identities and positivity of $K_\eta(z)$ hold for every real order η and $z > 0$, so no condition $\nu > 1$ is needed for the directional contrast. $\square$

E Proof of the finite-library certificate

Proof. Write $X_i = \Sigma_0^{1/2} Z_i$, where the Z_i are independent $\mathcal{N}_p(0, I_p)$ vectors, and stack them into $Z = (Z_1^\top, \dots, Z_N^\top)^\top$. For a fixed candidate θ ,

$$\sum_{i=1}^N X_i^\top \Sigma_\theta^{-1} X_i = Z^\top (I_N \otimes A_\theta) Z, \quad \mathbb{E} Z^\top (I_N \otimes A_\theta) Z = N \operatorname{tr}(A_\theta). \quad (71)$$

The two-sided Gaussian quadratic-form inequality gives, for every $t > 0$,

$$\Pr \left[\left| Z^\top (I_N \otimes A_\theta) Z - N \operatorname{tr}(A_\theta) \right| > 2\sqrt{N} \|A_\theta\|_F \sqrt{t} + 2\|A_\theta\|_{\text{op}} t \right] \leq 2e^{-t}. \quad (72)$$

Here $\|I_N \otimes A_\theta\|_F = \sqrt{N} \|A_\theta\|_F$ and $\|I_N \otimes A_\theta\|_{\text{op}} = \|A_\theta\|_{\text{op}}$. The log-determinant term in $\widehat{L}_N(\theta) - L(\theta)$ is deterministic. Dividing (72) by $2Np$, setting $t = \log(2M/\delta)$, and applying a union bound over the deterministic library proves (47) [13, 15].

On the simultaneous event,

$$\begin{aligned} L(\widehat{\theta}) - L(\theta^*) &= [L - \widehat{L}_N](\widehat{\theta}) + [\widehat{L}_N(\widehat{\theta}) - \widehat{L}_N(\theta^*)] + [\widehat{L}_N - L](\theta^*) \\ &\leq u_{\widehat{\theta}} + u_{\theta^*} \leq 2 \max_{\theta \in \mathcal{G}} u_\theta, \end{aligned} \quad (73)$$

which proves (48). Finally,

$$D_{\text{KL}}\{\mathcal{N}_p(0, \Sigma_0) \parallel \mathcal{N}_p(0, \Sigma_\theta)\} = \frac{1}{2} \left[\operatorname{tr}(\Sigma_\theta^{-1} \Sigma_0) - p + \log \frac{\det \Sigma_\theta}{\det \Sigma_0} \right], \quad (74)$$

which proves (49). Under (50), inversion and congruence give $0 \prec A_\theta \preceq c^{-1} I_p$; hence the stated operator- and Frobenius-norm bounds and (51) follow. $\square$

F Exact exponential benchmark

For $d = 1$, $\nu = 1/2$, and the Epanechnikov density $k(u) = \frac{3}{4}(1 - u^2)\mathbf{1}\{|u| \leq 1\}$, the density of $D = U - V$ is

$$f_D(w) = \frac{3}{5} - \frac{3}{4}|w|^2 + \frac{3}{8}|w|^3 - \frac{3}{160}|w|^5, \quad |w| \leq 2. \quad (75)$$

With $x = \alpha h$, define

$$s(x) = \mathbb{E}e^{-x|D|}, \quad M_k(x) = \mathbb{E}e^{xU} = \frac{3(x \cosh x - \sinh x)}{x^3}, \quad q(x) = M_k(x)^2. \quad (76)$$

Then $C_h(0) = \sigma^2 s(x)$, and for $R \geq 2h$,

$$C_h(R) = \sigma^2 q(x) e^{-\alpha R}. \quad (77)$$

The exact naive pair target is

$$(\sigma_h^2)^\dagger = \sigma^2 s(x), \quad \alpha_h^\dagger = \alpha - \frac{1}{R} \log \frac{q(x)}{s(x)}. \quad (78)$$

For $x > 0$, compact support and nondegeneracy give

$$1 < q(x)/s(x) < e^{2x}.$$

Indeed, condition on $A = |U|$ and $B = |V|$, and put $p = A+B$ and $m = |A-B|$. Symmetry of the signs gives conditional numerator and denominator proportional to $\cosh(xp) + \cosh(xm)$ and $e^{-xp} + e^{-xm}$, respectively. The lower inequality follows from $\cosh(t) \geq e^{-t}$, with strictness off $A = B = 0$. Since $p + m = 2 \max(A, B) \leq 2$, $\cosh(xp) < e^{xp} \leq e^{2x-xm}$ and $\cosh(xm) \leq e^{xm} \leq e^{2x-xp}$, with strictness on a set of positive probability. Averaging proves the upper inequality. Since $R \geq 2h$, this implies

$0 < \alpha - R^{-1} \log\{q(x)/s(x)\} < \alpha$. Variance and decay are therefore both strictly underestimated. The small-bandwidth expansions are

$$s(x) = 1 - \frac{18}{35}x + \frac{1}{5}x^2 - \frac{4}{63}x^3 + \frac{3}{175}x^4 + O(x^5), \quad (79)$$

$$q(x) = 1 + \frac{1}{5}x^2 + \frac{3}{175}x^4 + O(x^6), \quad (80)$$

$$\log\{q(x)/s(x)\} = \frac{18}{35}x + \frac{162}{1225}x^2 + O(x^3). \quad (81)$$

This benchmark is used as an exact software regression test, not as the paper’s novelty claim.

G Reproducibility contract

The public code at <https://github.com/Sasha-Cui/HighDimSpatialStatistics> records the exact configuration and commit identifier in each promoted run’s metadata; the frozen paper artifact is tagged `supportshift-geosim-v1.3.2`. The release includes a synthetic-data card, scoped reuse terms, a citation file, and a one-command verifier for the 23 promoted data/provenance inputs, 33 generated paper artifacts, 8,400 finite-grid fits, 12,800 replicated fits, and 168 reported numerical claims. The finite-grid full-likelihood track uses an immutable JSON manifest, SHA-256 configuration hashes, deterministic seed derivation, one task-local atomic shard, and an **afterany** reducer that refuses to mark a run complete when a shard is missing or invalid. The replicated-field track records its complete candidate grids, matrix-norm radii, derived seeds, dependency versions, result hashes, and validation gates. Those gates check the exact finite-grid ERM inequality, agreement with a public likelihood implementation, population-target interiority, finite-to-continuous objective resolution, the candidatewise simultaneous event, and exact reapplication of the saved observation operator. The promoted batch aborts unless the Git worktree is clean. The zero-bandwidth selector, row sums, row rank, corrected population target, grid interpolation, Gaussian certificate, driver shakedown, artifact hash manifest, and closed exponential formulas are tested. The reported implementation adds no adaptive diagonal jitter. All paper figures and tables are regenerated from audited aggregate CSV files.

References

- [1] Takeshi Amemiya and Roland Y. Wu. The effect of aggregation on prediction in the autoregressive model. *Journal of the American Statistical Association*, 67(339):628–632, 1972. doi: 10.1080/01621459.1972.10481264.
- [2] François Bachoc. Asymptotic analysis of covariance parameter estimation for Gaussian processes in the misspecified case. *Bernoulli*, 24(2):1531–1575, 2018. doi: 10.3150/16-BEJ906.
- [3] Claude Bellehumeur and Pierre Legendre. Aggregation of sampling units: An analytical solution to predict variance. *Geographical Analysis*, 29(3):258–266, 1997. doi: 10.1111/j.1538-4632.1997.tb00961.x.
- [4] Moreno Bevilacqua, Carlo Gaetan, Jorge Mateu, and Emilio Porcu. Estimating space and space-time covariance functions for large data sets: A weighted composite likelihood approach. *Journal of the American Statistical Association*, 107(497):268–280, 2012. doi: 10.1080/01621459.2011.646928.
- [5] Erick A. Chacón-Montalván, Peter M. Atkinson, Christopher Nemeth, Benjamin M. Taylor, and Paula Moraga. Spatial latent Gaussian modelling with change of support, 2024.
- [6] Isobel Clark. Regularization of a semivariogram. *Computers & Geosciences*, 3(2):341–346, 1977. doi: 10.1016/0098-3004(77)90010-3.

- [7] David Clifford, Alex B. McBratney, James Taylor, and Brett M. Whelan. Generalized analysis of spatial variation in yield monitor data. *The Journal of Agricultural Science*, 144(1):45–51, 2006. doi: 10.1017/S0021859606005892.
- [8] Victor De Oliveira and Zifei Han. On information about covariance parameters in Gaussian Matérn random fields. *Journal of Agricultural, Biological and Environmental Statistics*, 27(4):690–712, 2022. doi: 10.1007/s13253-022-00510-5.
- [9] Maria Myrto Folia and Magnus Rattray. Trajectory inference and parameter estimation in stochastic models with temporally aggregated data. *Statistics and Computing*, 28:1053–1072, 2018. doi: 10.1007/s11222-017-9779-x.
- [10] Montserrat Fuentes. Approximate likelihood for large irregularly spaced spatial data. *Journal of the American Statistical Association*, 102(477):321–331, 2007. doi: 10.1198/016214506000000852.
- [11] Alan E. Gelfand, Li Zhu, and Bradley P. Carlin. On the change of support problem for spatio-temporal data. *Biostatistics*, 2(1):31–45, 2001. doi: 10.1093/biostatistics/2.1.31.
- [12] Carol A. Gotway and Linda J. Young. Combining incompatible spatial data. *Journal of the American Statistical Association*, 97(458):632–648, 2002. doi: 10.1198/016214502760047140.
- [13] Daniel Hsu, Sham M. Kakade, and Tong Zhang. A tail inequality for quadratic forms of subgaussian random vectors. *Electronic Communications in Probability*, 17:1–6, 2012. doi: 10.1214/ECP.v17-2079.
- [14] Phaedon C. Kyriakidis and Eun-Hye Yoo. Geostatistical prediction and simulation of point values from areal data. *Geographical Analysis*, 37(2):124–151, 2005. doi: 10.1111/j.1538-4632.2005.00633.x.
- [15] Béatrice Laurent and Pascal Massart. Adaptive estimation of a quadratic functional by model selection. *The Annals of Statistics*, 28(5):1302–1338, 2000. doi: 10.1214/aos/1015957395.
- [16] Christopher J. Paciorek and Mark J. Schervish. Spatial modelling using a new class of nonstationary covariance functions. *Environmetrics*, 17(5):483–506, 2006. doi: 10.1002/env.785.
- [17] Frederik J. Simons, Olivia L. Walbert, Arthur P. Guillaumin, Gabriel L. Eggers, Kevin W. Lewis, and Sofia C. Olhede. Maximum-likelihood estimation of the Matérn covariance structure of isotropic spatial random fields on finite, sampled grids. *Geophysical Journal International*, 245(2):ggag044, 2026. doi: 10.1093/gji/ggag044.
- [18] Michael L. Stein. *Interpolation of Spatial Data: Some Theory for Kriging*. Springer, New York, 1999. doi: 10.1007/978-1-4612-1494-6.
- [19] Daniel O. Stram and William W. S. Wei. Temporal aggregation in the ARIMA process. *Journal of Time Series Analysis*, 7(4):279–292, 1986. doi: 10.1111/j.1467-9892.1986.tb00495.x.
- [20] Cristiano Varin, Nancy Reid, and David Firth. An overview of composite likelihood methods. *Statistica Sinica*, 21(1):5–42, 2011.
- [21] Hao Zhang. Inconsistent estimation and asymptotically equal interpolations in model-based geostatistics. *Journal of the American Statistical Association*, 99(465):250–261, 2004. doi: 10.1198/016214504000000241.