

LP_Practice_digitSumOdd

Sasi | 09 Feb 2023



Finish State: Normal

Test Taken on: February 09, 2023 09:40:21 AM IST



Sasi

sasidevi.s.2020.cse@ritchennai.edu.in


Overall Summary

40 Marks Scored
out of 40

100 % 100 percentile
out of 27333 Test Takers

4m 58s Time taken
of 1hr 5mins

Marks Scored



Attempt Summary

Distribution of questions attempted in a total of 1 question(s).



This shows the correctness of questions attempted by the test taker

Correct	1 Ques	40/40 Marks
Incorrect	0 Ques	0/0 Marks
Partially Correct	0 Ques	0/0 Marks
Not Attempted	0 Ques	0/0 Marks

Section-Wise Details

▼ Section 1 Program	question(s) 1 Q.	Time taken 4m 58s (Untimed)	Marks Scored 40 / 40
---------------------------	---------------------	--------------------------------	-------------------------

Marks Scored



Attempt Summary

Distribution of questions attempted in a total of 1 question(s).



■ Correct	1 Ques	40/40 Marks
-----------	--------	-------------

This shows the correctness of questions attempted by the test taker

About the Report

This Report is generated electronically on the basis of the inputs received from the assessment takers. This Report including the AI flags that are generated in case of availing of proctoring services, should not be solely used/relied on for making any business, selection, entrance, or employment-related decisions. Mettl accepts no liability from the use of or any action taken or refrained from or for any and all business decisions taken as a result of or reliance upon anything, including, without limitation, information, advice, or AI flags contained in this Report or sources of information used or referred to in this Report.



1. Program

Question 1

Revisit Later

How to Attempt?

Odd Digits' Sum:

In mathematics, the "digit sum" of a given integer is the sum of all its digits. e.g.
the digit sum of 84001 is calculated as $8+4+0+0+1 = 13$,
the digit sum of 158 is $1+5+8 = 14$.

Rohan's teacher has asked him to write a function (method) that takes as input a positive number and performs digitSum of only the odd digits in the given number.

Example 1: If the given number is 9625, we must add only the odd digits. i.e. $9+5 = 14$. Thus, the OddDigitsSum for the number 9625 is 14.

Example 2: If the given number is 2134, the OddDigitsSum will be $1+3 = 4$.

Assumption: The input number will be a positive integer number ≥ 1 and ≤ 25000 .

JAVA8

Compiler: Java - 1.8

```
1 import java.io.*;
2 import java.util.*;
3
4 // Read only region start
5 class UserMainCode
6 {
7
8     public int OddDigitsSum(int input1){
9         // Read only region end
10         int sum=0;
11         while(input1!=0)
12         {
13             int n=input1%10;
14             if(n%2!=0)
15                 sum+=n;
16             input1/=10;
17         }
18         return sum;
19     }
20 }
```

☐ Use Custom Input

?

Compile and Test

Submit Code