

* Total prime numbers between 1-50 $\rightarrow$ 15

50-100 $\rightarrow$ 10

100-150 $\rightarrow$ 10

150-200 $\rightarrow$ 11

200-250 $\rightarrow$ 7

250-300 $\rightarrow$ 10

Co - prime numbers :

- * Two numbers are co-prime if their HCF is 1.
- * Two prime numbers are always co-prime to each other.
- * Any two consecutive integers are always co-prime.

* Face value of 3 in 56738 is 3.

* Place value of 3 in 56738 is 30.

Operation	Rational, Rational	Rational, Irrational	Irrational, Irrational
Addition	Always rational $2 + 3 = 5$	Always irrational $2 + \sqrt{3}$	Can be rational or irrational $\sqrt{2} + (-\sqrt{2}) = 0$ $\sqrt{2} + \sqrt{2} = 2\sqrt{2}$
Subtraction	Always rational $3 - 2 = 1$	Always irrational $\sqrt{3} - \sqrt{2}$	Can be rational or irrational $\sqrt{3} - \sqrt{3} = 0$ $\sqrt{3} - \sqrt{2}$
Multiplication	Always rational $2 \times 3 = 6$	Always irrational Except if rational number is 0. $2 \times \sqrt{3} = 2\sqrt{3}$ $0 \times \sqrt{3} = 0$	Can be rational or irrational $2\sqrt{2} \times \sqrt{2} = 4$ $\sqrt{2} \times \sqrt{3} = \sqrt{6}$
Division	Always rational (if divisor $\neq 0$) $\frac{3}{4} = 0.75$	Always irrational Except if dividend is 0. $\frac{2}{\sqrt{3}}$ $\frac{0}{\sqrt{3}} = 0$	Can be rational or irrational $\frac{2\sqrt{2}}{\sqrt{2}} = 2$ $\frac{\sqrt{2}}{\sqrt{3}} = \sqrt{\frac{2}{3}}$

Successive Division

* A number 219 is successively divided by 6, 2, 4, 3

$$\begin{array}{r} 6 \overline{) 219} (36 \\ \underline{18} \\ 39 \\ \underline{36} \\ 3 \end{array}$$

$$\begin{array}{r} 2 \overline{) 36} (18 \\ \underline{2} \\ 16 \\ \underline{16} \\ 0 \end{array}$$

$$\begin{array}{r} 4 \overline{) 18} (4 \\ \underline{16} \\ 2 \end{array}$$

$$\begin{array}{r} 3 \overline{) 4} (1 \\ \underline{3} \\ 1 \end{array}$$

$$\begin{array}{r} 6 \swarrow + 3 \\ 2 \swarrow + 0 \\ 4 \swarrow + 2 \\ 3 \swarrow + 1 \\ \times 1 \end{array}$$

↖ last quotient

$$3 \times 1 + 1 = 4$$

$$4 \times 4 + 2 = 18$$

$$18 \times 2 + 0 = 36$$

$$36 \times 6 + 3 = 219$$

* If they don't give the last quotient, we can assume it as 0

Divisibility Rule of 2, 4, 8, 16

* $2 (2^1)$: Last digit is divisible by 2 or is 0.

* $4 (2^2)$: Last 2 digits are divisible by 4 or are 00.

* $8 (2^3)$: Last 3 digits are divisible by 8 or are 000.

* $16 (2^4)$: Last 4 digits are divisible by 16 or are 0000.

Divisibility Rule of 3, 9

* 3 : Sum of digits is divisible by 3.

* 9 : Sum of digits is divisible by 9.

Divisibility Rule of 5, 25, 125

* $5 (5^1)$: Last digit is 0 or 5.

* $25 (5^2)$: Last 2 digits are 00 or divisible by 25.

* $125 (5^3)$: Last 3 digits are 000 or divisible by 125.

Divisibility Rule of Numbers having known

- * 6 : Number is divisible by both 2 and 3.
- * 24 : Number is divisible by both 8 and 3.
- * 72 : Number is divisible by both 8 and 9.
- * 88 : Number is divisible by both 8 and 11.

Note - We have to take only co-primes.

Divisibility Rule of 7, 11, 13

Check 88440077061

$$88 \overline{) 440} \quad 077 \overline{) 061}$$

$$\begin{array}{r} 061 \\ 440 \\ \hline 171 \\ 154 \\ \hline 17 \end{array}$$

$$\begin{array}{r} 077 \\ 88 \\ \hline 11 \\ 11 \\ \hline 0 \end{array}$$

501

165

(-)

336

Now we need to check whether 336 is divisible by 7, 11 or 13.

Divisibility Rule of 11

Check 5453987

5 | 4 | 5 | 3 | 9 | 8 | 7

$$7 + 9 + 5 + 5 = 26$$

$$8 + 3 + 4 = 15$$

(-)

11

The result is divisible 11 or not.

Note :- Number in the form of $abcabc$ are always divisible by 7, 11, 13 and 1001.

Remainder Theorem

$$\frac{17}{3} \rightarrow 2$$

$$\frac{17^2}{3} = \frac{17 \times 17}{3} \rightarrow \frac{2^2}{3} \rightarrow \frac{4}{3} \rightarrow 1$$

$$\frac{35}{37} \rightarrow -2 \rightarrow 35$$

$$\frac{35^2}{37} = \frac{35 \times 35}{37} \rightarrow \frac{(-2)^2}{37} \rightarrow \frac{4}{37} \rightarrow 4$$

$$\frac{35^3}{37} = \frac{35 \times 35 \times 35}{37} \rightarrow \frac{(-2)^3}{37} \rightarrow \frac{-8}{37} \rightarrow 29$$

$$\frac{35 \times 38 \times 39}{37} \rightarrow \frac{-4}{37} \rightarrow 33$$

$$\frac{35 + 38 + 39}{37} \rightarrow \frac{1}{37} \rightarrow 1$$

$$* \frac{a^{p-1}}{p} \rightarrow \text{Remainder} = 1 \text{ -- where } p \text{ is prime number,}$$

a and p are co-primes.

$$\text{Ex: } \frac{8^{36}}{37} \rightarrow 1$$

$$\frac{8^{73}}{37} = \frac{8^{72} \cdot 8}{37} = \frac{8^{36} \cdot 8^{36} \cdot 8}{37} \rightarrow \frac{8}{37} \rightarrow 8$$

$$\frac{5^{50}}{17} = \frac{5^{48} \cdot 5^2}{17} = \frac{5^{16} \cdot 5^{16} \cdot 5^{16} \cdot 5^2}{17} \Rightarrow \frac{5^2}{17} \Rightarrow \frac{25}{17} \rightarrow 8$$

Number of zeroes

* Let $N = 2^p \times x^a \times y^b \times z^c \times 5^q$

Number of zeroes in $N = \min(p, q)$

* To find the number of zeroes in $n!$, we divide n by 5 repeatedly until the result is less than 5 and then add all the quotients. $(\lfloor \frac{n}{5} \rfloor + \lfloor \frac{n}{25} \rfloor + \lfloor \frac{n}{125} \rfloor + \dots)$

Ex: $\frac{50!}{5} \rightarrow 10 + 2 \rightarrow 12 \text{ zeroes}$

Factors

* $a^m - b^m$ (m is even) $\rightarrow (a+b), (a-b)$

* $a^m + b^m$ (m is odd) $\rightarrow a+b$

* $a^m - b^m$ (m is odd) $\rightarrow a-b$

* $a^m + b^m$ (m is even)

$$a^{12} + b^{12} \rightarrow (a^4)^3 + (b^4)^3 \rightarrow a^4 + b^4$$

$$(a^m + b^m \text{ (} m \text{ is odd)})$$

Number of factors

$$120 = 2^3 \times 3^1 \times 5^1$$

* Total number of factors (including 1 and itself)

$$= (3+1)(1+1)(1+1) = 4 \times 2 \times 2 = 16$$

* Total number of even factors = $(3)(1+1)(1+1)$

$$= 3 \times 2 \times 2 = 12$$

* Total number of odd factors = $(1)(1+1)(1+1) = 1(2)(2) = 4$

* Total number of prime factors = $3 + 1 + 1 = 5$

* Sum of all factors = $(2^0 + 2^1 + 2^2 + 2^3) \cdot (3^0 + 3^1) \cdot (5^0 + 5^1)$

* Sum of even factors (ignore 2^0)

$$= (2^1 + 2^2 + 2^3) \cdot (3^0 + 3^1) \cdot (5^0 + 5^1)$$

* Sum of odd factors = $(3^0 + 3^1) \cdot (5^0 + 5^1)$

Units Digit of powers

0 → Always 0

1 → Always 1

2 → 2, 4, 8, 6, ...

3 → 3, 9, 7, 1, ...

4 → 4, 6, 4, 6, ...

5 → Always 5


6 → Always 6

7 → 7, 9, 3, 1, ...

8 → 8, 4, 2, 6, ...

9 → 9, 1, 9, 1, ...

Divisibility by a or b, a and b



$x+y \rightarrow$ divisible by a

$y+z \rightarrow$ divisible by b

$x+y+z \rightarrow$ divisible by a or b

$(x+y) + (y+z) - (2y) = x+z \rightarrow$ divisible by a or b but not by a & b.

Recurring Decimal

$$0.\overline{23} = \frac{23}{99}$$

$$0.5\overline{29} = \frac{529 - 5}{990} = \frac{524}{990}$$

$$0.8\overline{351} = \frac{8351 - 8}{9990} = \frac{8343}{9990}$$

$$2.5\overline{32} = 2 + 0.5\overline{32} = 2 + \frac{532 - 53}{990} = 2 + \frac{479}{990}$$

Identification of Perfect Square

- * Last digit is 1, 4, 5, 6, 9 or 0, then the number may or may not be a perfect square.
- * Last digit is 2, 3, 7, 8, then the number is not a perfect square.
- * Last two digits of the number should match with the last two digits of the numbers 1^2 to 24^2 .
- * Digital sum should be 1, 4, 9 or 7 ($1+9+4=7$).