N Niet

Gerd Potent. It originated as part of a consignment of *P. callosum* imported from Thailand by the Karge Nursery in Dahlenburg in Germany. It has flowers that are similar to those of typical *P. callosum* but with a somewhat smaller dorsal sepal and no marginal warts on the petals. Gruss & Roth suggested that it might be a natural hybrid of *P. callosum* and another unnamed species. I have seen plants from the original introduction and consider that the minor differences from *P. callosum* do not warrant its recognition at specific rank.

DESCRIPTION. A terrestrial herb. Leaves 3–5, narrowly elliptic, oblong-elliptic or obovate, tridenticulate at acute apex, 10–20 cm long, 3.2–4.8 cm wide, ciliate at base, tessellated pale and dark above, sometimes purple at base on lower surface. Inflorescence 1- (or rarely 2-) flowered, 20–40 cm tall; peduncle purple, purple-pubescent, 12–25 cm long; bract ovate to elliptic, acute to subacute, 1.5–2.8 cm long, 1.5–2 cm wide, green, sometimes marked with purple, ciliate. Flower large, 8–11 cm across; sepals white flushed with purple in lower half, veined with purple and green; petals white to yellow-green with a purple apical third, spotted with maroon on upper margin and sometimes in basal half; lip green, heavily flushed deep maroon; staminode pale green with darker veins, flushed purple; pedicel and ovary 3–6.5 cm long, green, flushed with purple, purple-pubescent. Dorsal sepal broadly ovate to subcircular, apiculate, 4–5.5 cm long, 4.2–6 cm wide, with recurved ciliate margins. Synsepal concave, elliptic or lanceolate, acute, 2.7–3.2 cm long, 1.6–2.5 cm wide. Petals sometimes reflexed, subsigmoid, ligulate,



Fig. 104. Paphiopedilum callosum var. sublaeve, growing in forest, southern Thailand. (Photo.: H. Bänziger)