

# Assignment - 1

Gautam Singh

**Abstract**—This document contains the solution to Exercise 3.18 (a) of Oppenheim.

**Problem 1.** A causal LTI system has the system function

$$H(z) = \frac{1 + 2z^{-1} + z^{-2}}{\left(1 + \frac{1}{2}z^{-1}\right)(1 - z^{-1})} \quad (1)$$

Find the impulse response  $h(n)$  of the system.

**Solution:** We substitute  $x := z^{-1}$  and carry out long division

$$\begin{array}{r} -2 \\ -\frac{1}{2}x^2 - \frac{1}{2}x + 1 \overline{) x^2 + 2x + 1} \\ \underline{-x^2 \quad -x + 2} \phantom{1} \\ x + 3 \end{array}$$

Thus, on expanding using partial fractions,

$$H(z) = -2 + \frac{z^{-1} + 3}{\left(1 + \frac{1}{2}z^{-1}\right)(1 - z^{-1})} \quad (2)$$

$$= -2 + \frac{1}{3} \left( \frac{1}{1 + \frac{1}{2}z^{-1}} \right) + \frac{8}{3} \left( \frac{1}{1 - z^{-1}} \right) \quad (3)$$

We know that

$$a^n u(n) \stackrel{Z}{\rightleftharpoons} \frac{1}{1 - az^{-1}} \quad (4)$$

and

$$\delta(n) \stackrel{Z}{\rightleftharpoons} 1 \quad (5)$$

Thus, using (4) and (5) in (3),

$$h(n) = -2\delta(n) + \frac{1}{3} \left( 8 + \left( -\frac{1}{2} \right)^n \right) u(n) \quad (6)$$

The plot of  $h(n)$  has been generated using the Python code `codes/1_1.py`.




Fig. 1: Plot of  $h(n)$  against  $n$