

CSCI-GA.1170-001 Homework 2

Ankit Sati

TOTAL POINTS

27 / 36

QUESTION 1

1 Different Methods for Recurrences 8 / 14

✓ + 1 pts **Part a)** Correct values for c_1 and n (for Big-O)

✓ + 1 pts **Part a)** Correct values for c_2 and n (for Ω)

✓ + 1 pts **Part c)** Correct Level 0

✓ + 1 pts **Part c)** Correct Level 1

✓ + 0.5 pts **Part c)** Correct Level $d-1$

✓ + 1 pts **Part c)** Correct Level d

✓ + 0.5 pts **Part c)** Correct depth

+ 2 Point adjustment

Part a is wrong.

1 On what assumption n is being replaced with $n \log n$?

2 Incorrect $T(n)$ and asymptotic bound

QUESTION 2

2 Fun with Recurrences 14 / 15

✓ + 1 pts (c) correct substitution

✓ + 1 pts (c) correct relation for step 2

✓ + 1 pts (c) correct relation for step 3

✓ + 1 pts (a) Extra credit for correct constant

✓ + 5 pts (B) All Correct

✓ + 5 pts (a) all correct



Incorrect/missing steps for part c. Final answer missing/incorrect

QUESTION 3

3 Expected Running Time 5 / 7

✓ - 0 pts Correct

✓ - 1 pts Incorrect/Missing recurrence relation for $n > 5$ in part (a)

✓ - 1 pts Incorrect/Missing base case for $n \leq 5$ in part (a)

✓ - 0.5 pts Incorrect/Missing $T(n)$ in part (a)

✓ + 0.5 pts Correct final answer in part (c)

3 Missing justification for why $\text{Bla}(\text{Foo})$ is constant time

4 Missing base case ($n \leq 1$)

Problem 2-1.

$$a) T(n) = 2T\left(\frac{n}{2} + 1\right) + n - 2$$

$$\text{given } T(3) = 0$$

To prove. $T(n) = \Theta(n-2) \log(n-2) \therefore$ Tight bound.

\rightarrow Prove by master theorem for Cheat value.

$$T(n) = aT\left(\frac{n}{b}\right) + 1.$$

$$f(n) = \Theta(n^k \cdot \log^p n)$$

$$\log_b a = 1; k = 1; p = 0.$$

$$\text{Case 2 } f(n) = \Theta(n^k \cdot \log^{p+1} n)$$

$$= \Theta(n \cdot \log n) \rightarrow \text{best guess.}$$

Cheat Value by Master Theorem.

Guess then Verify

$$T(n) = 2T\left(\frac{n+2}{2}\right) + n - 2.$$

Substituting the values

$$= 2c\left(\frac{n \log n + 2}{2}\right) + n \log n - 2.$$

$$= c(n \log n + 2) + n \log n - 2.$$

$$\approx \Theta(n \log n).$$

Since $T(3) = 0$.

$$\therefore (n-2) \log(n-2) \quad \forall \quad n \geq 3.$$

which means for every number $(n \geq 3)$ the below expression holds true

$$c_1 n \log n \leq \Theta(n-2) \log(n-2) \leq c_2 n \log n$$

where $c_1, c_2 > 0$

①

$$\therefore C_1 \underbrace{n \log n} \leq \underbrace{(n-2) \log(n-2)} \leq C_2 n \log n$$

which means that our function is a tight bound.

$$\forall C_1, C_2 > 0 \Rightarrow f(n) = \Theta g(n)$$

$$\boxed{= \Theta \cdot (n-2) \log(n-2)}$$

← Proven by guess then verify.

b) Domain Range substitution for a Reducing Recurrence.

$$T(n) = 2T\left(\frac{n}{2} + 1\right) + (n-2)$$

Assuming that the function is reducing, we will solve towards $T(n) = 1$.

$$T_n = \begin{cases} 0 & ; n=3 \\ 2T\left(\frac{n}{2} + 1\right) + n-2 & ; n > 1 \end{cases}$$

$$T(n) = 2T\left(\frac{n}{2} + 1\right) + n-2 \quad \text{--- (1)}$$

Solving for $\left(\frac{n}{2}\right)$

~~$$T\left(\frac{n}{2}\right) = 2 \left[2T\left(\frac{n+1}{2^2}\right) + \frac{n}{2} - 2 \right] + n-2$$~~

$$T\left(\frac{n}{2}\right) = 2 \left[2T\left(\frac{n+1}{2^2}\right) + \frac{n}{2} - 2 \right] + n-2$$

$$2^2 T \frac{n+1}{2^2} + n-2 + n-2 \quad \text{--- (2)}$$

Relation to function in reduced form. Total work for full credit (Reduced form)

$$\textcircled{1} f(n) = \boxed{T_n = 2^k \cdot T\left(\frac{n+1}{2^k}\right) + k(n-1)} \rightarrow \text{Main equation}$$

$$T\left(\frac{n+1}{2^k}\right) = T(1)$$

$$\frac{n+1}{2^k} = 1$$

$$k = \log(n-2)$$

$$\therefore f(3) = 0$$

min value $(n-2)$

Cont. ②

Since

$$T(n) = 2^k \frac{T(n-2)}{2^k} + K(n-2)$$

$$\& K = \log(n-2) \text{ --- (4)}$$

Substituting the values

$$\begin{aligned} T(n) &= 2^k T(3) + K(n-2) \\ T(n) &= K(n-2) \\ T(n) &= (n-2) \cdot \log(n-2) \end{aligned}$$

← Answer, * Approach 1.

Alternative Approach:

$$T(n) = \Omega((n-2) \log(n-2))$$

$$\Rightarrow T(n) \geq C(n-2) \log(n-2)$$

Solution for sub problems with $\log(\frac{n}{2} + 1)$

$$\begin{aligned} \text{here, } T(\frac{n}{2} + 1) &\geq C\left(\left(\frac{n}{2} + 1\right) - 2\right) \log\left(\frac{n}{2} + 1 - 2\right) \\ &\geq C\left(\frac{n}{2} - 1\right) \log\left(\frac{n}{2} - 1\right) \end{aligned}$$

$$T(n) = 2T\left(\frac{n}{2} + 1\right) + (n-2)$$

$$T(n) \geq 2 \cdot \left[C\left(\frac{n}{2} - 1\right) \log\left(\frac{n}{2} - 1\right) \right] + (n-2)$$

$$T(n) \geq C(n-2) \log\left(\frac{n-2}{2}\right) + n-2$$

$$T(n) \geq C(n-2) [\log(n-2) - \log 2] + (n-2)$$

$$T(n) \geq C(n-2) \log(n-2) + (n-2) [1 - C \log 2]$$

$$T(n) \geq C(n-2) \log(n-2) \quad \& \quad \boxed{C \geq \frac{1}{\log 2}}$$

$$\therefore T(n) = \Omega((n-2) \log(n-2))$$

$$\text{Since } T(n) = O((n-2) \log(n-2))$$

→ Approach 2.

$$\therefore T(n) = \Theta((n-2) \log(n-2)) \quad \& \quad C_1, C_2 > 0 \quad \textcircled{3}$$

2-(c) Recursion Tree Method


$$T(n) = 2T(n-1) + n, \quad T(0) = 1.$$

$$T(n) = aT\left(\frac{n}{b}\right) + f(n)$$

$$f(n) = \Theta(n^k)$$

$$f(n) = \Theta(n^k \cdot \log n)$$

Recursion Tree



$$T(0) = 1.$$

Total Cost.

$$n$$

$$2(n-1)$$

$$2^2(n-2)$$

$$2^3(n-3)$$

$$2^k(n-k)$$

For the exact value of $T(n)$

$$T(n) = n + 2(n-1) + 2^2(n-2) + \dots + 2^{n-1}(n-(n-1)) + 2^n \cdot T(0)$$

$$T(n) = n + 2n + 2^2n + \dots + 2^{n-1}n - (1 \cdot 2 + 2 \cdot 2^2 + 3 \cdot 2^3 + \dots + (n-1)2^{n-1}) + 2^n$$

$$T(n) = n[1 + 2 + 2^2 + \dots + 2^{n-1}] - \sum_{i=1}^{n-1} i 2^i + 2^n$$

Per hint

$$= \frac{n \cdot (2^n - 1)}{2 - 1} + \frac{(n-1)2^{n+1} - 2^n \cdot n + 2 + 2^n}{(2-1)^2}$$

$$= n \cdot 2^n - n + n \cdot 2^{n+1} - 2^{n+1} - n \cdot 2^n + 2 + 2^n$$

(4)
Cost.

$$= 2 \cdot m 2^m - 2^m - m + 2$$

$$T(m) = 2 \cdot m 2^m - 2^m - m + 2$$

Correct answer as per the question

$$T(m) = \Theta(m \cdot 2^m)$$

Asymptotic Notation.

Table.

Level	Size of pr.	No of pr.	Non recursive	Total cost
0	m	1	m	m
1	(m-1)	2	2(m-1)	2m-2
d-1	(m-(d-1))	2^{d-1}	$2^{d-1}(m-(d-1))$	$2^{d-1} \cdot (m-(d-1))$
d	(m-d)	2^d	1	$2^d \cdot 1$

Let level d

$m-d=0$
 $\therefore m=d$ (as mentioned above)

Cost Asymptotically equal.

$m \rightarrow C.E.O$

$2 \rightarrow V.P.$

Cost at level d = 2^m
 cost at level d+1 = $2^{m+1} (m - (m-1))$
 $= 2^{m+1} \cdot 1$

~~Master Theorem~~

5

1 Different Methods for Recurrences 8 / 14

- ✓ + 1 pts **Part a)** Correct values for c_1 and n (for Big-O)
- ✓ + 1 pts **Part a)** Correct values for c_2 and n (for Ω)
- ✓ + 1 pts **Part c)** Correct Level 0
- ✓ + 1 pts **Part c)** Correct Level 1
- ✓ + 0.5 pts **Part c)** Correct Level $d-1$
- ✓ + 1 pts **Part c)** Correct Level d
- ✓ + 0.5 pts **Part c)** Correct depth
- + 2 Point adjustment

Part a is wrong.

- 1 On what assumption n is being replaced with $n \log n$?
- 2 Incorrect $T(n)$ and asymptotic bound

Problem 2-2. (Fun with Recurrences)

(a)

$$T(n, 1) = 3n$$

$$T(1, m) = 3m$$

$$T(n, m) = 3n + T\left(\frac{n}{3}, \frac{m}{3}\right)$$

$$T(n^2, m^2) = T\left(\frac{n}{3}, \frac{m^2}{3^2}\right) + 3n$$

$$= T\left(\frac{3n}{3^2}, \frac{m^2}{9}\right) + 3n + 3 \cdot \frac{m}{3}$$

$$= T\left(\frac{n}{3^3}, \frac{m^2}{3^5}\right) + 3n + 3 \cdot \frac{m}{3} + 3 \cdot \frac{m}{3^2}$$

$$= \cancel{T\left(\frac{n}{3^k}, \frac{m^2}{3^k}\right)} + 3n$$

$$T\left(\frac{n}{3^k}, \frac{m^2}{3^k}\right) + 3n \left[1 + \frac{1}{3} + \frac{1}{3^2} + \frac{1}{3} + \dots + \frac{1}{3^{k-1}} \right]$$

Put $\frac{m}{3^k} = 1 \Rightarrow 3^k = m \Rightarrow k = \log_3 m$

$$\Rightarrow T(n, m^2) = T\left(1, \frac{m^2}{m}\right) + 3n \left[\frac{1 \cdot (1 - 1/3)^k}{(1 - 1/3)} \right]$$

$$= T(1, m) + 3n \times \frac{3}{2} \left(1 - \frac{1}{3^k}\right)$$

$$= 3n + \frac{9n}{2} \left(1 - \frac{1}{m}\right)$$

$$= 3n + \frac{9n}{2} - \frac{9}{2}$$

$$= 15 \frac{n}{2} + \frac{9}{2} = O(n)$$

Answer.

$$\therefore T(n, m^2) = O(n)$$

Another Gedi:

$$T(n) = 15 \frac{n}{2} + \frac{9}{2}$$

$$= 15 \frac{n}{2} + O(1) \Rightarrow 15 \frac{n}{2} + O(1)$$

2-2 Recurrence Tree


Unbalanced

$$T(n) = T(0.01n) + T(0.99n) + Cn$$

\hookrightarrow longest branch \rightarrow VP.
 \hookrightarrow Shortest branch VP.

$$T(n) = T\left(\frac{n}{100}\right) + T\left(\frac{99n}{100}\right) + Cn$$

Recursion tree



depth of the Smallest branch $\boxed{n = \log_{100} n}$

$$\frac{n}{100} = 1$$

$$100^n = n$$

$$n = \log_{100} n$$

depth of the largest branch. $\therefore n \cdot \left(\frac{99}{100}\right)^k = 1$

$$n \approx \left(\frac{100}{99}\right)^k$$

$$k = \log_{\frac{100}{99}} n$$

$T(n) < cn + cn + cn \dots$ upto $\log_{\frac{100}{99}} n$ levels

$$T < cn \cdot \log_{\frac{100}{99}} n$$

$$\boxed{T(n) = O\left(n \log_{\frac{100}{99}} n\right)} \quad \text{--- I}$$

2. $T(n) > cn + cn + cn \dots$ upto $\log_{100} n$ level

$$T(n) > cn \log_{100} n$$

$$\boxed{T(n) = \Omega\left(n \log_{100} n\right)} \quad \text{--- II}$$

From equation I & II, we can conclude

Asymptotically

$$T(n) = \Theta(n \log n)$$

2-2.

(c) Domain Range Substitution

$$T(m) = 9 + \left(\frac{m}{3} + \frac{m^2}{\log m} \right)$$

$$\text{let } m = 3^k \Rightarrow k = \log_3 m.$$

$$\Rightarrow m^2 = (3^k)^2 = 9^k.$$

$$T(3^k) = 9T\left(\frac{3^k}{3}\right) + \frac{9^k}{k}.$$

$$T(3^k) = 9T(3^{k-1}) + \frac{9^k}{k}.$$

$$f(k) = T(3^k)$$

$$f(k-1) = T(3^{k-1})$$

$$f(k) = 9f(k-1) + \frac{9^k}{k}.$$

$$= 9 \left[9f(k-2) + \frac{9^{k-1}}{(k-1)} \right] + \frac{9^k}{k}.$$

$$= 9^2 \left[9f(k-3) + \frac{9^{k-2}}{(k-2)} \right] + \frac{9^k}{(k-1)} + \frac{9^k}{k}.$$

$$f(k) = 9^k f(0) + \frac{9^k}{1} + \frac{9^k}{2} + \dots + \frac{9^k}{k-1} + \frac{9^k}{k}.$$

$$= 9^k \cdot \Theta(1) + 9^k \left[1 + \frac{1}{2} + \dots + \frac{1}{k-1} + \frac{1}{k} \right].$$

$$T(k) = 9^k \cdot \Theta(1) + 9^k \log k.$$

$$T(3^k) = 9^k \cdot \Theta(1) + 9^k \log k.$$

$$\text{Since } 3^k = m \Rightarrow 9^k = m^2.$$

$$k = \log_3 m.$$

cont.

(9)

2 Fun with Recurrences 14 / 15

✓ + 1 pts (c) correct substitution

✓ + 1 pts (c) correct relation for step 2

✓ + 1 pts (c) correct relation for step 3

✓ + 1 pts (a) Extra credit for correct constant

✓ + 5 pts (B) All Correct

✓ + 5 pts (a) all correct

Incorrect/missing steps for part c. Final answer missing/incorrect

$$T(n) = n^2 : \Theta(1) + n^2 \log \log n.$$

$$= \Theta(n^2 \log \log n) \quad \text{Lightly bound.}$$

2-3 solution

$$T(n) = T\left(\frac{n}{3^{3/4}}\right) + \Theta(1).$$

$$T(n) = T\left(\frac{n}{3^{3/4}}\right) + C.$$

Compare with $T(n) = a T\left(\frac{n}{b}\right) + f(n).$

$$a=1, b=3, f(n)=C.$$

$$n \log^a n = n \log^3 n = n^0 = 1. \quad f(n) = \underline{\underline{C}}.$$

log. Master theorem Case 2.

$$T(n) = \Theta(1 \cdot \log_{3^{3/4}} n)$$

$$= \Theta(\log_{3^{3/4}} n)$$

(b) The function BLA returns value 1.


(c) Expecting running time of Foo. 3

Deriving the Recurrence.

$$S(n) = S\left(\frac{n}{3^{3/4}}\right) + \Theta(n)$$

$$= S\left(\frac{n}{3^{3/4}}\right) + \underline{\underline{C}}(n) \quad \text{4}$$

cont.



← Base case.

$$S(n) = C_m + \frac{C_m}{3^{3/4}} + \frac{C_m}{(3^{3/4})^2} + \dots + \frac{C_m}{(3^{3/4})^{k-1}} + S(1)$$

$$= C_m \left[1 + \frac{1}{3^{3/4}} + \frac{1}{(3^{3/4})^2} + \dots + \frac{1}{(3^{3/4})^{k-1}} \right] + C'$$

$$= C(n) \cdot \frac{1 - \frac{1}{(3^{3/4})^k}}{\left(1 - \frac{1}{3^{3/4}}\right)} \quad \because \left(\frac{1}{3^{3/4}}\right)^k = \frac{1}{n} \quad \text{(base case)}$$

$$k = \log_{3^{3/4}} n$$

$$S_m = C_m \left(\frac{3^{3/4}}{3^{3/4} - 1} \right) \cdot \left(1 - \frac{1}{n} \right)$$

$$S_m = \frac{3^{3/4}}{(3^{3/4} - 1)} \cdot (C_m - C)$$

$$S_m = \Theta(n)$$

3 Expected Running Time 5 / 7

✓ - 0 pts Correct

✓ - 1 pts Incorrect/Missing recurrence relation for $n > 5$ in part (a)

✓ - 1 pts Incorrect/Missing base case for $n \leq 5$ in part (a)

✓ - 0.5 pts Incorrect/Missing $T(n)$ in part (a)

✓ + 0.5 pts Correct final answer in part (c)

3 Missing justification for why $\text{Bla}(\text{Foo})$ is constant time

4 Missing base case ($n \leq 1$)