

# CSCI-GA.1170-001 Homework 12

Ankit Sati

TOTAL POINTS

42 / 35

QUESTION 1

1 Zero-Weight Cycles and Shortest Paths

22 / 14

- ✓ - 0 pts *a,b,c* Correct
- ✓ + 3 pts 12-1) *e)* extra credit correct
- ✓ + 2 pts 12-1) *d)* (extra credit) algo correct
- ✓ + 3 pts 12-1) *d)* (extra credit) correctness correct

QUESTION 2

2 Modified Dijkstra - Too Slow? 8 / 10

- ✓ - 0 pts Correct
- ✓ - 2 pts Partial points to proof in part (c)

1 You are explaining how the algorithm works.

Your proof needs to be more rigorous.

QUESTION 3

3 Testing the Tester! 12 / 11

- (a)
  - ✓ - 0 pts Correct
- (b)
  - ✓ - 0 pts Correct Case-1
  - ✓ + 3 pts Correct Case-2 - Follows from Check 2,4
  - ✓ + 2 pts Partially correct Case-3 (proper reason for "why shorter path exists mathematically" not specified, etc.)
  - ✓ + 2 pts Partially correct Case-4 (reasoning, etc.)
- (c)

✓ - 1.5 pts Incorrect/Missing  $\$w'_e$  blank

✓ - 1.5 pts Incorrect/Missing Reasoning

(d)

✓ - 1 pts Incorrect/Missing  $\$w'(P)$  blank

✓ - 1 pts Incorrect/Missing Derivation of the Equation (telescopic sum is main)

✓ - 1 pts Incorrect/Missing Reasoning behind statement "min-cost path in  $G$  is not affected by the new weights"

(e)

✓ - 0 pts Incorrect/Missing solution



2020

अक्टूबर  
October

19 सोमवार  
Monday

Problem 12 - 1

(a)

Base Case:

Let us consider a graph with 2 vertices as shown:



A zero weight cycle can be

$$1 \rightarrow 2 \rightarrow 3 \rightarrow 2 \rightarrow 1$$

Now, it can be further broken

into two simple zero-weight cycles

$$1 \rightarrow 2 \rightarrow 1 \quad \text{and} \quad 2 \rightarrow 3 \rightarrow 2.$$

अक्टूबर/October

Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25



अक्टूबर  
October

2020

20 मंगलवार  
Tuesday

Induction Hypothesis:

Let us consider a graph with  $k$  vertices and assume that it follows the given statement that a zero-weight cycle can be decomposed into further <sup>simple</sup> zero-weight cycles.

Induction step:

Let us now consider the graph has  $k+1$  vertices.

$$\text{So, } V_{k+1} = V_k \cup v_{k+1}$$

Basically, the graph with  $k$  vertices could be decomposed into

अक्टूबर/October

Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Fr	Sa	Su	Mo	Tu	We	
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24


2020



अक्टूबर  
October

21 बुधवार  
Wednesday

simple cycles.



This is the region where some vertices are there which are present in more than one cycle

Now, Since the  $(k+1)$ th vertex is a new node added to the graph with  $K$  vertices. So, this new graph formed also can decomposed further

अक्टूबर/October

Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25




अक्टूबर  
October

2020

22 बृहस्पतिवार  
Thursday

into simple cycles.

Ex:



The cycle  $1 \rightarrow 2 \rightarrow 3 \rightarrow 2 \rightarrow 4 \rightarrow 3 \rightarrow 1$

can be decomposed into

$1 \rightarrow 2 \rightarrow 4 \rightarrow 3 \rightarrow 1$ ,  $2 \rightarrow 3 \rightarrow 2$ ,  $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$

and  $2 \rightarrow 4 \rightarrow 3 \rightarrow 2$

अक्टूबर/October

Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa							
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31

2020



अक्टूबर  
October

23 शुक्रवार  
Friday

(b)

Given: R is relation such that

$(u, v) \in R$  if both u and v lie

on some cycle with zero weight

Reflexive:

$(a, a) \in R \quad \forall a \in V$

The shortest weight from a node to itself is 0 which is found by moving to no other vertex from a.

So, R is reflexive.

Symmetric:

अक्टूबर/October

Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17



अक्टूबर  
October

2020

24 शनिवार  
Saturday

Let us consider  $(a, b) \in R$

This means  $a$  and  $b$  lie

on some zero weight cycle of  
graph.

So, we can say that  $b$  and  $a$   
also lie on the same zero  
weight cycle.

So,  $(b, a) \in R$

Since  $(a, b) \in R$  and  $(b, a) \in R$

So,  $R$  is symmetric

Transitive:

Let  $(a, b) \in R$  and  $(b, c) \in R$

So,  $a$  and  $b$  lie on some zero-

अक्टूबर/October

Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25

2020




अक्टूबर  
October

25 रविवार  
Sunday

weight cycle. Also, b and c lie on some zero weight cycle. So, a, b and c all lie on some zero weight cycle.

So,  $(a, c) \in R$



$(a, b) \in R$  for the cycle  $a \rightarrow b \rightarrow a$

$(b, c) \in R$  for the cycle  $b \rightarrow c \rightarrow b$

$(a, c) \in R$  for the cycle  $a \rightarrow b \rightarrow c \rightarrow b \rightarrow a$

Therefore R is transitive.

Since R is reflexive, symmetric, and transitive. So, R is an equivalence relation.

अक्टूबर/October

Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa							
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31

अक्टूबर  
October



2020

26 सोमवार  
Monday

(c)

For the adjacency matrix of the graph, we will put  $\text{Adj}[u, u] = \infty$

which means the distance with node  $u$  to itself is  $\infty$  initially.

Now, we will run the

Floyd - Warshall Algorithm

and check if  $\text{Adj}[u, u]$  is 0

for some  $u \in V$

Algorithm  $\rightarrow$

FLOYD-WARSHALL( $W$ )

$n \leftarrow \text{rows}(W)$

$D^0 \leftarrow W$

अक्टूबर/October

Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa							
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31



2020

अक्टूबर

October

27 मंगलवार  
Tuesday

{ for  $k \leftarrow 1$  to  $n$

{ do for  $i \leftarrow 1$  to  $n$

{ { do for  $j \leftarrow 1$  to  $n$

if  $d_{ik}^{K-1} + d_{kj}^{K-1} < d_{ij}^K$ :

{  $d_{ij}^K = d_{ik}^{K-1} + d_{kj}^{K-1}$

$\Pi_{ij} = K$

{ }  
}

{ for  $i = 1$  to  $n$

{ if  $d_{ii}^n = 0$ :

{ Take a vector  $v$

$v = \text{NODES-IN-CYCLE}(i, \Pi)$

{

}

$\text{NODES-IN-CYCLE}(i, \Pi)$

Backtrack from node  $i$  over parent matrix  $\Pi$  until you again get  $i$

अक्टूबर/October

Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17

अक्टूबर  
October



2020

28 बुधवार  
Wednesday

Running Time :-

This algorithm is mainly Floyd-Warshall algorithm with slight changes. So, runtime is  $O(NV^3)$ .

Correctness -

We did a <sup>run of</sup> Floyd - Warshall - Algorithm to know if distance between a node u to itself is becoming 0. In this case, we call the function NODES-IN-CYCLE which backtracks from current node

अक्टूबर/October

Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25

2020



अक्टूबर  
October

29 बृहस्पतिवार  
Thursday

until it gets back to that node

again. So, in this way we are keeping track of all the vertices found in the zero weight cycle.

And hence the algorithm is

justified.

(d)

Algorithm →

## FLOYD - WARSHALL(W)

$m \leftarrow \text{rows}(w)$

$D^o \leftarrow W$

for K=1 to n

for  $i = 1$  to  $n$

for j = 1 to n

## अक्टूबर/October

Th Fr Sa Su Mo Tu We Th Fr Sa Su Mo Tu We Th Fr Sa Su Mo Tu We Th Fr Sa  
01 02 03 04 05 06 07 08 09 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31



अक्टूबर

October

2020

30 मंगलवार  
Tuesday

if  $d_{ik}^K + d_{kj}^{K-1} < d_{ij}^K$ :

$$d_{ij}^K = d_{ik}^{K-1} + d_{kj}^{K-1}$$

$$\pi_{ij} = K$$

for  $i = 1$  to  $n$

if  $d_{ii}^n = 0$

vector<int>  $v = \text{NODES\_IN\_CYCLE}(i, \pi)$

$m = \text{INT\_MAX}$

for  $i = 0$  to  $v.size()$

$$m = \min(m, v[i])$$

$i \cdot \text{equivalence} = m$

Approach →

Our algorithm is similar to the

अक्टूबर/October

Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa							
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31

2020



अक्टूबर  
October

31 बुधवार  
Wednesday

algorithm used in previous part.

Here after finding the vector of vertices which are involved in the cycle, we find the minimum of all those and store in equivalence property of every vertex.

In this way, if equivalence will store the minimum node  $v^*$  value for any node  $v$  which is present in the same zero weight cycle in which  $u$  is also present.

अक्टूबर/October

Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa							
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31

2020



नवम्बर

November

01 रविवार  
Sunday

In this way the algorithm is justified.

(e)

Since there is some zero-weight cycle which includes vertices  $t$  and  $i$ . Also, there is one another ~~zero~~-weight cycle which includes vertices  $j$  and  $k$ .

Since there would be some balancing edges whose weights must be cancelling each other's weight, so it will be optimal.

नवंबर/November

Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	
01	02	03	14	15	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24

2020



नवम्बर

November

07 शनिवार  
Saturday

to consider the shortest path

from ~~h~~ to i and shortest path  
from j to k.

So, for the shortest path from  
h to k, we can go via

shortest path from h to i,

then via shortest path from i to

j and then <sup>via</sup> the shortest path

from j to k.

This will provide us the  
shortest path between ~~h~~ and ~~k~~.

So,  $\delta(h, k) = \delta(h, i) + \delta(i, j) + \delta(j, k)$ .

नवम्बर/November

Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24

1 Zero-Weight Cycles and Shortest Paths 22 / 14

- ✓ - 0 pts *a,b,c* Correct
- ✓ + 3 pts *12-1) e) extra credit correct*
- ✓ + 2 pts *12-1) d) (extra credit) algo correct*
- ✓ + 3 pts *12-1) d) (extra credit) correctness correct*

नवम्बर

November




2020

02 सोमवार  
Monday

### Problem 12-2

(a) The required example is:



If we remove step 12, then  
the shortest path from 1 to 3 will  
come 2, which is wrong as  
the path is -1 which comes  
on going  $1 \rightarrow 2 \rightarrow 3$ .

नवम्बर/November

Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo							
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30

# 2020



नवम्बर

November

03 मंगलवार  
Tuesday

(b)

In the worst case, the shortest path from a source node  $s$  to a node  $v$  will pass through all the edges. So, the number of iterations will be bounded by the number of edges on.

Hence the algorithm terminates in a finite number of steps.

(c)

In the usual Dijkstra algorithm, we donot visit a vertex more

नवम्बर/November

Su	Mo	Tu	We	Th	Fr	Sa	Su	M.	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su								
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30

नवम्बर

November



2020

04 बुधवार  
Wednesday

than once. This was so because all the weights were considered to be positive in that case.

So, as soon as the vertex is visited, its distance from source node is the minimum.

But, when we consider negative weights also, then we may have to revisit some node to find even more shorter path. This thing is maintained by line 12, where we are.

नवम्बर/November

Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo							
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30

2020



नवम्बर

November

05 बुधसप्तविंशति  
Thursday

inserting the node  $v$  back to  
the priority-queue  $Q$  so that it  
may be visited again in the  
coming iterations to find more  
shorter paths.

In this way MODIFIED DIJKSTRA  
well works for the negative  
weight edge case such that there  
are no negative weight cycles.

Hence the ~~above~~ above algorithm  
is justified.

नवम्बर/November

Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	Tu	We	Th	Fr	Sa	Su	Mo	
01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24

## 2 Modified Dijkstra - Too Slow? 8 / 10

✓ - 0 pts Correct

✓ - 2 pts Partial points to proof in part (c)

- 1 You are explaining how the algorithm works. Your proof needs to be more rigourous.

### ~~Notes~~ Problem 3

(a) Runtime for each step of Algo.

- 1  $O(1)$
- 2  $O(|V| + |E|) + O(|E|) + O(|V| + |E|)$
- 3  $O(|V|)$
- 4  $O(|V| + |E|)$  {. Step 4 of Bellman Ford}
- 5  $O(1)$

$$\therefore \text{Runtime} = O(|V| + |E|)$$

The algorithm will accept this, if this answer is complete.

Note  $\Rightarrow$  Any right answer will be a single source of shortest path and it should have:

1.  $s.d = 0$ ,  $s.\pi = \text{NIL}$  for source vertex.
2. The given shortest path is reachable from  $s$ .
3.  $v.d$  is correctly calculated.
4. This graph cannot be relaxed any further.

If all the above arguments are met & verified then the algorithm will return correct.

Never, all the eight verification are made by the algorithm & will return correct for a right answer.

(a)  $S(s, v) = \infty$  but  $v.d < \infty$

→  $v$  is actually not reachable but  $v.d < \infty$  indicates it is still reachable.

Mence this will fail in:

(i) 2. a → there will be no edge  $(v, \pi, v)$  as.  $v$  is unreachable

(ii) 2. b → On running DFS, we fail to reach this node with  $v.d < \infty$

→  $S(s, v) < \infty$  but  $v.d = \infty$

$v$  is actually reachable but  $v.d = \infty$  indicates it is not.

It will be caught in:

→ 2. c. → On running DFS, it will reach node  $v$  with  $v.d = \infty$

→ 4. → On the Bellman-Ford pass, graph will relax further, as  $v.d$  shouldn't be  $\infty$ .

→  $S(s, v) < v.d < \infty$

As.  $S(s, v) < v.d$ , it means we can further relax  $v$ .

∴ never it will fail in.

→ 4. - on the bellman ford. pass, we will observe relaxations.

→  $v.d < S(s, v) < \infty$

This can indicate wrong calculation of  $v.d$ .

To catch this specific case, we need to find the  $v$  where  $v.d$  is not matching with  $v.\pi.d + w(v, \pi, v)$ .

Where this case will be caught in?

3 Testing the Tester! 12 / 11

(a)

✓ - 0 pts Correct

(b)

✓ - 0 pts Correct Case-1

✓ + 3 pts Correct Case-2 - Follows from Check 2,4

✓ + 2 pts Partially correct Case-3 (proper reason for "why shorter path exists mathematically" not specified, etc.)

✓ + 2 pts Partially correct Case-4 (reasoning, etc.)

(c)

✓ - 1.5 pts Incorrect/Missing \$\$w'\_e\$\$ blank

✓ - 1.5 pts Incorrect/Missing Reasoning

(d)

✓ - 1 pts Incorrect/Missing \$\$w'(P)\$\$ blank

✓ - 1 pts Incorrect/Missing Derivation of the Equation (telescopic sum is main)

✓ - 1 pts Incorrect/Missing Reasoning behind statement "min-cost path in  $G$  is not affected by the new weights"

(e)

✓ - 0 pts Incorrect/Missing solution