



Optical/Infrared Observations of the Extraordinary GRB 250702B: A Highly Obscured Afterglow in a Massive Galaxy Consistent with Multiple Possible Progenitors

Jonathan Carney¹ , Igor Andreoni¹ , Brendan O'Connor^{2,21} , James Freeburn¹ , Hannah Skobe² , Lewi Westcott³ , Malte Busmann⁴ , Antonella Palmese² , Xander J. Hall² , Ramandeep Gill^{5,6} , Paz Beniamini^{6,7,8} , Eric R. Coughlin⁹ , Charles D. Kilpatrick¹⁰ , Akash Anumalapudi¹ , Nicholas M. Law¹ , Hank Corbett¹ , Tomas Ahumada¹¹ , Ping Chen^{12,13} , Christopher Conselice³ , Guillermo Damke¹⁴ , Kaustav K. Das¹¹ , Avishay Gal-Yam¹⁵ , Daniel Gruen^{4,16} , Steve Heathcote¹⁴ , Lei Hu² , Viraj Karambelkar¹¹ , Mansi Kasliwal¹¹ , Kathleen Labrie¹⁷ , Dheeraj Pasham^{18,19} , Arno Riffeser⁴ , Michael Schmidt⁴ , Kritti Sharma¹¹ , Silona Wilke⁴ , and Weicheng Zang²⁰

¹ Department of Physics and Astronomy, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599-3255, USA; jcarney@unc.edu

² McWilliams Center for Cosmology and Astrophysics, Department of Physics, Carnegie Mellon University, Pittsburgh, PA 15213, USA

³ Jodrell Bank Centre for Astrophysics, Alan Turing Building, University of Manchester, Oxford Road, Manchester M13 9PL, UK

⁴ University Observatory, Faculty of Physics, Ludwig-Maximilians-Universität München, Scheinerstrasse 1, 81679 Munich, Germany

⁵ Instituto de Radioastronomía y Astrofísica, Universidad Nacional Autónoma de México, Antigua Carretera a Pátzcuaro # 8701, Ex-Hda. San José de la Huerta, Morelia, Michoacán, C.P. 58089, México

⁶ Astrophysics Research Center of the Open university (ARCO), The Open University of Israel, P.O. Box 808, Ra'anana 4353701, Israel

⁷ Department of Natural Sciences, The Open University of Israel, P.O. Box 808, Ra'anana 4353701, Israel

⁸ Department of Physics, The George Washington University, 725 21st Street NW, Washington, DC 20052, USA

⁹ Department of Physics, Syracuse University, Syracuse, NY 13210, USA

¹⁰ Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA), Northwestern University, 1800 Sherman Avenue, Evanston, IL 60201, USA

¹¹ Cahill Center for Astrophysics, California Institute of Technology, MC 249-17, 1216 E. California Boulevard, Pasadena, CA 91125, USA

¹² Institute for Advanced Study in Physics, Zhejiang University, Hangzhou 310027, People's Republic of China

¹³ Institute for Astronomy, School of Physics, Zhejiang University, Hangzhou 310027, People's Republic of China

¹⁴ Cerro Tololo Inter-American Observatory/NSF NOIRLab, Casilla 603, La Serena, Chile

¹⁵ Department of Particle Physics and Astrophysics Weizmann Institute of Science, 234 Herzl Street, Rehovot, Israel

¹⁶ Excellence Cluster ORIGINS, Boltzmannstrasse 2, 85748 Garching, Germany

¹⁷ Gemini Observatory/NSF's NOIRLab, 670 N. A'ohoku Place, Hilo, HI 96720, USA

¹⁸ Eureka Scientific, 2452 Delmer Street, Suite 100, Oakland, CA 94602-3017, USA

¹⁹ Department of Physics, The George Washington University, Washington, DC 20052, USA

²⁰ Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA

Received 2025 September 25; revised 2025 October 30; accepted 2025 November 1; published 2025 November 26

Abstract

GRB 250702B was the longest gamma-ray burst ever detected, with a duration that challenges standard collapsar models and suggests an exotic progenitor. We collected a rich set of optical and infrared follow-up observations of its rapidly fading afterglow using a suite of telescopes including the W. M. Keck Observatory, the Gemini telescopes, the Magellan Baade Telescope, the Victor M. Blanco 4 m telescope, and the Fraunhofer Telescope at Wendelstein Observatory. Our analysis reveals that the afterglow emission is well described by forward shock emission from a highly obscured relativistic jet. Deep photometric observations of the host galaxy reveal a massive ($10^{10.66} M_{\odot}$), dusty, and extremely asymmetric system that is consistent with two galaxies undergoing a major merger. The galactocentric offset, host galaxy properties, and jet characteristics disfavor a jetted tidal disruption event (TDE) around a supermassive black hole but do not definitively distinguish between competing progenitor scenarios. We find that the afterglow and host are consistent with a range of progenitors, including an atypical collapsar, a merger between a helium star and a stellar-mass black hole, the disruption of a star by a stellar-mass compact object (micro-TDE), and the tidal disruption of a star by an off-nuclear intermediate-mass black hole.

Unified Astronomy Thesaurus concepts: Relativistic jets (1390); Black holes (162); Tidal disruption (1696); Gamma-ray bursts (629); Time domain astronomy (2109); X-ray transient sources (1852); Gamma-ray transient sources (1853); Infrared astronomy (786)

1. Introduction

Relativistic jets can illuminate the entire electromagnetic spectrum, reaching very high luminosities from gamma rays to the radio when the observer's line of sight is fortuitously

aligned with the jet axis. Among the plethora of high-energy transients, gamma-ray bursts (GRBs) are the most common. Progenitors of the majority of GRBs include collapsars and compact object mergers (see e.g., T. Piran 2004, for a review). The prompt emission from GRBs is usually found to last from ~ 0.01 s up to several minutes (C. Kouveliotou et al. 1993), but a rare population of so-called ultralong GRBs exists (A. J. Levan et al. 2014), where prompt emission can last for several thousand seconds (see also Y. Y. Tikhomirova & B. E. Stern 2005). The progenitors of these systems are believed to be massive stars with particularly large radii, which is supported by the emergence of bright supernovae after some

²¹ McWilliams Fellow.



ultralong GRBs (e.g., G. Stratta et al. 2013; A. J. Levan et al. 2014). A helium star merger progenitor (C. L. Fryer & S. E. Woosley 1998) has also been proposed, for instance, in the case of ultralong GRB 101225A (C. C. Thöne et al. 2011).

Although exceptionally rare, the disruption of a star by a massive black hole may also produce relativistic jets, in which case the gamma-ray activity may last for a few days. This extended emission from a relativistic tidal disruption event (TDE) was observed for Sw J1644+57 (J. S. Bloom et al. 2011; D. N. Burrows et al. 2011; A. J. Levan et al. 2011; B. A. Zauderer et al. 2011), Sw J2058+05 (S. B. Cenko et al. 2012; D. R. Pasham et al. 2015), Sw J1112–82 (G. C. Brown et al. 2015, 2017), and AT 2022cmc (I. Andreoni et al. 2022; D. R. Pasham et al. 2023). The high-energy spectrum of these sources is softer than typical GRBs; thus, the emission is long-lasting predominantly in the hard X-rays.

On 2025 July 2, the Gamma-ray Burst Monitor (GBM) on board the Fermi satellite detected multiple GRBs likely originating from the same source (E. Neights et al. 2025c). They were observed between 2025 July 2 13:56:00 UT and 2025 July 2 16:21:33 UT and designated GRB 250702B (Fermi GBM Team 2025a), GRB 250702C (later retracted; Fermi GBM Team 2025b; E. Neights et al. 2025b), GRB 250702D (Fermi GBM Team 2025c), and GRB 250702E (Fermi GBM Team 2025d). Henceforth we will refer to the event as GRB 250702B, in accordance with standard GRB nomenclature. The transient was later reported to have been concurrently observed by other facilities, including Neil Gehrels Swift Observatory (hereafter Swift; N. Gehrels et al. 2004) Burst Array Telescope’s Gamma-ray Urgent Archiver for Novel Opportunities (J. DeLaunay et al. 2025; B. O’Connor et al. 2025), the Monitor of All-sky X-ray Image (MAXI; Y. Kawakubo et al. 2025), Konus-Wind (D. Frederiks et al. 2025), and the Space Variable Objects Monitor (SVOM; Svom/Grm Team et al. 2025). Konus-Wind detected multiple overlapping X-ray emission episodes over hours from 2025 July 2 12:40 UT to 2025 July 2 16:50 UT (D. Frederiks et al. 2025), suggesting an extreme duration for the transient. A complete gamma-ray analysis combining Konus-Wind and Fermi finds a gamma-ray duration of ~ 25 ks (E. Neights et al. 2025a). This extraordinary duration was further extended by retroactive analysis of stacked Einstein Probe (EP; W. Yuan et al. 2025) Wide-field X-ray Telescope (WXT) images, which revealed emission as early as July 1 (H. Q. Cheng et al. 2025), giving it a central engine duration of ~ 100 ks (E. Neights et al. 2025a). This makes GRB 250702B the longest-duration GRBs ever detected—the previous record holder, GRB 111209A, lasted ~ 15 ks in the gamma rays and had a central engine duration of ~ 25 ks (B. Gendre et al. 2013). The exceptionally long duration of GRB 250702B complicates explanation from the known population of GRBs (T. Piran 2004; S. E. Woosley & J. S. Bloom 2006).

The EP mission detected a soft X-ray counterpart (designated EP 250702a) at 2025 July 2 02:53 UT, with a localization accuracy of 2.4 . Follow-up observations with the Swift X-ray Telescope (XRT) further improved the localization to a radius of 2.0 centered on J2000 coordinates R.A. = $18^{\text{h}} 58^{\text{m}} 45^{\text{s}}.61$, decl. = $-07^{\circ} 52' 26.9''$ (J. A. Kennea et al. 2025). Following these initial reports, ultraviolet (UV), optical, and near-infrared (NIR) follow-up observations ensued. No UV or optical counterparts were identified in these initial searches (see Table 1). At 0.746 days after the first Fermi detection, Very Large Telescope (VLT) HAWK-I

observations identified an NIR counterpart (A. J. Levan et al. 2025) at coordinates R.A. = $18^{\text{h}} 58^{\text{m}} 45^{\text{s}}.57$, decl. = $-07^{\circ} 52' 26.2''$. Subsequent follow-up observations confirmed that the NIR counterpart was rapidly fading and coincident with an extended NIR galaxy (A. J. Levan et al. 2025). Neither the transient nor the host galaxy was initially detected in any optical photometric band during follow-up, suggesting significant extinction along the line of sight.

In this work, we focus on optical and NIR observations of GRB 250702B and its host galaxy. The Letter is organized as follows: We describe our observations in Section 2 and in Appendix A. Modeling of both the host galaxy and the transient emission is presented in Section 3. We discuss our results and their implications for the progenitor in Section 4. We present our conclusions in Section 5.

Throughout this work we adopt a standard Λ CDM cosmology with $H_0 = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.315$, and $\Omega_\Lambda = 0.685$ (Planck Collaboration et al. 2020). All magnitudes are reported in the AB system.

2. Observations

Following the initial Fermi and EP detections of GRB 250702B, we began an optical to NIR follow-up campaign, beginning with *griz* observations using the Dark Energy Camera (DECam; B. Flaugher et al. 2015) mounted on the Victor M. Blanco 4 m Telescope at Cerro Tololo Inter-American Observatory (Program ID 2025A-729671; PI: Palmese). After the XRT localization and counterpart detection by VLT became available, we continued to monitor the afterglow with the Fraunhofer Telescope at Wendelstein Observatory (FTW) using the Three Channel Imager (3KK; F. Lang-Bardl et al. 2016) for simultaneous optical and NIR imaging. We also acquired deep *z*-band imaging with both Gemini Multi Object Spectrographs on Gemini South (GMOS-S; program ID GS-2025A-Q-117; PI: Andreoni) and Gemini North (GMOS-N; program ID GN-2025A-Q-213; PI: Andreoni) and deep NIR monitoring with the Multi-Object Spectrograph for Infrared Exploration (MOSFIRE; I. S. McLean et al. 2012) on the W. M. Keck Observatory telescope I (PI: Kasliwal; program ID C348), the NOAO Extremely Wide Field Infrared Imager (NEWFIRM; R. G. Autry et al. 2003) on the Victor M. Blanco 4 m Telescope at Cerro Tololo Inter-American Observatory, and the FourStar infrared camera (S. E. Persson et al. 2013) mounted on the 6.5 m Magellan Baade Telescope. Observations extended from ~ 15 hr to ~ 41 days after the initial Fermi detection. Details for these observations can be found in Appendix A.

To measure the transient flux, we performed image subtraction using our late-time Magellan/FourStar observations as templates for the FTW/3KK, Keck/MOSFIRE, and NEWFIRM images. We also reprocessed publicly available data from the Hubble Space Telescope (HST; program ID 17988; PI: Levan; previously published in A. J. Levan et al. 2025). To reduce host contamination in HST photometry, we employed the morphological models described in Section 3.3 as templates for image subtraction. The extreme obscuration along the line of sight led to nondetections at all UV/optical wavelengths for the transient, with the NIR counterpart to GRB 250702B only detected in *H*, *F160W*, and *K_s* bands. The host galaxy was also found to be highly obscured, but our observations revealed significant detections in *z*, *H*, *F160W*, and *K_s* bands (Figure 1). This dataset was combined with publicly available UV/optical/IR data from the literature for our subsequent analysis (see Table 1 for the full

Table 1
Photometry Table of UVOIR Observations of GRB 250702B

Datetime	Instrument	Filter	Transient Mag	References
2025-07-03 02:06:10	GOTO	<i>L</i>	>20.35	A. Kumar et al. (2025)
2025-07-03 05:12:58	Blanco/DECam	<i>g</i>	>23.56	This work
2025-07-03 05:18:18	Blanco/DECam	<i>r</i>	>23.04	This work
2025-07-03 05:22:05	Blanco/DECam	<i>i</i>	>22.26	This work
2025-07-03 05:24:03	Blanco/DECam	<i>z</i>	>22.21	This work
2025-07-03 05:26:10	Blanco/DECam	<i>z</i>	>22.08	This work
2025-07-03 07:03:03	VLT/HAWK-I	<i>H</i>	20.78 ± 0.05	A. J. Levan et al. (2025)
2025-07-03 07:39:40	VLT/HAWK-I	<i>K</i>	19.36 ± 0.02	A. J. Levan et al. (2025)
2025-07-03 08:19:00	COLIBRÍ/DDRAGO	<i>i</i>	>22.0	R. L. Becerra et al. (2025)
2025-07-03 10:54:08	SVOM/VT	<i>R</i>	>23.6	Svom/Grm Team et al. (2025)
2025-07-03 10:54:08	SVOM/VT	<i>B</i>	>23.8	Svom/Grm Team et al. (2025)
2025-07-03 17:26:10	WFST	<i>r</i>	>22.2	Y.-L. Hua et al. (2025)
2025-07-03 19:36:00	SYSU 80 cm	<i>J</i>	>17.5	X. Li et al. (2025b)
2025-07-03 21:52:47	FTW/3KK	<i>r</i>	>22.47	This work
2025-07-03 21:52:47	FTW/3KK	<i>i</i>	>22.04	This work
2025-07-03 21:52:47	FTW/3KK	<i>J</i>	>21.25	This work
2025-07-03 22:50:00	NOT/StaNCAM	<i>i</i>	>24.9	A. J. Levan et al. (2025)
2025-07-03 23:45:36	CAHA2.2/CAFOS	<i>z</i>	>21.4	I. Pérez-García et al. (2025)
2025-07-04 01:50:20	Swift/UVOT	<i>b</i>	>19.83	M. H. Siegel & Swift/UVOT Team (2025)
2025-07-04 01:48:55	Swift/UVOT	<i>u</i>	>19.73	M. H. Siegel & Swift/UVOT Team (2025)
2025-07-04 01:57:09	Swift/UVOT	<i>v</i>	>18.44	M. H. Siegel & Swift/UVOT Team (2025)
2025-07-04 01:46:11	Swift/UVOT	<i>uvw1</i>	>19.93	M. H. Siegel & Swift/UVOT Team (2025)
2025-07-04 01:51:45	Swift/UVOT	<i>uvw2</i>	>20.12	M. H. Siegel & Swift/UVOT Team (2025)
2025-07-04 03:20:59	VLT/HAWK-I	<i>K</i>	20.85 ± 0.03	A. J. Levan et al. (2025)
2025-07-04 03:36:09	VLT/HAWK-I	<i>H</i>	22.39 ± 0.06	A. J. Levan et al. (2025)
2025-07-04 03:43:54	Gemini/GMOS-S	<i>z</i>	>23.95	This work
2025-07-04 05:32:54	Swift/UVOT	<i>uvm2</i>	>19.33	M. H. Siegel & Swift/UVOT Team (2025)
2025-07-04 22:40:29	FTW/3KK	<i>r</i>	>22.06	This work
2025-07-04 22:40:29	FTW/3KK	<i>i</i>	>21.79	This work
2025-07-04 22:40:43	FTW/3KK	<i>K_s</i>	21.07 ± 0.16	This work
2025-07-05 01:44:52	GTC/HiPERCAM	<i>g</i>	>23.6	A. J. Levan et al. (2025)
2025-07-05 01:44:52	GTC/HiPERCAM	<i>r</i>	>23.0	A. J. Levan et al. (2025)
2025-07-05 01:44:52	GTC/HiPERCAM	<i>i</i>	>22.6	A. J. Levan et al. (2025)
2025-07-05 01:44:52	GTC/HiPERCAM	<i>z</i>	>22.3	A. J. Levan et al. (2025)
2025-07-05 03:07:37	VLT/HAWK-I	<i>K</i>	21.49 ± 0.05	A. J. Levan et al. (2025)
2025-07-05 08:56:18	SVOM/VT	<i>R</i>	>23.6	H. L. Li et al. (2025a)
2025-07-05 08:56:18	SVOM/VT	<i>B</i>	>23.8	H. L. Li et al. (2025a)
2025-07-05 10:14:00	Keck/MOSFIRE	<i>K_s</i>	21.96 ± 0.1	This work
2025-07-05 10:14:00	Keck/MOSFIRE	<i>H</i>	>22.12	This work
2025-07-05 10:14:00	Keck/MOSFIRE	<i>J</i>	>22.29	This work
2025-07-06 02:16:22	GTC/EMIR	<i>K_s</i>	>21.5	A. J. Levan et al. (2025)
2025-07-08 02:43:27	GTC/EMIR	<i>K_s</i>	>22.0	A. J. Levan et al. (2025)
2025-07-08 06:50:09	VLT/HAWK-I	<i>K</i>	22.53 ± 0.06	A. J. Levan et al. (2025)
2025-07-10 19:26:50	NEWFIRM	<i>H</i>	>20.50	This work
2025-07-11 21:33:30	NEWFIRM	<i>H</i>	>20.10	This work
2025-07-11 21:50:51	NEWFIRM	<i>K_s</i>	>20.62	This work
2025-07-15 01:40:27	VLT/HAWK-I	<i>K</i>	22.82 ± 0.08	A. J. Levan et al. (2025)
2025-07-15 02:39:23	HST/WFC3	F160W	26.52 ± 0.10	This work
2025-07-20 09:55:45	Gemini/GMOS-N	<i>z</i>	>24.18	This work
2025-08-12 06:00:00	Magellan/FourStar	<i>K_s</i>	>21.94	This work

Note. All magnitudes are provided in the AB system and are not corrected for Galactic extinction.

afterglow photometry dataset and Table B1 for all measured host photometry).

3. Analysis

3.1. Jet Modeling

We used the observed NIR detections obtained with Keck/MOSFIRE, FTW, VLT/HAWK-I, and HST for model fitting

(see Table 1 and Figure 2). We fit the *H*- and *K*-band data out to 3 days after discovery. We neglected the last two VLT/HAWK-I *K*-band detections, as they are likely contaminated by host galaxy flux (A. J. Levan et al. 2025) and require late-time host galaxy templates to fully capture the transient flux. We first fitted a power law,

$$F(\nu, t) = A\nu^\beta(t - t_0)^\alpha, \quad (1)$$




Figure 1. The host as detected by Gemini in z band (left), HST in the F160W filter (similar to H band; middle), and Magellan in K_s band (right). In all three images north is up and east is to the left. These are the host detections used for SED modeling (Section 3.2). In each image a crosshair marks the location of the afterglow.

where ν is the effective frequency associated with each filter, β is the spectral index, α is the temporal index, and A is the normalization. While the exact start time of the afterglow (the jet launching time) for the forward shock could have been coincident with the Fermi B, D, and E bursts (E. Neights et al. 2025a), here we assumed the time of the first burst, GRB 250702D, as the initial time t_0 . From this joint temporal and spectral fit, we obtained a temporal index $\alpha = -1.83^{+0.05}_{-0.06}$ and spectral index $\beta = -4.95^{+0.32}_{-0.32}$. In particular, we find that the late-time HST detection, after removing the majority of the host contamination (Section A.7), is consistent with the single power-law decay shown at earlier times. Overall, the NIR temporal index is consistent with the decay of the X-ray light curve (B. O’Connor et al. 2025), but the observed spectral index is clearly inconsistent with the initial X-ray reports ($\beta = -0.65 \pm 0.20$; J. A. Kennea et al. 2025).

Assuming that the observed NIR emission arises from synchrotron radiation produced by the forward shock (the afterglow) of a relativistic jet (J. Granot & R. Sari 2002), we can infer the properties of the blastwave and its surrounding environment. Using the standard afterglow closure relations (J. Granot & R. Sari 2002), we can infer the slope p of the electron’s power-law energy distribution $N(\gamma) \propto \gamma^{-p}$. Assuming that the NIR emission lies below the cooling frequency ν_c implies $p = 1 - 2\beta = 10.9 \pm 0.64$, whereas if the NIR is above the cooling frequency, we have $p = -2\beta = 9.90 \pm 0.64$. Either of these values is significantly inconsistent with both the theoretical and observed values of p , which lie closer to $p \sim 2.3$ (A. Achterberg et al. 2001; D. C. Ellison & G. P. Double 2002; A. Spitkovsky 2008; P. A. Curran et al. 2010; X.-G. Wang et al. 2015). This demonstrates that the spectral index is incredibly steep and is strongly indicative of intrinsic dust extinction (see also A. J. Levan et al. 2025; B. O’Connor et al. 2025), in addition to the large component in the Milky Way that was already corrected for (E. F. Schlafly & D. P. Finkbeiner 2011).

Therefore, we consider the closure relations utilizing the temporal decay index. This depends on the slope of the surrounding environment’s density profile of $\rho(r) \propto r^{-k}$, where $k = 0$ refers to a homogeneous medium and $k = 2$ refers to a wind environment (Z. G. Dai & T. Lu 1998; R. A. Chevalier & Z.-Y. Li 2000). We start by considering the case of a pre-jet-break afterglow. Assuming a homogeneous medium, if the ultraviolet, optical, and infrared (UVOIR) data satisfy $\nu_m < \nu < \nu_c$, where ν_m is the injection frequency, this value of α implies an electron power-law index of $p = 3.44^{+0.07}_{-0.08}$. In a

wind environment, this instead yields $p = 2.77^{+0.07}_{-0.08}$. If the UVOIR data lie above the cooling frequency, instead we have $p = 3.11^{+0.07}_{-0.08}$, independent of the surrounding environment. While these values of p are high, they are not inconsistent with the broad range of values found for GRB afterglows.

If we instead consider a post-jet-break afterglow from a nonspreading jet (e.g., A. J. Levan et al. 2025; B. O’Connor et al. 2025), for $\nu_m < \nu < \nu_c$, we find values of $p = 2.44^{+0.07}_{-0.08}$ and $p = 2.11^{+0.07}_{-0.08}$ for a homogeneous and wind environment, respectively. As it is unlikely that the NIR lies above the cooling frequency, we consider only this solution. As both of these values are typical of GRB afterglows, we cannot differentiate between these environments based on the simple temporal fit alone. Specifically, the inclusion of an intrinsic dust component is required to derive the reasonable range of intrinsic spectral slopes and therefore aid in constraining p .

Therefore, we used VegasAfterglow (Y. Wang et al. 2025) to fit the observed UVOIR data to assess whether it is consistent with synchrotron emission arising from the forward shock of a top-hat jet (R. Sari et al. 1998; J. Granot & R. Sari 2002). We do this using bilby (G. Ashton et al. 2019) with nested sampling using dynesty (J. S. Speagle 2020). We assumed a wind medium ($k = 2$), as favored in other recent works (A. J. Levan et al. 2025; B. O’Connor et al. 2025), and fit for A_* , the number density of the wind environment measured at a radius of 10^{18} cm. We assumed an on-axis viewing angle, $\theta_{\text{obs}} = 0$, because GRB 250702B’s multi-wavelength light curve does not exhibit features suggestive of off-axis viewing—the X-ray light curve decays monotonically without the delayed rebrightening characteristic of off-axis jets. We fit for the jet’s core half-opening angle, θ_c . The free parameters in the fit are the isotropic-equivalent kinetic energy at the jet’s core $E_{\text{kin,iso}}$, the initial bulk Lorentz factor at the jet’s core Γ_0 , density A_* , the jet’s half-opening angle θ_c , the slope of the electron’s energy distribution p , and the electron (ϵ_e) and magnetic field (ϵ_B) energy fractions. We included an additional intrinsic dust component in our modeling using the Milky Way dust law reported in E. L. Fitzpatrick & D. Massa (2007), which is parameterized by $A_{V,\text{host}}$ with fixed $R_V = 3.1$. This dust component $A_{V,\text{host}}$ was uniformly sampled between 0 and 10 mag. We did not include UVOIR upper limits in the fit, and we focused on the H - and K_s -band detections (Figure 2). We supplemented the UVOIR dataset (Table 1) with additional X-ray and radio data (A. J. Levan et al. 2025; B. O’Connor et al. 2025).

We performed three VegasAfterglow fits to the full multiwavelength (X-ray to NIR to radio) light curve using




Figure 2. Light curve of the observed infrared counterpart of GRB 250702B, including UV and optical upper limits. $t = 0$ is defined as the time from 2025 July 2 13:09:02 in the observer frame. Rest-frame time from the first Fermi trigger and absolute magnitudes are calculated at $z = 1.036$ (B. P. Gompertz et al. 2025) without additional K -correction and are corrected for Milky Way extinction using the dust maps by E. F. Schlafly & D. P. Finkbeiner (2011). Error bars represent 1σ confidence, and upper limits represent 5σ confidence. UV/optical observations are denoted in gray scale, with lighter colors representing longer wavelengths; NIR observations are colored blue to red according to their wavelengths. Data presented for the first time in this work, together with our host-model-subtracted photometry of the HST data first published in A. J. Levan et al. (2025), are represented by filled symbols, whereas data from the literature are represented by open symbols (see Table 1). Later VLT epochs with likely host contamination are denoted separately as squares and were excluded from modeling. For z , J , H , and K bands, the 1σ region in the posteriors for the best-fit forward shock model described in Section 3.1 is shown as a shaded region.

bilby (G. Ashton et al. 2019) with nested sampling using *dynesty* (J. S. Speagle 2020). The upper limits were not included because they are unconstraining for the fit: in the UV/optical bands this is due to the large amount of dust along the line of sight, and in the NIR this is because of lack of depth (see Figure 2). The first fit was run with the lateral jet spreading effects excluded, and the second fit included jet spreading (see Y. Wang et al. 2025, for details). A third fit for a nonspreading jet was performed with the added requirement that $\Gamma_0\theta_c > 1$ such that the spherical jet approximation holds during the deceleration phase (e.g., J. E. Rhoads 1997, 1999). The fit results are shown in the corner plots in Figures C1, C2, and C3. The model comparison to the observed multi-wavelength light curve is shown in Figure 3 for both the nonspreading (without the requirement of $\Gamma_0\theta_c > 1$) and spreading models. We find a Bayes factor (BF) strongly in favor of the model without jet spreading enabled of $\log_{10}\text{BF} = 26.8$. We discuss these solutions in Section 4.3.

There are significant departures from our models in radio wavelengths. However, at these observing frequencies, scintillation can induce high variability (see D. Dobie et al. 2020, and references therein). We therefore cannot conclude

that these are intrinsic outliers. Additionally, the early 1 keV light curve departs significantly from the fitted models, which is interpreted to be flaring activity in B. O’Connor et al. (2025).

3.2. Host Galaxy SED Modeling

The host galaxy of GRB 250702B is extremely red and was only detected in the zHK filters (Figure 1). Due to the diffuse nature of the host galaxy (Figure 1), the standard point-source upper limits derived in other filters are misleading; thus, we exclude the i - and J -band upper limits from the model fitting. We choose to include g - and r -band upper limits (Table 1), as they are half a mag deeper and thus more likely to still be constraining on a highly extinguished spectral energy distribution (SED). We modeled the SED (top panel of Figure 4) using the zHK host detection (Table B1) with the flexible stellar population (FSPS) code (C. Conroy et al. 2009). We utilized *prospector* (J. Leja et al. 2017; B. D. Johnson et al. 2019, 2021) in accordance with the methods previously outlined in B. O’Connor et al. (2022). The model fit was performed with *emcee* (D. Foreman-Mackey et al. 2013). We




Figure 3. Radio, X-ray, and infrared detections of the counterparts to GRB 250702B. The 1σ posteriors from *VegasAfterglow* fit to these data are shown as shaded regions. The left panel shows the fitting results without jet spreading enabled, and the right panel shows the results with jet spreading enabled.

adopted a G. Chabrier (2003) initial mass function, a D. Calzetti et al. (2000) dust law, and a delayed- τ star formation history. We fit for the total mass M_{gal} , age of the host galaxy t_{age} , e -folding timescale τ , metallicity Z , and intrinsic dust $A_{V,\text{host}}$. We adopt uniform bounded priors with ranges of 0.01–5.71 Gyr for t_{age} , 0.1–10.0 Gyr for τ , -1.8 to $+1.2$ dex (relative to solar) for $\log(Z/Z_{\odot})$, and 0–25 mag for dust optical depth. We then convert these into posterior distributions for total stellar mass, mass-weighted stellar age, and star formation rate (SFR; Figure 4, bottom panels). Due to the sparsity of our observations, we poorly constrained all parameters except for total stellar mass and dust, for which we find a massive ($M_{*} = 10^{10.66 \pm 0.15} M_{\odot}$) and dusty ($A_{V,\text{host}} = 0.88^{+0.47}_{-0.45}$ mag) host. A corner plot showing the posterior distributions is presented in Figure C4. We note that without the addition of redder (rest-frame) wavelengths in the fit the inferred mass may be lower than the true stellar mass of the galaxy.

3.3. Morphological Analysis of the Host Galaxy

HST observations of the host galaxy of GRB 250702B revealed a peculiar, disturbed morphology. We performed a full morphological analysis using both parametric and nonparametric fitting techniques to investigate the underlying features in the galaxy’s light.

We began with a parametric analysis using GALFIT (C. Y. Peng et al. 2002). We produced a bad-pixel segmentation map using Photutils (P. Bradley et al. 2024), to mask out nearby objects that are unrelated to the host galaxy, and a point-spread function (PSF) using the *Spike* software (A. Polzin 2025). As the morphology of the galaxy clearly shows two separate components, we performed a fit using two Sérsic profiles (Figure 5). We derive a half-light radius of the upper (north) Sérsic of $r_{50} = 0''.31 \pm 0''.08$ and a half-light radius of the lower (south) Sérsic of $r_{50} = 0''.30 \pm 0''.11$. The Sérsic indices are 0.56 ± 0.41 and 0.49 ± 0.44 , respectively, indicative of disk galaxies.

An additional fit was performed to investigate any possible underlying spiral structure. We applied two Sérsic profiles stacked onto the same coordinates, with applied azimuthal profile functions on the top profile. The primary Sérsic captures the central light, while the secondary Sérsic fits to the spiral structure. The ellipsoid shape of the galaxy light was perturbed using three functions: Fourier modes, bending modes, and a power-law spiral. The fit is shown in Figure 5.

For this model, the derived half-light radii for the central Sérsic and second Sérsic are $r_{50} = 0''.18 \pm 0''.19$ and $r_{50} = 0''.45 \pm 1''.18$, respectively. The Sérsic indices are 0.31 ± 0.95 and 0.41 ± 0.64 , respectively.

These two parametric fits yield the same goodness of fit and reveal similar residuals to the underlying structure. As such, we cannot distinguish between these two models, and we suggest that further imaging with JWST is required. In both fits, we find that after subtracting the GALFIT models there is a clear diffuse, low surface brightness emission that extends in the east–west direction above and below the galaxy’s center for $\sim 1''.75$. This may be due to the possible merger of the two separate Sérsic profiles, and it does offer some support to that scenario, though it is still unclear whether the galaxy is two separated, possibly merging Sérsic profiles or a single galaxy separated by a dust lane (A. J. Levan et al. 2025). Notably, a residual consistent with a point source is visible in both cases at the transient location.

To further investigate the host’s nature, we performed a nonparametric analysis using the *Morfometryka* software (F. Ferrari et al. 2015). This makes use of the three-dimensional “CAS” volume, representing the the concentration (C), asymmetry (A), and clumpiness (S) of a galaxy’s light profile, to classify a given galaxy’s morphology (e.g., C. J. Conselice et al. 2000; C. J. Conselice 2003). For this galaxy, we derive a concentration of $C = 2.56^{+0.05}_{-0.06}$ and an asymmetry of $A = 0.90^{+0.05}_{-0.05}$. The concentration is determined by $C = 5 \log_{10}(r_{80}/r_{20})$, and the asymmetry is determined by rotating the image about its center, subtracting the rotated image from the initial image, and then measuring the degree of residuals left behind. The classification regions defined by M. A. Bershadsky et al. (2000) determined that a high asymmetry of $A > 0.35$ indicates a galaxy undergoing a major merger. With an unusually large value of A , GRB 250702B was found to be highly asymmetrical, suggesting that a significant merger is potentially taking place.

4. Discussion

4.1. Constraints on Rebrightening

Continued NIR observations extending to 40.67 (19.98) days in the observer (rest) frame following Fermi detection show no evidence of rebrightening (Figure 2). Our reported HST F160W photometry is consistent with a continued power-law decay out to 12.53 (6.15) days in the observer (rest) frame




Figure 4. Top panel: observed host photometry (z , F160W, K_s with g and r upper limits) overlaid with 1000 randomly drawn galaxy SEDs from the *prospector* posterior distributions (Figure C4). The data are corrected for Milky Way extinction prior to modeling. Bottom left: posterior distribution of stellar mass from *prospector* modeling. Bottom middle: posterior distribution of mass-weighted age from *prospector* modeling. Here we show it in units of 5.71 Gyr (the age of the Universe at $z = 1.036$). Bottom right: posterior distribution for SFR from *prospector* modeling.

(Figure 2). This is in contrast to the non-power-law decay in the combined H and F160W light curve reported by A. J. Levan et al. (2025), where no attempt was made to remove contaminating host flux from the HST measurement. Similarly, the last two K -band VLT epochs in Figure 2 (denoted by squares) appear nonlinear in log-log scale but are similarly contaminated by host flux (see A. J. Levan et al. 2025 for a discussion about removing host flux from the VLT photometry). These points are not included in our jet modeling (Section 3.1). The final observation that we acquired, using the Magellan telescope, allows us to place only limited constraints on the rebrightening of the NIR transient. We are able to rule out rebrightening below an absolute magnitude of $K_s = -22.32$ AB mag at $\Delta t \approx 20$ days in the rest frame. This is insufficiently deep to rule out rebrightening akin to

either the well-studied optical jetted TDE AT 2022cmc (I. Andreoni et al. 2022; D. R. Pasham et al. 2023) or the prototypical GRB supernova SN 1998bw (T. J. Galama et al. 1998). AT 2022cmc was observed in K_s at rest frame $\Delta t = 3.02, 9.02$, and 13.57 days after the first Zwicky Transient Facility detections with observed absolute magnitudes of $K_s = -24.01$, $K_s = -21.88$, and $K_s = -21.54$ AB mag, respectively. Given GRB 250702B’s slightly lower redshift (AT 2022cmc occurred at $z = 1.19325$; I. Andreoni et al. 2022), and assuming that the SED of AT 2022cmc’s slow component takes roughly the form of a 30,000 K blackbody around a rest wavelength of 1000 Å, we would expect these values to be $\sim +0.3$ mag dimmer, making our late-time observations ~ 1.1 mag too shallow to constrain this potentially slowly fading component of the UVOIR light




Figure 5. Left panel: HST image of the host galaxy of GRB 250702B in the F160W filter. The location of the transient is marked by a red circle. Top middle panel: GALFIT model of best-fitting single Sérsic light profile and power-law spiral arms. Top right panel: the residual image of GALFIT after subtracting the spiral plus Sérsic model. Bottom middle panel: GALFIT model of best-fitting double-Sérsic light profiles. Bottom right panel: the residual image of GALFIT after subtracting the double-Sérsic model. In all images north is up and east is to the left.

curve. SN 1998bw ($z = 0.0085$) peaked in the I band (the reddest wavelength for which photometry is available) at an absolute magnitude of -18.82 AB mag (converted from Vega according to M. R. Blanton & S. Roweis 2007) at $\Delta t \approx 19$ days after the associated GRB. Accounting for redshift, but not extinction or K -correction, this is roughly equivalent to an H -band observation of $H \approx 25.5$ AB mag for an object at $z = 1.036$. Based on IR spectra taken within a few days of the I -band peak (F. Patat et al. 2001), we infer that a constraining K_s observation for an SN 1998bw-like supernova at the location of GRB 250702B would require observations deeper than $K_s \approx 25.5$ AB mag.

4.2. Dust Extinction from Jet Modeling

Our observations and analysis of the NIR transient further support its relativistic nature (Sections 3.1 and 4.3) and therefore its association with the high-energy detections of GRB 250702B (A. J. Levan et al. 2025; E. Neights et al. 2025a; B. O’Connor et al. 2025). The bright peak absolute magnitude, rapid fading, and color evolution not only are well fit by relativistic jet models (Section 3.1) but also are inconsistent with other common classes of extragalactic transients in the optical/NIR (see peak magnitude vs. evolution timescale plots for optical transients in, e.g., M. M. Kasliwal 2011; A. Y. Q. Ho et al. 2022).

The source is very reddened owing to high extinction, as also discussed by A. J. Levan et al. (2025), B. O’Connor et al. (2025), and B. P. Gompertz et al. (2025). After correcting for Galactic extinction, we find that both the host galaxy and the jet models need to account for a significant amount of dust. The estimate for the extinction that we infer from the jet modeling ($A_{V,GRB} = 5.43^{+0.53}_{-0.47}$ mag) is significantly larger than the value we obtain from host galaxy SED fitting

($A_{V,host} = 0.88^{+0.47}_{-0.45}$ mag). This discrepancy suggests that the GRB was produced in a dense, dusty circumstellar environment, or that a thick lane of dust is present in the host galaxy on the line of sight, between the observer and the transient (A. J. Levan et al. 2025). The line-of-sight extinction is comparable to the dustiest of those observed in a set of eight optically dark GRBs discussed in T. Krühler et al. (2011; $A_V \gtrsim 3$ mag) and to the heavily obscured relativistic TDE Sw J1644+57 ($A_V \approx 4.5$ mag; D. N. Burrows et al. 2011; see also B. O’Connor et al. (2025).

4.3. Opening Angle and Collimation-corrected Energetics

As discussed in Section 3.1, there exist two quantitatively different solutions for the afterglow model (see Figure 3) depending on the inclusion or exclusion of lateral jet spreading effects. The nonspreading jet fit (Figure C1) requires a narrow jet such that the light curve is already in a post-jet-break phase prior to the start of the earliest observations at ~ 0.5 days from discovery. This provides a better match to the observed temporal slope and the required value of p as noted above. This post-jet-break solution was also found by A. J. Levan et al. (2025), B. O’Connor et al. (2025), and B. P. Gompertz et al. (2025). However, when including lateral jet spreading, the post-jet-break decay of $\approx t^{-p}$ (J. E. Rhoads 1999; R. Sari et al. 1999) is too steep to describe the observed temporal decline of $\alpha \approx -1.8$ found for both the optical and X-ray data. Therefore, the solution identified in the presence of lateral jet spreading (Figure C2) requires that the jet break occur either during or after our latest data point. The pre-jet-break temporal slope of the model is therefore slightly shallower ($\alpha \approx -1.45$ for $p \approx 2.3$) than the observed steep X-ray and NIR decay $\alpha \approx -1.8$. Instead of identifying a steeper value of p and placing the jet break at >65 days, the fit using a spreading jet places the jet break earlier (at ~ 1 – 3 days) and underproduces

the late-time Chandra X-ray detections (B. O’Connor et al. 2025). In this case, while the jet is not extremely wide (as would be required for a jet break of >65 days; see B. O’Connor et al. 2023), it is significantly wider compared to the nonspreading jet solution and therefore also requires a higher collimation-corrected energy. That said, the inferred opening angles and collimation-corrected energies (Figures C1 and C2) are consistent with the range observed from classical long GRBs (S. B. Cenko et al. 2010; X.-G. Wang et al. 2018; B. O’Connor et al. 2023).

While at present the nonspreading post-jet-break solution provides a better fit to the data, there is no clear break observed in the light curve. As such, both solutions (wide jet vs. narrow jet) for the afterglow are potentially valid (Section 3.1). While our afterglow modeling with a spreading jet (Figures 3 and C2) does not identify the late jet-break solution, an additional fit requiring a late jet break (>65 days) was successfully able to model the light curve. This fit requires a steeper value of $p \approx 2.6$ (with higher $A_V \approx 8.5$ mag) but underpredicts the observed H -band flux, though a different dust law may account for this. Qualitatively the fit is viable, and since the correct afterglow solution (wide jet vs. narrow jet) has significant consequences for the progenitor’s requirements, additional late-time X-ray observations and continued radio monitoring can constrain the presence of the onset of any jet break in the light curve and possibly differentiate between these two possible solutions.

4.4. Constraints on the Jet’s Lorentz Factor

An additional critical constraint on the progenitor of GRB 250702B comes from the inferred bulk Lorentz factor Γ_0 , which is found to be highly relativistic. Observing the end of the jet’s deceleration phase is required to determine the exact Lorentz factor (R. Sari & T. Piran 1999; E. Molinari et al. 2007; G. Ghisellini et al. 2010), but an upper limit to the deceleration time can also provide useful constraints (B. O’Connor et al. 2020, 2024). As we do not detect the deceleration of the afterglow (and therefore have an upper limit to the time of jet deceleration), the results of our fit can be considered a lower limit to the true Lorentz factor, as lower Lorentz factors will lead to later deceleration. The posterior distribution of GRB 250702B’s initial Lorentz factor ($\log \Gamma_0 = 1.25^{+0.16}_{-0.12}$) in Figure C1 favors an ultrarelativistic blast wave with Lorentz factor $\gtrsim 10$. Enforcing the condition $\Gamma_0 \theta_c > 1$ to prevent the jet from spreading laterally during the coasting phase requires a significantly larger Lorentz factor ($\log \Gamma_0 = 2.61^{+0.27}_{-0.41}$; see Figure C3). We note that the detection of high-energy >1 MeV photons by Fermi sets an independent requirement of $\Gamma_0 \gtrsim 50$ – 80 (E. Neights et al. 2025a). Measurements of Γ_0 observed from the relativistic jetted TDEs AT 2022cmc and Sw J1644+57 are ~ 8 – 20 (B. A. Zauderer et al. 2011; B. D. Metzger et al. 2012; I. Andreoni et al. 2022; L. Rhodes et al. 2023), which depart from the observed population of GRBs, which generally have far larger Lorentz factors (e.g., $\sim$ hundreds; E.-W. Liang et al. 2010; G. Ghirlanda et al. 2012, 2018). As we cannot rule out an earlier deceleration of the jet, the measurements can be treated as a lower bound to the Lorentz factor. Higher Lorentz factors clearly disfavor a TDE-like progenitor scenario and are more consistent with a GRB-like progenitor (Section 4.6).

4.5. Host Galaxy Properties and Population Comparisons

In Figure 6 we compare both the projected physical offset (left panel) and the host-normalized offset (right panel) of GRB 250702B with the population of classical long-duration GRBs (P. K. Blanchard et al. 2016). The location of GRB 250702B is displayed as a red vertical line and lies in the 90th percentile of long GRB physical offsets and the 80th percentile of host-normalized offsets. We find that the observed galactocentric offset of GRB 250702B also lies within the range of values observed for the population of candidate offset TDEs possibly arising from an intermediate-mass black hole (IMBH; D. Lin et al. 2018, 2020; C. C. Jin et al. 2025; Y. Yao et al. 2025). The offset alone does not distinguish between progenitor scenarios and is consistent with either a massive star or an offset IMBH progenitor.

Turning to the host parameters from SED modeling (Section 3.2), we compare the inferred stellar mass as a function of both redshift and extinction to long GRB hosts. In the left panel of Figure 7 we compare GRB 250702B to both long GRB hosts (S. Savaglio et al. 2009; D. A. Perley et al. 2013), two relativistic TDEs with well-studied host galaxies (G. C. Brown et al. 2015; A. J. Levan et al. 2016), and field galaxies from the CANDELS survey (A. Galametz et al. 2013). We find that while the host of GRB 250702B is inferred to be massive compared to most field galaxies at a similar redshift and to both relativistic TDE hosts, its inferred stellar mass is consistent with that of the most massive long GRB hosts. In the right panel of Figure 7 we find that the inferred stellar mass and extinction are consistent with the most massive moderately obscured long GRB hosts (S. Savaglio et al. 2009; D. A. Perley et al. 2013) and with the most massive of the eight nonoptically selected “dark” GRBs discussed in T. Krühler et al. (2011). The inferred total stellar mass, $\log(M_*/M_\odot) = 10.66 \pm 0.15$, is similar to the $\log(M_*/M_\odot) \approx 10.9$ host mass of all three known off-nuclear TDEs (D. Lin et al. 2018, 2020; C. C. Jin et al. 2025; Y. Yao et al. 2025). We caution that these comparisons are limited by the small populations of both optically dark GRBs and relativistic TDEs for which there are measured host properties.

In Section 3.3 we derived the host’s concentration ($C = 2.56^{+0.05}_{-0.06}$) and asymmetry ($A = 0.90^{+0.05}_{-0.05}$) and determined that the galaxy exhibits morphological features consistent with a major merger. Using the CAS parameter space, we compared the host of GRB 250702B with the long GRB sample from J. D. Lyman et al. (2017), which only yields hosts with $A < 0.4$. The morphology of GRB 250702B’s host is a significant outlier compared to the limited sample of morphologically characterized GRB hosts. The morphology of GRB 250702B’s host also differs significantly from the host of the relativistic TDE Sw J1644+57 ($A \approx 0.1$; A. J. Levan et al. 2016). However, the relativistic TDE Sw J1112.2–8238 occurred in a highly asymmetric and possibly merging galaxy (G. C. Brown et al. 2017). If confirmed, this merger scenario would be consistent with a TDE origin, as galaxy mergers are known to both enhance TDE rates (H. Pfister et al. 2021) and produce populations of wandering massive black holes (M. Tremmel et al. 2018; A. Ricarte et al. 2021).

4.6. Summary of the Possible Progenitors

The gamma-ray duration of GRB 250702B suggests a similar origin to previously studied ultralong GRBs. While the core




Figure 6. Left: the projected galactocentric offset of GRB 250702B compared to offset TDEs (Y. Yao et al. 2025) and the cumulative distribution for long GRBs (P. K. Blanchard et al. 2016). Right: same as the left panel, but for the host-normalized offset (r/r_{50}). The offset TDE candidates 3XMM J2150 (D. Lin et al. 2018, 2020), EP 240222a (C. C. Jin et al. 2025), and AT 2024tvd (Y. Yao et al. 2025) are believed to be associated with an IMBH.




Figure 7. Left: Stellar mass vs. redshift for long GRB host galaxies (blue circles; S. Savaglio et al. 2009; D. A. Perley et al. 2013) and field galaxies from the CANDELS survey (gray circles; A. Galametz et al. 2013). Also shown are the two relativistic TDEs (orange squares): Sw J1644+57 (Y. Yoon et al. 2015; A. J. Levan et al. 2016) and Sw J1112.2–8238 (G. C. Brown et al. 2015, 2017). GRB 250702B is shown as a red star. Right: stellar mass vs. the average host extinction inferred from galaxy SED modeling (D. A. Perley et al. 2013).

collapse of a massive star could explain a ~ 25 ks (E. Neights et al. 2025a) prompt emission, the observed long-lived, multiday X-ray emission is incompatible with the standard collapsar models (B. O’Connor et al. 2025). The off-center location of GRB 250702B is naturally consistent with stellar-mass black hole or IMBH engines from several proposed scenarios: collapsars, compact object–helium star mergers (E. Neights et al. 2025a), micro-TDEs (P. Beniamini et al. 2025)

or IMBH-TDEs (J. Granot et al. 2025; A. J. Levan et al. 2025). An association with a supernova would be expected in collapsars and some versions of micro-TDE origins involving a natal-kick-induced TDE after stellar collapse in one component of a field binary star system. An IMBH-TDE would also potentially require a supernova association. In this scenario, in the classic “frozen-in” approximation (J. H. Lacy et al. 1982; M. J. Rees 1988) energy dissipation

takes place primarily at the tidal radius,²² and the duration of fallback from the most inner-bound orbit is $t_{\text{fb}} \sim 10^6 (M_{\text{BH}}/10^4 M_{\odot})^{1/2} R_*^{3/2} M_*^{-1} \text{ s}$. As a result, matching the duration with that of GRB 250702B requires a smaller physical size of the disrupted star, which can be realized if the star is a white dwarf. In this situation, the compression of a white dwarf during the TDE process is expected to be sufficient to trigger a thermonuclear supernova (S. Rosswog et al. 2009). Helium star mergers are also expected to have associated supernovae, though significantly dimmer than those associated with collapsars (E. Neights et al. 2025a). Finally, the absence of a supernova would support a dynamically induced micro-TDE. At the time of writing, the large redshift and significant dust extinction in GRB 250702B preclude a strong conclusion regarding potential supernova association.

5. Conclusion

We presented deep optical and NIR observations of the high-energy transient GRB 250702B—the longest-duration GRB ever observed—whose extreme multiwavelength properties cannot be placed firmly among known transient classes (B. P. Gompertz et al. 2025; A. J. Levan et al. 2025; E. Neights et al. 2025a; B. O’Connor et al. 2025). Deep observations from a suite of optical/NIR facilities—including Blanco, Gemini, Wendelstein, Keck, and Magellan telescopes—combined with additional photometry from A. J. Levan et al. (2025), revealed a highly obscured, rapidly fading transient detectable only in the NIR. Our late-time observations constrain a luminous rebrightening ~ 20 days after the GRB in the rest frame, but deeper observations are needed to rule out rebrightening akin to the relativistic TDE AT 2022cmc’s slow thermal component (I. Andreoni et al. 2022) or SN 1998bw (F. Patat et al. 2001) given the high redshift and large extinction in the line of sight.

Using a redshift of $z = 1.036$ obtained from JWST observations (B. P. Gompertz et al. 2025), we modeled the decline of the NIR transient and found it to be consistent with forward shock emission originating from a nonspreading top-hat jet with a Lorentz factor of $\log \Gamma_0 = 1.25^{+0.16}_{-0.12}$. After correcting for Galactic extinction, we estimated the intrinsic line-of-sight extinction to be $A_{V,\text{GRB}} = 5.67^{+0.3}_{-0.33} \text{ mag}$ at $z = 1.036$. While this high extinction is consistent with both the most obscured GRBs (T. Krühler et al. 2011) and TDEs (D. N. Burrows et al. 2011), the Lorentz factor measurement remains model dependent: the value derived both without lateral spreading and with lateral spreading ($\Gamma_0 \approx 18$) aligns with relativistic TDEs (B. A. Zauderer et al. 2011; B. D. Metzger et al. 2012; I. Andreoni et al. 2022), while enforcing additional constraints ($\Gamma_0 \theta_c > 1$) on a nonspreading jet yields a value ($\Gamma_0 \approx 400$) more typical of GRB collapsars (E.-W. Liang et al. 2010; G. Ghirlanda et al. 2018).

Keck, Gemini, and HST images enabled the detection of the host galaxy in the z , H , K_s , and F160W filters. GRB 250702B’s galactocentric offset is consistent with both the population of observed long GRBs and the three known off-nuclear TDEs. Using the host galaxy SED modeling tools within Prospector, we found the galaxy to be massive ($10^{10.66 \pm 0.15} M_{\odot}$), dusty

($A_{V,\text{host}} = 0.88^{+0.47}_{-0.45} \text{ mag}$), and highly asymmetric ($A = 0.90$ in CAS space). The host’s inferred stellar mass is consistent with the masses of long GRB hosts and reminiscent of the $\sim 10^{10.9} M_{\odot}$ host masses observed in all three known off-nuclear TDEs (D. Lin et al. 2018, 2020; C. C. Jin et al. 2025; Y. Yao et al. 2025). The differential between the extinction of the NIR counterpart and the host suggests that GRB 250702B originated in a dense circumstellar environment or was obscured by a prominent dust cloud within the host. The galaxy’s high asymmetry makes it an outlier among long GRB hosts and suggests the possibility of a major merger that could support the TDE hypothesis.

Multiple scenarios remain viable to explain the nature of GRB 250702B. The jet modeling highlights the NIR emission accompanying GRB 250702B as consistent with both collapsar and TDE scenarios. Our data are also consistent with exotic models that could explain the origin of GRB 250702B, such as micro-TDEs (H. B. Perets et al. 2016; P. Beniamini et al. 2025), TDEs from IMBHs (J. Granot et al. 2025), and helium star merger (C. L. Fryer & S. E. Woosley 1998; E. Neights et al. 2025a). The multiwavelength signatures of the longest-duration GRB 250702B make it a unique puzzle that may originate from a novel progenitor.

Acknowledgments

We thank the anonymous referee for their thoughtful feedback and suggestions that strengthened this work.

We thank Brian Lemaux, staff scientist at Gemini North, for his exceptional assistance throughout this project. His expertise, dedication, and willingness to provide rapid advice across often inconvenient time zones were invaluable to the success of this work. We also thank Chris Simpson for his timely assistance with edge case GMOS reductions. J.C., I.A., and B.O. acknowledge useful discussions with Eliza Neights and Eric Burns and thank Ben Gompertz, Andrew Levan, and Antonio Martin-Carrillo for sharing the redshift derived by JWST.

The Andreoni Transient Astronomy Lab is supported by the National Science Foundation award AST 2505775, NASA grant 24-ADAP24-0159, and the Discovery Alliance Catalyst Fellowship Mentors award 2025-62192-CM-19. B.O. is supported by the McWilliams Postdoctoral Fellowship in the McWilliams Center for Cosmology and Astrophysics at Carnegie Mellon University. M.B. is supported by a Student grant from the Wübben Stiftung Wissenschaft. W.Z. acknowledges the support from the Harvard-Smithsonian Center for Astrophysics through the CfA Fellowship. E.R.C. acknowledges support from NASA through the Astrophysics Theory Program, grant 80NSSC24K0897. P.B.’s work was funded by a grant (No. 2020747) from the United States–Israel Binational Science Foundation (BSF), Jerusalem, Israel, and by a grant (No. 1649/23) from the Israel Science Foundation. P.C. acknowledges the support from the Zhejiang provincial top-level research support program. C.D.K. gratefully acknowledges support from the NSF through AST-2432037, from the HST Guest Observer Program through HST-SNAP-17070 and HST-GO-17706, and from JWST Archival Research through JWST-AR-6241 and JWST-AR-5441. A.P. is supported by NSF grant No. 2308193.

This Letter includes data gathered with the 6.5 m Magellan Telescopes located at Las Campanas Observatory, Chile. This Letter contains data obtained at the Wendelstein Observatory

²² If, instead, energy dissipation is dominated at the pericenter (for comparison of this with the “frozen-in” assumption, see E. Steinberg et al. 2019) and the penetration factor β is rather large ($\beta \gtrsim 5$), then the fallback time shortens significantly (as β^{-3} ; see P. Beniamini et al. 2025) and disruption of a main-sequence star by an IMBH becomes possible (albeit geometrically fine-tuned).

of the Ludwig-Maximilians University Munich. Funded in part by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy —EXC-2094—390783311. This work is based on observations made with the NASA/ESA Hubble Space Telescope. The data were obtained from the Mikulski Archive for Space Telescopes at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-03127 for JWST. This research is based on data obtained from the Astro Data Archive at NSF NOIRLab. NOIRLab is managed by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the US National Science Foundation. This project used data obtained with the Dark Energy Camera (DECam), which was constructed by the Dark Energy Survey (DES) collaboration. Funding for the DES Projects has been provided by the US Department of Energy, the US National Science Foundation, the Ministry of Science and Education of Spain, the Science and Technology Facilities Council of the United Kingdom, the Higher Education Funding Council for England, the National Center for Supercomputing Applications at the University of Illinois at Urbana-Champaign, the Kavli Institute of Cosmological Physics at the University of Chicago, the Center for Cosmology and Astro-Particle Physics at The Ohio State University, the Mitchell Institute for Fundamental Physics and Astronomy at Texas A&M University, Financiadora de Estudos e Projetos, Fundacao Carlos Chagas Filho de Amparo, Financiadora de Estudos e Projetos, Fundacao Carlos Chagas Filho de Amparo a Pesquisa do Estado do Rio de Janeiro, Conselho Nacional de Desenvolvimento Científico e Tecnológico and the Ministerio da Ciencia, Tecnologia e Inovacao, the Deutsche Forschungsgemeinschaft, and the Collaborating Institutions in the Dark Energy Survey.

Some of the data presented herein were obtained at the W. M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California, and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation. The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Maunakea has always had within the indigenous Hawaiian community. Based on observations at Cerro Tololo Inter-American Observatory, NSF’s NOIRLab (NOIRLab Prop. ID 2025A-729671; PI: Palmese), which is managed by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the National Science Foundation. Based on observations obtained at the international Gemini Observatory, a program of NSF’s OIR Lab, which is managed by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the National Science Foundation on behalf of the Gemini Observatory partnership: the National Science Foundation (United States), National Research Council (Canada), Agencia Nacional de Investigación y Desarrollo (Chile), Ministerio de Ciencia, Tecnología e Innovación (Argentina), Ministério da Ciência, Tecnologia, Inovações e Comunicações (Brazil), and Korea Astronomy and Space Science Institute (Republic of Korea). The data were acquired through the Gemini Observatory Archive at NSF NOIRLab and processed

using Data Reduction for Astronomy from Gemini Observatory North and South (DRAGONS).

The Collaborating Institutions are Argonne National Laboratory, the University of California at Santa Cruz, the University of Cambridge, Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas-Madrid, the University of Chicago, University College London, the DES-Brazil Consortium, the University of Edinburgh, the Eidgenössische Technische Hochschule (ETH) Zurich, Fermi National Accelerator Laboratory, the University of Illinois at Urbana-Champaign, the Institut de Ciències de l’Espai (IEEC/CSIC), the Institut de Física d’Altes Energies, Lawrence Berkeley National Laboratory, the Ludwig Maximilians Universität München and the associated Excellence Cluster Universe, the University of Michigan, NSF NOIRLab, the University of Nottingham, The Ohio State University, the University of Pennsylvania, the University of Portsmouth, SLAC National Accelerator Laboratory, Stanford University, the University of Sussex, and Texas A&M University.

The HST data presented in this Letter were obtained from the Mikulski Archive for Space Telescopes (MAST) at the Space Telescope Science Institute. The specific observations analyzed can be accessed via DOI: [10.17909/12kv-9c88](https://doi.org/10.17909/12kv-9c88).

Software: Astropy (Astropy Collaboration et al. 2013, 2018, 2022), matplotlib (J. D. Hunter 2007), SWarp (E. Bertin 2025), Source Extractor (E. Bertin & S. Arnouts 1996), Morfometryka (F. Ferrari et al. 2015), GALFIT (C. Y. Peng et al. 2002), DRAGONS (K. Labrie et al. 2019), Photutils (P. Bradley et al. 2024), VegasAfterglow (Y. Wang et al. 2025), astrometry.net (D. Lang et al. 2010), dynesty (J. S. Speagle 2020), DrizzlePac, emcee (D. Foreman-Mackey et al. 2013), bilby (G. Ashton et al. 2019), Claude (Anthropic 2024).

Appendix A Observation Details

A.1. DECam

We observed the location of GRB 250702B with the Dark Energy Camera (DECam; B. Flaugher et al. 2015) mounted on the Victor M. Blanco 4 m Telescope at Cerro Tololo Inter-American Observatory (program ID 2025A-729671; PI: Palmese), beginning approximately 15.3 hr after the first GBM detection. Observations spanned from 2025 July 3 05:12:58 UT to 2025 July 3 05:27:50 UT and consisted of exposures in four filters: 1×70 s in g band, 1×80 s in r band, 1×90 s in i band, and 2×100 s in z band. The images were reduced using the DESDM pipeline V5.5.3 (E. Morganson et al. 2018) with astrometric and photometric calibration against the United States Naval Observatory B catalog (D. G. Monet et al. 2003). No optical counterpart or host galaxy was detected in any filter, with 5σ limiting magnitudes of $g = 23.56$, $r = 23.04$, $i = 22.26$, and $z = 22.21$, 22.08 AB mag for the two z -band exposures, respectively.

A.2. Fraunhofer Telescope at Wendelstein Observatory

We observed the location of GRB 250702B with FTW/3KK (F. Lang-Bardl et al. 2016) for two epochs. The first epoch began at 2025 July 3 21:52:47 UT (~ 32 hr after the first GBM detection) and simultaneously imaged in the r , i , and J bands for 40×180 s. No sources were detected within the Swift/XRT

localization down to 5σ limiting magnitudes of $r = 22.47$, $i = 22.94$, $J = 21.25$ AB mag.

A second epoch was obtained at 2025 July 5 10:14:00 UT (~ 68 hr after the first GBM detection) with simultaneous imaging in the r , i , and K bands for 101×60 s. Using later imaging from Magellan (Section A.6) as a template for image subtraction, we detect the NIR counterpart at $K = 21.07 \pm 0.16$ AB mag. No detections were made in the optical bands down to 5σ limiting magnitudes of $r = 22.06$ and $i = 21.79$ mag. For both epochs we use the Pan-STARRS DR1 (PS1) catalog (K. C. Chambers et al. 2016) to photometrically calibrate r and i bands, and we use the Two Micron All Sky Survey catalog (M. F. Skrutskie et al. 2006) to photometrically calibrate J and K . We applied standard Vega-to-AB conversions (M. R. Blanton & S. Roweis 2007) to convert the calibrated J and K magnitudes to the AB system.

A.3. Gemini Telescopes

A.3.1. Gemini South

We observed the location of GRB 250702B in z band using GMOS-S (program ID GS-2025A-Q-117; PI. Andreoni) in imaging mode beginning at 2025 July 4 T03:43:54 UT (~ 38 hr after the first GBM detection). We obtained 120×60 s exposures for a total integration time of 2.0 hr. The images were reduced using the DRAGONS pipeline (K. Labrie et al. 2023), and photometric calibrations were obtained with the PS1 catalog (K. C. Chambers et al. 2016). No transient counterpart was detected to a 5σ limiting magnitude of $z = 23.95$ AB mag.

A.3.2. Gemini North

We observed the location of GRB 250702B in z band using the Gemini Multi Object Spectrograph on GMOS-N (program ID GN-2025A-Q-213; PI. Andreoni) in imaging mode beginning at 2025 July 20 T09:55:45 UT (~ 18 days after the first GBM detection). We obtained 52×120 s exposures for a total integration time of 1.73 hr. We did not coadd with the first epoch owing to differences in readout mode. The images were reduced using the DRAGONS pipeline (K. Labrie et al. 2023), and photometric calibrations were obtained with the PS1 catalog. No counterpart was detected to a 5σ limiting magnitude of $z = 24.19$ AB mag. We detected extended emission from the host galaxy using a $2''.25 \times 1''.0$ elliptical aperture, measuring $z = 23.20 \pm 0.13$ mag (Figure 1).

A.4. Keck

We observed the counterpart to GRB 250702B with MOSFIRE (I. S. McLean et al. 2012) on the W. M. Keck Observatory telescope I (PI: Kasliwal; program ID C348), obtaining J -, H -, and K_s -band images of the afterglow. Observations began at 2025 July 5 10:05:00 UT (~ 68 hr after the first GBM detection) and consisted of two box-9 dither sequences: 8 s exposures with four coadds in K_s , 4 s exposures with eight coadds in H , and one box-9 dither of 11 s exposures with three coadds in J . The data were reduced using standard procedures, and photometric calibration was performed against the VISTA Hemisphere Survey (VHS; R. G. McMahon et al. 2013) DR4.1 catalog for K_s and J bands and photometrically calibrated using the UKIRT Infrared Deep Sky Survey

(A. Lawrence et al. 2007) catalog in H band. We clearly detect the host galaxy in the K_s and H bands: $K_s = 20.50 \pm 0.07$ AB mag (using a $2''.88 \times 1''.08$ aperture) and $H = 21.55 \pm 0.17$ AB mag (using a $2''.26 \times 0''.9$ aperture). We perform image subtraction using K_s images taken ~ 38 days later with the Magellan telescope (Section A.6) and measure the transient to be $K_s = 21.96$ AB mag.

A.5. NEWFIRM

We observed the location of GRB 250702B at two epochs with the recommissioned NEWFIRM (R. G. Autry et al. 2003) on the Victor M. Blanco 4 m Telescope at Cerro Tololo Inter-American Observatory. The first epoch began at 2025 July 10 19:26:50 UT (~ 8 days after the first GBM detection) and consisted of 51×60 s exposures in H band. No transient emission was detected down to a 5σ limiting magnitude of $H = 20.50$ AB mag. The second epoch began the following night at 2025 July 11 21:33:30 UT (~ 9 days after the first GBM detection) with an additional 32×60 s exposures in H band, followed by 120×60 s exposures in K_s band beginning at 2025 July 11 21:50:51 UT. All data were processed using the photpipe imaging and photometry pipeline (A. Rest et al. 2005). Due to poor seeing conditions and high air mass during both nights, the combined H -band dataset achieved only a 5σ limiting magnitude of $H = 20.10$ AB mag. The K_s -band observations reached $K_s = 20.62$ AB mag (5σ). No transient was detected in either band, even after image subtraction with a later K_s epoch (Section A.6). We detected the host galaxy in K_s band at $K_s = 21.34 \pm 0.48$ AB mag using a $2''.4 \times 1''.2$ elliptical aperture. Photometric calibration was performed using VHS DR4.1 (R. G. McMahon et al. 2013) for K_s band and UKIDSS (A. Lawrence et al. 2007) for H band owing to limited VHS H -band coverage in this field.

A.6. Magellan

We obtained one epoch of K_s -band imaging with the NIR imager FourStar (S. E. Persson et al. 2013) mounted on the 6.5 m Magellan Baade Telescope. The observations were conducted at 2025 August 12 06:00:00 UT (~ 41 days post-burst) by executing a sequence of exposures at 12 random dithered positions, with each position consisting of 12 loops of 5.822 s integration. The total exposure time on target was 1397.28 s. The FourStar images were reduced using custom Python scripts following the standard procedures: dark subtraction, flat-fielding, sky background subtraction, astrometric registration of the science images using *astrometry.net* (D. Lang et al. 2010), and image stacking with *SWarp* (E. Bertin et al. 2002). No transient emission was detected down to a 5σ limiting magnitude of $K_s = 21.94$ AB mag. The host galaxy was detected at $K_s = 21.32 \pm 0.21$ AB mag using a $2''.4 \times 1''.12$ elliptical aperture (Figure 1). This template was used as a reference for image subtraction for the K_s images discussed in Sections A.2, A.4, and A.5.

A.7. Hubble Space Telescope

We reprocessed publicly available HST imaging obtained from the Mikulski Archive for Space Telescopes (MAST).²³ The HST data (program ID 17988; PI: Levan) were previously published in A. J. Levan et al. (2025). The images were

²³ <https://archive.stsci.edu/index.html>

obtained with the Wide Field Camera 3 (WFC3) in the F160W filter at 2025 July 15 02:39:23 (~ 12.5 days after the first GBM detection). The data were reduced using standard procedures within the `DrizzlePac` software package. We reprocessed the data to a final pixel scale of $0''.06 \text{ pixel}^{-1}$ using `pixfrac`=0.8. Using a $1''.75 \times 0''.65$ elliptical aperture, we measure a host magnitude with transient contamination of $F160W = 21.50 \pm 0.1$ mag (Figure 1), where the error bars have been increased to include systematics. In Section 3.3 we model the host morphology either as a spiral galaxy or as two galaxies undergoing a merger. We subtracted both of these models from the reprocessed image and performed photometry on the residuals. Using a $0''.30$ -diameter circular aperture, we measure the transient magnitude to be

$F160W = 26.52 \pm 0.1$ AB mag in the residual of the spiral galaxy model and $F160W = 26.75 \pm 0.1$ AB mag in the residual of the merger model, again with our error bars increased to include systematics. The residual flux after subtraction using the galaxy merger model is more PSF-like, so we employ a magnitude of $F160W = 26.52 \pm 0.1$ AB mag in our model fitting.

Appendix B Host Photometry

Host photometry presented in this work is displayed in Table B1.

Table B1
Host Galaxy Photometry of GRB 250702B

Datetime	Instrument	Filter	Host Mag	Aperture
2025-07-05 10:14:00	Keck/MOSFIRE	K_s	20.50 ± 0.07	$2''.88 \times 1''.08$
2025-07-11 21:50:51	NEWFIRM	K_s	21.34 ± 0.48	$2''.4 \times 1''.2$
2025-08-12 06:00:00	Magellan/FourStar	K_s	21.32 ± 0.21	$2''.4 \times 1''.12$
2025-07-05 10:14:00	Keck/MOSFIRE	H	21.55 ± 0.17	$2''.26 \times 0''.9$
2025-07-15 02:39:23	HST/WFC3	F160W	21.50 ± 0.1	$1''.75 \times 0''.65$
2025-07-20 09:55:45	Gemini/GMOS-N	z	23.20 ± 0.13	$2''.25 \times 1''.0$

Note. All magnitudes are provided in the AB system and are not corrected for Galactic extinction. The aperture column denotes the semimajor $\times$ semiminor axis of the elliptical aperture used for forced photometry.

Appendix C

Additional Model Fitting Results

The results of our jet modeling are displayed in Figures C1, C2, and C3. The results of our host galaxy SED fitting are displayed in Figure C4.




Figure C1. Posterior distributions of the jet parameters derived using VegasAfterglow to fit the available multiwavelength data with a nonspreading top-hat jet.




Figure C2. Posterior distributions of the jet parameters derived using *VegasAfterglow* to fit the available multiwavelength data with the inclusion of lateral jet spreading effects.




Figure C3. Posterior distributions of the jet parameters derived using VegasAfterglow to fit the available multiwavelength with a nonspreading top-hat jet constrained such that $\Gamma_0\theta_c > 1$.




Figure C4. Posterior distributions of our host galaxy SED fitting results using *prospector*. We note that M_{gal} refers to the total mass formed and not the galaxy's stellar mass. See also Figure 4.

ORCID iDs

Jonathan Carney <https://orcid.org/0000-0001-8544-584X>
 Igor Andreoni <https://orcid.org/0000-0002-8977-1498>
 Brendan O'Connor <https://orcid.org/0000-0002-9700-0036>
 James Freeburn <https://orcid.org/0009-0006-7990-0547>
 Hannah Skobe <https://orcid.org/0000-0003-0516-3485>
 Lewi Westcott <https://orcid.org/0009-0008-8642-5275>
 Malte Busmann <https://orcid.org/0009-0001-0574-2332>
 Antonella Palmese <https://orcid.org/0000-0002-6011-0530>
 Xander J. Hall <https://orcid.org/0000-0002-9364-5419>
 Ramandeep Gill <https://orcid.org/0000-0003-0516-2968>

Paz Beniamini <https://orcid.org/0000-0001-7833-1043>
 Eric R. Coughlin <https://orcid.org/0000-0003-3765-6401>
 Charles D. Kilpatrick <https://orcid.org/0000-0002-5740-7747>
 Akash Anumalapudi <https://orcid.org/0000-0002-8935-9882>
 Nicholas M. Law <https://orcid.org/0000-0001-9380-6457>
 Hank Corbett <https://orcid.org/0000-0002-6339-6706>
 Tomas Ahumada <https://orcid.org/0000-0002-2184-6430>
 Ping Chen <https://orcid.org/0000-0003-0853-6427>
 Christopher Conselice <https://orcid.org/0000-0003-1949-7638>

Guillermo Damke  <https://orcid.org/0000-0002-2651-7038>
 Kaustav K. Das  <https://orcid.org/0000-0001-8372-997X>
 Avishay Gal-Yam  <https://orcid.org/0000-0002-3653-5598>
 Daniel Gruen  <https://orcid.org/0000-0003-3270-7644>
 Steve Heathcote  <https://orcid.org/0000-0002-0856-3663>
 Lei Hu  <https://orcid.org/0000-0001-7201-1938>
 Viraj Karambelkar  <https://orcid.org/0000-0003-2758-159X>
 Mansi Kasliwal  <https://orcid.org/0000-0002-5619-4938>
 Kathleen Labrie  <https://orcid.org/0000-0002-6633-7891>
 Dheeraj Pasham  <https://orcid.org/0000-0003-1386-7861>
 Arno Riffeser  <https://orcid.org/0000-0002-5466-3892>
 Michael Schmidt  <https://orcid.org/0009-0003-1323-9774>
 Kritti Sharma  <https://orcid.org/0000-0002-4477-3625>
 Weicheng Zang  <https://orcid.org/0000-0001-6000-3463>

References

- Achterberg, A., Gallant, Y. A., Kirk, J. G., & Guthmann, A. W. 2001, *MNRAS*, **328**, 393
- Andreoni, I., Coughlin, M. W., Perley, D. A., et al. 2022, *Natur*, **612**, 430
- Anthropic, 2024 Claude 4 Sonnet: AI Assistant for Research and Analysis, <https://claude.ai>
- Ashton, G., Hübner, M., Lasky, P. D., et al. 2019, *ApJS*, **241**, 27
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, *ApJ*, **935**, 167
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, **156**, 123
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, **558**, A33
- Autry, R. G., Probst, R. G., Starr, B. M., et al. 2003, *SPiE*, **4841**, 525
- Becerra, R. L., Dornic, D., Pereyra, M., et al. 2025, GCN, **40918**, 1
- Beniamini, P., Perets, H. B., & Granot, J. 2025, arXiv:2509.22779
- Bershady, M. A., Jangren, A., & Conselice, C. J. 2000, *AJ*, **119**, 2645
- Bertin, E., 2025 SWarp: Resampling and Co-adding FITS Images Together, Astrophysics Source Code Library, ascl:1010.068
- Bertin, E., & Arnouts, S. 1996, *A&AS*, **117**, 393
- Bertin, E., Mellier, Y., Radovich, M., et al. 2002, *ASPC*, **281**, 228
- Blanchard, P. K., Berger, E., & Fong, W.-f. 2016, *ApJ*, **817**, 144
- Blanton, M. R., & Roweis, S. 2007, *AJ*, **133**, 734
- Bloom, J. S., Giannios, D., Metzger, B. D., et al. 2011, *Sci*, **333**, 203
- Bradley, P., Sipőcz, B., Robitaille, T., et al. 2024, astropy/photutils: v2.0.2, Zenodo, doi:10.5281/zenodo.13989456
- Brown, G. C., Levan, A. J., Stanway, E. R., et al. 2015, *MNRAS*, **452**, 4297
- Brown, G. C., Levan, A. J., Stanway, E. R., et al. 2017, *MNRAS*, **472**, 4469
- Burrows, D. N., Kennea, J. A., Ghisellini, G., et al. 2011, *Natur*, **476**, 421
- Calzetti, D., Armus, L., Bohlin, R. C., et al. 2000, *ApJ*, **533**, 682
- Cenko, S. B., Frail, D. A., Harrison, F. A., et al. 2010, *ApJ*, **711**, 641
- Cenko, S. B., Krimm, H. A., Horesh, A., et al. 2012, *ApJ*, **753**, 77
- Chabrier, G. 2003, *PASP*, **115**, 763
- Chambers, K. C., Magnier, E. A., Metcalfe, N., et al. 2016, arXiv:1612.05560
- Cheng, H. Q., Zhao, G. Y., Zhou, C., et al. 2025, GCN, **40906**, 1
- Chevalier, R. A., & Li, Z.-Y. 2000, *ApJ*, **536**, 195
- Conroy, C., Gunn, J. E., & White, M. 2009, *ApJ*, **699**, 486
- Conselice, C. J. 2003, *ApJS*, **147**, 1
- Conselice, C. J., Bershady, M. A., & Jangren, A. 2000, *ApJ*, **529**, 886
- Curran, P. A., Evans, P. A., de Pasquale, M., Page, M. J., & van der Horst, A. J. 2010, *ApJL*, **716**, L135
- Dai, Z. G., & Lu, T. 1998, *MNRAS*, **298**, 87
- DeLaunay, J., Ronchini, S., Tohuavohu, A., et al. 2025, GCN, **40903**, 1
- Dobie, D., Kaplan, D. L., Hotokezaka, K., et al. 2020, *MNRAS*, **494**, 2449
- Ellison, D. C., & Double, G. P. 2002, *APh*, **18**, 213
- Fermi GBM Team 2025a, GCN, **40883**, 1
- Fermi GBM Team 2025b, GCN, **40885**, 1
- Fermi GBM Team 2025c, GCN, **40886**, 1
- Fermi GBM Team 2025d, GCN, **40890**, 1
- Ferrari, F., de Carvalho, R. R., & Trevisan, M. 2015, *ApJ*, **814**, 55
- Fitzpatrick, E. L., & Massa, D. 2007, *ApJ*, **663**, 320
- Flaugher, B., Diehl, H. T., Honscheid, K., et al. 2015, *AJ*, **150**, 150
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, **125**, 306
- Frederiks, D., Lysenko, A., Ridnaia, A., et al. 2025, GCN, **40914**, 1
- Fryer, C. L., & Woosley, S. E. 1998, *ApJL*, **502**, L9
- Galama, T. J., Vreeswijk, P. M., van Paradijs, J., et al. 1998, *Natur*, **395**, 670
- Galamez, A., Grazian, A., Fontana, A., et al. 2013, *ApJS*, **206**, 10
- Gehrels, N., Chincarini, G., Giommi, P., et al. 2004, *ApJ*, **611**, 1005
- Gendre, B., Stratta, G., Atteia, J. L., et al. 2013, *ApJ*, **766**, 30
- Ghisellini, G., Ghirlanda, G., Nava, L., & Celotti, A. 2010, *MNRAS*, **403**, 926
- Ghirlanda, G., Ghisellini, G., Nava, L., et al. 2012, *MNRAS*, **422**, 2553
- Ghirlanda, G., Nappo, F., Ghisellini, G., et al. 2018, *A&A*, **609**, A112
- Gompertz, B. P., Levan, A. J., Laskar, T., et al. 2025, arXiv:2509.22778
- Granot, J., Gill, R., Beniamini, P., Perets, H. B., & O'Connor, B. 2025, Submitted
- Granot, J., & Sari, R. 2002, *ApJ*, **568**, 820
- Ho, A. Y. Q., Perley, D. A., Yao, Y., et al. 2022, *ApJ*, **938**, 85
- Hua, Y.-L., Geng, J.-J., Wu, X.-F., et al. 2025, GCN, **40943**, 1
- Hunter, J. D. 2007, *CSE*, **9**, 90
- Jin, C. C., Li, D. Y., Jiang, N., et al. 2025, arXiv:2501.09580
- Johnson, B. D., Leja, J., Conroy, C., & Speagle, J. S. 2021, *ApJS*, **254**, 22
- Johnson, B. D., Leja, J. L., Conroy, C., & Speagle, J. S., 2019 Prospector: Stellar Population Inference from Spectra and SEDs, Astrophysics Source Code Library, ascl:1905.025
- Kasliwal, M. M. 2011, PhD thesis, California Institute of Technology
- Kawakubo, Y., Serino, M., Negro, H., et al. 2025, GCN, **40910**, 1
- Kennea, J. A., Siegel, M. H., Evans, P. A., et al. 2025, GCN, **40919**, 1
- Kouveliotou, C., Meegan, C. A., Fishman, G. J., et al. 1993, *ApJL*, **413**, L101
- Krühler, T., Greiner, J., Schady, P., et al. 2011, *A&A*, **534**, A108
- Kumar, A., Gompertz, B. P., Ramsay, G., et al. 2025, GCN, **40908**, 1
- Labrie, K., Anderson, K., Cárdenes, R., Simpson, C., & Turner, J. E. H. 2019, *ASPC*, **523**, 321
- Labrie, K., Simpson, C., Cardenes, R., et al. 2023, *RNAAS*, **7**, 214
- Lacy, J. H., Townes, C. H., & Hollenbach, D. J. 1982, *ApJ*, **262**, 120
- Lang, D., Hogg, D. W., Mierle, K., Blanton, M., & Roweis, S. 2010, *AJ*, **139**, 1782
- Lang-Bardl, F., Bender, R., Goessl, C., et al. 2016, *SPiE*, **9908**, 990844
- Lawrence, A., Warren, S. J., Almaini, O., et al. 2007, *MNRAS*, **379**, 1599
- Leja, J., Johnson, B. D., Conroy, C., van Dokkum, P. G., & Byler, N. 2017, *ApJ*, **837**, 170
- Levan, A. J., Martin-Carrillo, A., Laskar, T., et al. 2025, *ApJL*, **990**, L28
- Levan, A. J., Tanvir, N. R., Brown, G. C., et al. 2016, *ApJ*, **819**, 51
- Levan, A. J., Tanvir, N. R., Cenko, S. B., et al. 2011, *Sci*, **333**, 199
- Levan, A. J., Tanvir, N. R., Starling, R. L. C., et al. 2014, *ApJ*, **781**, 13
- Li, H. L., Xin, L. P., Qiu, Y. L., et al. 2025a, GCN, **41122**, 1
- Li, X., Chen, C., Dong, Z.-n., et al. 2025b, GCN, **40986**, 1
- Liang, E.-W., Yi, S.-X., Zhang, J., et al. 2010, *ApJ*, **725**, 2209
- Lin, D., Strader, J., Carrasco, E. R., et al. 2018, *NatAs*, **2**, 656
- Lin, D., Strader, J., Romanowsky, A. J., et al. 2020, *ApJL*, **892**, L25
- Lyman, J. D., Levan, A. J., Tanvir, N. R., et al. 2017, *MNRAS*, **467**, 1795
- McLean, I. S., Steidel, C. C., Epps, H. W., et al. 2012, *SPiE*, **8446**, 84460J
- McMahon, R. G., Banerji, M., Gonzalez, E., et al. 2013, *Msngr*, **154**, 35
- Metzger, B. D., Giannios, D., & Mimica, P. 2012, *MNRAS*, **420**, 3528
- Molinari, E., Vergani, S. D., Malesani, D., et al. 2007, *A&A*, **469**, L13
- Monet, D. G., Levine, S. E., Canzian, B., et al. 2003, *AJ*, **125**, 984
- Morganson, E., Gruendl, R. A., Menanteau, F., et al. 2018, *PASP*, **130**, 074501
- Neights, E., Burns, E., Fryer, C. L., et al. 2025a, *MNRAS*, Accepted
- Neights, E., Roberts, O. J., Burns, E., et al. 2025b, GCN, **40931**, 1
- Neights, E., Roberts, O. J., Burns, E., Veres, P. & Fermi-GBM Team 2025c, GCN, **40891**, 1
- O'Connor, B., Beniamini, P., & Gill, R. 2024, *MNRAS*, **533**, 1629–1648
- O'Connor, B., Beniamini, P., & Kouveliotou, C. 2020, *MNRAS*, **495**, 4782
- O'Connor, B., Gill, R., DeLaunay, J., et al. 2025, *ApJL*, **994**, L17
- O'Connor, B., Troja, E., Dichiara, S., et al. 2022, *MNRAS*, **515**, 4890
- O'Connor, B., Troja, E., Ryan, G., et al. 2023, *SciA*, **9**, eadi1405
- Pasham, D. R., Cenko, S. B., Levan, A. J., et al. 2015, *ApJ*, **805**, 68
- Pasham, D. R., Lucchini, M., Laskar, T., et al. 2023, *NatAs*, **7**, 88
- Patat, F., Cappellaro, E., Danziger, J., et al. 2001, *ApJ*, **555**, 900
- Peng, C. Y., Ho, L. C., Impey, C. D., & Rix, H.-W. 2002, *AJ*, **124**, 266
- Perets, H. B., Li, Z., Lombardi, J. C., Jr., & Milcarek, S. R., Jr. 2016, *ApJ*, **823**, 113
- Pérez-García, I., Castro-Tirado, A. J., Caballero-García, M. D., et al. 2025, GCN, **40929**, 1
- Perley, D. A., Levan, A. J., Tanvir, N. R., et al. 2013, *ApJ*, **778**, 128
- Persson, S. E., Murphy, D. C., Smee, S., et al. 2013, *PASP*, **125**, 654
- Pfister, H., Dai, J. L., Volonteri, M., et al. 2021, *MNRAS*, **500**, 3944
- Piran, T. 2004, *RvMP*, **76**, 1143
- Planck Collaboration, Aghanim, N., Akrami, Y., et al. 2020, *A&A*, **641**, A6
- Polzin, A. 2025, *JOSS*, **10**, 8200
- Rees, M. J. 1988, *Natur*, **333**, 523

- Rest, A., Stubbs, C., Becker, A. C., et al. 2005, [ApJ](#), **634**, 1103
- Rhoads, J. E. 1997, [ApJL](#), **487**, L1
- Rhoads, J. E. 1999, [ApJ](#), **525**, 737
- Rhodes, L., Bright, J. S., Fender, R., et al. 2023, [MNRAS](#), **521**, 389
- Ricarte, A., Tremmel, M., Natarajan, P., Zimmer, C., & Quinn, T. 2021, [MNRAS](#), **503**, 6098
- Rosswog, S., Ramirez-Ruiz, E., & Hix, W. R. 2009, [ApJ](#), **695**, 404
- Sari, R., & Piran, T. 1999, [ApJ](#), **520**, 641
- Sari, R., Piran, T., & Halpern, J. P. 1999, [ApJL](#), **519**, L17
- Sari, R., Piran, T., & Narayan, R. 1998, [ApJL](#), **497**, L17
- Savaglio, S., Glazebrook, K., & Le Borgne, D. 2009, [ApJ](#), **691**, 182
- Schlafly, E. F., & Finkbeiner, D. P. 2011, [ApJ](#), **737**, 103
- Siegel, M. H. & Swift/UVOT Team 2025, GCN, **40952**, 1
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, [AJ](#), **131**, 1163
- Speagle, J. S. 2020, [MNRAS](#), **493**, 3132
- Spitkovsky, A. 2008, [ApJL](#), **682**, L5
- Steinberg, E., Coughlin, E. R., Stone, N. C., & Metzger, B. D. 2019, [MNRAS](#), **485**, L146
- Stratta, G., Gendre, B., Atteia, J. L., et al. 2013, [ApJ](#), **779**, 66
- Svom/Grm Team, Wang, C.-W., Zheng, S.-J., et al. 2025, GCN, **40923**, 1
- Thöne, C. C., de Ugarte Postigo, A., Fryer, C. L., et al. 2011, [Natur](#), **480**, 72
- Tikhomirova, Y. Y., & Stern, B. E. 2005, [AstL](#), **31**, 291
- Tremmel, M., Governato, F., Volonteri, M., Pontzen, A., & Quinn, T. R. 2018, [ApJL](#), **857**, L22
- Wang, X.-G., Zhang, B., Liang, E.-W., et al. 2015, [ApJS](#), **219**, 9
- Wang, X.-G., Zhang, B., Liang, E.-W., et al. 2018, [ApJ](#), **859**, 160
- Wang, Y., Chen, C., & Zhang, B. 2025, [JHEAp](#), **50**, 100490
- Woosley, S. E., & Bloom, J. S. 2006, [ARA&A](#), **44**, 507
- Yao, Y., Chornock, R., Ward, C., et al. 2025, [ApJL](#), **985**, L48
- Yoon, Y., Im, M., Jeon, Y., et al. 2015, [ApJ](#), **808**, 96
- Yuan, W., Dai, L., Feng, H., et al. 2025, [SCPMA](#), **68**, 239501
- Zauderer, B. A., Berger, E., Soderberg, A. M., et al. 2011, [Natur](#), **476**, 425