




arXiv is now an independent nonprofit! [Learn more](#)

arxiv





[Search](#) [Submit](#) [Donate](#) [Log in](#)

Showing 51–59 of 59 results

Search v0.5.6 released 2020-02-24

Query: order: -announced_date_first; size: 50; page_start: 50;
include_cross_list: True; terms: AND all=0.139

[Simple Search](#)

[Refine query](#)

[New search](#)

51. [arXiv:1205.0580](#) [[pdf](#), [ps](#), [other](#)] [hep-ph](#) [hep-ex](#) [doi](#) [10.1103/PhysRevD.86.014014](#)

SU(3) symmetry breaking and CP violation in $D \rightarrow PP$ decays

Authors: [Hai-Yang Cheng](#), [Cheng-Wei Chiang](#)

Abstract: ...is about $-(0.139 \pm 0.004)$ and $-(0.151 \pm 0.004)$ for the two solutions of W -exchange amplitudes, respectively. We also find that the CP asymmetry of $D^0 \rightarrow K^0 \nu K^0$ dominated by the interference between W -exchange amplitudes ranges from -0.62×10^{-3} to -1.82×10^{-3} . We study phenomenological implications of two new physics sc... [▼ More](#)

Submitted 23 June, 2012; v1 submitted 2 May, 2012; originally announced May 2012.

Comments: 23 pages, three new paragraphs added in the beginning of Sec. III. Version to appear in PRD. arXiv admin note: substantial text overlap with arXiv:1201.0785

52. [arXiv:1204.0754](#) [[pdf](#), [ps](#), [other](#)] [nucl-ex](#) [hep-ex](#) [doi](#) [10.1103/PhysRevC.86.024909](#)

Nuclear-Modification Factor for Open-Heavy-Flavor Production at Forward Rapidity in Cu+Cu Collisions at $\sqrt{s_{NN}}=200$ GeV

Authors: [A. Adare](#), [S. Afanasiev](#), [C. Aidala](#), [N. N. Ajitanand](#), [Y. Akiba](#), [H. Al-Bataineh](#), [J. Alexander](#), [K. Aoki](#), [L. Aphecetche](#), [R. Armendariz](#), [S. H. Aronson](#), [J. Asai](#), [E. T. Atomssa](#), [R. Averbeck](#), [T. C. Awes](#), [B. Azmoun](#), [V. Babintsev](#), [G. Baksay](#), [L. Baksay](#), [A. Baldisseri](#), [K. N. Barish](#), [P. D. Barnes](#), [B. Bassalleck](#), [S. Bathe](#), [S. Batsouli](#), et al. (351 additional authors not shown)

Abstract: ...Results: The charm-production cross section in p+p collisions at $\sqrt{s}=200$ GeV, integrated over p_T and in the rapidity range $1.4 < y < 1.9$ is found to be $d\sigma_{c\bar{c}}/dy = 0.139 \pm 0.029$ (stat) $^{+0.051}_{-0.058}$ (syst) mb. This result is consistent with calculations and with expectations based on the corresponding midrapidity charm-production cross s... [▼ More](#)

Submitted 3 April, 2012; originally announced April 2012.

Comments: 376 authors, 18 pages, 14 figures, 5 tables. Submitted to Phys. Rev. C. Plain text data tables for the points plotted in figures for this and previous PHENIX publications are publicly available at http://www.phenix.bnl.gov/phenix/WWW/info/data/ppg123_data.html

Journal ref: Phys. Rev. C 86, 024909 (2012)

53. [arXiv:0912.3760](#) [[pdf](#), [other](#)] [cond-mat.other](#) [doi](#) [10.1103/PhysRevB.81.064516](#)

Bound states of edge dislocations: The quantum dipole problem in two dimensions

Authors: [K. Dasbiswas](#), [D. Goswami](#), [C. -D. Yoo](#), [Alan T. Dorsey](#)

Abstract: ...of the ground state energy. We then numerically solve the eigenvalue problem and calculate the energy spectrum. In our dimensionless units, we find a ground state energy of -0.139 , which is lower than any previous estimate. We also make successful contact with the behavior of the energy spectrum as derived from semiclassical considerations. [▼ More](#)

Submitted 18 December, 2009; originally announced December 2009.

Comments: 6 pages, 3 figures, submitted to PRB

Journal ref: Phys. Rev. B 81, 064516 (2010)

54. [arXiv:0807.4750](#) [pdf, ps, other] astro-ph doi 10.1086/592134

TRIS II: search for CMB spectral distortions at 0.60, 0.82 and 2.5 GHz

Authors: [M. Gervasi](#), [M. Zannoni](#), [A. Tartari](#), [G. Boella](#), [G. Sironi](#)

Abstract: ...GHz; $T_{cmb}^{th} = (2.516 \pm 0.139 \pm 0.284) K$ at $\nu = 2.5$ GHz. The first error bar is statistic (1σ) while the second one is systematic. These results represent a significant improvement with respect to the previous measurements. We have also set new limits to the free-free distortions, $-6.3 \times 10^{-6} < Y_{ff} < 12.6 \times 10^{-6}$, and sl... [▼ More](#)

Submitted 29 July, 2008; originally announced July 2008.

Comments: accepted for publication in The Astrophysical Journal

55. [arXiv:astro-ph/0209205](#) [pdf, ps, other] astro-ph doi 10.1086/367556

Chandra Observations of Abell 2029: The Dark Matter Profile Down to $<0.01 R_{vir}$ in an Unusually Relaxed Cluster

Authors: [Aaron D. Lewis](#), [David A. Buote](#), [John T. Stocke](#)

Abstract: ... > 100 beyond 200 kpc. The consistency with an NFW halo model and the large M/L suggest the cluster is DM-dominated down to $<\sim 0.005 r_{vir}$. We observe a gas mass fraction = 0.139 ± 0.004 at the limit of our observations, finding an upper limit to $\Omega_m \leq 0.29 \pm 0.03$ (abridged). [▼ More](#)

Submitted 6 December, 2002; v1 submitted 11 September, 2002; originally announced September 2002.

Comments: ApJ Accepted Version; to appear in 2003 March 20 Issue; 8 pages, emulateapj5, 3 figures (2 color)

Journal ref: Astrophys.J.586:135-142,2003

56. [arXiv:astro-ph/0105481](#) [pdf, ps, other] astro-ph doi 10.1046/j.1365-8711.2001.04655.x

Theoretical insights into the RR Lyrae K-band Period-Luminosity relation

Authors: [G. Bono](#), [F. Caputo](#), [V. Castellani](#), [M. Marconi](#)

Abstract: ...predictions for metal-deficient ($0.0001 < Z < 0.006$) horizontal branch models, we eventually find that the infrared Period-Luminosity- Metallicity (PLZK) relation is $M_K = 0.139 - 2.071(\log P + 0.30) + 0.167 \log Z$, with a total intrinsic dispersion of 0.037 mag. As a consequence, the use of such a PLZK relation should constrain within ± 0.04 mag the infrared distanc... [▼ More](#)

Submitted 28 May, 2001; originally announced May 2001.

Comments: 8 pages, including 5 postscript figures, accepted for publication on MNRAS

Journal ref: Mon.Not.Roy.Astron.Soc. 326 (2001) 1183

57. [arXiv:cond-mat/9805073](#) [pdf, ps, other] cond-mat.dis-nn q-bio

doi 10.1088/0305-4470/31/43/005

Phase Diagram and Storage Capacity of Sequence Processing Neural Networks

Authors: [A. During](#), [A. C. C. Coolen](#), [D. Sherrington](#)

Abstract: ..., compared to $\alpha \sim 0.139$ for Hopfield networks storing static patterns. Our results are tested against extensive computer simulations and excellent agreement is found. [▼ More](#)

Submitted 6 May, 1998; originally announced May 1998.

Comments: 17 pages Latex2e, 2 postscript figures

58. [arXiv:hep-lat/9211009](#) [[pdf](#), [ps](#), [other](#)] [hep-lat](#) [doi](#) [10.1016/0920-5632\(93\)90344-6](#)

The confined-deconfined interface tension, wetting, and the spectrum of the transfer matrix

Authors: [B. Grossmann](#), [M. L. Laursen](#), [T. Trappenberg](#), [U. -J. Wiese](#)

Abstract: ...we find $\sigma_{cd} = 0.139(4)T_c^2$ for $L_t = 2$. The interfaces show universal behavior because the deconfined-deconfined interfaces are completely wet by the confined phase. The critical exponents of complete wetting follow from the analytic interface solutions of a $\mathbb{Z}(3)$ -symmetric Φ^4 model in three dimensions. We find numerical evidence that the conf... [▽ More](#)

Submitted 4 November, 1992; **originally announced** November 1992.

Comments: Talk presented at the International Conference on Lattice Field Theory, Lattice 92, to be published in the proceedings, 4 pages, 4 figures, figures 2,3,4 appended as postscript files, figure 1 not available as a postscript file but identical with figure 2 of Nucl. Phys. B372 (1992) 703, special style file espcrc2.sty required (available from hep-lat), BUTP-92/47

59. [arXiv:hep-lat/9208020](#) [[pdf](#), [ps](#), [other](#)] [hep-lat](#) [doi](#) [10.1016/0550-3213\(93\)90664-B](#)

The confined-deconfined Interface Tension and the Spectrum of the Transfer Matrix

Authors: [B. Grossmann](#), [M. L. Laursen](#), [T. Trappenberg](#), [U. -J. Wiese](#)

Abstract: ...yields $\sigma_{cd} = 0.139(4)T_c^2$ for $L_t = 2$. We found numerical evidence that the deconfined-deconfined domain walls are completely wet by the confined phase, and that the confined-deconfined interfaces are rough. [▽ More](#)

Submitted 21 August, 1992; **originally announced** August 1992.

Comments: 22 pages, LaTeX file with 4 ps figures included, HLRZ 92-47, BUTP-92/34

Journal ref: Nucl.Phys. B396 (1993) 584-610

[Previous](#)

[1](#)

[2](#)

We gratefully acknowledge support from our **major funders**, **member institutions**, and all contributors.

[About](#) · [Help](#) · [Contact](#) · [Subscribe](#) · [Copyright](#) · [Privacy](#) · [Accessibility](#) · [Operational Status](#)

Major funding support from

SIM NS
FOUNDATION



**SIMONS
FOUNDATION
INTERNATIONAL**


Schmidt Sciences