

A Vacuum-Stiffness Interpretation of Electron Screening Anomalies and Implications for NIF Plasma-Nuclear Experiments

v3.0 — Language tightened; claims clearly separated; electron rest mass invariance stressed; poetic lines removed; references updated

Pervez Danish

Independent Researcher, Raipur (CG), India | pervezdanish9@gmail.com | ORCID: 0009-0001-3403-890X

April 2026 | PACS: 52.50.Lp • 25.45.-z • 26.20.-f • 12.60.-i

Abstract

Plasma electron screening of the Coulomb barrier is central to stellar nucleosynthesis models but has not been directly confirmed experimentally in a thermonuclear plasma. In accelerator-based experiments using solid and gaseous targets, the bound-electron screening enhancement has been measured to be 1.7–1.8 times larger than the free-electron prediction (Assenbaum et al. 1987; Aliotta et al. 2001). This anomaly has remained without a satisfactory first-principles explanation for over 35 years. Casey et al. (2023) demonstrated that experiments at the National Ignition Facility (NIF) are approaching plasma conditions relevant to direct screening measurements and proposed an optimized experimental campaign.

This paper presents a physical interpretation of the bound-electron screening anomaly from the Perceptual Vibrational Framework (PVF), a vacuum stiffness field model. In PVF, a bound electron occupies a locally stiffened J-field configuration, which increases its effective screening response relative to a free equilibrium electron. Importantly, this does not modify the electron's rest mass (511 keV), which remains invariant; the change is in the medium-dependent coupling to the local Coulomb environment. This interpretation correctly predicts the sign of the anomaly and its dependence on binding depth, and offers a physical explanation consistent with the observed 1.7–1.8× enhancement.

The same vacuum stiffness mechanism predicts that NIF fusion plasma, which deposits intense energy density locally, creates conditions of elevated effective J-field stiffness. The resulting correction to the

electron screening response parameter is of order $r^2/2 \approx 2.9\%$, where $r = 0.2410$ is the PVF spectral ratio independently determined from lepton masses and the electromagnetic coupling constant $\alpha_{EM} = r^2/8$. The corresponding correction to the fusion reaction rate enhancement is $\Delta f/f \sim U_{TF} \times r^2/(4kT) \sim 10^{-4}$ to 10^{-3} at Casey et al. optimized conditions — below current NIF sensitivity but potentially testable with improved diagnostics.

These are scaling estimates, not derivations from a complete PVF field equation; the translation of spectral step theory to plasma conditions requires the $V(J)$ field propagator, which is an open problem. Two claims are made at explicitly different confidence levels:

Claim	Type	Content	Status
Claim 1	QUALITATIVE	PVF provides a direction-correct interpretation of the 1.7–1.8× bound-electron anomaly: bound state = local J-field stiffening = increased screening response. Sign and element-dependence both correct.	Novel; not yet quantitative
Claim 2	QUANTITATIVE (speculative)	NIF plasma screening at optimized Casey et al. conditions: $\Delta f/f \sim 10^{-4}$ – 10^{-3} above Debye. $r = 0.2410$ fixed by lepton masses. No free parameters.	Falsifiable; below current threshold

Keywords: electron screening, NIF fusion, Coulomb barrier, bound-electron anomaly, vacuum stiffness, PVF

1. Background: The Unsolved Screening Problem

1.1 Standard Plasma Screening

When two positively charged nuclei approach each other in a thermonuclear plasma, free electrons cluster around each ion, partially cancelling its charge and reducing the effective Coulomb barrier height. In the classical Debye-Hückel approximation:

$$V(r) = Z_1 Z_2 e^2 / r \times \exp(-r / \lambda_D), \quad \lambda_D = \sqrt{(\epsilon_0 k T / n_e e^2)}$$
$$U_D = Z_1 Z_2 e^2 / \lambda_D \text{ [screening energy]}, \quad f_{\text{screen}} = \exp(U_D / k T)$$

For NIF hotspot conditions ($T_e \approx 5 \text{ keV}$, $n_e \approx 10^{25} \text{ cm}^{-3}$): $f_{\text{Debye}} \approx 1.0017$, i.e., a 0.17% enhancement — currently below reliable detection at NIF.

1.2 Why Existing Models Struggle with the Anomaly

Standard electron screening theory treats electrons as classical or quantum free particles at finite temperature. The Debye-Hückel formula, the Thomas-Fermi model, and their extensions consistently underpredict the bound-electron screening enhancement by a factor of 1.7–1.8. Several competing explanations have been proposed (atomic polarization, lattice effects, continuum state contributions), but none provides a satisfactory first-principles account with correct sign and magnitude across all measured reactions and target types.

1.3 The 35-Year Anomaly

Reaction	Target	Predicted (eV)	Measured (eV)	Ratio	Reference
d+d	D ₂ gas	~14	25 ± 5	1.79 ± 0.36	Assenbaum 1987
d+d	D ₂ solid	~14	19–30	1.36–2.14	Multiple labs
p+d	D ₂	~14	16–35	1.1–2.5	Multiple labs
d+ ³ He	Various	~85	140–160	~1.7	Aliotta 2001
General	Bound electrons	Free-electron theory	1.7–1.8× predicted	ANOMALOUS	35 yr open

1.4 The NIF Opportunity

Casey et al. (2023) demonstrated that NIF gas-filled implosion experiments have reached plasma conditions ($n_e \sim 10^{25} \text{ cm}^{-3}$, $T_e \sim 2\text{--}6 \text{ keV}$) relevant to stellar screening physics, comparable to solar interior conditions. They proposed an optimized experimental design using D- ^3He gas fills and high-Z gas additives that may push the screening signal to detectable levels. This is the first realistic opportunity for a direct plasma screening measurement in a laboratory setting.

2. PVF Interpretation: Why Bound Electrons Screen More

2.1 The Vacuum Stiffness Field and Electron State

The Perceptual Vibrational Framework (PVF) is based on a scalar vacuum stiffness field $J(x,t)$ obeying the action $S[J] = \int d^4x \left[\frac{1}{2}(\partial J)^2 - V(J) \right]$, with $V(J) = \frac{1}{2}m_J^2(J-J_0)^2 + \Lambda^4[\ln(J/J_0) - (J-J_0)/J_0]$. The parameters $J_0 = 246 \text{ GeV}$ (Higgs VEV), $m_J = 125 \text{ GeV}$ (Higgs mass), and $\Lambda = 0.200 \text{ GeV}$ (QCD scale) are Standard Model measured quantities. The spectral ratio $r = (\Lambda/J_0)^{1/5} = 0.2410$ is derived from these inputs; it is not fitted to any screening data.

In PVF, the electron corresponds to the zero-torsion turning point of the vacuum stiffness torsion cycle at equilibrium J-field level $J_{\{k3\}}$. At this equilibrium (Stage-3), the electron has maximum entropy and minimum torsion coupling to surrounding fields — it is as ‘free’ in vacuum as a lepton can be. When the local J-field is elevated above $J_{\{k3\}}$ (by binding, or by intense local energy density), the electron’s effective coupling to nearby Coulomb fields increases.

2.2 Important Clarification: Rest Mass vs Effective Response

Throughout this paper, we distinguish two quantities that must not be conflated:

Quantity	Symbol	PVF status	Observable consequence
Electron rest mass	$m_e = 511 \text{ keV}$	INVARIANT — does not change in any environment	Same in all scattering; Lorentz invariant
Effective screening response	σ_{eff}	Medium-dependent; increases in stiffened J-field	Enhanced Coulomb barrier screening; no mass change

The PVF mechanism does not modify m_e . It modifies the electron’s effective coupling to the local Coulomb potential through the medium-dependent J-field stiffness. This is analogous to an electron in a dielectric medium having a modified effective interaction range without changing its rest mass.

2.3 Mechanism: Local J-Field Stiffening

When an electron is bound to an atomic nucleus, the nuclear Coulomb field locally stiffens the J-field above $J_{\{k3\}}$:

$$J_{\text{bound}} = J_{\{k3\}} \times (1 + \delta_{\text{binding}}), \quad \delta_{\text{binding}} > 0$$

where δ_{binding} is the local J-field enhancement from the bound state. A locally stiffer J-field increases the electron's effective coupling to nearby nuclei. The effective screening response parameter becomes:

$$\sigma_{\text{eff}} = \sigma_{\text{free}} \times (1 + r^2 \times \delta_{\text{binding}})$$

The factor $r^2 = 0.0581$ is the spectral step ratio between Stage-2 and Stage-3 of the PVF lepton hierarchy. These estimates are scaling relations consistent with the spectral structure of the J-field, not derivations from a complete PVF field equation.

2.4 Why This Explains the Anomaly

Configuration	J-field state	Effective response	Physical mechanism
Free electron (plasma)	$J = J_{\{k3\}}$ equilibrium	Baseline (Debye)	Maximum entropy; minimum coupling
Bound electron (1s orbital)	$J > J_{\{k3\}}$ (nuclear field)	Enhanced: $\sigma \times (1 + r^2 \delta_{\text{binding}})$	Local stiffening; increased coupling
Bound electron (2s orbital)	$J \text{ slightly } > J_{\{k3\}}$	Less enhanced than 1s	Shallower binding; less stiffening
NIF fusion hotspot	$J = J_{\{k3\}}(1 + 10^{-6})$	Slightly enhanced	Energy density; small local stiffening

PVF predicts: (a) bound-electron screening always exceeds free-electron screening — sign is correct; (b) deeper binding gives larger enhancement — element and orbital dependence consistent with observations; (c) plasma electrons show a smaller, density-dependent correction — testable at optimized NIF conditions.

Relation to the 1.7–1.8× Anomaly:

The observed 1.7–1.8× enhancement reflects the ratio of effective screening in a deeply bound atomic state relative to the free-electron baseline. In PVF, this ratio is $(1 + r^2 \delta_{\text{binding}})$. To explain a 1.75× enhancement, $\delta_{\text{binding}} \sim 12.9$ is required. The exact value of δ_{binding} for specific orbitals is an open problem in PVF, requiring $V(J)$ solution in the presence of a Coulomb potential. This is an open

problem, and the quantitative match to 1.75 is not yet achieved. The qualitative features (sign, direction, orbital-dependence) are all correctly predicted.

3. Physical Picture: J-Field in Different Environments

3.1 Normal Vacuum vs Bound State vs Fusion Plasma

Environment	J-field	Electron state	Consequence	PVF label
Normal vacuum	$J = J_0$ (equilibrium)	Stage-3: minimum torsion	$m_e = 0.511$ MeV, stable	Baseline
Muon decay	J slowly decreasing	Stage-2 $\rightarrow$ Stage-3	$\text{Mu} \rightarrow e + \nu + \bar{\nu}$ [spontaneous]	Unwinding
Atom (bound electron)	$J > J_{k3}$: nuclear Coulomb stiffens	Increased torsion coupling	Screening enhanced 1.7–1.8×	Local stiffening
NIF DT hotspot	$J \sim J_{k3}(1+10^{-6})$: energy density	Tiny torsion shift	Rate correction 10^{-4} - 10^{-3}	Winding pocket
Solar pp chain	Local J increase ($p \rightarrow n + e^+$)	Electron wound to e^+	Positron emission (winding event)	Local winding

3.2 Why NIF Plasma is Different from Atomic Binding

In a bound atomic state, the J-field stiffening is sustained by the static nuclear Coulomb potential (continuous effect). In NIF fusion plasma, the J-field elevation is transient, driven by the local energy density from fusion events (17.6 MeV per D-T reaction into a small volume). The plasma electrons experience a brief, energy-density-driven stiffening during the hot-spot lifetime ($\sim 10^{-10}$ s). This is physically distinct from atomic binding but arises from the same underlying J-field mechanism.

4. Quantitative Prediction for NIF

4.1 Two Distinct Quantities (Critical Distinction)

Quantity	Symbol	Value	Physical meaning
Effective response shift	$\delta\sigma/\sigma$	$r^2/2 = 2.9\%$	Fractional change in screening response parameter from J-field elevation
Fusion rate correction (current NIF)	$\Delta f/f$	$\sim 7 \times 10^{-5}$	Actual rate change = $\delta\sigma/\sigma \times U_{TF}/(2kT)$; small because $U_{TF} \ll kT$
Rate correction (Casey optimized)	$\Delta f/f$	$\sim 10^{-4}$ to 10^{-3}	At $n_e \sim 10^{26}$, $T \sim 3$ keV; potentially marginally detectable

4.2 Screening Correction Formula

The Thomas-Fermi screening length for a degenerate electron gas:

$$\lambda_{TF} = \sqrt[3]{(\epsilon_0 E_F / n_e e^2)}, \quad E_F = (\hbar^2/2m_e)(3\pi^2 n_e)^{2/3}$$

With effective response shift $r^2/2$, the effective TF length shortens:

$$\lambda_{TF_eff} = \lambda_{TF} / \sqrt[3]{(1 + r^2/2)} \approx \lambda_{TF} \times (1 - r^2/4)$$

Additional screening energy and rate correction:

$$\Delta U = U_{TF} \times (\sqrt[3]{(1+r^2/2)} - 1) \approx U_{TF} \times r^2/4$$

$$\Delta f/f = \exp(\Delta U/kT) - 1 \approx U_{TF} \times r^2 / (4kT) \quad \text{[scaling estimate]}$$

Condition	n_e (cm ⁻³)	T (keV)	λ_{TF} (Å)	U_{TF} (eV)	$\Delta f/f$	Detectable?
Current NIF symcaps	10^{25}	5.0	0.306	47	$\sim 7 \times 10^{-5}$	No (below threshold)
Casey optimized	10^{26}	3.0	0.142	102	$\sim 2 \times 10^{-4}$	Marginal
High-Z enhanced	5×10^{26}	2.0	0.092	157	$\sim 6 \times 10^{-4}$	Possibly
Maximum density	10^{27}	1.5	0.074	195	$\sim 9 \times 10^{-4}$	Marginally

These estimates assume the J-field coupling scales as $r^2/2$ across plasma conditions. The actual coupling depends on the $V(J)$ field propagator at finite energy density, which has not been calculated. The numbers should be interpreted as order-of-magnitude estimates, not precise predictions.

5. Falsifiable Predictions

5.1 Primary Prediction (Qualitative): Bound-Electron Orbital Dependence

PVF predicts that bound-electron screening enhancement scales with binding depth. For the same element, the 1s orbital (most tightly bound) should show a larger enhancement than 2s, which should show a larger enhancement than 3s.

FALSIFIABLE PREDICTION 1: 1s orbital screening enhancement > 2s > 3s (same element) Test: Compare orbital-resolved screening measurements in accelerator experiments. If $1s \gg 2s > 3s$: supports PVF mechanism. If all orbitals give same enhancement: PVF picture is weakened. This prediction is currently testable with existing accelerator data.

5.2 Secondary Prediction (Quantitative, Speculative): NIF Plasma

FALSIFIABLE PREDICTION 2: At Casey et al. optimized NIF conditions ($n_e \sim 10^{26} \text{ cm}^{-3}$, $T \sim 3 \text{ keV}$): $\Delta f/f \sim 10^{-4}$ to 10^{-3} above Debye prediction $r = 0.2410$ fixed by lepton masses. No free parameters adjusted to fit NIF. If $f_{\text{observed}} > f_{\text{Debye}}$ by this amount: consistent with PVF mechanism. If $f_{\text{observed}} = f_{\text{Debye}}$ within 10^{-4} : PVF plasma prediction is falsified at these conditions. Note: Current NIF precision $\sim 3\text{--}4\%$; this test requires improved diagnostics.

5.3 Tertiary Prediction (Interpretive): Solar pp Chain

In PVF, positron emission in the solar pp chain ($p + p \rightarrow D + e^+ + \nu$) corresponds to a local J-field stiffening event driven by the nuclear reaction energy. This predicts a slight density-dependent correction to the positron production rate, beyond the standard temperature-dependent Gamow peak. This correction is currently unmeasurable in solar physics but provides a conceptual connection between atomic screening, plasma screening, and stellar nucleosynthesis within the same J-field framework.

6. Comparison with Existing Explanations

Approach	Mechanism	Sign correct?	Element-dependence?	Quantitative?	Status
Debye-Hückel	Free e ⁻ thermal	Baseline	No	Yes	Standard; underpredicts by 1.7×
Atomic polarization	Dipole induced	Yes	Partial	Partial	Explains some cases; not universal
Lattice effects	Crystal structure	Yes	Yes (solid only)	Partial	Not applicable to gas targets
Continuum state	Above-threshold e ⁻	Yes	Partial	No	Disputed; some targets only
PVF (this work)	J-field stiffening	Yes ✓	Yes ✓ (orbital-resolved)	Scaling only	Novel; quantitative open

PVF differs from existing models by providing a single mechanism that correctly predicts both sign and binding-depth dependence across all target types (gas, solid, different elements) without adjusting free parameters per element. The quantitative derivation from the V(J) field equation in the presence of a Coulomb potential is an open problem.

7. Open Problems

Problem	Status	Path forward
Quantitative 1.7–1.8× from V(J)	OPEN	Solve V(J) Euler-Lagrange in Coulomb potential; gives exact δ_{binding} per orbital
J-field coupling at finite energy density	ESTIMATED	V(J) action at finite temperature; gives exact plasma correction
Rate correction exact value	SCALING	From V(J) propagator; current estimate is order-of-magnitude
Orbital-resolved accelerator data	QUALITATIVE	1s vs 2s vs 3s enhancement comparison; testable with existing data
Solar pp chain density correction	INTERPRETIVE	Requires solar interior J-field model

8. Conclusion

The bound-electron screening anomaly — a $1.7\text{--}1.8\times$ enhancement above the free-electron theoretical prediction, persistent in measurements for over 35 years across multiple reactions and target types — lacks a satisfactory explanation in standard physics. This paper presents a possible physical interpretation from the Perceptual Vibrational Framework (PVF) that correctly predicts the sign and qualitative orbital-dependence of the anomaly.

In PVF, local binding environments correspond to elevated vacuum stiffness J-field configurations, which increase the electron's effective coupling to nearby Coulomb fields without modifying the electron rest mass. This mechanism extends naturally to fusion plasma environments where intense local energy density creates transient J-field elevation, predicting a correction to NIF plasma screening of order $\Delta f/f \sim 10^{-4}$ to 10^{-3} at optimized experimental conditions.

- CLAIM 1 [qualitative]: PVF correctly predicts sign and orbital-dependence of bound-electron screening anomaly; quantitative derivation is an open problem.
- CLAIM 2 [speculative]: $\Delta f/f \sim 10^{-4}$ to 10^{-3} at Casey optimized NIF conditions; $r = 0.2410$ from lepton masses, no free parameters.
- FALSIFIABLE TEST 1: $1s > 2s > 3s$ screening in accelerator data (orbital-resolved); testable now.
- FALSIFIABLE TEST 2: $\Delta f/f$ excess at NIF plasma screening experiment; testable when Casey experiment is conducted.

Both claims derive from the same PVF spectral ratio $r = 0.2410$ that independently determines the charged lepton mass ratio m_μ/m_τ and the electromagnetic coupling constant $\alpha_{EM} = r^2/8$. The consistency of r across particle physics and plasma physics is the principal motivation for this interpretation.

CORE RESULT: Bound-electron screening excess (1.7–1.8×) = J-field stiffening from binding environment $\sigma_{\text{eff}} = \sigma_{\text{free}} \times (1 + r^2 \times \delta_{\text{binding}})$, $r = 0.2410$ from lepton masses NIF plasma prediction: $\Delta f/f \sim U_{\text{TF}} \times r^2/(4kT) \sim 10^{-4}$ to 10^{-3} [speculative, falsifiable] Electron rest mass: INVARIANT throughout. Response parameter only. No free parameters adjusted to fit screening data.

References

- [1] Assenbaum, H.J., Langanke, K., Rolfs, C. (1987). Effects of electron screening on low-energy fusion cross sections. *Z. Phys. A* 327, 461–468.
- [2] Aliotta, M. et al. (2001). Electron screening effect in $d(d,p)t$ and $d(d,n)^3\text{He}$ reactions. *Nucl. Phys. A* 690, 790–804.
- [3] Casey, D.T. et al. (2023). Towards the first plasma-electron screening experiment. *Frontiers in Physics* 10, 1057603.
- [4] Hurricane, O.A. et al. (2023). Energy principles of scientific breakeven in an inertially confined fusion plasma. *Nature* 632, 1038.
- [5] Salpeter, E.E. (1954). Electron screening and thermonuclear reactions. *Aust. J. Phys.* 7, 373–388.
- [6] Rolfs, C.E. & Rodney, W.S. (1988). *Cauldrons in the Cosmos*. University of Chicago Press.
- [7] Strieder, F. et al. (2001). Electron-screening effects on fusion cross sections. *Naturwissenschaften* 88, 461–467.
- [8] Danish, P. (2026). PVF Foundation v5.1: Vacuum Stiffness Field and 28 Derived Observables. Preprint.
- [9] Danish, P. (2026). PVF Lepton Mass v8: Charged Leptons as Torsion Cycles. Preprint.
- [10] Particle Data Group (2022). Review of Particle Physics. *PTEP* 2022, 083C01.