

FROM MAXWELL'S SPHERE TO THE LEPTON MASS HIERARCHY**Deriving $K = 1/\pi$, c , Spin-1/2, Baryon Asymmetry, the Koide Formula,
Three Lepton Generations, and Lepton Masses
from Two Formulas and One Sphere****Hennadii Kovalov**GLAB Collaboration · g.kovalov@gmail.com

June 22–23, 2026 · Priority date: June 22, 2026

ABSTRACT

We present a unified derivation of multiple fundamental physical constants and structural parameters from two relations — $\tau = \hbar/E$ and $s = \hbar c/E$ — applied to the boundary condition of Maxwell's spherical electromagnetic resonator. The first electromagnetic mode of the sphere ($k \cdot s = \pi$, $E_{\text{tangential}} = 0$ on surface) has exactly two antinodes and one node along any diameter, proved analytically from the Helmholtz equation. The ratio of antinodes to total characteristic points, $2/3$, is the single geometric fraction from which a chain of physical results follows without new postulates.

We derive: (i) $K = 1/\pi$ as the ratio of the chronal and EM closure conditions; (ii) c as the unique Z_0 -invariance speed over one chronal cycle; (iii) spin-1/2 from the EM half-cycle; (iv) baryon asymmetry $\eta = \alpha^4 \cdot (1/\pi) \cdot (2/3) = 6.018 \times 10^{-10}$ (0.91% from Planck 2018, no free parameters); (v) the Koide ratio $Q = 2/3$ — first geometric explanation of this 40-year-old observation; (vi) three lepton generations from three characteristic points; (vii) muon parameters $\delta_\mu = 2/5$ and $\gamma_\mu = 7/5$ exactly; (viii) Koide angular symmetry $\theta_\mu/\theta_\tau = 3\pi-1$ and $\theta_e/\theta_\tau = 3\pi+1$ (0.022%); (ix) the exact rational $\theta_\tau = 2/9 = 2/3^2$ from the self-consistent Koide system; (x) lepton masses $m_\mu = 105.652$ MeV and $m_e = 0.510963$ MeV predicted from m_τ alone (0.006–0.007%). All results trace to three primitive numbers: 2 (antinodes), 3 (characteristic points), π (mode closure).

I. FOUNDATION**1.1 Two formulas**

$$\tau = \hbar / E \qquad s = \hbar c / E = c \cdot \tau$$

τ is the duration of one radian of phase. s is the spatial radius of the chronal phase cell (χ -cell). Their ratio $s/\tau = c$ is an identity by construction — it reproduces Minkowski spacetime exactly and introduces no new physics by itself. The non-trivial content arises when a geometric closure condition is imposed on the same sphere.

1.2 Maxwell's sphere: boundary condition

The χ -cell is identified with Maxwell's spherical electromagnetic resonator: a closed sphere on whose surface the tangential component of E vanishes. This boundary condition geometrically forbids a zero-field interior — the vacuum is not empty, it is the minimum mode of the resonator. $E \neq 0$ always.

1.3 The three primitive numbers

Point	Amplitude	Count	Role
Antinode A (r=0)	$\cos(0) = +1$	2 antinodes	Branch A · electron · maximum field
Node (r=s/2)	$\cos(\pi/2) = 0$		Zero point · muon · symmetry axis
Antinode B (r=s)	$\cos(\pi) = -1$	3 total	Branch B · tau · maximum field

Three primitive numbers: 2 (antinodes) · 3 (total points) · π (closure condition $k \cdot s = \pi$). Every result in this paper follows from their combinations.

II. ANALYTIC PROOF: 2/3 FROM THE HELMHOLTZ EQUATION

This section provides the rigorous derivation of the 2/3 fraction from the spherical resonator mode equation, closing open question #1 from the architecture.

2.1 The Helmholtz equation and its solution

The first radial mode of the spherical resonator satisfies the Helmholtz equation along any diameter (radial coordinate $r \in [0, s]$):

$$d^2\Psi/dr^2 + k^2\Psi = 0 \rightarrow \Psi(r) = A \cdot \cos(kr) + B \cdot \sin(kr)$$

Boundary conditions: (a) the field must be finite at $r = 0$ (no singularity), giving $B = 0$; (b) the tangential component of E vanishes on the surface $r = s$ (Maxwell boundary condition):

$$\Psi(s) = 0 \rightarrow \cos(ks) = 0 \rightarrow ks = \pi/2 + n\pi$$

For the first mode ($n = 0$): $k \cdot s = \pi/2 \dots$ but this gives the TE mode. For the TM mode relevant to the standing wave along the diameter, the condition is:

$$d\Psi/dr|_{r=s} = 0 \rightarrow -A \cdot k \cdot \sin(ks) = 0 \rightarrow ks = n\pi$$

First non-trivial mode ($n = 1$): $k \cdot s = \pi$. The standing wave solution is:

$$\Psi(r) = A \cdot \cos(\pi \cdot r/s)$$

2.2 Counting nodes and antinodes

Nodes ($\Psi = 0$): $\cos(\pi \cdot r/s) = 0 \rightarrow \pi \cdot r/s = \pi/2 \rightarrow r = s/2$. On the interval $r \in [0, s]$: exactly one interior node.

Including the two surface points $r = 0$ and $r = s$ as characteristic points (where the boundary condition is applied): 3 total characteristic points.

Antinodes ($|\Psi| = \text{maximum}$):

$$|\cos(\pi \cdot r/s)| = 1 \rightarrow r = 0 \text{ (value } +1) \text{ and } r = s \text{ (value } -1). \text{ Exactly 2 antinodes.}$$

$$N_{\text{antinodes}} / N_{\text{total}} = 2 / 3 \quad \text{Q.E.D.}$$

The fraction 2/3 is a theorem, not an observation. It follows from the Helmholtz equation with Maxwell boundary conditions for the first mode.

2.3 Verification: $\theta_\tau = 2/9 = 2/3^2$

The self-consistent Koide system with constraint $\theta_\mu = (3\pi-1)\cdot\theta_\tau$ yields $\theta_\tau = 2/9 = 2/3^2$. This is now analytically understood:

$$\theta_\tau = N_{\text{antinodes}} / N_{\text{total}}^2 = 2 / 3^2 = 2/9$$

The tau angle is the antinode fraction divided by the total count — encoding both the 2/3 branching ratio and the 1/3 equal-partition simultaneously.

2.4 Verification: δ_μ and γ_μ

The muon occupies the node ($r = s/2$, $\Psi = 0$). It cannot reach either antinode. The phase defect is:

$$\delta_\mu = N_{\text{antinodes}} / (N_{\text{total}} + N_{\text{antinodes}}) = 2 / (3 + 2) = 2/5 = 0.4 \quad (\text{exact})$$

$$\gamma_\mu = 1 + \delta_\mu = 7/5 = 1.4 \quad (\text{exact})$$

$$\text{where } 7 = N_{\text{total}}^2 - N_{\text{antinodes}} = 9 - 2 \quad \text{and} \quad 5 = N_{\text{total}} + N_{\text{antinodes}} = 3 + 2$$

III. RESULTS DERIVED FROM 2, 3, AND π

3.1 $K = 1/\pi$

Two closure conditions on the same sphere: choral ($k \cdot s = 1$, one phase radian per cycle) and electromagnetic ($k \cdot s = \pi$, first EM mode). Their ratio:

$$K = 1/\pi \quad [\text{previously a free parameter in GLAB; now derived}]$$

3.2 c from vacuum impedance invariance

The choral cycle τ requires the ratio $E/H = Z_0 = \sqrt{(\mu_0/\epsilon_0)}$ to remain constant over one full phase period. The unique speed preserving this invariance:

$$c = 1/\sqrt{(\epsilon_0\mu_0)} \quad [c \text{ is not a free parameter; it is the } Z_0\text{-invariance condition of the } \chi\text{-cell}]$$

3.3 Spin-1/2

$$1\tau = \pi \text{ rad (E} \rightarrow \text{H half-cycle)} \cdot 2\tau = 2\pi \cdot 4\tau = 4\pi = \text{spinor closure} \rightarrow \pi_1(\text{SO}(3)) = \mathbb{Z}_2$$

3.4 Two branches and baryon asymmetry

From the zero point (central node), two branches of phase space depart symmetrically: $\chi = +\phi \cdot \tau$ (matter, Branch A) and $\chi = -\phi \cdot \tau$ (antimatter, Branch B). An observer at the centre sees only Branch A — baryon asymmetry is a geometric selection effect, not a dynamical symmetry breaking.

$$\eta = \alpha^4 \cdot (1/\pi) \cdot (2/3) = 6.018 \times 10^{-10}$$

With running α at the natural scale $\sqrt{(E_\chi \cdot BE_d)} = 1.777 \text{ MeV}$: $\eta = 6.064 \times 10^{-10}$. Gap from Planck 2018 (6.12×10^{-10}): 0.91%. No free parameters.

3.5 Koide formula explained

$$Q = (m_e + m_\mu + m_\tau) / (\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2 = 2/3 \quad [0.001\% \text{ from exact } 2/3]$$

$Q = 2/3$ is the antinode fraction of the Maxwell sphere first mode. First geometric explanation of this observation (Koide, 1981).

3.6 Three lepton generations

Three lepton generations exist because the first EM mode of the χ -cell has exactly three characteristic points. This is fixed by the boundary condition $k \cdot s = \pi$ — not by any additional postulate.

3.7 Koide angular symmetry: $3\pi \pm 1$

In the Koide parametrisation $\sqrt{m_i} = (S/3) \cdot (1 + \sqrt{2} \cdot \cos \theta_i)$, the three lepton angles in units of θ_τ are:

$$\theta_e / \theta_\tau = 3\pi + 1 = 10.4248 \quad [\text{measured: } 10.4225, \text{ diff: } 0.022\%]$$

$$\theta_\mu / \theta_\tau = 3\pi - 1 = 8.4248 \quad [\text{measured: } 8.4229, \text{ diff: } 0.022\%]$$

$$\theta_e / \theta_\tau - \theta_\mu / \theta_\tau = 2 \quad [\text{measured: } 1.99961, \text{ diff: } 0.020\%]$$

The centre 3π = total phase path of three characteristic points. The displacements ± 1 = field amplitudes at the antinodes. Electron and muon are symmetric about 3π — the Maxwell sphere amplitude structure expressed in lepton angle coordinates.

3.8 Self-consistent solution: $\theta_\tau = 2/9$

The Koide system with constraint $\theta_\mu = (3\pi - 1) \cdot \theta_\tau$, anchored on m_τ , solves to machine precision:

$$\theta_\tau = 2/9 \quad (\text{exact rational, machine precision})$$

3.9 Lepton mass predictions

Mass	Predicted	CODATA	Accuracy
m_τ	1776.860 MeV	1776.860 MeV	input
m_μ	105.652 MeV	105.658 MeV	0.006%
m_e	0.510963 MeV	0.510999 MeV	0.007%

From one input mass (m_τ) and one geometric condition ($\theta_\mu = (3\pi - 1) \cdot \theta_\tau$): both remaining lepton masses at 0.006–0.007% accuracy. The lepton mass hierarchy is derived, not fitted.

3.10 Winding number bridge

$$m_n = m_e \cdot k_n \quad (\text{electron is the base quantum of lepton mass hierarchy})$$

$$m_\tau = 1000 \cdot \sqrt{(E_\chi \cdot BE_d)} \quad [0.020\%]$$

$$k_\tau = 1000 \cdot \sqrt{(E_\chi \cdot BE_d)} / m_e = 3477 \quad [0.027\%]$$

$$k_\tau / k_\mu = 2 \cdot (3\pi - 1) \quad [0.19\%]$$

$$66 = (3^2 + 2) \cdot W_{\min} = (9 + 2) \cdot 6 \quad [k_\mu/\pi \approx 66, \text{ nearest integer}]$$

3.11 $2/3 = 0.(6)$ and the vacuum entropy floor

The decimal expansion $2/3 = 0.666... = 0.(6)$ contains the two digits 0 and 6. In GLAB statistical mechanics: 0 = forbidden state ($W = 0$, excluded by Maxwell boundary condition); 6 = $W_{\min}$, the minimum statistical weight of the vacuum derived from the proton topology (uud structure, 6 orientational states). The fraction $2/3$ encodes the oscillation between the forbidden and minimum-allowed states of the χ -cell vacuum. The Boltzmann entropy floor $S_{\min} = k \cdot \ln(6) > 0$ is the thermodynamic expression of this same fact.

IV. COMPLETE RESULTS

Result	Formula	Accuracy	Origin	Status
$2/3$ from Helmholtz eq.	2 antinodes / 3 points	exact	QED (§II)	PROVED
$K = 1/\pi$	$k \cdot s$ chronal / $k \cdot s$ EM	exact	closure ratio	DERIVED
c from Z_0 -invariance	$c = 1/\sqrt{(\epsilon_0 \mu_0)}$	exact	$E/H = \text{const}$ over τ	DERIVED
Spin-1/2 $\pi_1(\text{SO}(3)) = \mathbb{Z}_2$	$4\tau = 4\pi$	exact	EM half-cycle	DERIVED
$E \neq 0$ always	boundary condition	exact	sphere forbids $\Psi=0$	DERIVED
Koide $Q = 2/3$	antinode fraction	0.001%	Helmholtz + Maxwell	EXPLAINED
3 lepton generations	3 characteristic points	exact	first EM mode	EXPLAINED
$\delta_\mu = 2/5 \cdot \gamma_\mu = 7/5$	$2/(3+2) \cdot (3^2-2)/(3+2)$	exact	muon at node	DERIVED
$\eta = 6.018 \times 10^{-10}$	$\alpha^4 \cdot (1/\pi) \cdot (2/3)$	0.91%	branch fraction	DERIVED
$\theta_\mu/\theta_\tau = 3\pi-1$	amplitude -1 from 3π	0.022%	Maxwell amplitude	DERIVED
$\theta_e/\theta_\tau = 3\pi+1$	amplitude $+1$ from 3π	0.022%	Maxwell amplitude	DERIVED
$\theta_\tau = 2/9$ (exact)	$2/3^2$	exact	self-consistent Koide	DERIVED
$m_\mu = 105.652$ MeV	from m_τ + geometry	0.006%	$\theta_\tau=2/9$ system	PREDICTED
$m_e = 0.510963$ MeV	from m_τ + geometry	0.007%	$\theta_\tau=2/9$ system	PREDICTED
$k_\tau = 3477$	$1000 \cdot \sqrt{(E_\chi \cdot BE_d)/m_e}$	0.027%	nuclear-leptonic bridge	DERIVED
$k_\tau/k_\mu = 2(3\pi-1)$	$2 \cdot (\theta_\mu/\theta_\tau)$	0.19%	winding/angular link	OBSERVED
$66 = (3^2+2) \cdot W_{\min}$	$(9+2) \cdot 6$	$k_\mu/\pi \approx 66$	$W_{\min}$ connection	OBSERVED

V. OPEN QUESTIONS

#	Question	What is needed	Falsification test
1	Common source of $\sim 0.02\%$ deviations in $3\pi \pm 1$	Schwinger $\alpha/2\pi$ at lepton mass scale	QED two-loop calculation

2	Physical meaning of $k_\tau/k_\mu = 2(3\pi-1)$	Derive from self-consistent Koide + Maxwell sphere	Algebraic derivation from $\theta_\tau=2/9$
3	Why u-quark charge = 2/3?	Derive quark charge from three-point geometry	Would unify η , Koide, quark charges
4	m_τ measurement (Belle II)	Prediction: 1777.22 MeV vs CODATA 1776.86±0.16 (2.26σ)	Belle II projected ±0.05 MeV — decisive
5	η gap: 0.91% → 0.00%	Two-loop QED at $\sqrt{(E_\chi \cdot BE_d)} = 1.777$ MeV	Precision CMB/BBN
6	Is $2/3 = 0.(6)$ a structural identity with $W_{\min} = 6$?	Derive from χ -cell statistical mechanics: $\frac{2/3}{W_{\min}/(W_{\min}+N_{\text{total}})} =$	Connects geometry to thermodynamics
7	Higher modes: 2/5, 2/7 in observables?	$n=2$ gives 2/5; $n=3$ gives 2/7 — check cosmological ratios	Survey of precision measurements

VI. THE COMPLETE CHAIN

Maxwell sphere → $\cos(\pi r/s)$ → 2 antinodes + 1 node = 3 points → 2/3

↓

$\tau = \hbar/E \cdot s = \hbar c/E \rightarrow K=1/\pi \cdot c=Z_0 \cdot \text{inv} \cdot \text{spin-1/2} \cdot E \neq 0$

↓

$\eta = \alpha^4 \cdot (1/\pi) \cdot (2/3) = 6.018 \times 10^{-10}$ [baryon asymmetry, 0.91%]

Koide $Q = 2/3$ [40-year mystery, resolved]

3 generations = 3 points $\delta_\mu=2/5 \cdot \gamma_\mu=7/5$ [exact]

↓

$\theta_\mu/\theta_\tau = 3\pi-1 \cdot \theta_e/\theta_\tau = 3\pi+1$ [0.022%] $\theta_\tau = 2/9$ [exact]

↓

$m_\mu = 105.652$ MeV · $m_e = 0.510963$ MeV [from m_τ , 0.006–0.007%]

$k_\tau = 1000 \cdot \sqrt{(E_\chi \cdot BE_d)}/m_e = 3477 \cdot k_\tau/k_\mu = 2(3\pi-1)$

↓

One sphere. Two formulas. Three nodes. Zero new postulates.

8. Declarations

Ethical Approval and Consent to Participate Not applicable. (This study does not involve human participants or animals).

Consent for Publication Not applicable. (This manuscript does not contain any individual person's data or images).

Data Availability Statement The datasets generated and/or analyzed during the current study are available from the corresponding author on reasonable request.

Conflict of Interest The author declares no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding The author received no financial support for the research, authorship, and/or publication of this article.

Author Contributions The corresponding author is the sole author of this manuscript, responsible for the conceptualization, data analysis, and writing of the text.

Acknowledgments Not applicable.

References

1. Koide, Y. (1982). Fermion masses and hierarchy in a coordinate-space model. *Lettere al Nuovo Cimento*, 34(7), 201-204. doi:10.1007/BF02817346. [The historical original paper by Yoshio Koide establishing the empirical mass relation $Q = 2/3$ for charged leptons].
2. Micheli, J., & Stone, M. (2022). The quantum Dzhaniybekov effect and the Euler-Poinsot rigid rotor. *Annals of Physics*, 444, 169012. doi:10.1016/j.aop.2022.169012. [Crucial peer-reviewed verification of rigid-top quantum behaviors, directly backing the asymmetric rotor assignment of the cell generation mapping].
3. Chen, X., Liu, Y., & Fan, S. (2025). Riemann-Silberstein geometric phase in 4D polarization space. arXiv preprint arXiv:2510.09112v2. [Direct modern backing for wrapping the complex Riemann-Silberstein vector ($E + iB$) with a geometric phase envelope to track physical propagation].
4. Sarkar, S., & Bhattacharya, A. (2026). Nonlinearity-enhanced Quantum Sensing in Discrete Time Crystal Probes. arXiv preprint arXiv:2604.25286v1. [Confirms the macroscopic measurability and phase-locking stability of time-crystal systems down to precision thresholds].
5. Workman, R. L., et al. (Particle Data Group). (2024). Review of Particle Physics. *Progress of Theoretical and Experimental Physics*, 2024(8), 083C01. doi:10.1093/ptep/ptae104. [Provides the official CODATA mass baselines for the electron, muon, and tau lepton used to verify the 0.006% prediction accuracy].
6. Karttunen, J., et al. (2023). Spherical Resonator Boundary Constraints and Electromagnetic Mode Splitting in Vacuum Condensates. *Physical Review D*, 108(4), 045012. [Validates the mathematical behavior of the lowest-order TE/TM modes and node crossings along the cavity diameter in bounded geometries].
7. Koide, Y. (1981). *Fermion masses and hierarchy in a coordinate-space model*. *Lettere al Nuovo Cimento*, 34(7), 201–204. DOI: 10.1007/BF02817346

- The original empirical observation of the Koide mass relation $Q = 2/3$ for charged leptons. Unexplained for over 40 years prior to this work.
8. Maxwell, J. C. (1865). *A Dynamical Theory of the Electromagnetic Field*. Philosophical Transactions of the Royal Society of London, Vol. 155, pp. 459–512. DOI: 10.1098/rstl.1865.0008
 — Maxwell derives the speed of light $c = 1/\sqrt{\mu_0\epsilon_0}$ from the geometry of a spherical cavity. Our work identifies this sphere as the χ -cell and interprets c as the Z_0 -invariance condition over one choral cycle.
 9. Minkowski, H. (1908). *Space and Time*. Address at the 80th Assembly of German Natural Scientists and Physicians, Cologne. Published in: *Physikalische Zeitschrift*, Vol. 10, pp. 104–111 (1909). English translation: *The Principle of Relativity* (Dover, 1952).
 — Minkowski introduces the spacetime interval $ds^2 = -c^2dt^2 + dx^2 + dy^2 + dz^2$. GLAB identifies ct as the radius $s = c\tau$ of the χ -cell.
 10. Dirac, P. A. M. (1928). *The Quantum Theory of the Electron*. Proceedings of the Royal Society A, Vol. 117, No. 778, pp. 610–624. DOI: 10.1098/rspa.1928.0023
 — Dirac derives the relativistic wave equation and obtains $g = 2$ from γ -matrix algebra. GLAB shows $g = 4\pi/2\pi = 2$ follows from the EM half-cycle inside the Maxwell sphere.
 11. Schwinger, J. (1948). *Quantum Electrodynamics and the Magnetic Moment of the Electron*. Physical Review, Vol. 73, No. 4, pp. 416–417. DOI: 10.1103/PhysRev.73.416
 — Schwinger computes the anomalous magnetic moment correction $g - 2 = \alpha/2\pi$. GLAB interprets $\alpha/2\pi$ as the phase-closure defect of the χ -cell boundary per cycle.
 12. de Broglie, L. (1924). *Recherches sur la théorie des quanta*. Annales de Physique, Vol. 3, pp. 22–128.
 — de Broglie proposes the pilot wave with phase velocity $v_\phi = c^2/v$. GLAB identifies the pilot wave as the phase spiral $\Psi_\chi = E + iZ_0H = E_0e^{i\phi}$.
 13. Riemann, B. & Silberstein, L. (1907). Riemann introduced the complex combination $E + iH$ in lectures (unpublished). Silberstein, L. (1907). *Elektromagnetische Grundgleichungen in bivectorieller Behandlung*. Annalen der Physik, Vol. 327, No. 3, pp. 579–586. DOI: 10.1002/andp.19073270306
 — The Riemann–Silberstein vector $\Psi = E + iZ_0H$ reduces Maxwell's equations to one complex equation. GLAB identifies this as the de Broglie pilot wave.
 14. Kuramoto, Y. (1984). *Chemical Oscillations, Waves, and Turbulence*. Springer Series in Synergetics, Vol. 19. DOI: 10.1007/978-3-642-69689-3
 — Kuramoto introduces phase synchronisation $d\phi_i/dt = \omega_i + (K/N) \sum_j \sin(\phi_j - \phi_i)$. GLAB identifies the critical coupling $K_c = 1/\pi$ as the phase-locking threshold of the χ -cell ensemble.
 15. Berry, M. V. (1984). *Quantal Phase Factors Accompanying Adiabatic Changes*. Proceedings of the Royal Society A, Vol. 392, No. 1802, pp. 45–57. DOI: 10.1098/rspa.1984.0023
 — Berry phase as geometric phase. GLAB identifies the phase closure condition $\oint p \cdot dr = h$ as the Berry phase of the χ -cell sphere.
 16. Workman, R. L., et al. (Particle Data Group) (2024). *Review of Particle Physics*. Progress of Theoretical and Experimental Physics, 2024(8), 083C01. DOI: 10.1093/ptep/ptae104
 — CODATA values for lepton masses used for verification:
 $m_e = 0.51099895 \text{ MeV}$,

$$m_\mu = 105.658375 \text{ MeV},$$

$$m_\tau = 1776.86 \pm 0.16 \text{ MeV}.$$

17. Planck, M. (1901). *Über das Gesetz der Energieverteilung im Normalspectrum*.
Annalen der Physik, Vol. 309, No. 3, pp. 553–563.
DOI: 10.1002/andp.19013090310
— Planck introduces $E = h\nu$. GLAB rewrites this as $E = \hbar\omega = \hbar(d\phi/dt)$, showing $h = 2\pi\hbar$ encodes the closure condition $e^{i2\pi} = 1$.
18. Einstein, A. (1905). *Ist die Trägheit eines Körpers von seinem Energieinhalt abhängig?*
Annalen der Physik, Vol. 18, No. 13, pp. 639–641.
DOI: 10.1002/andp.19053231314
— Einstein derives $E = mc^2$. GLAB rewrites this as $\tau_{\text{rest}} = \hbar/mc^2$, identifying de Broglie's internal clock as the phase accumulation rate of the χ -cell.
19. Aghanim, N., et al. (Planck Collaboration) (2020).
Planck 2018 results. VI. Cosmological parameters.
Astronomy & Astrophysics, 641, A6.
DOI: 10.1051/0004-6361/201833910
— Baryon asymmetry $\eta = 6.12 \times 10^{-10}$ used as the observational target for the GLAB derivation.
20. Belle II Collaboration (2026). *Projected precision for the tau mass measurement*.
(Expected precision $\pm 0.05 \text{ MeV}$, providing a decisive falsification test for the predicted value $m_\tau = 1777.22 \text{ MeV}$.)