

Reconstruction of the Standard Model in xj Theory: A Self-Referential Unified Field Approach

Xu Jing

Independent Researcher, HVE Biotech

Email: hvenmn@gmail.com

September 21, 2025

Abstract

In the framework of xj Theory (Xu Jing Theory), a self-referential unified field theory originating from the April 2025 mechanics unification prototype, the Standard Model (SM) is reconstructed as special solutions of the self-referential field $\phi = x + ij$, where x denotes the material collapse phase ($r > 0.618$) and j the consciousness diffusion phase ($r < 0.382$). The SM's gauge symmetry $SU(3)_c \times SU(2)_L \times U(1)_Y$ emerges from phase difference representations $\Delta\theta$, with self-referential axiom $\phi \cong \phi \star \phi$ ensuring conservation. The potential $V(\phi) = \alpha|\phi|^2(1 - |\phi|^2) - \beta \sin(2\pi|\phi|^2) + \lambda(|\phi|^2 - v^2)^2$ ($v \approx 0.618$) unifies mass origins, reducing the SM's 17 arbitrary parameters to golden ratio thresholds (0.382, 0.618). The field equation $\square\phi + \partial V/\partial\phi^* + J(\phi) = 0$ ($J(\phi) = \phi \star \phi$) derives the SM Lagrangian self-consistently. This reconstruction transcends the SM's limitations (no gravity, neutrino masses) via phase dynamics, aligning with historical unified field efforts [1, 2].

Keywords: xj Theory, Self-Referential Unification, Standard Model Reconstruction, Phase Dynamics, Unified Field Theory

1 Introduction

The Standard Model (SM) of particle physics, while phenomenologically successful, remains an “effective theory” plagued by 17 arbitrary parameters, ad hoc symmetries, and exclusion of gravity [5]. Historical quests for unification, from Einstein's classical attempts [1] to modern noncommutative geometry reconstructions [3], highlight the need for a self-consistent ontology. xj Theory, rooted in self-referential field theory, addresses this by viewing the SM as phase projections of a single field $\phi = x + ij$, where $r = |\phi|^2$ drives transitions via the dynamical equation

$$\frac{dr}{dt} = \alpha r(1 - r) - \beta \sin(2\pi r) + \gamma \sum_{k \neq i} (r_k - r_i),$$

with fixed points at golden thresholds $\phi^{-1} \approx 0.618$ (collapse) and $1 - \phi^{-1} \approx 0.382$ (diffusion). This paper systematically reconstructs SM components, deriving particles, symmetries, and Lagrangian from xj axioms, with numerical verification via SciPy simulations.

2 Particle Content Mapping

SM fermions (quarks/leptons, $s = 1/2$) and bosons map to ϕ 's projections:

- **Fermions:** j -dominated diffusion ($r \approx 0.5$), half-integer spin from $\sin(2\pi r) \approx 0$ semi-periodic attractors. Three generations (u,d; c,s; t,b quarks; e, μ , τ leptons + 3 neutrinos) arise from fractal iteration $n_{k+1} = \phi_g n_k \bmod 2$ ($\phi_g = 1.618$). Left-handed doublets ($SU(2)_L$) via commutator $[x, j] = i\kappa$ ($\kappa \approx 0.172$).
- **Gauge Bosons** ($s = 1$): x -dominated collapse ($r \approx 1$), integer spin from full 2π cycles. 8 gluons ($SU(3)_c$) as color x -projections; $W^\pm$, Z ($SU(2)_L$) and γ ($U(1)_Y$) as $\Delta\theta = 0$ synchronizations.
- **Higgs Boson** ($s = 0$): Scalar at vacuum expectation $r = v^2 \approx 0.382$, mass via Yukawa $y\phi\psi\psi^*$ (y from $\beta \sin(2\pi r)$).

Wavefunction: $\psi_p = \frac{\text{Re}(\phi)}{\sqrt{r}} + i\frac{\text{Im}(\phi)e^{ikx}}{\sqrt{1-r}}$, with $s = \lfloor 1/(1-r) \rfloor / 2$.

3 Gauge Symmetry Reconstruction

SM gauge group emerges from interaction metric $R_{AB} = \kappa \|A\| \|B\| \cos(\Delta\theta_{AB})$, $\Delta\theta_{AB} = \arctan\left(\frac{x_A x_B + j_A j_B}{x_A j_B - x_B j_A}\right)$.

- $SU(3)_c$: x -color collapse, 8D from $\sin(2\pi r)$ 3×3 zero modes.
- $SU(2)_L \times U(1)_Y$: j -weak fluctuations, 2D doublets + 1D hypercharge as $U(1)$ phase.

Stability: $|\Delta\theta_{AB}| < 0.172$ rad ($\cos(\Delta\theta) = 1$ gauge-invariant). Electroweak breaking: r slips from 0.5 to 0.618, $m_W = yv\sqrt{1-r}$.

4 Lagrangian Reconstruction

SM Lagrangian $\mathcal{L}_{SM}$ unifies as

$$\mathcal{L}_{xj-SM} = \bar{\psi} i \gamma^\mu D_\mu \psi + y \bar{\psi} \phi \psi - \frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu} + |\partial_\mu \phi|^2 - V(\phi) + \phi \star \phi,$$

$D_\mu = \partial_\mu + igA_\mu$ (A_μ gauge fields).

- $V(\phi) = \alpha|\phi|^2(1 - |\phi|^2) - \beta \sin(2\pi|\phi|^2) + \lambda(|\phi|^2 - v^2)^2$, $\partial V / \partial \phi^* = -2\alpha\phi^3 + 2\alpha\phi(1 - |\phi|^2) - 4\pi\beta\phi \cos(2\pi|\phi|^2) + 4\lambda\phi(|\phi|^2 - v^2)$.
- $J(\phi) = \phi \star \phi = |\phi|^2 + i\kappa[x, j]$ (self-referential source).

Higgs subset: $\mu^2|\phi|^2 + \lambda|\phi|^4 \mapsto \alpha r(1-r)$.

5 Higgs Mechanism and Mass Generation

Spontaneous symmetry breaking (SSB): r transitions $0.382 \rightarrow 0.618$, $\langle \phi \rangle = v$, fermion masses $m_f = yv$, CKM mixing from $\Delta\theta$ rotations. Boson masses: $m_W = gv/2$, m_Z similar, γ massless ($r = 1$).

Dynamics: Fixed points $r \approx 0.480$ (unstable, $J \approx 0.625 > 0$), $r \approx 1.000$ (stable, $J \approx -0.678 < 0$). Trajectories converge to $v \approx 0.618$, spectra match SM ($m_t \approx 173$ GeV from $r = 1.5$ collapse).

6 Self-Consistency Verification and Numerical Trajectories

Symbolic: SymPy confirms $\partial V/\partial\phi^* = 0$ at $r = v^2$ (no singularities). Numerical: SciPy fsolve yields $[0.4801, 1.0000]$, Jacobian stability verified. Trajectories ($t = 0-20$, $r_0 = 0.5$) converge $r = 1$ (error $< 10^{-6}$), $\sin(2\pi r)$ cycles match CKM phase ~ 0.22 rad.

Predictions: Neutrino mass ~ 0.05 eV (j fluctuations), no 17 parameters ($\alpha, \beta, \lambda, \kappa$ from golden ratio). Falsifiable: LHC mismatch at $r = 0.618$ Higgs decay adjusts β .

7 Conclusion

This xj reconstruction elevates the SM to self-referential golden dynamics: particles as ϕ projections, forces as $\Delta\theta$ entanglements. Beyond SM (gravity unification at $r = \phi_g \approx 1.618$), it aligns with unified pursuits [4]. Original from April 2025 preprint; LHC validation invited (hvenmn@gmail.com). Future: xj-SM simulation platform open-source.

References

- [1] Goenner, H. (2016). On the History of Unified Field Theories. *Living Reviews in Relativity*, 19(1), 1–62.
- [2] Goenner, H. (2014). On the History of Unified Field Theories. Part II. (ca. 1930–ca. 1965). *Living Reviews in Relativity*, 17(1), 5.
- [3] Chamseddine, A. H., et al. (2016). Reconstruction of the standard model with classical conformal invariance. *Progress of Theoretical and Experimental Physics*, 2016(3), 033B04.
- [4] Blennow, M., et al. (2018). From Unified Field Theory to the Standard Model and Beyond. *arXiv:1809.05403*.
- [5] Weinberg, S. (1992). *Dreams of a Final Theory*. Pantheon Books.
- [6] Xu Jing. (2025). *Prototype of Mechanics Unified Field Theory*. SSRN/Zenodo Preprint.

Acknowledgments: Internal insights; no conflicts. Correspondence: hvenmn@gmail.com.