

A Unified $SU(4)$ Qudit–Gauge Lattice Framework for Emergent Geometry and Interaction

Bryan E. W. Van Zandt

KILOWADS Studios, LLC
Averill Park, NY, USA

May 21, 2026

Abstract

We explore a unified field framework in which the fundamental degrees of freedom are four-level qudits residing on the sites of a discrete lattice, with $SU(4)$ gauge variables living on the links. The four basis states of each qudit correspond to a singlet “gravity-mode” and a triplet of gauge-modes. Effective symmetry breaking $SU(4) \rightarrow SU(3) \times U(1)$ naturally decomposes the qudit field into a geometric sector and a gauge sector. Spatial variations in the singlet amplitude generate a scalar notion of curvature, leading to metric-like behavior in the continuum limit, while excitations in the triplet sector give rise to gauge interactions.

Because the singlet and triplet sectors couple through the unified $SU(4)$ structure, the local geometric density can influence the effective mass of the triplet fields. At the effective level, this allows for the possibility of a curvature-driven symmetry breaking in the early universe, leading to a composite Higgs-like condensate and a corresponding *geometric electroweak epoch*. This mechanism is exploratory and relies on effective operators consistent with the symmetries of the model.

The framework is intended as a conceptual proposal illustrating how geometry and gauge interactions might emerge from a single discrete quantum substrate. It highlights several avenues for further development, including the treatment of fermions, quantitative predictions, and the derivation of effective couplings from the microscopic $SU(4)$ theory.

1 Introduction

The unification of gravity with the gauge interactions of the Standard Model remains one of the central open challenges in theoretical physics. General relativity describes gravitation as the curvature of a smooth spacetime manifold, while quantum field theory models the electromagnetic, weak, and strong interactions as gauge fields defined on that manifold. Despite their successes, these frameworks rest on fundamentally different principles, and a single description that encompasses both geometric and gauge phenomena has yet to be established. Traditional approaches—including string theory [1, 2], loop quantum gravity [3, 4], and grand unified theories [5, 6]—typically require extended mathematical structures, high unification scales, or a continuous spacetime background as a starting point.

In this work, we explore a different possibility: that geometry and gauge interactions may emerge from a *single discrete quantum substrate*. We consider a lattice whose sites host four-level qudits transforming under the fundamental representation of a unified $SU(4)$ gauge group. The four basis states of each qudit correspond to four interaction modes: a singlet “gravity-mode” and a triplet of gauge-modes. After effective symmetry breaking, the qudit field decomposes into a geometric sector and a gauge sector. Spatial variations in the singlet amplitude define a scalar notion of curvature, while excitations in the triplet sector give rise to electromagnetic-, weak-, and strong-like interactions. In this picture, spacetime geometry, gauge fields, and matter-like excitations arise as emergent phenomena of a single qudit–gauge system.

This framework draws on several converging developments. Lattice gauge theory demonstrates that discrete formulations can capture essential features of non-Abelian gauge dynamics [7, 8]. Qudit-based quantum systems naturally encode multi-state internal degrees of freedom [9, 10]. Approaches to quantum gravity—including spin networks, causal dynamical triangulations, and tensor network geometries—suggest that spacetime itself may be emergent from discrete quantum degrees of freedom [11, 12, 13]. Meanwhile, the decomposition of the $SU(4)$ fundamental representation into a triplet and a singlet under $SU(3) \times U(1)$ provides a natural mechanism for distinguishing gauge and geometric sectors within a unified representation.

A central feature of the model is the coupling between the singlet (geometry) and triplet (gauge/Higgs) sectors through the unified $SU(4)$ potential. At the effective level, this coupling allows the local geometric density to influence the mass of the triplet fields. We explore the possibility that the relaxation of curvature in the early universe could trigger a second-stage symmetry breaking, leading to a composite Higgs-like condensate and a corresponding *geometric electroweak epoch*. This mechanism is speculative and is introduced here as an effective description consistent with the symmetries of the model.

The remainder of this paper develops the formal structure of the model, examines its emergent geometric and gauge behavior, and illustrates how familiar physical phenomena may arise from the unified qudit–gauge substrate. Section III introduces the lattice construction and the $SU(4)$ qudit representation.

Section IV describes the symmetry breaking pattern and the composite Higgs mechanism at the effective level. Section V develops the notion of emergent curvature from the singlet field. Section VI analyzes the gauge interactions arising from the triplet sector. Section VII presents explicit examples, including gravitational lensing and Coulomb-like forces. Section VIII discusses the continuum limit and effective field theory behavior. Section IX outlines phenomenological consequences and predictions. Section X introduces the geometric electroweak epoch and its cosmological implications. Section XI compares the framework to other approaches, and Section XII discusses limitations and future directions.

This work is intended as a conceptual proposal rather than a complete theory. It suggests that the apparent diversity of physical forces may arise from a single discrete quantum substrate and highlights several avenues for further development.

2 Background and Motivation

The idea that spacetime and the fundamental interactions may emerge from a deeper quantum substrate has gained increasing attention across multiple areas of theoretical physics. While general relativity treats spacetime as a smooth differentiable manifold endowed with a metric, and quantum field theory describes matter and forces as excitations of fields defined on that manifold, several conceptual and technical challenges suggest that this continuum description may not be fundamental. Among these challenges are the non-renormalizability of perturbative quantum gravity, the black hole information problem, and the apparent discreteness implied by holographic entropy bounds [14, 15].

Discrete approaches to quantum gravity, including loop quantum gravity [11, 16], causal dynamical triangulations [12, 17], and spin foam models [18], propose that spacetime geometry arises from combinatorial or algebraic structures rather than a pre-existing manifold. Similarly, tensor network models have demonstrated that geometric notions such as distance, curvature, and even emergent gravitational dynamics can arise from entanglement patterns in many-body quantum systems [13, 19]. These developments suggest that geometry may be a derived concept, emerging from more fundamental quantum degrees of freedom.

Parallel to these efforts, lattice gauge theory has provided a powerful non-perturbative framework for studying gauge interactions [7, 8]. By discretizing spacetime into a lattice and placing gauge variables on the links, lattice gauge theory captures essential features of non-Abelian gauge dynamics, including confinement, chiral symmetry breaking, and phase transitions. Importantly, lattice formulations demonstrate that gauge fields can be defined without reference to a smooth underlying manifold, relying instead on group-valued variables associated with discrete structures.

The present work draws inspiration from both of these lines of research. We propose that the fundamental degrees of freedom of the universe are not fields defined on a manifold but rather *qudits*—finite-dimensional quantum systems—

located on the sites of a discrete lattice. Qudits generalize qubits by allowing d internal states, and have been extensively studied in quantum information theory, condensed matter physics, and quantum error correction [9, 10]. In particular, qudits naturally encode multi-state internal degrees of freedom, making them suitable for representing the internal structure of matter and interaction modes.

The choice of a four-level qudit is motivated by the structure of the fundamental interactions. The Standard Model contains three gauge interactions—electromagnetic, weak, and strong—while gravity is described separately by general relativity. A four-level system provides a natural way to encode these four interaction modes within a single quantum object. Moreover, the fundamental representation of $SU(4)$ is four-dimensional, and its decomposition under the subgroup $SU(3) \times U(1)$ yields a triplet and a singlet:

$$\mathbf{4} \rightarrow \mathbf{3} \oplus \mathbf{1}, \quad (1)$$

mirroring the division between gauge interactions and gravity-like behavior in our model.

The idea of unifying interactions through higher-dimensional gauge groups has a long history, including the Pati-Salam model [20], $SO(10)$ unification [21], and E_6 -based theories [22]. However, these approaches typically embed the Standard Model gauge group into a larger continuous group defined on a smooth spacetime. In contrast, our framework does not assume a pre-existing manifold or continuous geometry. Instead, geometry and gauge interactions both emerge from the dynamics of qudits and gauge variables on a discrete lattice.

The unified $SU(4)$ structure plays a central role in our approach. At high energies or short distances, the full $SU(4)$ symmetry governs the dynamics of the qudit field. At lower energies, effective symmetry breaking to $SU(3) \times U(1)$ separates the qudit into a gauge-mode triplet and a gravity-mode singlet. This breaking pattern is reminiscent of GUT symmetry breaking but differs in that the singlet is interpreted not as a sterile matter field but as a geometric degree of freedom. Spatial variations in the singlet amplitude define a discrete notion of curvature, while excitations in the triplet sector give rise to gauge interactions.

This perspective aligns with the broader idea that geometry may emerge from quantum information. Recent work has shown that entanglement entropy can encode geometric features of spacetime [23, 24], and that tensor networks can reproduce aspects of AdS/CFT duality [13]. In our model, the qudit field plays a dual role: it encodes both geometric and gauge information, with the unified $SU(4)$ structure ensuring that these sectors are not fundamentally distinct.

Finally, the lattice formulation provides a natural setting for defining dynamics without reference to a background geometry. The gauge variables on the links define parallel transport and interaction strengths, while the qudit field defines local internal states. Together, they form a self-contained quantum system from which geometry, gauge fields, and matter emerge in appropriate limits.

In summary, the motivation for this work arises from the convergence of several ideas: the discreteness suggested by quantum gravity, the success of lattice gauge theory, the versatility of qudits, and the unifying structure of $SU(4)$. By combining these elements, we aim to construct a framework in which geometry and interaction arise from a single quantum substrate, offering a new perspective on unification and the nature of spacetime.

3 Formal Construction of the $SU(4)$ Qudit–Gauge Lattice

The formal structure of the $SU(4)$ qudit–gauge lattice begins with a discrete hypercubic lattice whose sites carry four-level quantum systems and whose links carry $SU(4)$ gauge variables. This section develops the mathematical foundation of the model, from the definition of the qudit Hilbert space to the construction of link operators, plaquette curvature, and the unified action. The goal is to show how a single $SU(4)$ structure can encode both geometric and gauge degrees of freedom in a unified, discrete substrate.

3.1 Lattice Structure

We consider a d -dimensional hypercubic lattice with lattice spacing a . Each lattice site x hosts a four-level qudit, and each oriented link $(x, x + \hat{\mu})$ hosts an $SU(4)$ matrix.

The basic objects are:

- lattice sites x
- unit vectors $\hat{\mu}$
- link variables $U_{\mu}(x) \in SU(4)$
- qudit fields $\psi(x) \in \mathbb{C}^4$

The lattice is the fundamental structure; no continuum manifold is assumed.

Guided Link: Lattice structure

3.2 Qudit Hilbert Space

Each site carries a four-dimensional Hilbert space:

$$\mathcal{H}_x = \mathbb{C}^4.$$

We choose a basis:

$$\psi(x) = \begin{pmatrix} \psi_0(x) \\ \psi_1(x) \\ \psi_2(x) \\ \psi_3(x) \end{pmatrix}.$$

The key structural insight is that the $SU(4)$ fundamental representation decomposes as:

$$4 \rightarrow 1 \oplus 3,$$

corresponding to:

- a **singlet** $g(x) = \psi_0(x)$
- a **triplet** $\phi_i(x) = \psi_i(x)$, $i = 1, 2, 3$

The singlet becomes the geometric sector; the triplet becomes the gauge sector.

Guided Link: Qudit decomposition




Figure 1: Decomposition of the qudit field in the fundamental representation $\mathbf{4}$ of $SU(4)$ into a gravity-mode singlet and a gauge-mode triplet under $SU(3) \times U(1)$.

3.3 $SU(4)$ Gauge Variables

Each oriented link carries an $SU(4)$ matrix:

$$U_\mu(x) \in SU(4).$$

These link variables mediate interactions between qudits at neighboring sites. They are the discrete analog of gauge connections.

The $SU(4)$ generators T^A satisfy:

$$\text{Tr}(T^A T^B) = \frac{1}{2} \delta^{AB}.$$

Guided Link: $SU(4)$ gauge variables

3.4 Local Gauge Transformations

A local $SU(4)$ transformation acts as:

$$\psi(x) \rightarrow G(x)\psi(x), \quad G(x) \in SU(4),$$

and the link variables transform as:

$$U_\mu(x) \rightarrow G(x)U_\mu(x)G^\dagger(x + \hat{\mu}).$$

This ensures invariance under independent $SU(4)$ rotations at each site.

Guided Link: Local gauge transformations

3.5 Link Operators and Transport

The link variable defines forward parallel transport:

$$\psi(x + \hat{\mu}) \rightarrow U_\mu(x)\psi(x).$$

Backward transport uses the Hermitian conjugate:

$$\psi(x - \hat{\mu}) \rightarrow U_\mu^\dagger(x - \hat{\mu})\psi(x).$$

These rules define how qudit amplitudes propagate across the lattice.

Guided Link: Parallel transport

3.6 Plaquette Operators and Curvature

The elementary plaquette in the $\mu\nu$ plane is:

$$U_{\mu\nu}(x) = U_\mu(x)U_\nu(x + \hat{\mu})U_\mu^\dagger(x + \hat{\nu})U_\nu^\dagger(x).$$

This is the discrete curvature operator. If $U_{\mu\nu}(x) = \mathbb{I}$, the region is flat; deviations encode curvature.

Guided Link: Plaquette curvature

3.7 Unified Action

The gauge action is the Wilson action:

$$S_{\text{gauge}} = \frac{\beta}{4} \sum_{\mu\nu, x} \text{Re Tr}(\mathbb{I} - U_{\mu\nu}(x)).$$

The matter action is:

$$S_{\text{matter}} = \sum_{x, \mu} [\psi^\dagger(x)U_\mu(x)\psi(x + \hat{\mu}) + \psi^\dagger(x + \hat{\mu})U_\mu^\dagger(x)\psi(x)].$$

The unified action is:

$$S = S_{\text{gauge}} + S_{\text{matter}}.$$

Guided Link: Unified action

3.8 Gauge Invariance Proof

Under a local transformation:

$$\psi(x) \rightarrow G(x)\psi(x), \quad U_\mu(x) \rightarrow G(x)U_\mu(x)G^\dagger(x + \hat{\mu}),$$

the matter term transforms as:

$$\psi^\dagger(x)U_\mu(x)\psi(x + \hat{\mu}) \rightarrow \psi^\dagger(x)G^\dagger(x)G(x)U_\mu(x)G^\dagger(x + \hat{\mu})G(x + \hat{\mu})\psi(x + \hat{\mu}),$$

which leaves the expression invariant.

The gauge term is invariant because the trace of a closed loop is invariant under conjugation.

Guided Link: Gauge invariance

3.9 Operator Algebra

The $SU(4)$ generators satisfy:

$$[T^A, T^B] = if^{ABC}T^C,$$

and the link variables satisfy:

$$U_\mu(x)U_\mu^\dagger(x) = \mathbb{I}.$$

The qudit operators obey standard canonical commutation relations.

Guided Link: Operator algebra

3.10 Continuum Limit Considerations

As $a \rightarrow 0$, the link variables expand as:

$$U_\mu(x) = \exp(ia\mathcal{A}_\mu(x)),$$

and the plaquette becomes:

$$U_{\mu\nu}(x) = \exp(ia^2F_{\mu\nu}(x)).$$

The action reduces to a Yang–Mills action plus kinetic terms for the qudit fields.

Guided Link: Continuum limit

4 Symmetry Breaking and Representation Theory

The unified $SU(4)$ structure contains both geometric and gauge information. In this section we describe how the qudit field decomposes into a gravity-mode singlet and a gauge-mode triplet under the breaking $SU(4) \rightarrow SU(3) \times U(1)$, and how an additional $SU(3) \rightarrow SU(2)$ breaking may occur at the effective level. The latter is driven by a composite Higgs-like mode constructed from the triplet sector. Throughout this section, we emphasize that the symmetry-breaking potential is introduced as an *effective description* consistent with the symmetries of the model, rather than a derivation from the microscopic lattice action.

4.1 Structure of $SU(4)$

$SU(4)$ is a 15-generator Lie group with:

- rank 3,
- fundamental representation **4**,
- adjoint representation **15**.

The generators decompose naturally into:

- 8 generators of $SU(3)$,
- 1 generator of $U(1)$,
- 6 coset generators mixing singlet and triplet sectors.

This decomposition underlies the emergent geometric and gauge structure.

4.2 Decomposition of the Fundamental Representation

The qudit lives in the fundamental representation:

$$\mathbf{4} = \{0, 1, 2, 3\}.$$

Under symmetry breaking,

$$\mathbf{4} \rightarrow \mathbf{1} \oplus \mathbf{3},$$

corresponding to:

- gravity-mode singlet 0,
- gauge-mode triplet $\{1, 2, 3\}$.

4.3 Physical Interpretation of the Singlet and Triplet

4.3.1 Gravity-mode (Singlet)

The singlet transforms trivially under $SU(3)$:

$$0 \rightarrow e^{i\theta} 0.$$

Its amplitude

$$g(x) = \psi_0(x)$$

defines the geometric sector. Spatial variations in $g(x)$ encode curvature in the emergent scalar–tensor sense.

4.3.2 Gauge-mode (Triplet)

The triplet transforms as:

$$i \rightarrow U_{ij}^{(3)} j, \quad i, j = 1, 2, 3,$$

and carries color-like, electromagnetic-like, and weak-like structure.

4.4 Effective Symmetry Breaking Mechanism

The breaking $SU(4) \rightarrow SU(3) \times U(1)$ may be explicit, effective, or spontaneous. A convenient effective description is provided by a qudit condensate:

$$\langle \psi(x) \rangle = v 0,$$

which selects the singlet direction and leaves $SU(3) \times U(1)$ unbroken.

The subsequent breaking $SU(3) \rightarrow SU(2)$ is not derived from the microscopic lattice action. Instead, we introduce an *effective potential* for the singlet $g(x)$ and triplet $\phi_i(x)$ fields, consistent with the symmetries of the model, to explore how such a breaking might occur.

4.5 Effective Potential for the Singlet and Triplet Sectors

At the effective level, the most general renormalizable potential consistent with $SU(3) \times U(1)$ symmetry and the singlet–triplet decomposition is:

$$V_{\text{eff}}[g, \phi] = m_g^2 g^2 + \lambda_g g^4 + m_\phi^2 \phi^\dagger \phi + \lambda_\phi (\phi^\dagger \phi)^2 + \alpha g^2 \phi^\dagger \phi + \beta g |\phi_3|^2 + \gamma |\phi_3|^4. \quad (2)$$

We emphasize that:

- This potential is *not* derived from the microscopic $SU(4)$ lattice action.
- It is introduced as an *effective field theory* capturing the lowest-order couplings allowed by symmetry.
- The coefficients $(m_g^2, m_\phi^2, \alpha, \beta, \gamma)$ are phenomenological parameters.

This effective description allows us to explore how geometry and gauge sectors may interact at low energies.

4.6 Curvature-Triggered Effective Symmetry Breaking

To study the breaking $SU(3) \rightarrow SU(2)$, we restrict to the direction relevant for the composite Higgs-like mode by setting $\phi_1 = \phi_2 = 0$ and $\phi_3 \equiv h(x)$. The effective potential becomes:

$$V_{\text{eff}}(h; g) = m_{\text{eff}}^2(g) |h|^2 + (\lambda_\phi + \gamma) |h|^4,$$

with

$$m_{\text{eff}}^2(g) = m_\phi^2 + \alpha g^2 + \beta g.$$

High curvature ($|g|$ large) gives $m_{\text{eff}}^2(g) > 0$ and

$$\langle h(x) \rangle = 0,$$

leaving $SU(3)$ unbroken.

As curvature relaxes, $m_{\text{eff}}^2(g)$ may become negative, triggering a composite Higgs-like condensation:

$$\langle h(x) \rangle = v(x) = \sqrt{-\frac{m_{\text{eff}}^2(g(x))}{2(\lambda_\phi + \gamma)}}.$$

This breaks

$$SU(3) \rightarrow SU(2),$$

where the unbroken $SU(2)$ subgroup acts on (ϕ_1, ϕ_2) .

We stress that this mechanism is *exploratory* and depends on the effective parameters in Eq. (2). It is not claimed to reproduce the Standard Model electroweak sector.

4.7 Summary of the Breaking Pattern

$$SU(4) \rightarrow SU(3) \times U(1) \rightarrow SU(2) \times U(1).$$

Sector	Representation	Physical Role
Singlet	1	Geometry (gravity-mode)
Triplet	3	Gauge interactions; composite Higgs-like mode
Coset	$3 \oplus \bar{3}$	Gravity-gauge mixing (suppressed)

This effective symmetry-breaking structure provides a conceptual route by which geometry and gauge dynamics may interact within the unified $SU(4)$ qudit-gauge framework.

5 Emergent Geometry

The central claim of this framework is that **geometry is not fundamental**. Instead, it emerges from the **gravity-mode singlet** of the $SU(4)$ qudit field:

$$g(x) = \psi_0(x).$$

Spatial variations in $g(x)$ define curvature, metric behavior, and gravitational effects. This section formalizes how geometry arises from the qudit–gauge substrate.

5.1 The Gravity-Mode Field as a Geometric Scalar

The singlet field $g(x)$ transforms only under $U(1)$:

$$g(x) \rightarrow e^{i\theta(x)}g(x),$$

making it a natural candidate for a **geometric scalar**.

The magnitude $g(x)$ encodes **local geometric density**, while its phase encodes **geometric flux**.

Guided Links: Gravity-mode definition, Scalar geometric fields

5.2 Discrete Metric from Qudit Amplitudes

We define a **lattice metric weight**:

$$w_\mu(x) = g(x + \hat{\mu})^2 - g(x)^2.$$

Interpretation:

- If $w_\mu(x) = 0$: locally flat
- If $w_\mu(x) \neq 0$: curved

This gives a **direction-dependent metric distortion**.

We then define an **effective discrete metric**:

$$ds^2 = \sum_{\mu} (1 + \alpha w_\mu(x)) dx_\mu^2,$$

where α is a coupling constant.

Guided Links: Discrete metric, Metric emergence

5.3 Curvature from Plaquette Distortions

Curvature arises from **second-order differences** in the singlet field:

$$R_{\mu\nu}(x) \sim g(x + \hat{\mu} + \hat{\nu}) - g(x + \hat{\mu}) - g(x + \hat{\nu}) + g(x).$$

This is the discrete analog of:

$$R \sim \partial_\mu \partial_\nu g.$$

The curvature couples to the gauge sector through the unified action.

Guided Links: Discrete curvature, Lattice Ricci tensor

5.4 Geodesics from Qudit Transport

Particles follow **paths of extremal qudit amplitude overlap**.

Define the **transition amplitude**:

$$A(x \rightarrow y) = \psi(y) |U(x, y)| \psi(x).$$

A geodesic is the path that **maximizes**:

$$\prod_{\text{path}} A(x \rightarrow x + \hat{\mu}).$$

This reproduces gravitational lensing behavior.

Guided Links: Geodesics from amplitudes, Emergent lensing

5.5 Effective Gravitational Action

The geometric sector has an effective action:

$$S_{\text{geom}} = \sum_x [(\nabla g)^2 + V(g) + \lambda R[g]].$$

Where:

- $(\nabla g)^2$ gives kinetic curvature
- $V(g)$ stabilizes the vacuum
- $R[g]$ is the discrete curvature scalar

This is the analog of:

$$S_{\text{GR}} = \int d^4x \sqrt{-g} R.$$

Guided Links: Emergent gravitational action, Scalar-curvature coupling

5.6 Coupling Between Geometry and Gauge Modes

The unified $SU(4)$ structure ensures that geometry and gauge fields are not independent.

The coupling term:

$$S_{\text{mix}} = \sum_x g(x) \bar{\phi}(x) \phi(x)$$

means:

- gauge excitations distort geometry
- geometry influences gauge propagation

This is the discrete analog of:

- gravitational redshift
- curvature-dependent gauge propagation
- energy–geometry coupling

Guided Links: Gauge–geometry coupling, Backreaction

5.7 Small-Fluctuation Analysis and Newtonian-Like Behavior

To explore whether the emergent geometry of the model can reproduce aspects of weak-field gravity, we consider small fluctuations of the singlet field around a homogeneous background value,

$$g(x) = g_0 + \delta g(x), \quad |\delta g| \ll |g_0|.$$

The effective metric induced by the singlet amplitude is

$$g_{\mu\nu}(x) = \eta_{\mu\nu} + \alpha \partial_\mu g \partial_\nu g.$$

To leading order in the fluctuation δg , the metric perturbation is quadratic in derivatives,

$$h_{\mu\nu}(x) \approx \alpha (\partial_\mu \delta g)(\partial_\nu \delta g).$$

This structure differs from the linear weak-field expansion of general relativity and already indicates that the present construction yields only a schematic description of gravitational effects.

For a static, spherically symmetric fluctuation $\delta g = \delta g(r)$, the dominant component of the perturbation is

$$h_{00}(r) \approx \alpha \left(\frac{d\delta g}{dr} \right)^2.$$

In conventional weak-field gravity, the Newtonian potential $\Phi(r)$ is related to the metric perturbation by $h_{00} \approx -2\Phi$. One may therefore tentatively associate a Newtonian-like potential with the gradient of the fluctuation,

$$\Phi(r) \sim -\frac{\alpha}{2} \left(\frac{d\delta g}{dr} \right)^2.$$

However, because h_{00} depends on the *square* of the gradient rather than on δg itself, the resulting effective potential does not automatically reproduce the $1/r$ Newtonian form or satisfy the Poisson equation $\nabla^2\Phi = 4\pi G\rho$. A full demonstration that this framework recovers Newtonian gravity in the appropriate limit would require a more detailed analysis, including the back-reaction of matter sources on the singlet field and a proper identification of the stress-energy tensor. Such a derivation lies beyond the scope of the present work.

The above exercise serves only to illustrate that spatial variations in the geometry-mode field can, in principle, generate an effective gravitational potential. Whether this potential can be made quantitatively consistent with observed gravitational phenomenology remains an open question that merits further investigation.

5.8 Continuum Limit: Scalar–Tensor Behavior

As the lattice spacing $a \rightarrow 0$, the singlet field $g(x)$ becomes a smooth scalar field, and the discrete curvature operators converge to second-derivative structures of the form

$$R_{\mu\nu}(x) \sim \partial_\mu \partial_\nu g(x).$$

The resulting continuum description exhibits scalar–tensor behavior: the effective metric

$$g_{\mu\nu}(x) = \eta_{\mu\nu} + \alpha \partial_\mu g \partial_\nu g$$

is induced by gradients of the singlet field rather than introduced as a fundamental degree of freedom. This leads to metric-like and curvature-like effects, but does not reproduce the Einstein–Hilbert action or diffeomorphism invariance. Instead, the continuum limit yields a scalar–tensor model in which geometric responses arise from the dynamics of $g(x)$.

This behavior is consistent with the emergent nature of geometry in the qudit–gauge framework and provides a basis for exploring gravitational phenomena without assuming a fundamental spacetime manifold.

6 Emergent Gauge Interactions

The triplet sector of the $SU(4)$ qudit field gives rise to emergent gauge interactions. After symmetry breaking

$$SU(4) \rightarrow SU(3) \times U(1),$$

the triplet fields transform under the residual gauge symmetry and couple to the corresponding gauge fields. This section formalizes how gauge interactions arise from the qudit–gauge substrate.

6.1 Triplet Transformation Properties

The triplet fields are defined as:

$$\phi_i(x) = \psi_i(x), \quad i = 1, 2, 3.$$

Under an $SU(3)$ transformation:

$$\phi_i(x) \rightarrow U_{ij}^{(3)}(x) \phi_j(x),$$

where $U^{(3)}(x) \in SU(3)$.

Under the residual $U(1)$:

$$\phi_i(x) \rightarrow e^{i\theta(x)} \phi_i(x).$$

Thus the triplet carries both color-like and electromagnetic-like structure.

Guided Links: Triplet transformation, Gauge-mode structure

6.2 Gauge Fields from $SU(4)$ Link Variables

The $SU(4)$ link variable decomposes as:

$$U_\mu(x) = \exp \left[ia \left(A_\mu^{(3)}(x) + B_\mu(x) + X_\mu(x) \right) \right],$$

where:

- $A_\mu^{(3)}(x)$ are the $SU(3)$ gauge fields,
- $B_\mu(x)$ is the $U(1)$ gauge field,
- $X_\mu(x)$ are the coset fields (suppressed after breaking).

Guided Links: Gauge field decomposition, $SU(4)$ link structure

6.3 Covariant Derivative and Transport

The covariant derivative acting on the triplet is:

$$D_\mu \phi(x) = \frac{1}{a} [U_\mu(x) \phi(x + \hat{\mu}) - \phi(x)].$$

In the continuum limit:

$$D_\mu = \partial_\mu + ig_3 A_\mu^{(3)} + ig_1 B_\mu.$$

This defines how gauge interactions modify qudit propagation.

Guided Links: Covariant derivative, Gauge transport

6.4 Field Strength and Plaquette Dynamics

The plaquette operator:

$$U_{\mu\nu}(x) = U_\mu(x)U_\nu(x + \hat{\mu})U_\mu^\dagger(x + \hat{\nu})U_\nu^\dagger(x)$$

encodes the discrete field strength.

Expanding for small a :

$$U_{\mu\nu}(x) = \exp(ia^2 F_{\mu\nu}(x)) ,$$

where $F_{\mu\nu}$ contains both $SU(3)$ and $U(1)$ components.

Guided Links: Plaquette field strength, Lattice gauge curvature

6.5 Emergent $SU(3)$ Dynamics

The $SU(3)$ field strength is:

$$F_{\mu\nu}^{(3)} = \partial_\mu A_\nu^{(3)} - \partial_\nu A_\mu^{(3)} + ig_3[A_\mu^{(3)}, A_\nu^{(3)}].$$

The corresponding lattice action is:

$$S_{SU(3)} = \frac{\beta_3}{4} \sum_{x, \mu\nu} \text{Tr} \left(\mathbb{I} - U_{\mu\nu}^{(3)}(x) \right) .$$

This reproduces color-like interactions.

Guided Links: $SU(3)$ dynamics, Color interactions

6.6 Emergent $U(1)$ Dynamics

The $U(1)$ field strength is:

$$F_{\mu\nu}^{(1)} = \partial_\mu B_\nu - \partial_\nu B_\mu.$$

The lattice action is:

$$S_{U(1)} = \frac{\beta_1}{4} \sum_{x, \mu\nu} \left(1 - \cos F_{\mu\nu}^{(1)}(x) \right) .$$

This reproduces electromagnetic-like behavior.

Guided Links: $U(1)$ dynamics, Electromagnetic emergence

6.7 Triplet Self-Interactions

The triplet fields interact through the gauge fields:

$$S_{\text{int}} = \sum_x \bar{\phi}(x) \left(g_3 A_\mu^{(3)} + g_1 B_\mu \right) \phi(x).$$

This generates:

- color exchange,
- phase rotation,
- mixing between triplet components.

Guided Links: Gauge self-interactions, Triplet dynamics

6.8 Coupling to Geometry

Because the unified $SU(4)$ structure ties the singlet and triplet sectors together, the gauge fields couple to geometry through the singlet amplitude $g(x)$.

The coupling term:

$$S_{\text{mix}} = \sum_x g(x) \bar{\phi}(x) \phi(x)$$

implies:

- gauge excitations distort geometry,
- geometric gradients affect gauge propagation.

This is the discrete analog of curvature-dependent gauge dynamics.

Guided Links: Gauge-geometry coupling, Backreaction effects

7 Worked Phenomenological Examples

This section demonstrates how the unified $SU(4)$ qudit-gauge lattice reproduces familiar physical phenomena. Each example shows how geometry and gauge dynamics emerge from the underlying discrete substrate.

7.1 Gravitational Lensing

Gravitational lensing arises from variations in the singlet field $g(x)$, which modify the effective metric:

$$ds^2 = \sum_{\mu} (1 + \alpha w_{\mu}(x)) dx_{\mu}^2, \quad w_{\mu}(x) = g(x + \hat{\mu})^2 - g(x)^2.$$

Lightlike excitations follow geodesics defined by maximizing the transition amplitude:

$$A(x \rightarrow y) = \psi(y) |U(x, y)| \psi(x).$$

Paths with larger cumulative amplitude correspond to curved trajectories, reproducing lensing behavior.

Guided Links: [Geodesic lensing](ca://s?q=Explain_emergent_lensing), [Metric distortion](ca://s?q=Explain_metric_distortion)

7.2 Coulomb-Like Forces

Triplet excitations interact through the emergent $U(1)$ gauge field $B_\mu(x)$.

The lattice field strength is:

$$F_{\mu\nu}^{(1)}(x) = \partial_\mu B_\nu(x) - \partial_\nu B_\mu(x).$$

In the static limit, the potential between two triplet excitations becomes:

$$V(r) \sim \frac{1}{r},$$

matching the Coulomb form.

Guided Links: [Emergent electromagnetism](ca://s?q=Explain_emergent_electromagnetism), [Coulomb potential](ca://s?q=Explain_Coulomb_potential)

7.3 Scattering Processes

Two triplet excitations ϕ_i and ϕ_j scatter through the exchange of $SU(3)$ and $U(1)$ gauge fields.

The interaction term:

$$S_{\text{int}} = \sum_x \bar{\phi}(x) \left(g_3 A_\mu^{(3)} + g_1 B_\mu \right) \phi(x)$$

produces:

- color exchange,
- phase rotation,
- channel mixing.

The lattice scattering amplitude is:

$$\mathcal{A}(p \rightarrow p') = \sum_x e^{i(p-p')x} \phi'(x) |U(x, x + \hat{\mu})| \phi(x).$$

Guided Links: [Gauge scattering](ca://s?q=Explain_gauge_scattering), [Triplet interactions](ca://s?q=Explain_triplet_interactions)

7.4 Wave Propagation in Curved Geometry

A wave packet in the triplet sector satisfies:

$$D_\mu D^\mu \phi(x) = 0,$$

where:

$$D_\mu = \partial_\mu + ig_3 A_\mu^{(3)} + ig_1 B_\mu.$$

In regions where $g(x)$ varies, the effective metric changes, producing:

- refraction,
- time delay,
- frequency shift.

This reproduces gravitational redshift and Shapiro delay.

Guided Links: [Wave propagation](ca://s?q=Explain_wave_propagation_in_curved_space), [Redshift](ca://s?q=Explain_gravitational_redshift)

7.5 Summary of Phenomenology

The $SU(4)$ qudit–gauge lattice reproduces:

- gravitational lensing from singlet gradients,
- Coulomb-like forces from $U(1)$ dynamics,
- color-like scattering from $SU(3)$ exchange,
- wave propagation in curved geometry,
- unified coupling between geometry and gauge fields.

These examples demonstrate that familiar physical behavior emerges naturally from the discrete qudit substrate.

Guided Links: [Emergent phenomena](ca://s?q=Explain_emergent_phenomena), [Unified dynamics](ca://s?q=Explain_unified_dynamics)

8 Continuum Limit and Effective Field Theory

The $SU(4)$ qudit–gauge lattice is fundamentally discrete, but in the limit of small lattice spacing a , it reproduces continuum field theory. This section formalizes how gauge fields, geometry, and matter fields emerge as smooth continuum objects and how the effective action reduces to familiar forms.

8.1 Scaling Toward the Continuum

As $a \rightarrow 0$, the link variables expand as:

$$U_\mu(x) = \exp(ia \mathcal{A}_\mu(x)),$$

where $\mathcal{A}_\mu(x)$ is the continuum $SU(4)$ gauge potential. The qudit field becomes a smooth function:

$$\psi(x) \rightarrow \psi(x) + a \partial_\mu \psi(x) + \mathcal{O}(a^2).$$

The singlet and triplet components scale as:

$$g(x) = \psi_0(x), \quad \phi_i(x) = \psi_i(x).$$

Guided Links: [Continuum scaling](ca://s?q=Explain_continuum_scaling), [Lattice-to-fieldmapping](ca://s?q=Explain_lattice_to_ffield_mapping)

8.2 Continuum Gauge Fields

The $SU(4)$ gauge field decomposes as:

$$\mathcal{A}_\mu(x) = A_\mu^{(3)}(x) + B_\mu(x) + X_\mu(x),$$

where:

- $A_\mu^{(3)}$ are the $SU(3)$ gauge fields,
- B_μ is the $U(1)$ gauge field,
- X_μ are the coset fields (suppressed at low energies).

The continuum field strengths are:

$$F_{\mu\nu}^{(3)} = \partial_\mu A_\nu^{(3)} - \partial_\nu A_\mu^{(3)} + ig_3[A_\mu^{(3)}, A_\nu^{(3)}],$$

$$F_{\mu\nu}^{(1)} = \partial_\mu B_\nu - \partial_\nu B_\mu.$$

Guided Links: [Continuum gauge fields](ca://s?q=Explain_continuum_gauge_fields), [Fieldstrengthtensors](ca://s?q=Explain_field_strengh_tensors)

8.3 Continuum Geometry

The singlet field $g(x)$ becomes a smooth scalar field. Metric distortions arise from gradients:

$$g_{\mu\nu}(x) = \eta_{\mu\nu} + \alpha \partial_\mu g \partial_\nu g.$$

Curvature emerges from second derivatives:

$$R_{\mu\nu}(x) \sim \partial_\mu \partial_\nu g(x).$$

This produces a scalar-tensor geometric structure.

Guided Links: [Emergent metric](ca://s?q=Explain_emergent_metric), [Scalargeometry](ca://s?q=Explain_scalar_geometry)

8.4 Effective Continuum Action

The lattice action:

$$S = S_{\text{gauge}} + S_{\text{matter}} + S_{\text{geom}}$$

reduces to the continuum effective action:

$$S_{\text{eff}} = \int d^4x \left[-\frac{1}{4} F_{\mu\nu}^{(3)} F_{(3)}^{\mu\nu} - \frac{1}{4} F_{\mu\nu}^{(1)} F_{(1)}^{\mu\nu} + D_\mu \phi^2 + (\partial_\mu g)^2 + V(g) + \lambda R[g] \right].$$

This contains:

- Yang–Mills terms,
- scalar kinetic terms,
- scalar potential,
- curvature coupling.

Guided Links: [Effective action](ca://s?q=Explain_effective_action), [Continuum reduction](ca://s?q=Explain_continuum_reduction)

8.5 Recovery of Known Physics

In the continuum limit, the model reproduces:

- $SU(3)$ gauge theory (color-like interactions),
- $U(1)$ gauge theory (electromagnetic-like interactions),
- scalar–tensor gravity (geometry from $g(x)$),
- geodesic motion from amplitude maximization.

The unified $SU(4)$ structure ensures that these sectors are not independent but emerge from a single discrete substrate.

Guided Links: [Standard interactions](ca://s?q=Explain_standard_interactions), [Emergent gravity](ca://s?q=Explain_emergent_gravity)

8.6 Breakdown of the Continuum Approximation

The continuum description fails when:

- gradients of $g(x)$ become large,
- gauge fields vary on the scale of the lattice spacing,
- high-energy excitations probe the discrete structure,
- coset fields X_μ become relevant.

In this regime, the full lattice theory must be used.

Guided Links: [Continuum breakdown](ca://s?q=Explain_continuum_breakdown), [Lattice corrections](ca://s?q=Explain_lattice_corrections)

9 Predictions and Phenomenology

The unified $SU(4)$ qudit–gauge lattice makes concrete predictions that arise from its discrete structure, symmetry breaking pattern, and emergent continuum behavior. These predictions span gravitational, gauge, and cosmological phenomena and provide potential observational signatures of the underlying substrate.

9.1 Discreteness Signatures

Because the model is fundamentally discrete, it predicts observable effects when physical processes probe the lattice scale.

Possible signatures include:

- **Modified dispersion relations** at high energies,
- **anisotropic propagation** due to lattice directions,
- **frequency-dependent lensing** from discrete curvature,
- **small violations of Lorentz symmetry** near the cutoff scale.

These effects vanish smoothly as $a \rightarrow 0$.

Guided Links: [Discreteness effects](ca://s?q=Explain_discreteness_effects), [Lorentz violation](ca://s?q=Explain_Lorentz_violation)

9.2 Scalar–Tensor Deviations

The emergent metric:

$$g_{\mu\nu}(x) = \eta_{\mu\nu} + \alpha \partial_\mu g \partial_\nu g$$

predicts deviations from pure Einstein gravity.

Observable consequences include:

- **scalar gravitational waves**,
- **modified post-Newtonian parameters**,
- **fifth-force–like corrections** at intermediate scales,
- **environment-dependent gravitational strength**.

These arise because geometry is encoded in a scalar field rather than a fundamental metric.

Guided Links: [Scalar gravity](ca://s?q=Explain_scalar_gravity), [Modified gravity](ca://s?q=Explain_modified_gravity)

9.3 Gauge Sector Unification Patterns

The symmetry breaking:

$$SU(4) \rightarrow SU(3) \times U(1)$$

predicts specific relationships between gauge couplings:

$$g_4 \rightarrow (g_3, g_1),$$

with constraints:

- g_3 and g_1 unify at the lattice scale,
- running couplings diverge from Standard Model predictions at high energies,
- coset fields X_μ may appear as heavy resonances.

These features provide potential collider signatures.

Guided Links: [Gauge unification](ca://s?q=Explain_{gauge_unification}), [Running couplings](ca://s?q = Explain_{running_couplings})

9.4 Discrete Spectral Features

The lattice structure produces quantized spectral features:

- **band-like energy levels** for qudit excitations,
- **momentum quantization** near the Brillouin zone edges,
- **suppressed high-frequency modes**,
- **nonlinear dispersion** for both gauge and geometric waves.

These effects could appear in:

- gravitational wave spectra,
- high-energy cosmic rays,
- early-universe relic radiation.

Guided Links: [Discrete spectra](ca://s?q=Explain_{discrete_spectra}), [Brillouin zone physics](ca://s?q = Explain_{Brillouin_zone_physics})




Figure 2: Flowchart of the geometric electroweak epoch. High curvature suppresses triplet condensation, keeping $SU(2)$ unbroken. As the geometry-mode field relaxes, the effective mass of the triplet becomes negative, triggering $SU(3) \rightarrow SU(2)$ breaking and generating a position-dependent Higgs VEV $v(x)$ that stabilizes as curvature settles.

9.5 Cosmological Implications

The singlet field $g(x)$ acts as a scalar degree of freedom with potential cosmological roles:

- **inflation-like behavior** from $V(g)$,
- **dark-energy-like acceleration** from slow-roll dynamics,
- **dark-matter-like clustering** from oscillatory modes,
- **suppressed small-scale structure** from lattice discreteness.

The unified action naturally couples geometry and gauge fields, producing nontrivial early-universe dynamics.

Guided Links: [Emergent cosmology](ca://s?q=Explain_{emergent}_{cosmology}), [Scalar-field cosmology](ca://s?q=Explain_{scalar field}_{cosmology})

The corresponding local electroweak scale is set by $v(x)$, and the $SU(2)$ gauge boson masses become explicitly geometry-dependent:

$$m_W(x) = \frac{1}{2}g_2 v(x), \quad m_Z(x) = \frac{1}{2}\sqrt{g_2^2 + g_Y^2} v(x), \quad (3)$$

so that curvature, via $g(x)$, directly controls the onset and magnitude of electroweak symmetry breaking.

9.6 Summary of Predictions

The $SU(4)$ qudit-gauge lattice predicts:

- discrete corrections to high-energy propagation,
- scalar-tensor deviations from GR,
- unified gauge coupling relationships,
- quantized spectral features,
- cosmological signatures from the singlet field.

These predictions provide multiple avenues for testing the model across gravitational, gauge, and cosmological regimes.

Guided Links: [Unified predictions](ca://s?q=Explain_{unified}_{predictions}), [Phenomenological tests](ca://s?q=Explain_{phenomenological}_{tests})

10 Cosmological Implications: The Geometric Electroweak Epoch

The coupling between the geometry-mode singlet $g(x)$ and the triplet fields $\phi_i(x)$ suggests that curvature and symmetry breaking may not be independent phenomena in the early universe. In this section we explore, at the effective level, how the relaxation of the singlet field could influence the mass of the triplet sector and potentially trigger a second-stage symmetry breaking. This mechanism is speculative and is introduced as a conceptual possibility rather than a derived prediction of the microscopic lattice model.

10.1 Early Universe: High Curvature and Unbroken $SU(2)$

In the earliest stages of cosmic evolution, the geometry-mode field $g(x)$ is expected to exhibit large amplitude and strong spatial gradients, reflecting a high-curvature, pre-geometric phase of the qudit-gauge substrate. Within the effective potential

$$V_{\text{eff}}(h; g) = m_{\text{eff}}^2(g) |h|^2 + (\lambda_\phi + \gamma) |h|^4, \quad (4)$$

with

$$m_{\text{eff}}^2(g) = m_\phi^2 + \alpha g^2 + \beta g, \quad (5)$$

large values of $|g|$ keep $m_{\text{eff}}^2(g)$ positive. The composite Higgs-like mode $h(x) = \phi_3(x)$ therefore remains uncondensed:

$$\langle h(x) \rangle = 0. \quad (6)$$

In this regime, the triplet sector retains its full $SU(3) \times U(1)$ symmetry, with the $SU(2)$ subgroup acting on (ϕ_1, ϕ_2) unbroken. We emphasize that this structure is internal to the qudit representation and is not identified with the Standard Model electroweak group.

10.2 Relaxation of Geometry and the Onset of Instability

As the universe expands, the singlet field $g(x)$ may relax toward a smoother, lower-curvature configuration. Because the effective mass of the triplet depends on $g(x)$, this relaxation can drive the system toward an instability:

$$m_{\text{eff}}^2(g(x)) \rightarrow 0. \quad (7)$$

If $m_{\text{eff}}^2(g)$ becomes negative in extended regions, the symmetric point $h = 0$ becomes unstable. The composite Higgs-like mode may then condense:

$$\langle h(x) \rangle = v(x) = \sqrt{-\frac{m_{\text{eff}}^2(g(x))}{2(\lambda_\phi + \gamma)}}. \quad (8)$$

This breaks the internal symmetry

$$SU(3) \rightarrow SU(2), \quad (9)$$

with the unbroken $SU(2)$ subgroup acting on (ϕ_1, ϕ_2) . We stress that this breaking pattern is an effective description and is not intended to reproduce the Standard Model electroweak sector.

10.3 The Geometric Electroweak Epoch

The period during which $g(x)$ relaxes and drives the condensation of the composite Higgs-like mode constitutes what we refer to as the *geometric electroweak epoch*. In this exploratory scenario, the Higgs vacuum expectation value becomes explicitly geometry-dependent:

$$v(x) = \sqrt{-\frac{m_\phi^2 + \alpha g(x)^2 + \beta g(x)}{2(\lambda_\phi + \gamma)}}. \quad (10)$$

The corresponding gauge boson masses in the internal $SU(2)$ sector become local functions of curvature:

$$m_W(x) = \frac{1}{2}g_2 v(x), \quad m_Z(x) = \frac{1}{2}\sqrt{g_2^2 + g_Y^2} v(x), \quad (11)$$

where g_2 and g_Y are effective couplings associated with the residual symmetry. These expressions are included to illustrate the structure of the effective theory and are not claimed to match Standard Model values.

Regions with different curvature may undergo the transition at different times, producing domains of broken and unbroken $SU(2)$. Interfaces between these regions act as geometry–symmetry boundaries where both $g(x)$ and $v(x)$ vary rapidly. Such features are reminiscent of domain walls in conventional phase transitions, though here they arise from the unified qudit–gauge substrate.

10.4 Late Universe: Stabilized Geometry and Constant Effective Scale

As $g(x)$ approaches a nearly homogeneous equilibrium value g_0 , the Higgs-like VEV stabilizes to a uniform value:

$$v_0 = \sqrt{-\frac{m_\phi^2 + \alpha g_0^2 + \beta g_0}{2(\lambda_\phi + \gamma)}}. \quad (12)$$

In this effective picture, the internal electroweak-like scale is set by the asymptotic geometry of the universe. Residual fluctuations in $g(x)$ produce small spatial variations in $v(x)$, implying that particle masses in the internal sector may be slightly geometry-dependent. These effects are suppressed in the late universe but could be relevant in regions of extreme curvature.

10.5 Summary and Caveats

The geometric electroweak epoch provides a conceptual illustration of how curvature and symmetry breaking might interact within the unified $SU(4)$ qudit–gauge framework. The key ideas are:

- curvature influences the effective mass of the triplet sector,
- a composite Higgs-like mode may emerge from the qudit triplet,
- symmetry breaking can be triggered by geometric relaxation,
- the effective electroweak-like scale is tied to the asymptotic value of $g(x)$.

We emphasize that this mechanism is exploratory and relies on effective operators rather than a derivation from the microscopic lattice action. Its cosmological implications—such as reheating, baryogenesis, or consistency with observational constraints—remain open questions for future investigation.

11 Conceptual Implications and Comparison to Other Approaches

The $SU(4)$ qudit–gauge lattice provides a unified substrate from which both geometry and gauge interactions emerge. This section compares the framework

to other major approaches to quantum gravity and unification, highlighting both conceptual similarities and key differences.

11.1 Relation to Loop Quantum Gravity

Loop Quantum Gravity (LQG) quantizes geometry directly using spin networks. The present model differs in several fundamental ways:

- Geometry arises from a **scalar singlet field** $g(x)$, not from quantized area/volume operators.
- The underlying structure is a **qudit lattice**, not a spin network.
- Gauge fields and geometry share a **common SU(4) origin**.
- The model naturally includes **gauge interactions**, which LQG typically adds separately.

However, both frameworks share:

- a fundamentally discrete substrate,
- background independence in the continuum limit,
- geometric emergence from non-geometric degrees of freedom.

Guided Links: [Loop Quantum Gravity](ca://s?q=Explain_L*QG_comparison*), [Spinnetworks](ca://s?q=Explain_s*pin_networks*)

11.2 Relation to String Theory

String theory posits that all particles arise from vibrational modes of fundamental strings. The present model differs conceptually:

- No extended objects; only **qudits and link variables**.
- No requirement for extra dimensions.
- Geometry is **emergent**, not built into the background.
- Gauge fields arise from **SU(4) decomposition**, not from string excitations.

Yet there are parallels:

- unified gauge and geometric sectors,
- scalar fields playing geometric roles,
- effective low-energy actions resembling scalar–tensor gravity.

Guided Links: [String theory comparison](ca://s?q=Explain_s*string_theory_comparison*), [Emergent geometry](ca://s?q=Explain_e*emergent_geometry_theory*)

11.3 Relation to Tensor Networks

Tensor network approaches (MERA, PEPS, etc.) model spacetime as an emergent structure from entanglement patterns.

Connections include:

- geometry emerges from **local quantum degrees of freedom**,
- curvature corresponds to **distortions in local structure**,
- gauge fields arise from **symmetry constraints**.

Differences:

- The qudit lattice is a **physical substrate**, not a computational ansatz.
- Gauge fields are **explicit SU(4) link variables**, not emergent from entanglement alone.

Guided Links: [Tensor networks](ca://s?q=Explain_tensor_networks), [Entanglement geometry](ca://s?q=Explain_entanglement_geometry)

11.4 Relation to Lattice Gauge Theory

The model is structurally closest to lattice gauge theory (LGT), but with key extensions:

- LGT includes only gauge fields; here, **geometry also emerges**.
- The qudit field plays a dual role: matter + geometric scalar.
- The gauge group is **SU(4)**, not chosen arbitrarily.
- The singlet–triplet decomposition produces **gravity + gauge** from one structure.

Thus the model can be viewed as a **geometric extension of LGT**.

Guided Links: [Lattice gauge theory](ca://s?q=Explain_LGT_comparison), [Wilson action](ca://s?q=Explain_Wilson_action)

11.5 Relation to Quantum Information Approaches

Quantum information approaches often treat spacetime as emergent from:

- entanglement entropy,
- quantum error correction,
- tensor factorization structure.

The present model aligns with this philosophy:

- qudits are the fundamental degrees of freedom,
- geometry is encoded in **local amplitudes**,
- gauge interactions arise from **unitary constraints**.

But it differs in that:

- the model is a **physical theory**, not an information-theoretic analogy,
- the $SU(4)$ structure is **dynamical**, not imposed.

Guided Links: [Quantum information geometry](ca://s?q=Explain_Q*I_ggeometry*), [Error-correctingspacetime](ca://s?q = Explain_Q*EC_spacetime*)

11.6 Philosophical Implications

The model suggests:

- spacetime is not fundamental but **emergent**,
- geometry is a **collective phenomenon** of qudit amplitudes,
- gauge forces and gravity share a **common origin**,
- discreteness is a **physical feature**, not a computational tool.

This places the theory in the broader category of **emergent physics**, where macroscopic laws arise from microscopic quantum structure.

Guided Links: [Emergent spacetime](ca://s?q=Explain_e*emergent_spacetime*), [Foundationsofgeometry](ca://s?q = Explain_f*foundations_of_ggeometry*)

11.7 Summary of Conceptual Positioning

The $SU(4)$ qudit-gauge lattice:

- shares discreteness with LQG,
- shares unification with string theory,
- shares entanglement structure with tensor networks,
- shares gauge dynamics with lattice gauge theory,
- shares information-theoretic intuition with QI approaches,

but remains distinct in its:

- unified $SU(4)$ origin of geometry and gauge fields,
- qudit-based substrate,
- scalar geometric field,
- emergent continuum behavior.

Guided Links: [Conceptual summary](ca://s?q=Explain_c*conceptual_ssummary*), [Unifiedframework](ca://s?q = Explain_u*nified_fframework*)

12 Conclusion

The $SU(4)$ qudit–gauge lattice provides a speculative but coherent framework in which geometry, gauge interactions, and matter-like excitations emerge from a single discrete quantum substrate. A four-level qudit at each lattice site, together with $SU(4)$ link variables, contains the structure needed to generate a geometric singlet sector, a gauge triplet sector, and effective couplings between them. While highly simplified, this construction illustrates how gravitational and gauge phenomena may arise from a unified microscopic origin.

12.1 Summary of the Framework

The model is built from a small set of ingredients:

- A discrete hypercubic lattice with spacing a .
- A four-level qudit field $\psi(x)$ at each site.
- $SU(4)$ link variables $U_\mu(x)$ on each oriented edge.
- A symmetry breaking pattern:

$$SU(4) \rightarrow SU(3) \times U(1) \rightarrow SU(2) \times U(1).$$

- A singlet field $g(x)$ encoding geometric information.
- A triplet field $\phi_i(x)$ encoding gauge interactions and a composite Higgs-like mode.

These elements yield an emergent scalar–tensor geometry and gauge dynamics in the continuum limit.

12.2 Emergent Geometry and Gauge Dynamics

Geometry arises from the singlet component

$$g(x) = \psi_0(x),$$

with metric-like behavior encoded in gradients of $g(x)$ and curvature arising from second differences. The resulting continuum theory resembles a scalar–tensor model rather than general relativity, and no spin-2 graviton or diffeomorphism invariance is recovered at this stage.

The triplet fields

$$\phi_i(x) = \psi_i(x), \quad i = 1, 2, 3,$$

transform under the residual $SU(3) \times U(1)$ symmetry. Gauge fields emerge from the decomposition of the $SU(4)$ link variables, and the continuum limit produces $SU(3)$ and $U(1)$ gauge dynamics with suppressed coset fields.

12.3 Effective Symmetry Breaking and Cosmology

At the effective level, couplings between the singlet and triplet sectors allow the local geometric density to influence the mass of the triplet fields. We explored the possibility that the relaxation of curvature in the early universe could trigger a composite Higgs-like condensation, leading to a geometric electroweak epoch. This mechanism is speculative and relies on effective operators consistent with the symmetries of the model rather than a derivation from the microscopic lattice action.

12.4 Limitations and Open Directions

Several important aspects of a complete physical theory are not yet addressed:

- The emergent geometry is scalar–tensor and does not reproduce the Einstein–Hilbert action or diffeomorphism invariance.
- Fermions, chiral structure, and Standard Model quantum numbers are not included.
- The effective potential used for symmetry breaking is not derived from the microscopic $SU(4)$ action.
- The $SU(3) \rightarrow SU(2)$ breaking pattern does not reproduce the Standard Model electroweak sector.
- Quantitative predictions (e.g., Newtonian limit, lensing angles, coupling constants) remain to be developed.
- Cosmological implications such as reheating, baryogenesis, and consistency with observational constraints have not been analyzed.

These limitations highlight the speculative nature of the framework and point toward several directions for future work.

12.5 Final Perspective

Despite its simplifications, the $SU(4)$ qudit–gauge lattice offers a conceptually unified setting in which geometry and gauge interactions arise from the same microscopic degrees of freedom. It provides a concrete example of how spacetime and forces might emerge from a discrete quantum substrate and suggests new avenues for exploring the quantum structure of geometry and interaction. Further development is required to assess whether this framework can be extended toward a more complete description of fundamental physics.

13 Discussion

The $SU(4)$ qudit–gauge lattice explored in this work provides a concrete example of how geometry and gauge interactions might emerge from a common microscopic substrate. While highly simplified, the model illustrates several conceptual themes that appear across a wide range of approaches to quantum gravity and unification.

13.1 Emergent Geometry from Non-Geometric Degrees of Freedom

A central feature of the framework is that geometric behavior arises from the dynamics of a scalar singlet field rather than from a fundamental metric. This places the model within the broader class of emergent geometry proposals, where spacetime is not assumed but reconstructed from more primitive quantum variables. The resulting continuum theory resembles a scalar–tensor model, highlighting both the promise and the limitations of this approach. In particular, the absence of diffeomorphism invariance and spin-2 excitations underscores the need for additional structure if one hopes to recover general relativity or a close analogue.

13.2 Gauge Interactions from Internal Structure

The triplet sector of the qudit field naturally supports non-Abelian and Abelian gauge interactions after the breaking $SU(4) \rightarrow SU(3) \times U(1)$. This demonstrates how gauge fields can emerge from internal degrees of freedom on a discrete substrate, consistent with insights from lattice gauge theory, tensor networks, and quantum information approaches. The model also suggests a route toward composite Higgs-like behavior, though the effective potential used here is not derived from the microscopic action.

13.3 Interplay Between Geometry and Symmetry Breaking

One of the more speculative aspects of the framework is the possibility that geometric relaxation in the early universe could influence the effective mass of the triplet sector and thereby trigger an internal symmetry-breaking transition. While this “geometric electroweak epoch” is introduced at the effective level, it illustrates how curvature and gauge dynamics might be coupled in a unified substrate. Further work is needed to determine whether such a mechanism can be derived from the underlying $SU(4)$ structure or whether alternative symmetry-breaking patterns yield closer connections to the Standard Model.

13.4 Position Within the Landscape of Emergent Frameworks

The model shares features with several major approaches:

- With loop quantum gravity and causal dynamical triangulations, it shares a fundamentally discrete substrate.
- With tensor network models, it shares the idea that geometry emerges from local quantum degrees of freedom.
- With lattice gauge theory, it shares the use of link variables and plaquette dynamics.
- With quantum information approaches, it shares the use of qudits as fundamental building blocks.

At the same time, the unified $SU(4)$ structure and the singlet–triplet decomposition give the model a distinctive character. It provides a compact setting in which geometry, gauge interactions, and symmetry breaking arise from a single representation.

13.5 Outlook

The present work should be viewed as a conceptual starting point. Many questions remain open, including the incorporation of fermions, the emergence of chiral structure, the derivation of effective couplings, and the recovery of Lorentz symmetry and spin-2 modes. Nonetheless, the model offers a coherent framework for exploring how spacetime and interactions might emerge from a unified quantum substrate, and it suggests several promising directions for future investigation.

14 Limitations and Future Work

The $SU(4)$ qudit–gauge lattice presented in this work is intended as a conceptual framework rather than a complete physical theory. Several important limitations remain, and addressing them will be essential for assessing the viability of the approach.

14.1 Limitations

Scalar–tensor nature of emergent geometry. The geometric sector arises from a single scalar field $g(x)$, leading to a scalar–tensor model rather than general relativity. The framework does not recover the Einstein–Hilbert action, diffeomorphism invariance, or a spin-2 graviton.

Absence of fermions and chiral structure. The model currently contains only bosonic degrees of freedom. A realistic unified theory must incorporate fermions, chiral asymmetry, anomaly cancellation, and Standard Model quantum numbers.

Effective nature of symmetry breaking. The composite Higgs-like potential is introduced at the effective level and is not derived from the microscopic $SU(4)$ lattice action. The parameters $(m_\phi^2, \alpha, \beta, \gamma)$ remain phenomenological.

Non-Standard-Model breaking pattern. The internal breaking $SU(3) \rightarrow SU(2)$ does not reproduce the Standard Model electroweak group. The relationship between the internal $SU(2)$ and physical weak interactions remains unclear.

Lack of quantitative predictions. While qualitative phenomena such as lensing and Coulomb-like forces can be illustrated, quantitative predictions (e.g., Newtonian limit, coupling constants, cosmological constraints) remain undeveloped.

Cosmological uncertainties. The geometric electroweak epoch is exploratory. Its implications for reheating, baryogenesis, dark matter, and nucleosynthesis have not been analyzed.

14.2 Future Work

Several directions appear promising:

- Deriving effective couplings from the microscopic $SU(4)$ lattice action.
- Incorporating fermions and exploring mechanisms for chiral emergence.
- Developing quantitative predictions for gravitational and gauge observables.
- Studying the continuum limit more systematically, including possible emergent spin-2 modes.
- Exploring cosmological dynamics, including reheating and domain wall evolution.
- Investigating whether alternative symmetry-breaking patterns within $SU(4)$ yield closer connections to the Standard Model.

These open directions highlight the speculative nature of the present framework while identifying concrete steps toward a more complete theory.

Appendices

15 Appendix A: SU(4) Algebra and Representation Theory

This appendix summarizes the algebraic structure of $SU(4)$ used throughout the main text.

15.1 A.1 Generators

The generators T^A of $SU(4)$ satisfy:

$$[T^A, T^B] = if^{ABC}T^C, \quad \text{Tr}(T^AT^B) = \frac{1}{2}\delta^{AB}.$$

They decompose under the breaking:

$$SU(4) \rightarrow SU(3) \times U(1)$$

into:

- 8 generators of $SU(3)$,
- 1 generator of $U(1)$,
- 6 coset generators.

Guided Links: [SU(4) algebra](ca://s?q=Explain_{SU4algebra}), [Representation theory](ca://s?q=Explain_{representationtheory})

15.2 A.2 Fundamental Representation

The fundamental representation is:

$$\mathbf{4} = 1 \oplus 3,$$

corresponding to the singlet $g(x)$ and triplet $\phi_i(x)$ fields.

Guided Links: [Fundamental rep](ca://s?q=Explain_{fundamentalrepresentation}), [Singlet-triplet split](ca://s?q=Explain_{singlettripletsplit})

16 Appendix B: Lattice Conventions

16.1 B.1 Coordinates and Directions

We use a hypercubic lattice with spacing a and unit vectors $\hat{\mu}$.

Forward and backward shifts:

$$x + \hat{\mu}, \quad x - \hat{\mu}.$$

16.2 B.2 Link Variables

Link variables live on oriented edges:

$$U_\mu(x) \in SU(4).$$

Backward links:

$$U_{-\mu}(x) = U_\mu^\dagger(x - \hat{\mu}).$$

16.3 B.3 Plaquette Definition

The elementary plaquette is:

$$U_{\mu\nu}(x) = U_\mu(x)U_\nu(x + \hat{\mu})U_\mu^\dagger(x + \hat{\nu})U_\nu^\dagger(x).$$

Guided Links: [Lattice basics](ca://s?q=Explain_{lattice}basics), [Plaquetteoperator](ca://s?q=Explain_plaquetteoperator)

17 Appendix C: Continuum Limit Expansions

17.1 C.1 Link Expansion

For small a :

$$U_\mu(x) = \exp(ia \mathcal{A}_\mu(x)) = 1 + ia \mathcal{A}_\mu(x) - \frac{a^2}{2} \mathcal{A}_\mu^2(x) + \dots$$

17.2 C.2 Plaquette Expansion

$$U_{\mu\nu}(x) = \exp(ia^2 F_{\mu\nu}(x)) = 1 + ia^2 F_{\mu\nu}(x) + \mathcal{O}(a^3).$$

Guided Links: [Continuum expansion](ca://s?q=Explain_continuumexpansion), [Fieldstrength](ca://s?q=Explain_fieldstrengthexpansion)

18 Appendix D: Numerical Simulation Methods

18.1 D.1 Update Algorithms

Typical update schemes include:

- heatbath updates for $SU(4)$ links,
- overrelaxation steps,
- Metropolis acceptance for qudit fields.

18.2 D.2 Observables

Key observables include:

- plaquette expectation values,
- correlation functions of $g(x)$,
- triplet propagators,
- curvature estimators.

Guided Links: [Lattice numerics](ca://s?q=Explain_{lattice_numerics}), [MonteCarlo methods](ca://s?q=Explain_{MonteCarlo_methods})

19 Appendix E: Symbol Glossary

Symbol	Meaning
$\psi(x)$	Qudit field
$g(x)$	Singlet (gravity-mode) component
$\phi_i(x)$	Triplet (gauge-mode) components
$U_\mu(x)$	$SU(4)$ link variable
$U_{\mu\nu}(x)$	Plaquette operator
$A_\mu^{(3)}$	$SU(3)$ gauge field
B_μ	$U(1)$ gauge field
X_μ	Coset fields
a	Lattice spacing
$F_{\mu\nu}$	Field strength tensor

Table 1: Glossary of symbols used throughout the text.

Guided Links: [Symbol glossary](ca://s?q=Explain_{symbol_glossary}), [Notation reference](ca://s?q=Explain_{notation_reference})

References

- [1] M. B. Green, J. H. Schwarz, and E. Witten. *Superstring Theory, Vol. 1: Introduction*. Cambridge University Press, 1987.
- [2] J. Polchinski. *String Theory, Vol. 1: An Introduction to the Bosonic String*. Cambridge University Press, 1998.
- [3] C. Rovelli. Loop quantum gravity. *Living Reviews in Relativity*, 1:1, 1998.
- [4] A. Ashtekar and J. Lewandowski. Background independent quantum gravity: A status report. *Classical and Quantum Gravity*, 21:R53–R152, 2004.

- [5] H. Georgi and S. L. Glashow. Unity of all elementary particle forces. *Physical Review Letters*, 32:438–441, 1974.
- [6] H. Georgi. The state of the art—gauge theories. *AIP Conference Proceedings*, 23:575–582, 1974.
- [7] K. G. Wilson. Confinement of quarks. *Physical Review D*, 10:2445–2459, 1974.
- [8] M. Creutz. *Quarks, Gluons and Lattices*. Cambridge University Press, 1983.
- [9] T. Durt, B.-G. Englert, I. Bengtsson, and K. Życzkowski. On mutually unbiased bases. *International Journal of Quantum Information*, 8:535–640, 2010.
- [10] A. B. Klimov, L. L. Sánchez-Soto, and H. de Guise. Generalized coherent states for $su(n)$ systems. *Journal of Physics A: Mathematical and Theoretical*, 42:485303, 2009.
- [11] C. Rovelli and L. Smolin. Spin networks and quantum gravity. *Physical Review D*, 52:5743–5759, 1995.
- [12] J. Ambjørn, J. Jurkiewicz, and R. Loll. Dynamically triangulating lorentzian quantum gravity. *Nuclear Physics B*, 610:347–382, 2001.
- [13] B. Swingle. Entanglement renormalization and holography. *Physical Review D*, 86:065007, 2012.
- [14] T. Jacobson. Thermodynamics of spacetime: The einstein equation of state. *Physical Review Letters*, 75:1260–1263, 1995.
- [15] R. Bousso. The holographic principle. *Reviews of Modern Physics*, 74:825–874, 2002.
- [16] A. Ashtekar and J. Lewandowski. Background independent quantum gravity: A status report. *Classical and Quantum Gravity*, 21:R53–R152, 2004.
- [17] J. Ambjørn, J. Jurkiewicz, and R. Loll. Reconstructing the universe. *Physical Review D*, 72:064014, 2005.
- [18] A. Perez. The spin foam approach to quantum gravity. *Living Reviews in Relativity*, 16:3, 2013.
- [19] X.-L. Qi. Exact holographic mapping and emergent space-time geometry. *Nature Physics*, 9:284–291, 2013.
- [20] J. C. Pati and A. Salam. Lepton number as the fourth color. *Physical Review D*, 10:275–289, 1974.
- [21] H. Georgi. The state of the art—gauge theories. *AIP Conference Proceedings*, 23:575–582, 1974.

- [22] F. Gürsey and P. Ramond. E6 gauge theory and the standard model. *Physical Review D*, 24:447–448, 1981.
- [23] S. Ryu and T. Takayanagi. Holographic derivation of entanglement entropy from ads/cft. *Physical Review Letters*, 96:181602, 2006.
- [24] M. Van Raamsdonk. Building up spacetime with quantum entanglement. *General Relativity and Gravitation*, 42:2323–2329, 2010.