

Geometric Foundations of the Selection-Stitch Model: Deriving c , G_N , l_P , Lorentz Invariance, and the Decoherence Threshold from $K = 12$ Lattice Topology

Raghu Kulkarni

SSMTheory Group, IDrive Inc., Calabasas, CA 91302, USA

Abstract

We present the geometric foundations of the Selection-Stitch Model (SSM) [1], in which the vacuum is a discrete Face-Centered Cubic ($K = 12$) tensor network. From this single structure, we derive five fundamental quantities that the Standard Model and General Relativity treat as independent empirical inputs: (1) the speed of light, $c = 4v_{\text{lattice}}$, from the structure tensor eigenvalue of the cuboctahedral unit cell [15]; (2) the lattice spacing $L = \sqrt{2\sqrt{6}\ln 2}l_P \approx 1.84l_P$, from the intersection of the vacuum's elastic energy density with the Planck density; (3) Newton's gravitational constant, $G_N = \sqrt{2/3}L^2/(4\ln 2) \approx 1.0l_P^2$, from the exact Ryu-Takayanagi relation on the hexagonal boundary [7, 8]; (4) exact Lorentz invariance $\text{SO}(3,1)$, inherited holographically from the continuous $\text{SO}(2)$ symmetry of the $K = 6$ boundary sheets via four $\{111\}$ stacking planes [4, 5]; and (5) a two-step quantum decoherence threshold at $m_{\text{soft}} \approx 11.8\mu\text{g}$ and $m_{\text{hard}} \approx 20.5\mu\text{g}$, from the intersection of the Compton wavelength with the lattice spacing and the metric wall [6]. We compare all predictions against experimental data in a comprehensive table spanning 15 observables, including bounding recent macroscopic optomechanical superpositions [9]. No free parameters are introduced.

Keywords: Discrete Spacetime, Lorentz Invariance, Planck Scale, Quantum Decoherence, Holographic Principle, Tensor Networks

1. Introduction

The Standard Model of particle physics and General Relativity together contain approximately 25 free parameters—coupling constants, mass ratios, and cosmological constants that must be measured experimentally. The Selection-Stitch Model (SSM) [1] proposes that these parameters are not free: they are geometric invariants of a discrete vacuum lattice.

In this paper, we consolidate the foundational derivations of the SSM into a single, self-contained presentation. The starting point is a single structural assumption:

Email address: raghu@idrive.com (Raghu Kulkarni)

The vacuum is a Face-Centered Cubic (FCC) tensor network with coordination number $K = 12$, where each bond is a maximally entangled Bell pair.

From this assumption alone—with no additional inputs, no continuous fields, and no free parameters—we derive the five foundational scales of physics listed in the abstract. We then compare every prediction against experimental data.

Interactive 3D visualizations. To immediately ground the geometry discussed in this paper, readers can explore the foundational mappings of the SSM through two interactive WebGL applications:

1. Vacuum Phase Transitions: The $K = 6 \rightarrow K = 4 \rightarrow K = 12$ topological relaxation, explicitly illustrating the emergence of the bulk tensor network:

https://raghu91302.github.io/ssmtheory/ssm_regge_deficit.html

2. Holographic Emergence and Symmetry: An interactive 3D construction demonstrating the stacking of 2D hexagonal boundary sheets to form the 3D $K = 12$ FCC bulk. It visualizes the emergence of the 12-coordinated cuboctahedron and the four $\{111\}$ continuous stacking planes:

https://raghu91302.github.io/ssmtheory/ssm_lorentz_holographic.html

2. The Vacuum Lattice

Definition 1 (The SSM Vacuum). *The vacuum is a three-dimensional tensor network with the following properties:*

- **Lattice type:** *Face-Centered Cubic (FCC)—the unique 3D lattice saturating the Kepler bound [3].*
- **Coordination number:** $K = 12$ nearest neighbours per node.
- **Bond structure:** *Each bond is a maximally entangled Bell pair $|\Phi^+\rangle = (|00\rangle + |11\rangle)/\sqrt{2}$.*
- **Local geometry:** *The 12 neighbours of any node form a **cuboctahedron**—a polyhedron with 8 triangular faces and 6 square faces.*
- **Degrees of freedom:** *Each node has translational displacement u^i and microrotation θ^i (Cosserat elasticity).*
- **Bond decomposition:** $K = S_{\text{trans}} + S_{\text{tors}} = 4 + 8$, where $S_{\text{trans}} = 4$ translational bonds carry gravity and electromagnetism, and $S_{\text{tors}} = 8$ torsional bonds carry the dark sector. This decomposition follows from standard Cosserat (micropolar) elasticity applied to the FCC bond topology [11].

2.1. Origin: the $K = 6 \rightarrow K = 4 \rightarrow K = 12$ phase transition

The $K = 12$ FCC vacuum is not an *ad hoc* assumption. It arises from a topological phase transition in the entanglement network [1]. The sequence is:

Stage 1: $K = 6$ ground state. The 2D hexagonal sheet with $K = 6$ is the unique planar network satisfying the Euler characteristic constraint $\chi = V(1 - K/6) = 0$ for a flat, infinite surface. This is the topological ground state: zero curvature, maximum entropy, minimum free energy.

Stage 2: $K = 4$ tetrahedral nucleation. Under cooling (analogous to crystallization), the 2D sheet begins to buckle into 3D. The first stable 3D structure is the tetrahedron ($K = 4$), which has the minimum coordination for a rigid 3D framework. The Regge deficit angle of the tetrahedron is $\delta = 2\pi - 5 \arccos(1/3) \approx 0.128$ rad, which drives exponential inflation.

Stage 3: $K = 12$ saturation. As the crystallization wave propagates, tetrahedral clusters pack together. The densest possible packing of tetrahedra and octahedra fills 3D space with FCC symmetry, saturating the Kepler bound at $K = 12$. This is the vacuum we observe today.

The SSM thus has a *dynamical* origin: the vacuum is not placed on a lattice by hand; it crystallizes into the $K = 12$ FCC structure via a geometric phase transition from the $K = 6$ boundary.

Definition 2 (The Holographic Boundary). *The 3D FCC bulk is not fundamental. It is the holographic projection of a 2D boundary: a continuous hexagonal ($K = 6$) entanglement network—the pre-crystallization vacuum state. The Lift operator projects this 2D sheet into 3D via ABC stacking at height $h = \sqrt{2/3}L$, building the FCC structure layer by layer.*

The metric wall is the geometric exclusion limit of the lattice: under extreme compression, nodes cannot approach closer than $r_{\min} = L/\sqrt{3}$ without destroying the network topology. This exclusion radius is the geometric consequence of sphere packing at the Kepler limit [3], verified by direct kinematic simulation of the lattice framework [2].

The geometric relationship between the 2D boundary sheets and the emergent 3D bulk is illustrated in Figure 1.

3. The Speed of Light

3.1. The 12 FCC bond vectors

The FCC lattice has 12 nearest-neighbour bond vectors, each connecting a node to one of its 12 cuboctahedral neighbours. In units of the lattice parameter a (where the nearest-neighbour distance is $a/\sqrt{2}$), the normalized bond vectors are:

$$\hat{n}_j \in \left\{ \frac{1}{\sqrt{2}}(\pm 1, \pm 1, 0), \frac{1}{\sqrt{2}}(\pm 1, 0, \pm 1), \frac{1}{\sqrt{2}}(0, \pm 1, \pm 1) \right\}. \quad (1)$$

There are $\binom{3}{2} \times 2^2 = 12$ vectors, confirming $K = 12$. These vectors point toward the 12 vertices of the cuboctahedron.

3.2. The structure tensor

Consider a field $\psi(\mathbf{x})$ propagating through the lattice via nearest-neighbour hopping [15]. The discrete Dirac operator sums over the 12 FCC bond vectors:

$$D_{\text{SSM}}(\mathbf{k}) = \frac{1}{\tau} \sum_{j=1}^{12} (\gamma^\mu \hat{n}_{j,\mu}) e^{i\mathbf{k} \cdot \mathbf{n}_j}, \quad (2)$$

where τ is the discrete time step. In the long-wavelength limit ($|\mathbf{k}|L \ll 1$), expanding to first order in $\mathbf{k}$, the kinetic term depends on the second-rank **structure tensor**:

$$S^{\mu\nu} = \sum_{j=1}^{12} \hat{n}_j^\mu \hat{n}_j^\nu. \quad (3)$$

We compute this explicitly. For the xx component:

$$S^{xx} = \sum_{j=1}^{12} (\hat{n}_j^x)^2 = 4 \times \left(\frac{1}{\sqrt{2}}\right)^2 + 4 \times \left(\frac{1}{\sqrt{2}}\right)^2 + 4 \times 0 = 2 + 2 + 0 = 4. \quad (4)$$

By the cubic symmetry of FCC, $S^{yy} = S^{zz} = 4$. For the off-diagonal components:

$$S^{xy} = \sum_{j=1}^{12} \hat{n}_j^x \hat{n}_j^y = \frac{1}{2} \left[(+1)(+1) + (+1)(-1) + (-1)(+1) + (-1)(-1) \right] + 0 + 0 = 0. \quad (5)$$

Therefore:

$$\boxed{S^{\mu\nu} = 4\delta^{\mu\nu}.} \quad (6)$$

The factor of 4 is exact and unique to the FCC lattice. For comparison, the BCC lattice ($K = 8$) gives $S^{\mu\nu} = (8/3)\delta^{\mu\nu}$, and the simple cubic ($K = 6$) gives $S^{\mu\nu} = 2\delta^{\mu\nu}$. The FCC value of 4 is the largest possible for any Bravais lattice.

3.3. Speed of light as geometric renormalization

The long-wavelength dispersion relation extracted from D_{SSM} is:

$$\omega^2 = \frac{S^{\mu\nu} k_\mu k_\nu}{\tau^{-2}} = \frac{4|\mathbf{k}|^2}{\tau^{-2}} \implies \omega = 2 \frac{|\mathbf{k}|}{\tau^{-1}}. \quad (7)$$

The phase velocity is $v_{\text{phase}} = \omega/|\mathbf{k}| = 2/\tau^{-1}$. Identifying this with the macroscopic speed of light and using $v_{\text{lattice}} = L/\tau$:

$$\boxed{c = 4 \left(\frac{L}{\tau} \right) = 4 v_{\text{lattice}}.} \quad (8)$$

Light is a collective excitation moving exactly four times faster than the underlying lattice update rate. This factor of 4 is not arbitrary; it is the eigenvalue of the FCC structure tensor, which in turn is a direct consequence of $K = 12$ coordination.

3.4. Vanishing of Lorentz-violating corrections

The Taylor expansion of the discrete operator generates higher-order tensors:

First order (mass term): $\sum_{j=1}^{12} \hat{n}_j^\mu = 0$ by inversion symmetry of FCC. No lattice-induced mass is generated.

Third order (Lorentz violation): $\sum_{j=1}^{12} \hat{n}_j^\mu \hat{n}_j^\nu \hat{n}_j^\lambda = 0$ by the centrosymmetry of the cuboctahedron. This is the critical result: the leading Lorentz-violating correction vanishes identically.

Fourth order (first nonzero correction): The fourth-rank tensor $\sum \hat{n}_j^\mu \hat{n}_j^\nu \hat{n}_j^\lambda \hat{n}_j^\sigma$ is nonzero but suppressed by $(kL)^2 \sim (E/E_P)^2$. At optical frequencies ($E \sim 1$ eV, $E_P \sim 10^{28}$ eV), this gives corrections of order 10^{-56} , far below the experimental bound of 10^{-16} [4].

4. The Lattice Spacing and the Planck Scale

4.1. Maximum energy density

Each lattice node updates once per time step τ . The maximum energy a single node can transmit is therefore bounded by the uncertainty principle:

$$E_{\max} = \frac{\hbar}{\tau} = \frac{\hbar c}{4L}, \quad (9)$$

where we used $c = 4L/\tau$ from Eq. 8. The FCC lattice has 4 atoms per conventional cubic cell of side a . The volume per node is:

$$V_{\text{node}} = \frac{a^3}{4} = \frac{L^3}{\sqrt{2}}, \quad (10)$$

where $a = L\sqrt{2}$ is the conventional cell parameter (the nearest-neighbour distance in FCC is $a/\sqrt{2} = L$). The maximum energy density the lattice can support is:

$$\rho_{\text{lattice}} = \frac{E_{\max}}{V_{\text{node}}} = \frac{\hbar c/(4L)}{L^3/\sqrt{2}} = \frac{\sqrt{2} \hbar c}{4L^4}. \quad (11)$$

4.2. The metric wall

Under extreme gravitational compression, the lattice nodes are forced together. However, the FCC geometry imposes an absolute minimum approach distance: the **metric wall** at

$$r_{\min} = \frac{L}{\sqrt{3}}. \quad (12)$$

This is the inscribed sphere radius of the tetrahedron formed by four mutually touching spheres of diameter L . Below $r_{\min}$, the tetrahedral voids collapse and the network topology is destroyed. This is not a soft potential; it is a hard geometric exclusion. The lattice cannot be compressed further without shattering into a lower-dimensional structure (dimensional reduction to 2D sheets).

4.3. Saturation at the Planck density

Setting $\rho_{\text{lattice}} = \rho_P = c^7/(\hbar G^2)$:

$$\frac{\sqrt{2}\hbar c}{4L^4} = \frac{c^7}{\hbar G^2} \implies L = \left(\frac{\sqrt{2}}{4}\right)^{1/4} \sqrt{\frac{\hbar G}{c^3}}. \quad (13)$$

Combining this with the Ryu-Takayanagi boundary entanglement entropy (Section 5) gives the refined value:

$$L = \sqrt{2\sqrt{6}\ln 2} l_P \approx 1.84 l_P \approx 2.97 \times 10^{-35} \text{ m}. \quad (14)$$

The Planck scale is not an arbitrary UV cutoff; it is the physical limit where vacuum compression reaches the metric wall.

5. Newton's Constant from the Ryu-Takayanagi Relation

5.1. The Stitch operator and boundary state

Each bond of the hexagonal boundary is a maximally entangled Bell pair. Formally, the **Stitch operator** S_{ij} acting on adjacent nodes i and j (each carrying a local Hilbert space $\mathcal{H}_i \cong \mathbb{C}^d$ with orthonormal basis $\{|k\rangle_i\}_{k=0}^{d-1}$) is the projector:

$$S_{ij} = |\Phi^+\rangle\langle\Phi^+|_{ij}, \quad |\Phi^+\rangle_{ij} = \frac{1}{\sqrt{d}} \sum_{k=0}^{d-1} |k\rangle_i \otimes |k\rangle_j. \quad (15)$$

For the minimal SSM, $d = 2$ (qubit per node), so $|k\rangle \in \{|0\rangle, |1\rangle\}$ and $|\Phi^+\rangle = (|00\rangle + |11\rangle)/\sqrt{2}$. The complete boundary state is:

$$|\Psi_{\text{bdy}}\rangle = \prod_{(i,j) \in \text{edges}} S_{ij} |0\rangle^{\otimes N}, \quad (16)$$

i.e., start with all N nodes in $|0\rangle$, then project each adjacent pair onto $|\Phi^+\rangle$. The result is a highly entangled N -party state where every adjacent pair shares one ebit. This is a stabilizer tensor network in the same mathematical class as the HaPPY holographic error-correcting code [8].

5.2. Boundary entanglement entropy

Theorem 1 (Exact Entanglement Entropy). *For a connected boundary region A with perimeter P , the von Neumann entanglement entropy is exactly $S_A = n_{\text{cut}} \ln d$, where $n_{\text{cut}} = P/L$ is the number of severed stitches.*

Proof. A stitch entirely within A or entirely within $\bar{A}$ contributes zero entanglement across the cut. A stitch with one endpoint in A and the other in $\bar{A}$ is severed. By the Schmidt decomposition of a Bell pair, tracing out one qudit yields the maximally mixed state $\rho = I/d$ with entropy $\ln d$. Since the stitches are independent, $S_A = n_{\text{cut}} \ln d$. $\square$

On the hexagonal lattice with spacing L , $n_{\text{cut}} = P/L$. For qubits ($d = 2$):

$$S_A = \frac{P}{L} \ln 2. \quad (17)$$

This is the holographic area law (manifesting as a perimeter law in 2D). Figure 2 illustrates the bond-cutting geometry.

5.3. The minimal bulk surface

The Lift operator constructs the FCC bulk by stacking hexagonal sheets at height $h = \sqrt{2/3} L$.

Theorem 2 (Minimality of the One-Layer Curtain). *Among all bulk surfaces homologous to ∂A , the one-layer curtain with height $h = \sqrt{2/3} L$ achieves the minimum area: $\text{Area}(\gamma_A) = P \times \sqrt{2/3} L$.*

Proof. Any surface homologous to ∂A must form a closed wall separating the bulk interior of A from $\bar{A}$. Surfaces shallower than h fail the homology condition because inter-layer bonds at height h still connect the two regions. Surfaces deeper than h have area $\geq P \times nh$ ($n \geq 2$), which is strictly larger. Non-planar zigzag surfaces at height h have path lengths $> P$ along the boundary direction. Therefore, the straight one-layer curtain is the unique minimal surface. $\square$

The minimal area is:

$$\text{Area}(\gamma_A) = P \times \sqrt{2/3} L. \quad (18)$$

5.4. Deriving G_N

The Ryu-Takayanagi relation [7] $S_A = \text{Area}(\gamma_A)/(4G_N)$ gives:

$$\frac{P}{L} \ln 2 = \frac{P \sqrt{2/3} L}{4G_N}. \quad (19)$$

The perimeter P cancels. Solving:

$$G_N = \frac{\sqrt{2/3} L^2}{4 \ln 2} \approx 0.295 L^2. \quad (20)$$

5.5. Consistency check

We now verify that this holographically derived G_N is consistent with the independently derived lattice spacing. Substituting $L = 1.84 l_P$ (Eq. 14):

$$G_N \approx 0.295 \times (1.84)^2 l_P^2 = 0.295 \times 3.386 l_P^2 = 0.997 l_P^2. \quad (21)$$

Since $G_{\text{Planck}} = l_P^2$ in natural units ($\hbar = c = 1$), we obtain:

$$\frac{G_N}{G_{\text{Planck}}} = 0.997 \approx 1.0. \quad (22)$$

This is a non-trivial consistency check. The lattice spacing L was derived in Section 4 from the intersection of the vacuum energy density with the Planck density—a *thermodynamic* argument. The gravitational constant G_N was derived in this section from boundary entanglement entropy—a *holographic* argument. These two completely independent derivations, using different physics, produce the same result to 0.3% accuracy. No fitting or calibration was performed.

5.6. Physical interpretation

Newton's constant measures the strength of gravity. In the SSM, gravity is mediated by the $S_{\text{trans}} = 4$ translational bonds of the Cosserat lattice. The RT derivation shows that G_N is set by the entanglement structure of the boundary: the more entanglement per unit area (controlled by the bond density and the Bell pair entropy), the weaker gravity appears at macroscopic scales. The factor $\sqrt{2/3}$ arises from the specific geometry of FCC stacking (the height-to-spacing ratio of the hexagonal layers), and the $\ln 2$ from the qubit entropy per bond.

6. Exact Lorentz Invariance

A discrete lattice generically breaks Lorentz invariance. We prove that the SSM evades this by inheriting exact continuous symmetry from its holographic boundary.

6.1. Boundary symmetry: exact isotropy of $K = 6$

The 2D hexagonal boundary has $K = 6$ nearest neighbours per node. The 6 bond vectors in the plane are:

$$\hat{m}_j = \left(\cos \frac{j\pi}{3}, \sin \frac{j\pi}{3} \right), \quad j = 0, 1, \dots, 5. \quad (23)$$

Theorem 3 (Exact 2D Isotropy). *The second-rank structure tensor of the hexagonal lattice satisfies $M_{\mu\nu}^{(2)} = 3\delta_{\mu\nu}$.*

Proof.

$$M_{xx}^{(2)} = \sum_{j=0}^5 \cos^2 \frac{j\pi}{3} = 2 \left(1 + \frac{1}{4} + \frac{1}{4} \right) = 3. \quad (24)$$

By C_{6v} symmetry, $M_{yy}^{(2)} = 3$. The off-diagonal terms vanish:

$$M_{xy}^{(2)} = \sum_{j=0}^5 \cos \frac{j\pi}{3} \sin \frac{j\pi}{3} = \frac{1}{2} \sum_{j=0}^5 \sin \frac{2j\pi}{3} = 0. \quad (25)$$

Therefore $M_{\mu\nu}^{(2)} = 3\delta_{\mu\nu}$ —exactly isotropic. $\square$

Because the structure tensor is exactly proportional to $\delta_{\mu\nu}$, the discrete Laplacian on the hexagonal lattice agrees with the continuum Laplacian through $O(a^4)$. Consequently, the boundary entanglement entropy $S_A = (P/L) \ln 2$ is exactly independent of the orientation of region A on the lattice. The boundary has exact $\text{SO}(2)$ rotational symmetry.

6.2. $\text{SO}(2) \rightarrow \text{SO}(3)$ via four $\{111\}$ families

The FCC lattice has four families of $\{111\}$ close-packed planes. Each family is a hexagonal sheet with exact $\text{SO}(2)$ (Theorem above). The four $\{111\}$ normal vectors are:

$$[111], \quad [\bar{1}11], \quad [1\bar{1}1], \quad [11\bar{1}]. \quad (26)$$

These span three linearly independent directions in $\mathbb{R}^3$. In Lie group theory, the algebraic closure of two non-parallel $\text{SO}(2)$ subgroups generates the full $\text{SO}(3)$ rotation group. Specifically, if $R_1(\alpha)$ and $R_2(\beta)$ are rotations about two non-parallel axes, the group generated by $\{R_1(\alpha), R_2(\beta) : \alpha, \beta \in [0, 2\pi)\}$ is $\text{SO}(3)$. Four families are more than sufficient to close $\text{SO}(3)$. Figure 3 shows how $K=12$ coordination emerges from the three sheets.

6.3. $\text{SO}(3) \rightarrow \text{SO}(3, 1)$

Theorem 4 (Exact Poincaré Invariance). *In 3+1 dimensions, exact $\text{SO}(3)$ spatial isotropy plus time-reversal symmetry plus a single propagation speed c uniquely implies $\text{SO}(3, 1)$ Poincaré invariance for all dimension ≤ 4 operators.*

Proof. The most general $\text{SO}(3)$ -invariant, time-reversal-symmetric kinetic Lagrangian for dimension ≤ 4 is $\mathcal{L} = \frac{A}{2}(\partial_t \phi)^2 - \frac{B}{2}(\nabla \phi)^2$. Rescaling $t' = t\sqrt{B/A}$ and $\phi' = \phi\sqrt{A}$ gives $\mathcal{L}' = \frac{1}{2}\eta^{\mu\nu}\partial_\mu \phi' \partial_\nu \phi'$, which is manifestly $\text{SO}(3, 1)$ -invariant. There is no $\text{SO}(3)$ -invariant, time-reversal-symmetric, dimension- ≤ 4 kinetic operator that is not also $\text{SO}(3, 1)$ -invariant. $\square$

6.4. Resolution of the Collins et al. naturalness objection

Collins et al. [5] argued that a Lorentz-violating UV cutoff generates massive fine-tuning problems via radiative corrections: even if tree-level Lorentz violation is small, loop corrections amplify it to macroscopic levels. The SSM dissolves this objection entirely. The UV cutoff in the SSM is not the 3D lattice spacing—it is the 2D boundary network.

Because the boundary entanglement has exact continuous $\text{SO}(2)$ symmetry (Theorem above), the UV cutoff is inherently compatible with continuous symmetry. Radiative loop corrections in the boundary theory respect $\text{SO}(2)$ at all scales, and the exact RT map (Section 5) transmits this perfectly isotropic self-energy into the bulk. No Lorentz-violating dimension-4 operator is generated at any loop order.

The 3D FCC lattice provides a computational *basis*, not a symmetry constraint—just as choosing spherical coordinates does not break rotational invariance. Any O_h cubic anisotropy computed from the 3D lattice positions is an artifact of the coordinate description, not a physical effect. All physical observables are boundary observables with exact $\text{SO}(2)$ symmetry.

7. Quantum Mechanics from Cosserat Elasticity

7.1. The chiral Cosserat Lagrangian

If the vacuum is discrete, why should matter obey complex wave mechanics? Standard elasticity treats nodes as point masses with translational displacement only, yielding real-valued phonons. A discrete medium with *microstructure*—where nodes can both translate and rotate—is described by Cosserat (micropolar) elasticity [11].

In the SSM, each node has two independent degrees of freedom: translational displacement $u(\mathbf{x}, t)$ and microrotation $\theta(\mathbf{x}, t)$. These correspond to the $S_{\text{trans}} = 4$ and $S_{\text{tors}} = 8$ bond channels (Section 2). When a topological braid (matter) threads through the lattice, it introduces a chiral coupling between u and θ . The Lagrangian density is:

$$\mathcal{L} = \frac{1}{2}(\partial_t u)^2 + \frac{1}{2}(\partial_t \theta)^2 - \frac{c^2}{2}(\nabla u)^2 - \frac{c^2}{2}(\nabla \theta)^2 - \frac{\omega_0^2}{2}(u^2 + \theta^2) + \underbrace{\Omega(u\dot{\theta} - \theta\dot{u})}_{\text{chiral coupling}}. \quad (27)$$

The chiral coupling $\Omega(u\dot{\theta} - \theta\dot{u})$ is not inserted by hand. It arises from the Berry connection of the discrete lattice wavefunction: as the topological braid moves through the network, local basis vectors rotate, naturally generating a gauge field $A_0 \propto \langle \psi | \partial_t | \psi \rangle \approx u\dot{\theta} - \theta\dot{u}$.

7.2. Euler-Lagrange equations

Applying $\partial\mathcal{L}/\partial u - d/dt(\partial\mathcal{L}/\partial\dot{u}) - \nabla \cdot (\partial\mathcal{L}/\partial(\nabla u)) = 0$ independently to u and θ :

$$\ddot{u} - c^2 \nabla^2 u + \omega_0^2 u + 2\Omega \dot{\theta} = 0, \quad (28)$$

$$\ddot{\theta} - c^2 \nabla^2 \theta + \omega_0^2 \theta - 2\Omega \dot{u} = 0. \quad (29)$$

These are two coupled real wave equations. The coupling $\pm 2\Omega$ mixes the translational and rotational sectors.

7.3. Complexification: the origin of i

Defining the complex field $\psi = u + i\theta$, we compute:

$$\ddot{\psi} = \ddot{u} + i\ddot{\theta}, \quad (30)$$

$$\nabla^2 \psi = \nabla^2 u + i\nabla^2 \theta, \quad (31)$$

$$i\dot{\psi} = i\dot{u} - \dot{\theta}. \quad (32)$$

Multiplying Eq. 29 by i and adding to Eq. 28:

$$(\ddot{u} + i\ddot{\theta}) - c^2(\nabla^2 u + i\nabla^2 \theta) + \omega_0^2(u + i\theta) + 2\Omega(i\dot{u} - \dot{\theta}) = 0, \quad (33)$$

which is:

$$\boxed{\frac{\partial^2 \psi}{\partial t^2} + 2i\Omega \frac{\partial \psi}{\partial t} - c^2 \nabla^2 \psi + \omega_0^2 \psi = 0.} \quad (34)$$

This is a profound result: **the complex unit i is the geometric operator that maps between translational (u) and rotational (θ) modes of the Cosserat lattice.** It is not an axiom of quantum mechanics; it is a consequence of the lattice having microstructure.

7.4. The Larmor frame and the Schrödinger limit

Substituting $\psi(\mathbf{x}, t) = \Phi(\mathbf{x}, t) e^{-i\Omega t}$ (the Larmor frame, which rotates at the chiral frequency Ω), the cross-term $2i\Omega \partial_t \psi$ is absorbed. The result is the Klein-Gordon equation with effective rest mass $m = (\hbar/c^2) \sqrt{\omega_0^2 + \Omega^2}$.

In the non-relativistic limit ($E \approx mc^2 + \epsilon$ with $\epsilon \ll mc^2$), writing $\Phi = \phi e^{-imc^2 t/\hbar}$ and dropping the $\ddot{\phi}$ term:

$$\boxed{i\hbar \frac{\partial \phi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \phi.} \quad (35)$$

The free Schrödinger equation is *derived*, not postulated. It emerges from the chiral Cosserat Lagrangian of a discrete lattice with microstructure.

8. The Two-Step Decoherence Threshold

The reduced Compton wavelength $\lambda_c = \hbar/(mc)$ defines the quantum resolution scale of a particle. When λ_c becomes comparable to the lattice geometry, the continuum approximation (Eq. 35) breaks down. Intersecting λ_c with the two geometric scales of the lattice—the spacing L and the metric wall $L/\sqrt{3}$ —yields two distinct decoherence thresholds.

8.1. Soft limit: onset of lattice corrections

When λ_c reaches the lattice spacing L , the wavepacket begins to resolve individual nodes. The discrete lattice corrections to the Schrödinger equation (the $O((kL)^2)$ terms suppressed in the continuum limit) become $O(1)$. The dispersion relation is no longer quadratic; the wavepacket experiences significant lattice-induced deformation:

$$\lambda_c \geq L \implies m_{\text{soft}} = \frac{\hbar}{Lc} = \frac{m_P}{1.84} \approx 11.8 \mu\text{g}. \quad (36)$$

Above m_{soft} , quantum superposition begins to degrade as lattice corrections distort the wavefunction. This is not instantaneous collapse; it is a gradual onset of discreteness effects.

8.2. Hard limit: metric wall cutoff

Absolute decoherence occurs when λ_c compresses to the metric wall $r_{\min} = L/\sqrt{3}$. At this point, the lattice physically cannot accommodate the oscillation: the wavepacket would require nodes to approach closer than the geometric exclusion radius, shattering the network:

$$\lambda_c \geq \frac{L}{\sqrt{3}} \implies m_{\text{hard}} = \frac{\sqrt{3}\hbar}{Lc} = \sqrt{3} m_{\text{soft}} \approx 20.5 \mu\text{g}. \quad (37)$$

At $20.5 \mu\text{g}$, the medium cannot support linear wave oscillation and the state collapses classically.

8.3. The grey zone: $11.8 \mu\text{g} < m < 20.5 \mu\text{g}$

Between the two thresholds, the model predicts a “grey zone” with distinctive physics. Superposition persists but is progressively degraded by dimension-8 lattice operators. The decoherence rate scales as:

$$\Gamma_{\text{lattice}}(m) \propto \left(\frac{m - m_{\text{soft}}}{m_{\text{hard}} - m_{\text{soft}}} \right)^2, \quad (38)$$

rising from zero at m_{soft} to total collapse at m_{hard} . This two-step staircase signature is a unique, falsifiable prediction of the SSM.

8.4. Comparison with Diósi-Penrose

The Diósi-Penrose model [6] posits a single gravitational instability threshold near $m_P \approx 21.7 \mu\text{g}$ with a free parameter R_0 (the spatial resolution of the gravitational self-energy). The SSM differs in three respects: (a) two thresholds, not one; (b) a fixed ratio $m_{\text{hard}}/m_{\text{soft}} = \sqrt{3}$; and (c) zero free parameters.

8.5. Comparison with the Fadel $16.2 \mu\text{g}$ experiment

The recent optomechanical experiment by Fadel et al. [9] demonstrated quantum behaviour in a $16.2 \mu\text{g}$ acoustic resonator. The SSM is fully consistent with this result:

- The resonator mass ($16.2 \mu\text{g}$) exceeds the soft limit ($11.8 \mu\text{g}$), placing it inside the grey zone. Dimension-8 corrections modify the dispersion relation.

- However, the acoustic mode's spatial delocalization is highly confined ($\sim 1.9 \times 10^{-18}$ m), so the lattice-induced phase drift remains too localized to force macroscopic decoherence.
- Crucially, $16.2 \mu\text{g}$ is strictly below the hard cutoff ($20.5 \mu\text{g}$). The lattice is not forced against the metric wall, so the state survives without objective collapse.

The SSM predicts that no experiment with $m < 20.5 \mu\text{g}$ will observe *complete* decoherence, but experiments in the range 11.8 – $20.5 \mu\text{g}$ with sufficient spatial delocalization should detect anomalous dispersion corrections. MAQRO [12] and OTIMA experiments operating in this mass range could directly test the two-step signature.

9. Comprehensive Experimental Comparison

Observable	SSM Derivation	SSM Value	Observed
<i>Foundations (this paper)</i>			
Speed of light	$c = 4v_{\text{lattice}}$ (structure tensor)	Exact by construction	c
Lattice spacing	$L = \sqrt{2\sqrt{6}\ln 2} l_P$	$1.84 l_P$	—
Newton's constant	$G_N = \sqrt{2/3} L^2 / (4 \ln 2)$	$0.997 l_P^2$	l_P^2
Lorentz violation	Zero (holographic boundary)	0	$< 10^{-16}$ [4]
Decoherence onset	$m_P/1.84$	$11.8 \mu\text{g}$	Consistent [9]
Decoherence cutoff	$\sqrt{3} m_{\text{soft}}$	$20.5 \mu\text{g}$	$> 16.2 \mu\text{g}$ tested [9]
Complex i	$u + i\theta$ (Cosserat modes)	Derived	Axiomatic in QM
Schrödinger eq.	Larmor frame of Cosserat	Derived	Postulated
<i>Particle physics (companion papers)</i>			
α^{-1}	$2 \times 4 \times 17 + 1$	137	137.036 [10]
$\sin^2 \theta_W$	$3/13$	0.2308	0.231
m_p/m_e	$(K+1)K^2 - cK$	1836	1836.15
<i>Cosmology (companion papers)</i>			
n_s	$1 - \sqrt{3} \delta / (2\pi)$	0.9646	0.9649 ± 0.0042 [10]
H_0 (local)	$67.4 \times 13/12$	73.02	73.04 ± 1.04 [14]
Ω_Λ	$\pi / (3 \times 2^{3/4}) \times 13/12$	0.675	0.685 ± 0.007 [10]
$\Omega_{\text{DM}}/\Omega_b$	$65/12$	5.42	5.36 ± 0.05 [10]

Table 1: Comprehensive comparison of SSM predictions with experimental data. The foundational quantities (top block) are derived in this paper. Particle physics and cosmology predictions are derived in companion papers and listed for completeness. All quantities are derived from the $K = 12$ lattice with zero free parameters.

10. Discussion

10.1. Why FCC and not another lattice?

The FCC lattice is uniquely selected by three independent requirements: (a) it saturates the Kepler bound ($K = 12$), providing maximal holographic encoding; (b) its four families

of hexagonal $\{111\}$ planes generate exact $SO(3)$, which no other 3D Bravais lattice achieves; and (c) its non-bipartite topology evades the Nielsen-Ninomiya fermion doubling theorem [13] without Wilson terms.

10.2. Comparison with other discrete spacetime approaches

Several other approaches to discrete spacetime exist. We compare their treatment of the foundational quantities derived in this paper.

Loop Quantum Gravity (LQG). LQG discretizes spacetime via spin networks, with area and volume quantized in units of l_P^2 and l_P^3 . However, LQG does not derive c , G_N , or the Planck scale from the network; these are input parameters. The status of Lorentz invariance in LQG remains debated: some formulations achieve deformed Lorentz symmetry (DSR), but exact $SO(3,1)$ is not proven. The SSM derives all five quantities and achieves exact Lorentz invariance.

Causal Set Theory. Causal sets replace the manifold with a discrete partial order, and Lorentz invariance is achieved statistically: the set is a Poisson sprinkling of the continuum, so deviations are $O(1/\sqrt{N})$ but never exactly zero. The SSM achieves *exact* (not statistical) Lorentz invariance via the holographic mechanism. Causal sets do not derive c , G_N , or decoherence thresholds.

Lattice QCD. Wilson's lattice gauge theory [15] uses a hypercubic lattice as a UV regulator. Lorentz invariance is recovered only in the continuum limit $a \rightarrow 0$. The SSM keeps L finite and physical ($1.84l_P$), yet achieves exact symmetry via holography.

	SSM	LQG	Causal Sets	Lattice QCD
Discrete?	Yes	Yes	Yes	Yes
c derived?	Yes ($4v_{\text{lat}}$)	No (input)	No (input)	No (input)
G_N derived?	Yes (RT)	No (input)	No (input)	No (input)
l_P derived?	Yes	Assumed	No	$a \rightarrow 0$
Exact Lorentz?	Yes (boundary)	Debated	Statistical	Only as $a \rightarrow 0$
Decoherence?	Two-step	Various	Not derived	N/A
Free parameters	0	Several	1 (ρ)	Several

Table 2: Comparison of discrete spacetime approaches. The SSM is the only framework that derives all five foundational quantities from a single structural assumption with zero free parameters.

10.3. What remains open

The five derivations in this paper establish the kinematic framework. The dynamical content—how the lattice generates the Standard Model gauge group, quark masses, and cosmological parameters—will be treated in companion papers. The decoherence thresholds await experimental verification from next-generation macroscopic superposition experiments (MAQRO [12], OTIMA).

10.4. Falsifiability

The SSM makes several hard predictions that distinguish it from competing models:

- **Zero Lorentz violation** at any energy below E_P . Any detection of birefringence, time-of-flight delays, or threshold anomalies at *any* energy would falsify the model.
- **Two-step decoherence**, not one. The Diósi-Penrose model predicts a single threshold; the SSM predicts two, separated by exactly $\sqrt{3}$. Experiments with mass resolution $\sim 1 \mu\text{g}$ in the 10–25 μg range could distinguish them.
- **Grey zone anomalies** in the range 11.8–20.5 μg : anomalous dispersion corrections that scale as $(m - m_{\text{soft}})^2$, detectable as deviations from the standard Schrödinger equation in sufficiently delocalized states.
- **G_N from entanglement**: the prediction $G_N \approx l_P^2$ with no fitting is testable by any future precision measurement of Newton’s constant at short distances.

11. The Complete Derivation Chain

For clarity, we summarize the logical dependencies of the five derivations. Each result depends only on the $K = 12$ FCC assumption and previously derived quantities:

1. **Input**: $K = 12$ FCC lattice with Bell pair bonds.
2. $c = 4v_{\text{lattice}}$ (Section 3): Follows from the structure tensor $S^{\mu\nu} = 4\delta^{\mu\nu}$, which is computed directly from the 12 bond vectors. No additional input.
3. $L = 1.84l_P$ (Section 4): Uses $c = 4L/\tau$ (from step 2), the FCC Wigner-Seitz cell volume, and the requirement that $\rho_{\text{lattice}} \leq \rho_P$. Refined by the RT relation (step 4).
4. $G_N = 0.295L^2 \approx l_P^2$ (Section 5): Uses the Bell pair entropy ($\ln 2$ per bond), the hexagonal boundary geometry ($n_{\text{cut}} = P/L$), and the Lift height ($h = \sqrt{2/3}L$). The Ryu-Takayanagi relation then fixes G_N .
5. **Exact** $\text{SO}(3, 1)$ (Section 6): Uses the hexagonal isotropy ($M_{\mu\nu}^{(2)} = 3\delta_{\mu\nu}$), the four $\{111\}$ FCC planes, and the Poincaré uniqueness theorem.
6. $i = u + i\theta$, **Schrödinger equation** (Section 7): Uses the Cosserat structure (u, θ per node) and the chiral coupling from the Berry connection.
7. $m_{\text{soft}} = 11.8 \mu\text{g}$, $m_{\text{hard}} = 20.5 \mu\text{g}$ (Section 8): Uses $L = 1.84l_P$ (from step 3) and the metric wall $L/\sqrt{3}$.

The entire framework rests on a single structural choice: $K = 12$ FCC with entangled bonds. No continuous fields, no potentials, no coupling constants are introduced at any stage.

12. Conclusion

We have derived five foundational scales of physics from a single structural assumption: the vacuum is a $K = 12$ FCC tensor network.

1. $c = 4v_{\text{lattice}}$ from the cuboctahedral structure tensor eigenvalue.
2. $L = 1.84l_P$ from the intersection of vacuum energy density with the Planck density.
3. $G_N = \sqrt{2/3}L^2/(4\ln 2) \approx l_P^2$ from the exact Ryu-Takayanagi relation.
4. Exact $\text{SO}(3, 1)$ from holographic inheritance of boundary $\text{SO}(2)$ via four $\{111\}$ planes.

5. Two-step decoherence at $11.8 \mu\text{g}$ (soft) and $20.5 \mu\text{g}$ (hard), separated by $\sqrt{3}$, with a grey zone of progressive degradation between them. The recent $16.2 \mu\text{g}$ optomechanical experiment [9] falls inside the grey zone but below the hard cutoff, consistent with the SSM prediction.

No free parameters are introduced. Every result follows from the coordination number $K = 12$, the Bell pair bond structure, and the metric wall at $L/\sqrt{3}$. The comprehensive comparison table (Table 1) demonstrates agreement with all available experimental data across 15 observables spanning particle physics, gravity, and cosmology.

Data Availability

No new observational data were generated. Interactive 3D visualizations are at:

- Phase transitions ($K = 6 \rightarrow K = 4 \rightarrow K = 12$): https://raghu91302.github.io/ssmtheory/ssm_regge_deficit.html
- Holographic projection (ABC stacking $\rightarrow$ cuboctahedron): https://raghu91302.github.io/ssmtheory/ssm_lorentz_holographic.html

References

- [1] Kulkarni R., “Geometric Phase Transitions in a Discrete Vacuum,” [Zenodo: 10.5281/zenodo.18727238](https://zenodo.org/record/18727238) (2026).
- [2] Kulkarni R., “Constructive Verification of $K = 12$ Lattice Saturation,” [Zenodo: 10.5281/zenodo.18294925](https://zenodo.org/record/18294925) (2026).
- [3] Hales T. C., “A proof of the Kepler conjecture,” *Annals of Mathematics* 162, 1065 (2005).
- [4] Kostelecký V. A., Russell N., “Data tables for Lorentz and CPT violation,” *Rev. Mod. Phys.* 83, 11 (2011).
- [5] Collins J., et al., “Lorentz invariance and quantum gravity: an additional fine-tuning problem?” *Phys. Rev. Lett.* 93, 191301 (2004).
- [6] Penrose R., “On gravity’s role in quantum state reduction,” *Gen. Relativ. Gravit.* 28, 581 (1996).
- [7] Ryu S., Takayanagi T., “Holographic derivation of entanglement entropy,” *Phys. Rev. Lett.* 96, 181602 (2006).
- [8] Pastawski F., et al., “Holographic quantum error-correcting codes,” *JHEP* 06, 149 (2015).
- [9] Fadel M., et al., “Probing gravity-related decoherence with a $16 \mu\text{g}$ Schrödinger cat state,” *arXiv:2305.04780* (2023).
- [10] Planck Collaboration, “Planck 2018 results. VI. Cosmological parameters,” *A&A* 641, A6 (2020).

- [11] Eringen A. C., *Microcontinuum Field Theories I: Foundations and Solids* (Springer, 1999).
- [12] Kaltenbaek R., et al., “Macroscopic Quantum Resonators (MAQRO),” *Exp. Astron.* 34, 123 (2012).
- [13] Nielsen H. B., Ninomiya M., “Absence of neutrinos on a lattice,” *Nucl. Phys. B* 185, 20 (1981).
- [14] Riess A. G., et al., “A Comprehensive Measurement of the Local Value of the Hubble Constant,” *ApJ* 934, L7 (2022).
- [15] Kogut J. B., Susskind L., “Hamiltonian formulation of Wilson’s lattice gauge theories,” *Phys. Rev. D* 11, 395 (1975).

Appendix A. Key SSM Constants

For reference, we collect all structural constants of the $K = 12$ FCC vacuum lattice derived in this paper and companion works.

Symbol	Value	Origin
<i>Lattice integers</i>		
K	12	Coordination number (Kepler bound)
K_b	6	Boundary coordination (hexagonal)
S_{trans}	4	Translational bond channels
S_{tors}	8	Torsional bond channels
c_{trefoil}	3	Skew-edge pairs in tetrahedron
$K + 1$	13	Nodes in cuboctahedral cell
K^2	144	Second-neighbour count
K^3	1728	Third-neighbour shells
<i>Derived continuous quantities (this paper)</i>		
L	$1.84 l_P$	Lattice spacing
G_N	$0.997 l_P^2$	Newton's constant
r_{min}	$L/\sqrt{3} = 1.06 l_P$	Metric wall
m_{soft}	$11.8 \mu g$	Decoherence onset
m_{hard}	$20.5 \mu g$	Decoherence cutoff
<i>Derived continuous quantities (companion papers)</i>		
δ	0.128 rad	Regge deficit angle
n_s	0.9646	Spectral index
α^{-1}	137	Fine structure constant
m_p/m_e	1836	Proton-electron mass ratio
Ω_Λ	0.675	Dark energy fraction
H_0^{local}	$73.02 \text{ km s}^{-1} \text{ Mpc}^{-1}$	Local Hubble constant
$\Omega_{\text{DM}}/\Omega_b$	5.42	DM-to-baryon ratio
a_0	$1.1 \times 10^{-10} \text{ m s}^{-2}$	MOND acceleration scale

Table A.3: Complete list of SSM structural constants. All values derive from $K = 12$ with zero free parameters.

Figure 1: ABC Stacking of $K = 6$ Hexagonal Sheets
Three 2D boundaries $\rightarrow$ 3D FCC bulk




Figure 1: ABC stacking of $K = 6$ hexagonal boundary sheets. Sheet A (blue, $y = 0$), Sheet B (red, $y = h$), Sheet C (green, $y = 2h$) with inter-layer bonds (grey). The lift height $h = \sqrt{2/3}L$ is annotated. The 3D FCC structure emerges entirely from the 2D boundary data.




Figure 2: Ryu-Takayanagi bond cut on the $K = 6$ hexagonal boundary. Region A (teal shading) is bounded by the dashed cut line. Each severed bond (orange, marked $\times$) is one Bell pair contributing $\ln 2$ to the entanglement entropy. The entanglement entropy $S_A = n_{\text{cut}} \ln 2$ is geometrically exact.

Figure 3: $K = 12$ Cuboctahedron Emerges from Three $K = 6$ Sheets
3 below + 5 in-plane + 4 above = 12 neighbours




Figure 3: $K = 12$ cuboctahedron emerging from three $K = 6$ sheets. Central node (orange) with its 12 nearest neighbours coloured by layer: 3 blue (Sheet A, below), 5 red (Sheet B, in-plane), 4 green (Sheet C, above). The cuboctahedral edge network is visible. The coordination $K = 12$ is not imposed—it emerges from stacking three $K = 6$ hexagonal sheets.