

8.circle

ee24btech11027- satwikagv

- 6) The lines $2x-3y=5$ and $3x-4y=7$ are diameters of a circle having area as 154 sq.units. Then the equation of the circle is **(2003)**
- $x^2 + y^2 - 2x + 2y = 62$
 - $x^2 + y^2 + 2x - 2y = 62$
 - $x^2 + y^2 + 2x - 2y = 47$
 - $x^2 + y^2 - 2x + 2y = 47$
- 7) If a circle passes through the point (a, b) and cuts the circle $x^2 + y^2 = 4$ orthogonally, then the locus of its centre is **(2004)**
- $2ax - 2by - (a^2 + b^2 + 4) = 0$
 - $2ax + 2by - (a^2 + b^2 + 4) = 0$
 - $2ax - 2by + (a^2 + b^2 + 4) = 0$
 - $2ax + 2by + (a^2 + b^2 + 4) = 4$
- 8) A variable circle passes through the fixed point $A(p, q)$ and touches x -axis. The locus of the other end of the diameter through A is **(2004)**
- $(y - q)^2 = 4px$
 - $(x - q)^2 = 4py$
 - $(y - p)^2 = 4qx$
 - $(x - p)^2 = 4qy$
- 9) If the lines $2x + 3y + 1 = 0$ and $3x - y - 4 = 0$ lie along diameter of a circle of circumference 10π , then the equation of the circle is **(2004)**
- $x^2 + y^2 + 2x - 2y - 23 = 0$
 - $x^2 + y^2 - 2x - 2y - 23 = 0$
 - $x^2 + y^2 + 2x + 2y - 23 = 0$
 - $x^2 + y^2 - 2x + 2y - 23 = 0$
- 10) Intercept on the line $y = x$ by the circle $x^2 + y^2 - 2x = 0$ is AB . Equation of the circle on AB as a diameter is **(2004)**
- $x^2 + y^2 + x - y = 0$
 - $x^2 + y^2 - x + y = 0$
 - $x^2 + y^2 + x + y = 0$
 - $x^2 + y^2 - x - y = 0$
- 11) If the circles $x^2 + y^2 + 2ax + cy + a = 0$ and $x^2 + y^2 - 3ax + dy - 1 = 0$ intersect in two distinct points P and Q then the line $5x + by - a = 0$ passes through P and Q for **(2005)**
- exactly one value of a
 - no value of a
 - infinitely many values of a
 - exactly two values of a
- 12) A circle touches the x -axis and also touches the circle with centre at $(0, 3)$ and radius 2. The locus of the centre of the circle is **(2005)**
- an ellipse
 - a circle
 - a hyperbola
 - a parabola
- 13) If a circle passes through the point (a, b) and cuts the circle $x^2 + y^2 = p^2$ orthogonally, then the equation of the locus of its centre is **(2005)**
- $x^2 + y^2 - 3ax - 4by + (a^2 + b^2 - p^2) = 0$
 - $2ax + 2by - (a^2 - b^2 + p^2) = 0$
 - $x^2 + y^2 - 2ax - 3by + (a^2 - b^2 - p^2) = 0$
 - $2ax + 2by - (a^2 + b^2 + p^2) = 0$
- 14) If the pair of lines $ax^2 + 2(a + b)xy + by^2 = 0$ lie along diameters of a circle and divide the circle into four sectors such that the area of one of the sectors is thrice the area of another sector then **(2005)**
- $3a^2 - 10ab + 3b^2 = 0$
 - $3a^2 - 2ab + 3b^2 = 0$
 - $3a^2 + 10ab + 3b^2 = 0$
 - $3a^2 + 2ab + 3b^2 = 0$
- 15) If the lines $3x - 4y - 7 = 0$ and $2x - 3y - 5 = 0$ are two diameters of a circle of area 49π square units, the equation of the circle is **(2006)**
- $x^2 + y^2 + 2x - 2y - 47 = 0$
 - $x^2 + y^2 + 2x - 2y - 62 = 0$
 - $x^2 + y^2 - 2x + 2y - 62 = 0$
 - $x^2 + y^2 - 2x + 2y - 47 = 0$
- 16) Let C be the circle with centre $(0, 0)$ and radius 3 units. The equation of the locus of the mid points of the chords of the circle C that subtend an angle of $\frac{2\pi}{3}$ at its centre is **(2006)**
- $x^2 + y^2 = \frac{3}{2}$
 - $x^2 + y^2 = 1$
 - $x^2 + y^2 = \frac{27}{4}$
 - $x^2 + y^2 = \frac{9}{4}$
- 17) Consider a family of circles which are passing through the point $(-1, 1)$, and are tangent to x -

axis. If (h, k) are the coordinate of the centre of the circles, then the set of values of k is given by the interval **(2007)**

- (a) $-\frac{1}{2} \leq k \leq \frac{1}{2}$ (c) $0 \leq k \leq \frac{1}{2}$
 (b) $k \leq \frac{1}{2}$ (d) $k \geq \frac{1}{2}$

18) The point diametrically opposite to the point **P** (1, 0) on the circle $x^2 + y^2 + 2x + 2y - 3 = 0$ is **(2008)**

- (a) (3, -4) (c) (-3, -4)
 (b) (-3, 4) (d) (3, 4)

19) The differential equation of the family of circles with fixed radius 5 units and centre on the line $y = 2$ is

- (a) $(x - 2)y'^2 = 25 - (y - 2)^2$
 (b) $(y - 2)y'^2 = 25 - (y - 2)^2$
 (c) $(y - 2)^2 y'^2 = 25 - (y - 2)^2$
 (d) $(x - 2)^2 y'^2 = 25 - (y - 2)^2$

20) If **P** and **Q** are the points of intersection of the circles $x^2 + y^2 + 3x + 7y + 2p - 5 = 0$ and $x^2 + y^2 + 2x + 2y - p^2 = 0$ then there is a circle passing through **P, Q** and (1, 1) for: **(2009)**

- (a) all except one value of p
 (b) all except two values of p
 (c) exactly one value of p
 (d) all value of p