Unit 2: Optics

1. Basic Optical Laws: Reflection, Refraction & Refractive Index

Definition: Optics is the branch of physics that studies the behavior and properties of light. Reflection is the bouncing back of light in the same medium, while refraction is the bending of light when it passes from one transparent medium to another.

Light behaves predictably when it interacts with different surfaces and media. This behavior is governed by fundamental laws of optics.

Laws of Reflection:

When light strikes a polished surface (like a mirror), it bounces back. This follows two rules:

- The incident ray, the reflected ray, and the normal to the surface at the point of incidence all lie in the same plane.
- The angle of incidence (∠i) is always equal to the angle of reflection (∠r). **[∠i = ∠r]**

Laws of Refraction (Snell's Law):

When light travels from one transparent medium to another (e.g., air to glass), its speed changes, causing it to bend.

- The incident ray, the refracted ray, and the normal at the point of incidence all lie in the same plane.
- **Snell's Law:** The ratio of the sine of the angle of incidence to the sine of the angle of refraction is a constant for a given pair of media.

Snell's Law: $\mu = \sin(i) / \sin(r)$

Refractive Index (μ or n):

The refractive index is a measure of how much a medium slows down light. It is defined as the ratio of the speed of light in a vacuum (c) to the speed of light in that medium (v).

Formula: $\mu = c / v$ (where $c \approx 3 \times 10^8$ m/s).

Higher refractive index means the medium is optically denser (light travels slower, bends more towards the normal).

```
REFLECTION:                    REFRACTION (Rarer to Denser):
      Normal                         Normal
         |                              |
 Incident|  Reflected           Incident |
  Ray \   |   / Ray              Ray \    |
       \ | /                          \   |
________\|/________Surface             \_|_ Rarer (Air)
                                       | \_
```

```
                                        |    \ Refracted Ray
                                        |     \ (Bends towards normal)
                                              Denser (Glass)
```

👉 **Hinglish Explanation:** Reflection matlab light ka kisi surface se takra kar wapas aana (angle i = angle r). Refraction matlab light ka ek medium se dusre medium me jate waqt mud jana. Refractive index (μ) batata hai ki koi medium light ko kitna slow karega. Snell's law (sin i / sin r) exam me confirm pucha jata hai!

2. Image Formation, Lens Formula, Power & Defects

Definition: An image is formed when light rays originating from an object converge (real image) or appear to diverge (virtual image) after reflection from a mirror or refraction through a lens.

Lenses and Image Formation:

A lens is a transparent optical medium bounded by two spherical surfaces. There are two main types:

- **Convex Lens (Converging):** Thicker in the middle. It brings parallel rays of light to a focus at a real focal point. It can form both real/inverted and virtual/erect images depending on the object's distance.
- **Concave Lens (Diverging):** Thinner in the middle. It spreads out parallel rays of light, causing them to appear as if they originate from a virtual focal point. Always forms virtual, erect, and diminished images.

Lens Formula & Magnification:

The relationship between object distance (u), image distance (v), and focal length (f) is given by the lens formula. Proper sign conventions must be used (e.g., distances measured against the direction of incident light are negative).

Lens Formula: $1/f = 1/v - 1/u$
Magnification (m) = v/u = Height of image (h') / Height of object (h)

Power of Lens (P):

The ability of a lens to converge or diverge light rays is its power. It is the reciprocal of focal length expressed in meters. The SI unit of power is **Diopter (D)**.
P = 1 / f(in meters). Convex lens has positive power (+D), Concave has negative power (-D).

Defects of Lenses (Aberrations):

- **Spherical Aberration:** The inability of a spherical lens to focus all marginal and paraxial rays at a single point, causing a blurred image. Fixed by using stops or aspheric lenses.
- **Chromatic Aberration:** The lens acts like a prism and disperses white light into its constituent colors, causing colored fringes around the image. Fixed by using achromatic doublets (combining convex and concave lenses of different materials).

👉 **Hinglish Explanation:** Lens formula (1/f = 1/v - 1/u) aur Power (P = 1/f) par direct numericals aate hain. Diopter yaad rakhna! Magnification batata hai image kitni badi bani. Lenses me 2 main defects hote hain: Spherical (image blur hona kyunki edges ki light pehle focus ho jati hai) aur Chromatic (image me colors/rainbow dikhna kyunki alag colors alag jagah focus hote hain).

3. Total Internal Reflection (TIR) & Optical Fibers

Definition: Total Internal Reflection (TIR) occurs when a light ray traveling from a denser to a rarer medium strikes the interface at an angle greater than the critical angle, causing the ray to reflect entirely back into the denser medium.

Critical Angle (C):

When light travels from an optically denser medium to a rarer medium, it bends away from the normal. As the angle of incidence increases, the angle of refraction also increases. The specific angle of incidence for which the angle of refraction becomes exactly 90° is called the Critical Angle.

$\sin(C) = 1 / \mu$ (where μ is the refractive index of denser medium w.r.t rarer)

Conditions for TIR:

1. Light must travel from an **optically denser medium to an optically rarer medium** (e.g., from glass/water to air).
2. The angle of incidence in the denser medium must be **greater than the critical angle** ($i > C$).

Applications of TIR in Optical Fibers:

An optical fiber is a thin, flexible strand of high-quality glass or plastic used to transmit light signals over long distances with minimal loss of energy. It consists of two main parts:

- **Core:** The inner part with a higher refractive index (μ_1).
- **Cladding:** The outer covering with a slightly lower refractive index (μ_2).

Working: Light enters the core at a specific angle and strikes the core-cladding boundary at an angle greater than the critical angle. Because the core is denser than the cladding, TIR occurs. The light undergoes multiple total internal reflections, zigzagging its way down the fiber without escaping.

```
Optical Fiber Principle (TIR):

   Cladding (Lower μ)
_____________________________________________
  \       /\       /\       /\       /
   \ TIR /  \ TIR /  \ TIR /  \ TIR /    <-- Light Ray
    \   /    \   /    \   /    \   /         trapped in Core
_____\ /_______\ /_______\ /_______\ /_______
   Core (Higher μ)
```

Key Uses:

- **Telecommunications:** High-speed internet (broadband) and telephone signal transmission without data loss.
- **Medical (Endoscopy):** Doctors use fiber-optic bundles to look inside the human body (e.g., stomach, lungs) without major

surgery.

👉 **Hinglish Explanation:** TIR bahut important topic hai. Jab light denser (kaanch/paani) se rarer (hawa) me jati hai, aur angle of incidence Critical angle se bada ho jaye, to light bahar jane ke bajay andar hi wapas reflect ho jati hai. Optical Fiber (Jio Fiber/Airtel Xstream) me yahi TIR use hota hai light signals ko bina loss ke hazaro kilometer door bhejne ke liye.

4. Simple & Compound Microscope

Definition: Microscopes are optical instruments used to produce highly magnified images of tiny, closely situated objects that cannot be seen clearly with the naked eye.

1. Simple Microscope:

A simple microscope is just a single convex lens of short focal length. When an object is placed between the optical center and the focus of the lens, it forms a virtual, erect, and magnified image on the same side as the object.

- **Magnifying Power (M):** It is the ratio of the angle subtended by the image at the eye to the angle subtended by the object at the least distance of distinct vision (D = 25 cm).
- **Formula:** M = 1 + (D/f) [when final image is at D].
- **Uses:** Reading small print, watchmakers for tiny gears, jewelers.

2. Compound Microscope:

To get a much higher magnification than a simple microscope, a compound microscope uses two convex lenses:

- **Objective Lens:** Placed near the object. It has a short focal length and a small aperture. It forms a real, inverted, and magnified intermediate image.
- **Eyepiece:** Placed near the eye. It has a slightly larger focal length and larger aperture. It acts like a simple microscope and magnifies the intermediate image further, forming a highly magnified, virtual, and inverted final image.

Magnifying Power of Compound Microscope:

$$M = m(\text{objective}) \times m(\text{eyepiece}) = (v_o / u_o) \times (1 + D/f_e)$$

👉 **Hinglish Explanation:** Simple microscope ek normal magnifying glass hota hai (1 lens). Par agar bacteria ya cells dekhne hain to 1 lens kafi nahi, isliye Compound microscope use hota hai jisme 2 lenses (Objective aur Eyepiece) hote hain. Pehla lens thoda zoom karta hai, dusra lens us zoom ki hui image ko aur zoom kar deta hai (M = mo × me).

5. Astronomical Telescope & Resolving Power

Definition: An astronomical telescope is an optical instrument used to see magnified images of distant heavenly bodies like stars, planets, and galaxies. Resolving power is its ability to separate two closely placed distant objects.

Astronomical Telescope (Normal Adjustment):

Like a compound microscope, it consists of two convex lenses mounted coaxially in a tube. However, their specifications are reversed:

- **Objective Lens:** Has a very large aperture (to gather maximum light from faint distant stars) and a large focal length (f_o).
- **Eyepiece:** Has a small aperture and a short focal length (f_e).

Working in Normal Adjustment: "Normal adjustment" means the final image is formed at infinity, which is the most relaxed state for the human eye. Parallel rays from a distant star fall on the objective, forming a real, inverted, and diminished image at its principal focus. The eyepiece is adjusted so that this image lies exactly at its focus. Thus, the final rays emerge parallel, creating the final image at infinity.

Magnifying Power in Normal Adjustment: $M = f_o / f_e$
Length of the tube: $L = f_o + f_e$

Resolving Power of Telescope:

Magnification alone is useless if the image is blurry. The resolving power is the ability of an instrument to produce distinct, separate images of two objects that are very close together (like binary stars). For a telescope, resolving power depends directly on the diameter (aperture) of the objective lens. Larger lenses give better resolving power.

> 👉 **Hinglish Explanation:** Telescope space ki cheezon (stars/planets) ko dekhne ke liye use hota hai. Iska aage wala lens (Objective) bahut bada hota hai taaki zyada light collect kar sake. Normal adjustment ka matlab hai final image infinity par banna, jisse aankhon par zor nahi padta. Magnifying power f_o/f_e hoti hai. Resolving power batati hai ki do aas-paas ke stars ek dikhenge ya alag-alag (HD quality).

6. Optical Projection Systems & Uses

Definition: Optical projection systems are devices designed to project a bright, magnified real image of a small object or slide onto a large screen for viewing by an audience.

A standard optical projection system (like a slide projector or cinema projector) consists of two main parts:

1. **Illumination System:** Consists of a powerful light source, a concave mirror (to reflect light forward), and condenser lenses. The condenser concentrates massive amounts of light evenly onto the transparent slide/object.
2. **Projection System (Projection Lens):** A combination of lenses (often an achromatic triplet) that takes the brightly illuminated slide and projects its real, magnified, and inverted image onto the screen. (The slide is placed upside down so the image appears erect on the screen).

Summary of Uses of Optical Instruments:

- **Microscopes:** Used in biology to study cells/microorganisms, in metallurgy to study crystal structures of metals, in forensics for analyzing trace evidence, and in medical pathology for blood tests.
- **Telescopes:** Used in astronomy to observe planets/galaxies, in navigation, and in terrestrial observation (using an extra erecting lens to make images upright).
- **Projection Systems:** Used in education (overhead projectors, smart classrooms), entertainment (movie theaters), and business presentations.

👉 **Hinglish Explanation:** Projection system ka kaam ek choti photo ya slide ko bada karke screen par dikhana hai (jaise cinema hall me projector). Isme condenser lens light ko slide par focus karta hai, aur projection lens slide ki badi image deewar par banata hai.

🔁 Quick Revision Notes (Exam Saver)

- **Snell's Law:** $\mu = \sin(i) / \sin(r)$. Basic rule for refraction.
- **Lens Formula & Power:** $1/f = 1/v - 1/u$. Power $P = 1/f$ (in meters). Unit: Diopter. Convex is (+), Concave is (-).
- **TIR Conditions:** Light must go from Denser to Rarer medium, and angle i > Critical angle C. Used in Optical Fibers.
- **Microscope vs Telescope:** Microscope has small objective, Telescope has huge objective.
- **Magnifying Powers:** Microscope (Compound) $M = m_o \times m_e$. Telescope (Normal Adj.) $M = f_o / f_e$.
- **Defects of Lenses:** Spherical aberration (blurred focus) and Chromatic aberration (color separation).

📝 **Powered & Inspired by: Rgpvnotes.tech**