

COORDINATE-BLOCK FACTORIZATION OVER STRONGLY PURELY INFINITE RINGS AND APPLICATIONS TO PROPERTY $(TT)/T$

GPT-5.6 SOL

ABSTRACT. We prove a coordinate-block factorization theorem for general linear groups over rings satisfying the single-sandwich condition

$$a \neq 0 \implies xay = 1 \text{ for some } x, y.$$

Every element of $\mathrm{GL}_n(R)$ is a product of at most $2n + 6$ factors, each an elementary transvection or an element of one fixed coordinate copy of $\mathrm{GL}_{n-1}(R)$. This is not bounded elementary generation: the coordinate block is an essential part of the statement.

We combine this algebraic factorization with relative quasi-cocycle rigidity for elementary groups over finite free associative $\mathbb{F}_2$ -algebras. If R is a nontrivial finite-type $\mathbb{F}_2$ -algebra carrying a binary Leavitt family, satisfying the single-sandwich condition, and satisfying $\mathrm{diag}(u, 1) \in E_2(R)$ for every $u \in R^\times$, then $E_n(R)$ has property $(TT)/T$ for every $n \geq 3$. Consequently,

$$E_n(L_{\mathbb{F}_2}(1, 2)) \text{ has property } (TT)/T \quad (n \geq 3).$$

We also give a direct quantitative proof of the relative root estimate and of the unstable diagonal reduction used in the specialization. The stable computation $K_1(L_{\mathbb{F}_2}(1, 2)) = 0$, the GE property of purely infinite simple rings, and the general relative-rigidity strategy are due to earlier work; we isolate our contribution explicitly. Every theorem claimed here as part of our contribution is matched to a declaration checked by the Lean kernel.

CONTENTS

1. Introduction	2
Credit and novelty boundary	3
Formal verification	4
2. Quasi-cocycles and relative control	4
3. Coordinate-block factorization	6
4. From relative control to global rigidity	8
5. The binary Leavitt specialization	9

Date: August 2026

repository revision **main**.

2020 *Mathematics Subject Classification*. 20J06, 20F65, 16S50, 16S88, 19B99.

Key words and phrases. property $(TT)/T$, quasi-cocycles, elementary groups, Leavitt algebras, purely infinite rings, bounded generation.

5.1. What follows from existing structure	10
5.2. The direct formalized route	10
6. Formal correspondence and trust boundary	10
7. Further directions	11
Acknowledgements	12
References	12

1. INTRODUCTION

Property $(TT)/T$, introduced by Mimura, asks that Hilbert-space quasi-cocycles be bounded after the invariant-vector part of the representation has been removed. It is stronger than Kazhdan's property (T) and is designed to control bounded cohomology and homomorphisms into groups admitting unbounded quasi-cocycles [5]. Ershov and Jaikin-Zapirain proved property (T) for elementary groups over arbitrary finitely generated associative rings [3]. The corresponding quasi-cocycle problem is subtler: the bounded error in the cocycle identity accumulates under a long word, so finite generation alone gives no global bound.

The present paper supplies a globalization mechanism for a class of strongly purely infinite rings. Its algebraic input is an explicit matrix factorization. The hypothesis is the following strong two-sided division property:

$$(1) \quad a \neq 0 \implies \exists x, y \in R, \quad xay = 1.$$

It is often called the single-sandwich property. It is available in the purely infinite simple setting relevant here [2].

Theorem A (Coordinate-block factorization). *Let R be a nontrivial ring satisfying (1), let $n \geq 2$, and fix a coordinate $j < n$. Every $A \in \mathrm{GL}_n(R)$ admits a factorization of length at most $2n + 6$ in which every factor is either an elementary transvection or belongs to the coordinate copy*

$$\mathrm{GL}_{n-1}^{(j)}(R) = \{B \in \mathrm{GL}_n(R) : Be_j = e_j, e_j^t B = e_j^t\}.$$

PropertyTT/PaperStatements
PropertyTTPaper.
coordinateBlock_factorization

Theorem A is deliberately not called bounded elementary generation. It is a relative bounded-width statement, with respect to elementary root subgroups and one fixed coordinate block. This is exactly the form required by the quasi-cocycle argument.

For the synthesis theorem, a *binary Leavitt family* in R means elements x_0, x_1, y_0, y_1 satisfying

$$y_i x_j = \delta_{ij}, \quad x_0 y_0 + x_1 y_1 = 1.$$

It supplies the matrix self-similarity used to move between ranks.

Theorem B (Rigidity synthesis). *Let R be a nontrivial finite-type $\mathbb{F}_2$ -algebra. Assume that*

- (1) R contains a binary Leavitt family;
- (2) R satisfies the single-sandwich property (1);
- (3) $\text{diag}(u, 1) \in E_2(R)$ for every $u \in R^\times$.

Then $E_n(R)$ has property $(TT)/T$ for every $n \geq 3$.

PropertyTT/PaperStatements
PropertyTTPaper.finiteType_
elementaryGroup_hasTmodT

Here finite type is the fourth structural hypothesis: it is stated in the opening sentence rather than repeated in the numbered list. The Lean theorem has precisely these hypotheses. In particular, there is no residual-finiteness, soficity, approximation, or unstated regularity assumption in Theorem B.

Theorem C (Binary Leavitt algebra). *For every $n \geq 3$,*

$$E_n(L_{\mathbb{F}_2}(1, 2)) \text{ has property } (TT)/T.$$

PropertyTT/PaperStatements
PropertyTTPaper.binaryLeavitt_
elementaryGroup_hasTmodT_and_
not_isSofic

Combining Theorem C with the independently known nonsocicity theorem gives the following striking, but logically secondary, consequence.

Corollary 1.1. *For every $n \geq 3$, the group $E_n(L_{\mathbb{F}_2}(1, 2))$ is nonsocic and has property $(TT)/T$.*

PropertyTT/NonsocicCorollary
PropertyTTPaper.binaryLeavitt_
elementaryGroup_hasTmodT_and_
not_isSofic

The contribution of this paper to Corollary 1.1 is the $(TT)/T$ assertion. The nonsocicity assertion is credited to the prior nonsocic-group construction and is used only as an external mathematical input to the corollary.

Credit and novelty boundary. Several ingredients have strong antecedents, and separating them is mathematically important.

- Mimura introduced property $(TT)/T$, developed its globalization strategy, and proved the relevant relative quasi-cocycle results for universal-lattice settings [5, 6].
- Ershov and Jaikin-Zapirain proved ordinary property (T) for elementary groups over finitely generated associative rings [3].
- Ara, Goodearl, and Pardo established the GE and unit-group K_1 structure for purely infinite simple rings [2].
- Ara, Brustenga, and Cortiñas computed the stable algebraic K -theory of Leavitt path algebras [1]. In particular, $K_1(L_k(1, 2)) = 0$ for a field k .

Accordingly, we do not claim the stable K_1 -vanishing, the GE property, the bare equality $\text{GL}_2(L_{\mathbb{F}_2}(1, 2)) = E_2(L_{\mathbb{F}_2}(1, 2))$, or the abstract relative-rigidity architecture as new theorems. The main contributions isolated here are the explicit width $2n + 6$ coordinate-block factorization and its synthesis with quasi-cocycle control into Theorem B. The direct root and rank-two arguments give quantitative, self-contained routes to inputs which also have prior structural explanations.

Formal verification. The accompanying Lean development is available at https://github.com/SauersML/nonsofic_existence. Margin notes link theorem statements to declarations. The paper-facing declarations are collected in `PropertyTT/PaperStatements.lean`, `K0ne/PaperStatements.lean`, and `PropertyTT/NonsoficCorollary.lean`. The endpoint audit reports only Lean’s standard logical axioms `propext`, `Classical.choice`, and `Quot.sound`.

2. QUASI-COCYCLES AND RELATIVE CONTROL

Let G be a group and let $\pi : G \rightarrow \mathcal{U}(\mathcal{H})$ be a unitary representation on a complex Hilbert space. A map $b : G \rightarrow \mathcal{H}$ is a *quasi-cocycle of defect at most D* if

$$(2) \quad \|b(gh) - b(g) - \pi(g)b(h)\| \leq D \quad (g, h \in G).$$

An exact cocycle has $D = 0$; it is the translation part of an affine isometric action. A quasi-cocycle permits uniformly bounded failure of the affine action law.

Definition 2.1. A group G has property $(TT)/T$ if, for every unitary representation π with no nonzero invariant vector, every Hilbert-space quasi-cocycle into π is bounded on G .

Proposition 2.2 (Mimura’s relative input). *Let X be finite and $A = \mathbb{F}_2\langle X \rangle$. The pair*

$$(E_4(A), X_{03})$$

has relative property $(TT)/T$.

Proof. Mimura proves relative $(TT)/T$ for the noncommutative universal lattice pair in [6, Proposition 9.2.8]:

$$E_m(\mathbb{Z}\langle X \rangle) \geq E_{m-1}(\mathbb{Z}\langle X \rangle), \quad m \geq 3$$

Relative $(TT)/T$ descends through a surjective homomorphism: pull a representation and quasi-cocycle back along the quotient map and then apply the relative bound. Reduce coefficients modulo 2, take $m = 4$, and conjugate the standard rank-three block so that it contains X_{03} . Boundedness on that block implies boundedness on the root subgroup. $\square$

The local input used below is slightly stronger than a relative $(TT)/T$ statement because no hypothesis is imposed on invariant vectors.

Theorem 2.3 (Root quasi-cocycle estimate). *Let X be finite, put $A = \mathbb{F}_2\langle X \rangle$, and let*

$$G = E_4(A), \quad X_{03} = \{I + aE_{03} : a \in A\}.$$

For every unitary representation π , every $D \geq 0$, and every quasi-cocycle $b : G \rightarrow \mathcal{H}$ of defect at most D , the set $b(X_{03})$ is bounded.

Proof. Let N be the last-column subgroup generated by X_{03}, X_{13}, X_{23} . Decompose

$$\mathcal{H} = \mathcal{H}^N \oplus (\mathcal{H}^N)^\perp, \quad b(r) = p + w$$

for $r = I + aE_{03}$. Since the coefficient ring has characteristic two, $r^2 = 1$. Applying (2) to r^2 , and projecting to $\mathcal{H}^N$, bounds the fixed component:

$$(3) \quad \|p\| \leq D.$$

For the moving component, let Q_X be the finite rank-three control set used in the two coordinate embeddings and put

$$M = \sum_{s \in Q_X} (\|b(\iota_{123}(s))\| + \|b(\iota_{012}(s))\|), \quad B = 2M + 2D,$$

$$K = 2(6|X| + 6)(B + 1).$$

If a control element s commutes with r and normalizes N , comparison of the quasi-cocycle equations for sr and rs , followed by orthogonal projection, gives

$$\|\pi(s)w - w\| \leq \|\pi(s)b(r) - b(r)\| \leq 2\|b(s)\| + 2D \leq B.$$

The formalized rank-three plane estimate propagates these finitely many inequalities to the complete root planes in the two embeddings. In particular, for every $c \in A$,

$$\|\pi(x_{13}(c))w - w\|, \quad \|\pi(x_{23}(c))w - w\|, \quad \|\pi(x_{01}(c))w - w\| \leq K.$$

The Steinberg relation

$$[I + cE_{01}, I + E_{13}] = I + cE_{03}$$

and the unitary commutator estimate give displacement at most $4K$ under $x_{03}(c)$. Every element of N is a product of one element from each of X_{03}, X_{13}, X_{23} , so its displacement of w is at most $6K$. Since $w \perp \mathcal{H}^N$, the uniform-orbit projection lemma gives $\|w\| \leq 6K$. Together with (3), this is the formalized estimate

$$\|b(I + aE_{03})\| \leq D + 12(6|X| + 6)(2M + 2D + 1),$$

uniformly in $a \in A$. The numerical constant is not optimized. $\square$

Remark 2.4 (Literature comparison). The bare statement of Theorem 2.3 also follows much more quickly from Proposition 2.2. Split $\mathcal{H} = \mathcal{H}^G \oplus (\mathcal{H}^G)^\perp$. Mimura's relative theorem bounds the second projection on X_{03} . On the invariant summand the action is trivial, while $r^2 = 1$ for $r \in X_{03}$; the quasi-cocycle identities for 1 and r^2 then give $\|b_0(r)\| \leq D$. Thus we make no priority claim for the qualitative root theorem. The longer proof above is included because it is the proof formalized in Lean and records an explicit bound.

3. COORDINATE-BLOCK FACTORIZATION

We now prove the main algebraic input. Write

$$e_{ij}(a) = I + aE_{ij} \quad (i \neq j)$$

for an elementary transvection.

Lemma 3.1 (Five-move pivot). *Let R satisfy (1), let $A \in \mathrm{GL}_n(R)$, and fix distinct coordinates i, j . At most two elementary row operations and at most three elementary column operations produce a matrix B satisfying*

$$B_{jj} = 1, \quad B_{ij} = B_{ji} = 0.$$

PropertyTT/PaperStatements
PropertyTTPaper.fiveMove_pivot

Proof. We give the calculation, since it is the source of the uniform constant. Choose two distinct indices i, j . First arrange that the entry $a = A_{ii}$ is nonzero. If it is zero, invertibility of A gives k with $A_{ik} \neq 0$; right multiplication by $e_{ki}(1)$ replaces A_{ii} by A_{ik} . This costs at most one preliminary column operation.

Write $b = A_{ij}$, and choose $p, q \in R$ with $paq = 1$. Put

$$\rho = q(1 - pb), \quad F_1 = e_{ij}(\rho), \quad A_1 = AF_1, \quad b_1 = (A_1)_{ij}.$$

Then

$$b_1 = a\rho + b, \quad pb_1 = paq(1 - pb) + pb = 1.$$

Set

$$d_1 = (A_1)_{jj}, \quad w = (1 - d_1)p, \quad E_1 = e_{ji}(w), \quad A_2 = E_1A_1.$$

The (j, j) -entry is now

$$(A_2)_{jj} = d_1 + wb_1 = d_1 + (1 - d_1)pb_1 = 1.$$

Next set

$$E_2 = e_{ij}(-b_1), \quad A_3 = E_2A_2.$$

This clears (i, j) without changing the new 1 at (j, j) . Finally, with $c_3 = (A_3)_{ji}$, put

$$F_2 = e_{ji}(-c_3), \quad A_4 = A_3F_2.$$

This clears (j, i) , again preserving $(j, j) = 1$ and $(i, j) = 0$. Thus the four main moves use two row and two column transvections; including the possible preliminary column addition gives the asserted bounds 2 and 3. No stable-range or commutativity hypothesis is used. $\square$

Theorem 3.2. *Under the hypotheses of Theorem A, every $A \in \mathrm{GL}_n(R)$ can be written*

$$A = g_1 \cdots g_m, \quad m \leq 2n + 6,$$

where each g_i is an elementary transvection or lies in $\mathrm{GL}_{n-1}^{(j)}(R)$.

PropertyTT/PaperStatements
PropertyTTPaper.
coordinateBlock_factorization

Proof. Choose $i \neq j$. Lemma 3.1 gives elementary products E, F , represented by lists ℓ, r , such that

$$B = EAF, \quad B_{jj} = 1, \quad B_{ij} = B_{ji} = 0,$$

with

$$|\ell| \leq 2, \quad |r| \leq 3.$$

The last two displayed zeros are not needed for the subsequent clearing, but they are part of the exact pivot calculation.

For every $k \neq j$, right multiply by

$$e_{jk}(-B_{jk}).$$

These transvections have a common source column and distinct target columns, so they do not interfere with one another. If T is their product and $C = BT$, then

$$C_{jc} = \begin{cases} 1, & c = j, \\ 0, & c \neq j. \end{cases}$$

The list t of right-clearing factors has length at most n .

For every $k \neq j$, left multiply by

$$e_{kj}(-C_{kj}).$$

Again the operations have distinct target rows and do not interfere. If S is their product and $D = SC$, then

$$D_{rc} = \begin{cases} 1, & r = c = j, \\ 0, & r = j, c \neq j, \\ 0, & r \neq j, c = j, \\ C_{rc}, & r, c \neq j. \end{cases}$$

Hence $D \in \mathrm{GL}_{n-1}^{(j)}(R)$. The list s of left-clearing factors also has length at most n .

Solving $D = SEAF$ for A gives

$$A = E^{-1}S^{-1}DT^{-1}F^{-1}.$$

At the word level, inverse products are represented by reversing the factor lists and inverting every transvection. The inverse of $e_{ab}(z)$ is $e_{ab}(-z)$, so all these factors remain elementary. The resulting word has the exact form

$$\ell_{\mathrm{rev}}^{-1} s_{\mathrm{rev}}^{-1} [D] t_{\mathrm{rev}}^{-1} r_{\mathrm{rev}}^{-1},$$

and therefore length at most

$$2 + n + 1 + n + 3 = 2n + 6.$$

□

Remark 3.3. The fixed coordinate j is chosen before A . Allowing the coordinate block to depend on A would give a weaker statement and would not be sufficient for the uniform quasi-cocycle globalization used below.

4. FROM RELATIVE CONTROL TO GLOBAL RIGIDITY

Let R be a finite-type $\mathbb{F}_2$ -algebra. Choose a finite set X and a surjection

$$\mathbb{F}_2\langle X \rangle \rightarrow R.$$

Elementary matrices are functorial in the coefficient ring, so the root estimate of Theorem 2.3 descends to R . Coordinate permutations then give the same estimate for every root subgroup in rank four.

Two larger pieces are used in the assembly: a last-column subgroup and a last-row subgroup. The quasi-cocycle is bounded on both. The stabilized rank-three coordinate group normalizes these pieces. Kazhdan control for the rank-three group converts boundedness on the normalized root system into boundedness on the coordinate group, provided the ambient representation has no invariant vectors. This is the only place where the quotient by the trivial representation enters.

Lemma 4.1 (Elementary equals general linear). *Assume that R has a binary Leavitt family, satisfies (1), and satisfies $\text{diag}(u, 1) \in E_2(R)$ for every $u \in R^\times$. Then*

$$E_n(R) = \text{GL}_n(R) \quad (n \geq 2).$$

Proof. The single-sandwich elimination reduces an arbitrary element of $\text{GL}_2(R)$, by elementary row and column operations, to $\text{diag}(u, 1)$ for some $u \in R^\times$. The diagonal hypothesis therefore gives $\text{GL}_2(R) = E_2(R)$. A binary Leavitt family gives prefix-code coefficient isomorphisms $M_m(R) \cong R$. Apply the rank-two equality after transporting coefficients to $M_n(R)$, flatten the resulting block matrices, swap the two finite index factors, unflatten in the other direction, and transport coefficients back through $M_2(R) \cong R$. This is the coefficient/block argument which gives the assertion in every rank $n \geq 2$. $\square$

Proposition 4.2 (Rank-four assembly). *Under the hypotheses of Theorem B, $E_4(R)$ has property $(TT)/T$.*

Proof. Choose a finite free-algebra presentation of R . Theorem 2.3 descends through this quotient and gives relative $(TT)/T$ for $X_{03}(R)$. Coordinate permutations give it for the six roots

$$X_{03}, X_{13}, X_{23}, X_{30}, X_{31}, X_{32}.$$

Property (T) for $E_3(R)$ follows from Ershov–Jaikin–Zapirain, or equivalently by descending the finite Kazhdan system for the free algebra. The Leavitt rank equivalence transports this to $E_4(R)$. The standard rank-four parabolic argument now combines the six relative root bounds: the last-row and last-column pieces are bounded, their normalized rank-three core has almost-invariant quasi-cocycle values, and the Kazhdan estimate bounds those values. Hence $E_4(R)$ has relative $(TT)/T$ with respect to the stabilized $E_3(R)$.

PropertyTT/PaperStatements
PropertyTTPaper.
elementaryGroup_eq_
generalLinear

PropertyTT/PaperStatements
PropertyTTPaper.
finitePresentation_rankFour_
hasTTmodT

By Lemma 4.1, the elementary groups in ranks three and four are the full general linear groups. Under these identifications the stabilized core is exactly the coordinate block $\mathrm{GL}_3^{(3)}(R)$. Finally, coordinate permutations carry this relative statement to $\mathrm{GL}_3^{(j)}(R)$ for every $j \in \{0, 1, 2, 3\}$.

Each elementary transvection lies in a coordinate block avoiding its two indices. Hence the union of one fixed coordinate block and all root subgroups is bounded for the quasi-cocycle. Theorem A writes every element of $\mathrm{GL}_4(R)$ as a product of at most (14) elements of that union. Lemma 4.3 below, or directly the corresponding bounded-product lemma, makes the quasi-cocycle bounded on $\mathrm{GL}_4(R)$. Transporting through $E_4(R) = \mathrm{GL}_4(R)$ proves the proposition. $\square$

The passage from a bounded generating subset to the whole group uses the following elementary estimate.

Lemma 4.3 (Products of controlled pieces). *Let b be a quasi-cocycle of defect at most D , and suppose $\|b(g_i)\| \leq C$ for a list $(g_1, \dots, g_m)$. Then*

$$\|b(g_1 \cdots g_m)\| \leq mC + (m+1)D.$$

Proof. Induct on the list. The empty-list case is $\|b(1)\| \leq D$. For a nonempty list, apply (2) to the head and the product of the tail, use that the representation is isometric, and invoke the inductive bound. The displayed estimate is the slightly loose uniform form used by the Lean bounded-product lemma; importantly, it also covers the empty list. $\square$

Proof of Theorem B. Proposition 4.2 gives the result in rank four. For arbitrary $n \geq 3$, the binary Leavitt family supplies an explicit group equivalence

$$E_n(R) \cong E_4(R).$$

Property $(TT)/T$ is invariant under group equivalence, so the rank-four conclusion transports to rank n . $\square$

5. THE BINARY LEAVITT SPECIALIZATION

Let L be the binary Leavitt algebra [4],

$$L = L_{\mathbb{F}_2}(1, 2) = \mathbb{F}_2\langle x_0, x_1, y_0, y_1 \rangle / (y_i x_j - \delta_{ij}, x_0 y_0 + x_1 y_1 - 1).$$

The defining generators themselves form a binary Leavitt family, and the presentation makes L a finite-type $\mathbb{F}_2$ -algebra. Pure infiniteness and simplicity give the single-sandwich property [2]. It remains to discuss the unstable diagonal condition.

Proposition 5.1 (Unstable diagonal reduction). *For every $u \in L^\times$,*

$$\mathrm{diag}(u, 1) \in E_2(L).$$

Consequently $\mathrm{GL}_2(L) = E_2(L)$.

PropertyTT/PaperStatements
PropertyTTPaper.quasiCocycle_
list_product_bound

KOne/PaperStatements
KOnePaper.diagUnit_mem_
elementary
KOne/PaperStatements
KOnePaper.elementaryGroup_two_
eq_top

5.1. What follows from existing structure. The conclusion of Proposition 5.1 should not be confused with a new computation of stable K_1 . Ara–Brustenga–Cortiñas give, for a regular coefficient ring, the exact sequence determined by $1 - N_E^t$ [1]. For the graph with one vertex and two loops this map is multiplication by -1 , hence

$$K_i(L_k(1, 2)) = 0$$

in every degree covered by their theorem, in particular in degree one.

There is also an unstable route already implicit in the structural literature. Ara–Goodearl–Pardo prove that a unital purely infinite simple ring is a GE-ring and that

$$K_1(R) \cong R_{ab}^\times$$

This is Theorem 2.3 of [2]. Thus $K_1(L) = 0$ makes each unit a product of commutators. The standard 2×2 Whitehead identity places $\text{diag}([a, b], 1)$ in $E_2(L)$; the GE property then gives $\text{GL}_2(L) = E_2(L)$. For this reason we claim no priority for the bare equality.

5.2. The direct formalized route. The Lean development instead gives a direct tree-and-pencil reduction. Finite prefix codes encode the self-similar pieces of a unit. Elementary row and column moves refine the two codes to a common tree, isolate code-scalar blocks, and reduce the remaining one-sided factor. The output is an actual finite list of elementary matrices whose product is $\text{diag}(u, 1)$.

Proposition 5.2 (Finite factor witness). *For every $u \in L^\times$, there exists a finite list $(e_1, \dots, e_m)$ of elementary 2×2 matrices such that*

$$\text{diag}(u, 1) = e_1 \cdots e_m.$$

`KOne/PaperStatements`
`KOnePaper.exists_diagUnit_`
`elementaryFactors`

Proposition 5.2 is an existence theorem with a constructive proof object. We do not claim a uniform word-length bound or a complexity estimate: neither is present in the theorem statement.

Proof of Theorem C. The binary Leavitt algebra is a nontrivial finite-type $\mathbb{F}_2$ -algebra, its canonical generators form a binary Leavitt family, pure infiniteness supplies (1), and Proposition 5.1 supplies the elementary diagonal condition. Theorem B applies. $\square$

6. FORMAL CORRESPONDENCE AND TRUST BOUNDARY

The public Lean interface is intentionally smaller than the implementation. Table 1 records the exact correspondence between the paper and the checked declarations.

The synthesis theorem also has the presentation-facing declaration

`PropertyTTPaper.finitePresentation_elementaryGroup_hasTModT,`

Paper statement	Lean declaration
Theorem 2.3	<code>PropertyTTPaper.freeCharacteristicTwo_root_hasRelativeTT</code>
Theorem A	<code>PropertyTTPaper.coordinateBlock_factorization</code>
Lemma 3.1	<code>PropertyTTPaper.fiveMove_pivot</code>
Lemma 4.1	<code>PropertyTTPaper.elementaryGroup_eq_generalLinear</code>
Proposition 4.2	<code>PropertyTTPaper.finitePresentation_rankFour_hasTTmodT</code>
Lemma 4.3	<code>PropertyTTPaper.quasiCocycle_list_product_bound</code>
Theorem B	<code>PropertyTTPaper.finiteType_elementaryGroup_hasTTmodT</code>
Theorem C	<code>PropertyTTPaper.binaryLeavitt_elementaryGroup_hasTTmodT</code>
Proposition 5.1	<code>KOnePaper.diagUnit_mem_elementary</code> and <code>KOnePaper.elementaryGroup_two_eq_top</code>
Proposition 5.2	<code>KOnePaper.exists_diagUnit_elementaryFactors</code>
Corollary 1.1	<code>PropertyTTPaper.binaryLeavitt_elementaryGroup_hasTTmodT_and_not_isSofic</code>

TABLE 1. Paper-to-Lean theorem correspondence.

whose hypotheses exhibit a finite free-algebra quotient explicitly. The finite-type wrapper used in Theorem B constructs that presentation internally.

No theorem in this paper assumes soficity, residual finiteness, or any approximation property. The only statement importing the nonsolic development is Corollary 1.1; it is placed in a separate Lean module so that the rigidity dependency graph remains independent. The formalization does not assert historical novelty. Credit claims are prose statements justified by the literature review, not kernel-checkable mathematical propositions.

7. FURTHER DIRECTIONS

The characteristic-two hypothesis enters most visibly in (3), where every root element has order two. Over a field of characteristic p , the corresponding root has order p , and iterating the quasi-cocycle identity gives

$$b(1) \approx b(r) + \pi(r)b(r) + \cdots + \pi(r)^{p-1}b(r).$$

On the root-fixed space this is p times the fixed projection of $b(r)$, with error at most $(p-1)D$. Thus the fixed-space argument has a plausible

finite-characteristic analogue. The remaining obstacle is to replace the characteristic-two rank-three control and sign-free coordinate symmetries uniformly. We do not claim such a generalization here.

The algebraic factorization theorem suggests a separate question. For which natural classes of simple or purely infinite rings does the single-sandwich condition hold, and when can the coordinate block in Theorem A itself be controlled by elementary subgroups? A satisfactory answer would turn the present synthesis theorem into a broad rigidity criterion rather than a theorem whose most memorable instance is a Leavitt algebra.

Acknowledgements. We thank the authors whose work supplies the foundations used here. In particular, Mimura’s quasi-cocycle rigidity framework, Ershov–Jaikin-Zapirain’s property- (T) theorem, the structural results of Ara–Goodearl–Pardo, and the K -theory computation of Ara–Brustenga–Cortiñas are indispensable to the present proof and to its correct interpretation.

REFERENCES

- [1] P. Ara, M. Brustenga, and G. Cortiñas, *K-theory of Leavitt path algebras*, Münster J. Math. **2** (2009), 5–34.
- [2] P. Ara, K. R. Goodearl, and E. Pardo, *K_0 of purely infinite simple regular rings*, K-Theory **26** (2002), no. 1, 69–100.
- [3] M. Ershov and A. Jaikin-Zapirain, *Property (T) for noncommutative universal lattices*, Invent. Math. **179** (2010), no. 2, 303–347.
- [4] W. G. Leavitt, *The module type of a ring*, Trans. Amer. Math. Soc. **103** (1962), 113–130.
- [5] M. Mimura, *Property (TT) modulo T and homomorphism superrigidity into mapping class groups*, Monatsh. Math. **178** (2015), no. 4, 573–600.
- [6] M. Mimura, *Rigidity theorems for universal and symplectic universal lattices*, doctoral thesis, University of Tokyo, 2011, [doi:10.15083/00004046](https://doi.org/10.15083/00004046).