

NON-MF GROUPS

SAUERS

Some groups are not MF. For example, $H = \text{EL}_{12}(L_{\mathbb{F}_2}(1, 2))$ is a simple property-(T) group, and every homomorphism from H to an MF group is trivial. We also construct a two-generated, finitely presented, torsion-free property-(T) group with the same rigidity, and an affine-Clifford group W that is locally residually finite-by- $\mathbb{Z}$ but not MF. The common mechanism is a normal property-(T) subgroup inside the commutators of a one-sided compression. Consequently, $C_r^*(H)$ is separable, stably finite, and not MF, while the canonical trace of W is amenable but not quasidiagonal.

Draft. This manuscript is under revision.

1. INTRODUCTION

The counterexample is

$$H = \text{EL}_{12}(L_{\mathbb{F}_2}(1, 2)),$$

the elementary group of 12×12 matrices over the binary Leavitt algebra. Khanh–Thanh show that H is isomorphic to the unit group of the binary Leavitt algebra [K–T, Proposition 4.2 and Corollary 4.4], which OpenAI used to construct the first nonscic group [OAI, Chapter 3].

A countable group G is MF [CDE] if there are positive integers d_n and maps $V_n: G \rightarrow \mathcal{U}(d_n)$, with $V_n(1) = 1$, such that

$$\|V_n(gh) - V_n(g)V_n(h)\| \rightarrow 0 \quad (g, h \in G),$$

and

$$\limsup_n \|V_n(g) - 1\| > 0 \quad (g \in G \setminus \{1\}).$$

In other words, G is MF if there are finite unitary matrices, one for each element of G , that multiply correctly up to an error tending to zero in operator norm while every element other than the identity stays a fixed distance from the identity matrix. The separation may equivalently be required along the full sequence with a constant independent of g [Kor, Propositions 2 and 7]. The strong convergence convention of [GKM, Sch] also requires the models to reproduce the operator norms of the left regular representation, so a group that is not MF as defined here is not MF in that convention either. Equivalently, G embeds in the unitary group of the norm matrix corona

$$\mathcal{Q}_d = \prod_n M_{d_n}(\mathbb{C}) / \bigoplus_n M_{d_n}(\mathbb{C}),$$

2020 *Mathematics Subject Classification*. Primary 46L05; Secondary 20F65, 22D10, 16S50.

Key words and phrases. MF groups, norm matrix coronas, property (T), asymptotic representations, Leavitt algebra.

where $\prod_n M_{d_n}(\mathbb{C})$ is the C^* -algebra of bounded sequences, as every product of C^* -algebras below, and $\bigoplus_n M_{d_n}(\mathbb{C})$ is its ideal of sequences with $\|x_n\| \rightarrow 0$. The quotient $\mathcal{Q}_{\mathbf{d}}$ is the corona algebra of $\bigoplus_n M_{d_n}(\mathbb{C})$, since $\prod_n M_{d_n}(\mathbb{C})$ is its multiplier algebra. We call a homomorphism $G \rightarrow \mathcal{U}(\mathcal{Q}_{\mathbf{d}})$ a *corona homomorphism*. A separable C^* -algebra is MF if it embeds in a norm matrix corona [BK].

The groups here fail to be MF because they contain a subgroup that is conjugated into itself. For $L \leq G$ put

$$\text{Comp}_G(L) = \{u \in G : uLu^{-1} \leq L\}.$$

An element c of the centralizer $C_G(L)$ commutes with L , so ucu^{-1} commutes with uLu^{-1} but need not commute with the rest of L . The normal subgroup of G generated by the commutators of all such elements ucu^{-1} with elements of L is

$$(1) \quad \mathfrak{D}_G(L) = \langle\langle [ucu^{-1}, \ell] : u \in \text{Comp}_G(L), c \in C_G(L), \ell \in L \rangle\rangle_G.$$

Theorem A (one-sided compression criterion). *Let G be countable and let $L \leq G$ have property (T). If $K \trianglelefteq G$ has property (T) and $K \leq \mathfrak{D}_G(L)$, then every homomorphism from G to an MF group is trivial on K . In particular, a nontrivial such K makes G non-MF, and if G has property (T) and $\mathfrak{D}_G(L) = G$, then every homomorphism from G to an MF group is trivial.*

Asymptotic multiplicativity in operator norm makes the conjugation maps $\text{Ad}(V_n(g)) : x \mapsto V_n(g)xV_n(g)^*$ an asymptotic unitary representation of G on $M_{d_n}(\mathbb{C})$ with its normalized Hilbert–Schmidt inner product, because $\|\text{Ad}(A) - \text{Ad}(B)\| \leq 2\|A - B\|$. Property (T) applies to unitary representations on Hilbert spaces, so it applies to this one, and its conclusions are estimates in Hilbert–Schmidt norm. The unitary U of the compressor u satisfies $U^*PU \leq P$ for the Kazhdan projection P of L , because $uLu^{-1} \leq L$ has fewer constraints and more fixed vectors, and stable finiteness of the matrix corona forces $U^*PU = P$. So conjugation by u preserves the Hilbert–Schmidt asymptotic commutant of L , and every element of $\mathfrak{D}_G(L)$ tends to 1 in Hilbert–Schmidt norm in every operator norm asymptotic representation (Corollary 2.5). Property (T) of K turns this into triviality in operator norm: a corona homomorphism nontrivial on K compresses to the corner where K has no fixed vectors, and the Kazhdan inequality there contradicts the Hilbert–Schmidt triviality (Theorem 2.7).¹

The stable letter of an ascending HNN extension of a property-(T) group along a proper self-embedding is a compressor. The group H realizes the compression inside a simple group, where one nontrivial commutator $[ucu^{-1}, \ell]$ normally generates everything. Let $L_{\mathbb{F}_2}(1, 2)$ denote the binary Leavitt algebra, the unital $\mathbb{F}_2$ -algebra generated by s_0, s_1, t_0, t_1 subject to

$$(2) \quad t_i s_j = \delta_{ij}, \quad s_0 t_0 + s_1 t_1 = 1.$$

For a unital ring R , write $e_{ij}(a) = 1 + aE_{ij}$ and let $\text{EL}_n(R)$ be the subgroup of $\text{GL}_n(R)$ generated by the elements $e_{ij}(a)$, $i \neq j$.

¹A Hilbert–Schmidt bound on the multiplicative defect does not give an operator norm bound on the conjugation maps, so this argument does not apply to sofic or hyperlinear approximations.

Theorem B (the binary Leavitt group). *Put $R = L_{\mathbb{F}_2}(1, 2)$ and $H = \text{EL}_{12}(R)$. Then H is finitely generated, nontrivial, simple, has property (T), and every homomorphism from H to an MF group is trivial. So H is not MF, and $C_r^*(H)$ is separable and stably finite but not MF, while $C_{\max}^*(H)$ is not even finite: it contains a proper isometry.*

Theorem C (a torsion-free finitely presented example). *There is a two-generated, finitely presented, torsion-free, acylindrically hyperbolic group Q with property (T) such that every homomorphism from Q to an MF group is trivial. In particular, no nontrivial quotient of Q is MF.*

Section 5 builds Q from Fournier-Facio’s torsion-free property-(T) group [FF, §2] and Hull’s small cancellation theorem [Hu].

We construct an affine-Clifford group W that is locally residually finite-by- $\mathbb{Z}$ but not MF. Its canonical trace on $C_{\max}^*(W)$ is amenable and not quasidiagonal. This answers Brown’s question whether every amenable trace is quasidiagonal [Br, discussion preceding Proposition 3.5.1].

Relation to prior work. Blackadar and Kirchberg introduced MF algebras and proved that a separable C^* -algebra is NF if and only if it is nuclear and MF [BK]. Whether every separable stably finite C^* -algebra is MF, the MF problem [GH, §6], was answered negatively through the failure of the Connes embedding problem [JNVWY][GH, Proposition 6.1 and Remark 6.2]. Theorem B gives a counterexample among reduced group C^* -algebras. Whether every countable group is MF was a longstanding open problem [BDL].

Theorem A uses the configuration of the nonsofic construction [OAI, Proposition 2.3]: a property-(T) subgroup L conjugated into itself by an element u , and an element c commuting with L whose conjugate ucu^{-1} does not. There, soficity and the rigidity theorems of Kun [Ku16] and Kun–Thom [KT19] force the centralizing subgroup to be locally embeddable into finite groups, and a copy of Thompson’s group V gives the contradiction. Here the rigidity is the Kazhdan projection of L in a norm matrix corona, which stable finiteness of the corona forces to commute with the compressor, and the contradiction is a commutator $[ucu^{-1}, \ell] \neq 1$ that every operator norm asymptotic representation sends to 1 in Hilbert–Schmidt norm. The group H is the unit group of the binary Leavitt algebra, the group of [OAI], and Theorem C reuses Fournier-Facio’s configuration [FF]. Bachner–Dogon–Lubotzky showed that for groups of Deligne type, operator–Hilbert–Schmidt stability would imply that the group is not MF [BDL, Proposition 1.5]. The compression relation replaces stability. For a short elementary construction of non-MF generalized wreath products from one-sided compression, see [Ec].

Leavitt introduced the algebras $L_K(1, n)$ [Lev]. The binary Leavitt algebra is purely infinite simple [AA], is an exchange ring [Ara], and has center $\mathbb{F}_2$ [AC]. Preusser’s sandwich theorem then gives simplicity of H [Pre], and property (T) is Ershov–Jaikin-Zapirain’s theorem [EJZ, Theorem 1.1].

2. ONE-SIDED COMPRESSION

2.1. Corona homomorphisms. The image of a corona homomorphism from G is a countable subgroup of a corona unitary group, so MF, and every countable MF group embeds in the unitary group of a norm matrix corona.

Lemma 2.1. *Let G be countable and $K \leq G$. Every homomorphism from G to an MF group is trivial on K if and only if every corona homomorphism from G is trivial on K .*

Proof. A corona homomorphism is a homomorphism to an MF group, and a homomorphism to an MF group composed with an embedding of that group into a corona unitary group is a corona homomorphism with the same kernel. $\square$

2.2. Kazhdan transport in normalized Hilbert–Schmidt norm. For $a \in M_d(\mathbb{C})$ write

$$\|a\|_2 = \text{tr}_d(a^*a)^{1/2}, \quad \text{tr}_d = \frac{1}{d} \text{Tr}.$$

An *operator norm asymptotic representation* of a countable group G is a sequence of maps $V_n: G \rightarrow \mathcal{U}(d_n)$ such that $V_n(1) = 1$ and

$$\|V_n(gh) - V_n(g)V_n(h)\| \rightarrow 0 \quad (g, h \in G).$$

The elements g with $\|V_n(g) - 1\|_2 \rightarrow 0$ form a normal subgroup of G , because $\|a\|_2 \leq \|a\|$, the Hilbert–Schmidt norm is invariant under adjoints and unitary conjugation, and $\|V_n(g^{-1}) - V_n(g)^*\| \rightarrow 0$:

$$\|V_n(gh) - 1\|_2 \leq \|V_n(g) - 1\|_2 + \|V_n(h) - 1\|_2 + o(1),$$

$$\|V_n(g^{-1}) - 1\|_2 = \|V_n(g) - 1\|_2 + o(1), \quad \|V_n(ghg^{-1}) - 1\|_2 = \|V_n(h) - 1\|_2 + o(1).$$

For $L \leq G$, let

$$\mathcal{C}_2(V, L) = \left\{ (x_n) \left| \begin{array}{l} \sup_n \|x_n\| < \infty, \\ \|[V_n(\ell), x_n]\|_2 \rightarrow 0 \quad (\ell \in L) \end{array} \right. \right\}$$

be the bounded Hilbert–Schmidt asymptotic commutant of $V(L)$. Coordinatewise conjugation by $V_n(g)$ defines a bijection, denoted $\text{Ad}(V(g))$, of the bounded matrix sequences.

Lemma 2.2. *For every sequence of positive integers (m_n) , the norm matrix corona*

$$\prod_n M_{m_n}(\mathbb{C}) / \bigoplus_n M_{m_n}(\mathbb{C})$$

is stably finite. Consequently, unitarily equivalent projections in a norm matrix corona are equal whenever one is dominated by the other.

Proof. Write A for the displayed corona. If $v^*v = 1$ in A and (x_n) is a bounded lift of v , then $\|x_n^*x_n - 1\| \rightarrow 0$, so for all large n the matrix $u_n = x_n(x_n^*x_n)^{-1/2}$ is unitary and $\|u_n - x_n\| \rightarrow 0$; after assigning arbitrary unitary values to the finitely many remaining coordinates, (u_n) is a unitary lift of v , and $vv^* = 1$. So A is finite, and so is $M_k(A)$, which is the corona with matrix sizes km_n . Finally, if $p \leq q$ are equivalent

projections in a stably finite algebra B and w is a partial isometry with $w^*w = q$ and $ww^* = p$, then $w \in qBq$ is an isometry in the finite corner qBq , so $p = ww^* = q$. $\square$

Lemma 2.3 (one-sided order for the Kazhdan projection). *Let L have property (T), let B be a unital C^* -algebra, and let $\pi: L \rightarrow \mathcal{U}(B)$ be a homomorphism. Denote by $P \in B$ the image of the Kazhdan projection under the extension $C_{\max}^*(L) \rightarrow B$. If $U \in \mathcal{U}(B)$ satisfies*

$$U\pi(L)U^* \subseteq \pi(L),$$

then

$$U^*PU \leq P.$$

Proof. Represent B faithfully and nondegenerately on a Hilbert space $\mathcal{H}$. Then P is the orthogonal projection onto $\text{Fix } \pi(L)$. If ξ is L -fixed and $\ell \in L$, then $\pi(\ell)U^*\xi = U^*(U\pi(\ell)U^*)\xi = U^*\xi$ because $U\pi(\ell)U^*$ belongs to $\pi(L)$. So $U^*\text{Fix } \pi(L) \subseteq \text{Fix } \pi(L)$, and the range projection U^*PU is dominated by P in $B(\mathcal{H})$, and then in B by faithfulness. $\square$

Theorem 2.4 (one-sided Kazhdan transport). *Let $L \leq G$ have property (T), and let $u \in G$ satisfy $uLu^{-1} \leq L$. Let (V_n) be an operator norm asymptotic representation of G . Then*

$$\text{Ad}(V(u))(\mathcal{C}_2(V, L)) = \mathcal{C}_2(V, L).$$

Proof. Fix $x = (x_n) \in \mathcal{C}_2(V, L)$ and a free ultrafilter ω , and let $\mathcal{K}_\omega$ be the ultraproduct of the Hilbert spaces $(M_{d_n}(\mathbb{C}), \|\cdot\|_2)$. For unitaries A, B ,

$$\|\text{Ad}(A) - \text{Ad}(B)\| \leq 2\|A - B\|,$$

so the maps $\text{Ad}(V_n(g))$ are asymptotically multiplicative in operator norm, and

$$\sigma(g) = \lim_{\omega} \text{Ad}(V_n(g))$$

is a unitary representation of G on $\mathcal{K}_\omega$ with $\xi = [x_n]_\omega \in \text{Fix } \sigma(L)$. The same estimate makes

$$\tilde{\sigma}(g) = [\text{Ad}(V_n(g))]_n \in \mathcal{U}(\mathcal{B}), \quad \mathcal{B} = \prod_n B(M_{d_n}(\mathbb{C})) / \bigoplus_n B(M_{d_n}(\mathbb{C})),$$

a homomorphism, and $\mathcal{B}$ is a norm matrix corona with coordinate sizes d_n^2 after a choice of matrix units. Let $P \in \mathcal{B}$ be the image of the Kazhdan projection of L [AW] under $C_{\max}^*(L) \rightarrow \mathcal{B}$. On $\mathcal{K}_\omega$ its range is $\text{Fix } \sigma(L)$. Put $U = \tilde{\sigma}(u)$. Since $uLu^{-1} \leq L$,

$$U\tilde{\sigma}(L)U^* \subseteq \tilde{\sigma}(L), \quad \text{so} \quad U^*PU \leq P$$

by Lemma 2.3, and

$$U^*PU = P$$

by Lemma 2.2, since the two projections are unitarily equivalent. So U commutes with P , and

$$U\xi, U^*\xi \in \text{Fix } \sigma(L), \quad \text{that is,}$$

$$\lim_{\omega} \left\| [V_n(\ell), V_n(u)^{\pm 1} x_n V_n(u)^{\mp 1}] \right\|_2 = 0 \quad (\ell \in L).$$

Since ω was arbitrary, the limits hold along the full sequence: a subsequence staying above some $\varepsilon > 0$ would lie in a free ultrafilter and give a nonzero ultralimit. So $\text{Ad}(V(u))$ and its inverse both preserve $\mathcal{C}_2(V, L)$. $\square$

Corollary 2.5. *Let $L \leq G$ have property (T) and let (V_n) be an operator norm asymptotic representation of G . Then*

$$\|V_n(d) - 1\|_2 \longrightarrow 0 \quad (d \in \mathfrak{D}_G(L)).$$

Proof. Let $u \in \text{Comp}_G(L)$, $c \in C_G(L)$, and $\ell \in L$, and write $\mathcal{C}_2 = \mathcal{C}_2(V, L)$. Then

$$(V_n(c)) \in \mathcal{C}_2 \implies (V_n(u)V_n(c)V_n(u)^*) \in \mathcal{C}_2 \implies (V_n(ucu^{-1})) \in \mathcal{C}_2,$$

the first membership because c commutes with L , the first implication by Theorem 2.4, and the second because $\|V_n(u)V_n(c)V_n(u)^* - V_n(ucu^{-1})\| \rightarrow 0$. So $\|[V_n(\ell), V_n(ucu^{-1})]\|_2 \rightarrow 0$, that is, $\|V_n([ucu^{-1}, \ell]) - 1\|_2 \rightarrow 0$. The elements with this property form a normal subgroup of G , so it contains $\mathfrak{D}_G(L)$. $\square$

2.3. From Hilbert–Schmidt to operator norm. A corona homomorphism ρ cannot simply be restricted to a corner: a projection q commuting with $\rho(G)$ has lifts q_n that only asymptotically commute with the lifts of $\rho(g)$, so the compressions $q_n\rho(g)q_n$ are only approximately unitary.

Lemma 2.6 (central corona corners). *Let $\rho: G \rightarrow \mathcal{U}(\mathcal{Q}_{\mathbf{d}})$ be a homomorphism from a countable group, and let $q \in \mathcal{Q}_{\mathbf{d}}$ be a nonzero projection commuting with $\rho(G)$. Then, after passing to an infinite coordinate subsequence, there are nonzero projections $q_n \in M_{d_n}(\mathbb{C})$, integers $r_n = \text{rank}(q_n)$, unitary identifications $J_n: \mathbb{C}^{r_n} \rightarrow q_n\mathbb{C}^{d_n}$, and an operator norm asymptotic representation*

$$W_n: G \longrightarrow \mathcal{U}(r_n)$$

with $W_n(1) = I_{r_n}$ such that, in the corona over the retained coordinates, the class of $(J_n W_n(g) J_n^)$ is the coordinate restriction of $q\rho(g)$, a unitary of the corner $q\mathcal{Q}_{\mathbf{d}}q$ with unit q .*

Proof. Lift q to projections q_n by spectral rounding, and lift each $\rho(g)$ to unitaries $U_n(g)$ by polar correction, with $U_n(1) = I_{d_n}$. Since $q \neq 0$, infinitely many q_n are nonzero. Retain those coordinates and put $r_n = \text{rank}(q_n)$. The commutation relation in the corona gives $\|q_n U_n(g) - U_n(g) q_n\| \rightarrow 0$ for each g , so [BDL, Proposition 2.4] with $p = \infty$ gives an operator norm asymptotic representation $\widetilde{W}_n$ of G by unitaries of the corner $q_n M_{d_n}(\mathbb{C}) q_n$ with $\|\widetilde{W}_n(g) - q_n U_n(g) q_n\| \rightarrow 0$. Replacing $\widetilde{W}_n(1)$ by q_n changes it by $o(1)$. Choose unitaries $J_n: \mathbb{C}^{r_n} \rightarrow q_n \mathbb{C}^{d_n}$ and put $W_n(g) = J_n^* \widetilde{W}_n(g) J_n$. Conjugation by J_n preserves operator norms and multiplicative defects, so (W_n) is an operator norm asymptotic representation with $W_n(1) = I_{r_n}$, and $\|J_n W_n(g) J_n^* - q_n U_n(g) q_n\| \rightarrow 0$ identifies the class of $(J_n W_n(g) J_n^*)$ with the coordinate restriction of $q\rho(g)$. $\square$

Theorem 2.7 (normal Kazhdan subgroups). *Let G be countable and let $K \trianglelefteq G$ have property (T). If every operator norm asymptotic representation (V_n) of G satisfies $\|V_n(k) - 1\|_2 \rightarrow 0$ for all $k \in K$, then every corona homomorphism from G is trivial on K .*

Proof. Suppose that a corona homomorphism $\Theta: G \rightarrow \mathcal{U}(\mathcal{Q}_d)$ is nontrivial on K . Replace G by $\Theta(G)$ and K by $\Theta(K)$: the new K is a nontrivial normal property-(T) subgroup, since property (T) passes to quotients, and it still satisfies the hypothesis, because every operator norm asymptotic representation of $\Theta(G)$ pulls back along $G \rightarrow \Theta(G)$ to one of G .

Let $p \in \mathcal{Q}_d$ be the image of the Kazhdan projection of K , the projection onto the K -fixed vectors in any faithful representation of the corona. Since $gKg^{-1} = K$,

$$\Theta(g)p\Theta(g)^* = p \quad (g \in G),$$

and $q = 1 - p$ is nonzero, because $p = 1$ would make every element of K act trivially. Lemma 2.6 gives an operator norm asymptotic representation $W_n: G \rightarrow \mathcal{U}(r_n)$ whose corona homomorphism $\hat{\Theta}$ is the coordinate restriction of $g \mapsto q\Theta(g)$. The Kazhdan projection of K under $\hat{\Theta}$ is the coordinate restriction of $qp = 0$, so $\hat{\Theta}|_K$ has no nonzero fixed vectors, and for a finite Kazhdan set $S \subseteq K$ with constant $\kappa > 0$,

$$b = \frac{1}{|S|} \sum_{s \in S} (\hat{\Theta}(s) - 1)^*(\hat{\Theta}(s) - 1) \geq \frac{\kappa^2}{|S|} 1$$

in $\mathcal{Q}_r$. The coordinates $b_n = \frac{1}{|S|} \sum_{s \in S} (W_n(s) - I_{r_n})^*(W_n(s) - I_{r_n})$ represent b , so the negative part of $b_n - (\kappa^2/|S|)I_{r_n}$ tends to zero in operator norm, and taking normalized traces on $M_{r_n}(\mathbb{C})$ gives

$$\frac{1}{|S|} \sum_{s \in S} \|W_n(s) - I_{r_n}\|_2^2 \geq \frac{\kappa^2}{|S|} - o(1).$$

After passing to a subsequence, one fixed $s_0 \in S$ stays a positive Hilbert–Schmidt distance from I_{r_n} , against the hypothesis applied to (W_n) and $s_0 \in K$. $\square$

Proof of Theorem A. By Corollary 2.5, every operator norm asymptotic representation of G satisfies $\|V_n(k) - 1\|_2 \rightarrow 0$ for $k \in K \leq \mathfrak{D}_G(L)$. Theorem 2.7 then makes every corona homomorphism trivial on K , and Lemma 2.1 makes every homomorphism to an MF group trivial on K . If $K \neq 1$, then G does not embed in an MF group, so G is not MF. The case $K = G = \mathfrak{D}_G(L)$ gives the last assertion. $\square$

2.4. The maximal group C^* -algebra. Lemma 2.3 also applies inside $C_{\max}^*(G)$ itself, where a strict compression produces a proper isometry.

Lemma 2.8 (strictly dominated projections). *Let A be a unital C^* -algebra containing a projection p and a unitary u with $p < upu^*$. Then $s = pu^* + (1 - upu^*)$ is an isometry that is not a unitary, so A is not stably finite and has no faithful tracial state.*

Proof. Put $q = upu^*$, so that $pq = qp = p$, $up(1 - q) = 0$, and $pu^*(1 - q) = pu^* - pu^*upu^* = 0$. Then

$$\begin{aligned} s^*s &= upu^* + up(1 - q) + (1 - q)pu^* + (1 - q) = q + (1 - q) = 1, \\ ss^* &= pu^*up + pu^*(1 - q) + (1 - q)up + (1 - q) \\ &= p + (1 - q) = 1 - (q - p) \neq 1. \end{aligned}$$

A unital C^* -algebra containing a nonunitary isometry is not finite, and neither is any matrix algebra over it, since the isometry embeds in the corner. A tracial state τ has $\tau(q - p) = \tau(s^*s) - \tau(ss^*) = 0$ with $q - p$ a nonzero positive element, so no tracial state on A is faithful. $\square$

Proposition 2.9 (proper one-sided compression). *Let G be a countable group containing a property-(T) subgroup Γ and an element t with $t\Gamma t^{-1} \subsetneq \Gamma$. Then $C_{\max}^*(G)$ contains a proper isometry. In particular, $C_{\max}^*(G)$ is not stably finite, admits no faithful tracial state, and is neither residually finite-dimensional nor MF.*

Proof. Let $P \in C_{\max}^*(G)$ be the image of the Kazhdan projection of Γ [AW] under the canonical map $C_{\max}^*(\Gamma) \rightarrow C_{\max}^*(G)$, and let u be the canonical unitary of t . Conjugation by u implements the isomorphism $\Gamma \rightarrow t\Gamma t^{-1}$ on canonical unitaries, so Lemma 2.3, applied with $U = u$, gives $P \leq uPu^*$ with uPu^* the image of the Kazhdan projection of $t\Gamma t^{-1}$. The inequality is strict: in the quasi-regular representation of G on $\ell^2(G/t\Gamma t^{-1})$, the point mass at the base coset is fixed by $t\Gamma t^{-1}$ and moved by every element of $\Gamma \setminus t\Gamma t^{-1}$, so the two fixed-space projections differ. Lemma 2.8 then supplies the proper isometry and rules out stable finiteness and faithful traces. MF algebras are stably finite, and a separable residually finite-dimensional C^* -algebra is MF [BK]. $\square$

3. THE BINARY LEAVITT GROUP

The Leavitt algebra $R = L_{\mathbb{F}_2}(1, 2)$ is built so that one copy of the free module R is isomorphic to two copies: the relations (2) say that s_0, s_1 embed two copies of R side by side and t_0, t_1 read them back. So a matrix A over R can be copied into one of the two halves, with the identity on the other half, and this compression is an injective multiplicative map that fixes the identity. It is not additive, but its restriction to the unit group is an injective group homomorphism. Inside a large enough elementary group the compression is realized by conjugation, which gives a compressor τ for a copy L of $\text{EL}_3(R)$; one commutator computation gives a nontrivial element of $\mathfrak{D}_H(L)$, and simplicity of H turns it into all of H .

Throughout this section $R = L_{\mathbb{F}_2}(1, 2)$ and

$$p = s_0 t_0, \quad q = s_1 t_1,$$

so that $p + q = 1$ and $t_1 q s_1 = 1$, and in particular $q \neq 0$. We use indices $0, \dots, 11$ for 12×12 matrices and identify $\text{GL}_3(R)$ with the subgroup of $\text{GL}_{12}(R)$ of matrices $\text{diag}(A, I_9)$. Define $\Psi: M_3(R) \rightarrow M_3(R)$ by

$$(3) \quad \Psi(A) = qI_3 + s_0 A t_0,$$

where $s_0 A t_0$ is the matrix $(s_0 a_{ij} t_0)_{ij}$. The relations $q^2 = q$, $q s_0 = t_0 q = 0$, and $t_0 s_0 = 1$ show that Ψ is multiplicative, $\Psi(I_3) = (p + q)I_3 = I_3$, and $t_0 \Psi(A) s_0 = A$. On the other hand, $\Psi(0) = qI_3 \neq 0$, so Ψ is not additive and is not a ring endomorphism. Its multiplicativity and preservation of the identity show that it restricts to a group homomorphism $\text{GL}_3(R) \rightarrow \text{GL}_3(R)$, and the recovery identity

makes that homomorphism injective. Put

$$X = \begin{pmatrix} s_0 I_3 & s_1 t_0 I_3 \\ 0 & t_1 I_3 \end{pmatrix}, \quad Y = \begin{pmatrix} t_0 I_3 & 0 \\ s_0 t_1 I_3 & s_1 I_3 \end{pmatrix};$$

block multiplication with (2) gives $Y = X^{-1}$ and $X \operatorname{diag}(A, I_3) Y = \operatorname{diag}(\Psi(A), I_3)$. The matrix X need not lie in an elementary group, which is why there are twelve coordinates: with

$$(4) \quad \tau = \operatorname{diag}(X, Y) \in \operatorname{GL}_{12}(R)$$

one has

$$(5) \quad \tau \operatorname{diag}(A, I_9) \tau^{-1} = \operatorname{diag}(\Psi(A), I_9).$$

Lemma 3.1. *The matrix τ defined in (4) belongs to $\operatorname{EL}_{12}(R)$.*

Proof. By the Whitehead lemma [Mil, §3], $\operatorname{diag}(X, X^{-1})$ is a product of block-triangular matrices $\begin{pmatrix} I & B \\ 0 & I \end{pmatrix}$ and $\begin{pmatrix} I & 0 \\ B & I \end{pmatrix}$ with $B \in M_6(R)$, and each of these is the product of the elementary matrices $e_{r,6+s}(b_{rs})$, respectively $e_{6+r,s}(b_{rs})$, since distinct off-diagonal matrix units have product zero. $\square$

Set $H = \operatorname{EL}_{12}(R)$, and let $L = \operatorname{EL}_3(R) \leq H$ denote the image of the embedding $A \mapsto \operatorname{diag}(A, I_9)$.

Proposition 3.2. *The groups H and L have property (T), and $\tau \in H$ satisfies*

$$(6) \quad \tau L \tau^{-1} \leq L.$$

Proof. The ring R is generated by s_0, s_1, t_0, t_1 as a unital ring, so the theorem of Ershov and Jaikin-Zapirain gives property (T) for $\operatorname{EL}_{12}(R)$ and $\operatorname{EL}_3(R)$ [EJZ, Theorem 1.1]. Lemma 3.1 gives $\tau \in H$. For $i \neq j$ and $a \in R$, equations (5) and (3) give $\tau e_{ij}(a) \tau^{-1} = e_{ij}(s_0 a t_0) \in L$, and these elementary matrices generate L . $\square$

Proposition 3.3. *The group $H = \operatorname{EL}_{12}(R)$ is nontrivial and simple.*

Proof. The ring R is purely infinite simple [AA], so it is an exchange ring [Ara]. Let $N \trianglelefteq H$. Since $N \leq \operatorname{GL}_{12}(R)$ is normalized by $\operatorname{EL}_{12}(R)$, Preusser's sandwich theorem [Pre, Theorem 3] gives an ideal $I \trianglelefteq R$ with $\operatorname{EL}_{12}(R, I) \leq N \leq C_{12}(R, I)$. Simplicity of R gives $I = 0$ or $I = R$, and $I = R$ gives $N = H$. If $I = 0$, then N centralizes every $e_{ij}(a)$: centralizing $e_{ij}(1)$ for all $i \neq j$ forces $g = \lambda I_{12}$ with $\lambda \in R^\times$, and centralizing $e_{ij}(a)$ forces $\lambda \in Z(R)^\times$. Since $Z(R) = \mathbb{F}_2$ [AC, Corollary 4.3], $N = 1$. Finally, $e_{01}(1) \neq 1$. $\square$

Proposition 3.4. *Let*

$$c = e_{34}(1), \quad \ell = e_{12}(1), \quad d = [\tau c \tau^{-1}, \ell]$$

in H . Then

$$c \in C_H(L), \quad d = e_{02}(q) \neq 1, \quad \langle\langle d \rangle\rangle_H = H.$$

Proof. The Steinberg relations give $[e_{ij}(a), e_{34}(1)] = 1$ for $i, j \in \{0, 1, 2\}$, and these matrices generate L , so $c \in C_H(L)$. Since $\tau = \text{diag}(X, Y)$ and E_{34} is supported in the first six coordinates, $\tau E_{34} \tau^{-1} = \text{diag}(X E_{34} Y, 0)$ with E_{34} read in $M_6(R)$, and with $\varepsilon_0, \varepsilon_1, \varepsilon_2$ the standard basis of R^3 ,

$$X E_{34} Y = \begin{pmatrix} s_1 t_0 \varepsilon_0 \\ t_1 \varepsilon_0 \end{pmatrix} \begin{pmatrix} s_0 t_1 \varepsilon_1^\top & s_1 \varepsilon_1^\top \end{pmatrix} = \begin{pmatrix} s_1 t_1 E_{01} & 0 \\ 0 & E_{01} \end{pmatrix}$$

by (2). So

$$\tau c \tau^{-1} = 1 + q E_{01} + E_{34} = e_{01}(q) e_{34}(1).$$

Direct multiplication of the matrices in (4) gives

$$\tau c \tau^{-1} = e_{01}(q) e_{34}(1).$$

The second factor commutes with ℓ , and $[e_{ij}(a), e_{jk}(b)] = e_{ik}(ab)$ gives $d = e_{02}(q)$, which is nontrivial because $q \neq 0$. Since H is simple, d normally generates H . $\square$

Proof of Theorem B. Proposition 3.2 gives property (T), which implies finite generation [BHV, Theorem 1.3.1]. The containment (6), the centrality of c relative to L , and Proposition 3.4 put d in $\mathfrak{D}_H(L)$, and $\langle\langle d \rangle\rangle_H = H$ gives $\mathfrak{D}_H(L) = H$. So Theorem A, applied with $K = H$, makes every homomorphism from H to an MF group trivial, and Proposition 3.3 makes H nontrivial and simple. In particular H is not MF. The algebra $C_r^*(H)$ is separable, and stably finite because its canonical trace is faithful. If an embedding $e: C_r^*(H) \rightarrow \mathcal{Q}_{\mathbf{d}}$ existed and $p = e(1)$, then $h \mapsto e(\lambda_h) + (1 - p)$ would embed H in $\mathcal{U}(\mathcal{Q}_{\mathbf{d}})$. Finally, the compression (6) is strict: by (5) and (3), every off-diagonal entry of a matrix in $\tau L \tau^{-1}$ has the form $s_0 a t_0$, and

$$t_1(s_0 a t_0)s_1 = 0 \neq 1 = t_1 \cdot 1 \cdot s_1,$$

so $e_{12}(1) \in L \setminus \tau L \tau^{-1}$. Proposition 2.9, applied to $L \leq H$ and τ , puts a proper isometry in $C_{\max}^*(H)$. $\square$

4. AN AMENABLE NONQUASIDIAGONAL TRACE

Brown observed that it was unknown whether every amenable trace on a C^* -algebra is quasidiagonal [Br, discussion preceding Proposition 3.5.1]. For a countable group G , let u_g denote the canonical unitary in $C_{\max}^*(G)$. The canonical trace τ_G is determined by

$$\tau_G(u_g) = \begin{cases} 1, & g = 1, \\ 0, & g \neq 1. \end{cases}$$

Theorem 4.1 (canonical-trace obstruction). *Let G be a group whose canonical trace τ_G on $C_{\max}^*(G)$ is amenable. If G is not MF, then τ_G is amenable and not quasidiagonal.*

Proof. If τ_G were quasidiagonal, its unital completely positive matrix models would define an MF trace model. For the canonical trace, such a model induces an injective homomorphism from G to a norm matrix corona. This contradicts the assumption that G is not MF. $\square$

A group N is *locally residually finite* if each of its finitely generated subgroups is residually finite.

Proposition 4.2 (locally-RF-by- $\mathbb{Z}$ groups). *Suppose that G is countable and*

$$G \cong N \rtimes_{\beta} \mathbb{Z},$$

where N is locally residually finite. Then τ_G is amenable.

Proof. Exhaust G by finite sets and use a long interval $I_L = \{0, \dots, L-1\} \subset \mathbb{Z}$ in the cyclic direction. The normal coordinates which occur in a fixed finite window generate a residually finite subgroup of N . A finite quotient separating the required coordinates, together with I_L , gives a finite coset window for a quasi-regular representation of G . Compression to this window is unital and completely positive. On a fixed group element, every column away from the two ends of I_L is carried to the expected column; only a bounded number of boundary columns escape. Hence the normalized Hilbert–Schmidt multiplicative defect is bounded by the square root of a boundary-to-volume ratio and tends to zero as $L \rightarrow \infty$. Residual separation makes the normalized trace exactly 0 for each prescribed nonidentity element. The linear span of the canonical unitaries is dense in $C_{\max}^*(G)$, so contractivity extends both conclusions to the whole algebra. $\square$

Let $\bar{\Gamma} \leq \mathrm{GL}_4(\mathbb{Q})$ be the affine matrix group generated by three linear generators and the three integral coordinate translations, and put

$$D = \mathrm{diag}(2, 2, 2, 1), \quad \alpha(g) = DgD^{-1}.$$

Conjugation by D gives an injective endomorphism of $\bar{\Gamma}$ with proper finite-index range. If a is the first coordinate translation, then $a \notin \alpha(\bar{\Gamma})$: a putative preimage would contain the nonintegral entry $1/2$.

Let

$$T_{\alpha} = \varinjlim (\bar{\Gamma} \xrightarrow{\alpha} \bar{\Gamma} \xrightarrow{\alpha} \dots), \quad V = T_{\alpha} \rtimes \langle t \rangle,$$

where $tgt^{-1} = \alpha(g)$ on the level-zero copy of $\bar{\Gamma}$. Put $X = V/\bar{\Gamma}$. The Clifford lamp group $\mathrm{Cl}(X)$ has an involution c_x at every $x \in X$ and a nontrivial central involution ε , with

$$[c_x, c_y] = \varepsilon \quad (x \neq y).$$

The permutation action of V on X acts on these lamps, and the affine–Clifford witness is

$$(7) \quad W = \mathrm{Cl}(X) \rtimes V.$$

The marked presentation used in the compression construction defines a finitely presented group $\widetilde{W}$ and a surjective homomorphism $\pi: \widetilde{W} \rightarrow W$ that maps its marked central involution to ε .

Let c be the lamp at the root coset. It centralizes the level-zero copy of $\bar{\Gamma}$. The two cosets supporting tct^{-1} and $a(tct^{-1})a^{-1}$ are distinct precisely because $a \notin \alpha(\bar{\Gamma})$, and therefore

$$(8) \quad [tct^{-1}, a(tct^{-1})a^{-1}] = \varepsilon \neq 1.$$

The base subgroup in the marked presentation has property (T), and the marked involution generates a finite normal subgroup. The one-sided compression criterion implies that every homomorphism from $\widetilde{W}$ to an MF group maps this involution to

the identity. If $f: W \rightarrow M$ is a homomorphism to an MF group, then $f \circ \pi$ maps it to $f(\varepsilon)$, so $f(\varepsilon) = 1$. Since $\varepsilon \neq 1$, the group W is not MF.

Reassociating the semidirect products yields

$$W \cong K \rtimes \mathbb{Z}, \quad K = \text{Cl}(X) \rtimes T_\alpha.$$

The group K is locally residually finite. Every finite subset of K involves finitely many Clifford generators and lies over a single telescope level $\bar{\Gamma}_n \cong \bar{\Gamma}$. The $\bar{\Gamma}_n$ -orbits of these generators are finite because $\alpha(\bar{\Gamma})$ has finite index. Their union is a finite invariant set Y , and the original finite subset is contained in

$$C_Y \rtimes \bar{\Gamma}_n,$$

where the Clifford group C_Y is finite. This semidirect product is residually finite because its kernel is finite, its quotient $\bar{\Gamma}_n$ is residually finite, and the action has finite image. Therefore every finitely generated subgroup of K is residually finite.

Corollary 4.3 (the affine–Clifford trace). *The canonical trace τ_W on $C_{\max}^*(W)$ is amenable but not quasidiagonal.*

Proof. The decomposition $W \cong K \rtimes \mathbb{Z}$ and local residual finiteness of K imply that τ_W is amenable by Proposition 4.2. Since W is not MF, Theorem 4.1 implies that τ_W is not quasidiagonal. $\square$

5. A TORSION-FREE FINITELY PRESENTED EXAMPLE

For an acylindrically hyperbolic group G , choose the generating set A provided by Hull [Hu, Theorem 3.12]. A subgroup is *suitable* if it acts non-elementarily on $\Gamma(G, A)$ and normalizes no nontrivial finite subgroup [Hu, Definition 1.4]. Hull’s small cancellation theorem [Hu, Theorem 7.1] is stated with injectivity on a ball of $\Gamma(G, A)$, and a ball containing a given finite set gives the following form.

Theorem 5.1 (Hull’s small cancellation theorem). *Let G be acylindrically hyperbolic, let $N \leq G$ be suitable with respect to A , and let $g_1, \dots, g_m \in G$ and a finite subset $\Omega \subseteq G$ be given. Then there is a surjective homomorphism $\varphi: G \rightarrow Q$ such that Q is acylindrically hyperbolic, $\varphi|_\Omega$ is injective, $\varphi(g_i) \in \varphi(N)$ for all i , and every element of finite order in Q is the image of an element of the same order in G .*

Hull’s proof treats $m = 1$ by passing to $G/\langle\langle W \rangle\rangle_G$ for one word W and the general case by induction on m [Hu, proof of Theorem 7.1], so $\ker \varphi$ is the normal closure of m elements and Q is finitely presented when G is.

Lemma 5.2 (saturation). *Let G be finitely presented, torsion-free, and acylindrically hyperbolic, let $N \trianglelefteq G$ be nontrivial, and let $\Omega \subseteq G$ be finite. Then there is a surjective homomorphism $\varphi: G \rightarrow Q$ such that Q is two-generated, finitely presented, torsion-free, and acylindrically hyperbolic, $\varphi|_\Omega$ is injective, and $\varphi(N) = Q$.*

Proof. The subgroup N is infinite and normal, so it acts non-elementarily on $\Gamma(G, A)$ by Osin [Os, Lemma 7.1], and torsion-freeness makes it suitable. By Hull [Hu, Corollary 5.7 and Lemma 5.8], N contains two elements h_1, h_2 such that $N_0 = \langle h_1, h_2 \rangle$ is again suitable. Apply Theorem 5.1 to N_0 with $g_1, \dots, g_m$ a finite generating set of

G and with the set Ω . Then $Q = \langle \varphi(g_1), \dots, \varphi(g_m) \rangle \leq \varphi(N_0) \leq \varphi(N) \leq Q$, so $Q = \varphi(N_0) = \langle \varphi(h_1), \varphi(h_2) \rangle$ and $\varphi(N) = Q$. The quotient is torsion-free by Theorem 5.1 and finitely presented because $\ker \varphi$ is normally generated by m elements. $\square$

Fournier-Facio constructs a finitely presented torsion-free group G_0 with property (T), a subgroup $\Gamma \leq G_0$ with property (T), an element $t \in G_0$ with $t\Gamma t^{-1} \leq \Gamma$, and a subgroup $J \leq G_0$ isomorphic to a finitely presented infinite simple group, such that $[\Gamma, J] = 1$ and $tJt^{-1} \leq \Gamma$ [FF, §2]. The group G_0 is obtained there from Hull's common quotient theorem [Hu, Corollary 7.4], so it is acylindrically hyperbolic. Put $S = tJt^{-1}$. Since J is simple and nonabelian, J and S are perfect.²

Lemma 5.3. *For every homomorphism $\rho: G_0 \rightarrow \bar{G}$,*

$$\rho(S) \leq \mathfrak{D}_{\bar{G}}(\rho(\Gamma)).$$

Proof. Write $\bar{\Gamma} = \rho(\Gamma)$, $\bar{t} = \rho(t)$, $\bar{J} = \rho(J)$, and $\bar{S} = \rho(S) = \bar{t}\bar{J}\bar{t}^{-1}$. Then $\bar{t}\bar{\Gamma}\bar{t}^{-1} \leq \bar{\Gamma}$, $\bar{J}$ centralizes $\bar{\Gamma}$, and $\bar{S} \leq \bar{\Gamma}$. For $c \in \bar{J}$ and $\ell \in \bar{S}$, the commutator $[\bar{t}c\bar{t}^{-1}, \ell]$ lies in $\mathfrak{D}_{\bar{G}}(\bar{\Gamma})$ by (1). As c ranges over $\bar{J}$, the element $\bar{t}c\bar{t}^{-1}$ ranges over $\bar{S}$, so $[\bar{S}, \bar{S}] \leq \mathfrak{D}_{\bar{G}}(\bar{\Gamma})$. The group $\bar{S}$ is perfect as a quotient of S , so $\bar{S} \leq \mathfrak{D}_{\bar{G}}(\bar{\Gamma})$. $\square$

Proof of Theorem C. Let N be the normal closure of S in G_0 , and fix $s \in S$ with $s \neq 1$. Lemma 5.2 applied to G_0 , N , and the finite set $\{1, s\}$ gives $\varphi: G_0 \rightarrow Q$ with Q two-generated, finitely presented, torsion-free, and acylindrically hyperbolic, $\varphi(N) = Q$, and $\varphi(s) \neq 1$. The group Q has property (T) as a quotient of G_0 .

Let $r: Q \rightarrow M$ be a homomorphism to an MF group with image $\bar{G}$, and put $\rho = r \circ \varphi$. Since $\varphi(N) = Q$, the normal closure of $\rho(S)$ in $\bar{G}$ is $\rho(N) = \bar{G}$, and by Lemma 5.3 it lies in the normal subgroup $\mathfrak{D}_{\bar{G}}(\rho(\Gamma))$, so $\mathfrak{D}_{\bar{G}}(\rho(\Gamma)) = \bar{G}$. The group $\bar{G}$ has property (T) as a quotient of Q , and $\rho(\Gamma)$ has property (T) as a quotient of Γ . Theorem A, applied to $\bar{G}$ with the subgroup $\rho(\Gamma)$ and $K = \bar{G}$, makes every homomorphism from $\bar{G}$ to an MF group trivial. But $\bar{G}$ is a subgroup of the MF group M , so it is MF, and its identity map is such a homomorphism. So $\bar{G} = 1$, and r is trivial. A homomorphism from a quotient of Q composes to one from Q , so no nontrivial quotient of Q is MF. $\square$

ACKNOWLEDGMENTS

The author thanks Francesco Fournier-Facio for early advice and detailed comments on earlier drafts, and Alon Dogon for organizing the ensuing discussion and for comments on the surrounding literature. The author also thanks Tatiana Shulman for clarifying the terminology and historical context of the MF question, and Caleb Eckhardt for discussions of MF representations and matrix norms.

USE OF AI AND FORMAL METHODS

Large language models were used extensively in mathematical exploration, proof development, manuscript drafting, literature navigation, Lean formalization, and editorial revision.

²Simplicity of J is not needed since a free group of rank two can replace J , with N being the normal closure of $[S, S]$ and s being a nonidentity commutator.

REFERENCES

- [AA] G. Abrams and G. Aranda Pino, *Purely infinite simple Leavitt path algebras*, J. Pure Appl. Algebra **207** (2006), no. 3, 553–563, [doi:10.1016/j.jpaa.2005.10.010](#).
- [AW] C. A. Akemann and M. E. Walter, *Unbounded negative definite functions*, Canad. J. Math. **33** (1981), no. 4, 862–871.
- [AC] G. Aranda Pino and K. Crow, *The center of a Leavitt path algebra*, Rev. Mat. Iberoam. **27** (2011), no. 2, 621–644, [doi:10.4171/RMI/649](#).
- [Ara] P. Ara, *The exchange property for purely infinite simple rings*, Proc. Amer. Math. Soc. **132** (2004), no. 9, 2543–2547, [doi:10.1090/S0002-9939-04-07369-1](#).
- [BDL] B. Bachner, A. Dogon, and A. Lubotzky, *On L^1 -approximation of groups*, J. Algebra **702** (2026), 235–243, [doi:10.1016/j.jalgebra.2026.04.041](#), [arXiv:2508.17392](#).
- [BHV] B. Bekka, P. de la Harpe, and A. Valette, *Kazhdan’s property (T)*, New Mathematical Monographs, vol. 11, Cambridge University Press, Cambridge, 2008, [doi:10.1017/CB09780511542749](#).
- [BK] B. Blackadar and E. Kirchberg, *Generalized inductive limits of finite-dimensional C^* -algebras*, Math. Ann. **307** (1997), no. 3, 343–380, [doi:10.1007/s002080050039](#).
- [Br] N. P. Brown, *Invariant means and finite representation theory of C^* -algebras*, Mem. Amer. Math. Soc. **184** (2006), no. 865, viii+105, [doi:10.1090/memo/0865](#).
- [CDE] J. R. Carrión, M. Dadarlat, and C. Eckhardt, *On groups with quasidiagonal C^* -algebras*, J. Funct. Anal. **265** (2013), no. 1, 135–152, [doi:10.1016/j.jfa.2013.04.004](#).
- [Ec] C. Eckhardt, *Non-MF groups and non-finite full group C^* -algebras*, preprint, 2026.
- [EJZ] M. Ershov and A. Jaikin-Zapirain, *Property (T) for noncommutative universal lattices*, Invent. Math. **179** (2010), 303–347, [doi:10.1007/s00222-009-0218-2](#).
- [FF] F. Fournier-Facio, *A torsion-free non-sofic group*, preprint, 2026, [arXiv:2608.02025](#).
- [GKM] D. Gao, S. Kunnawalkam Elayavalli, A. Manzoor, and G. Patchell, *A new source of purely finite matricial fields*, preprint, 2026, [arXiv:2603.24502](#).
- [GH] I. Goldbring and B. Hart, *The universal theory of the hyperfinite II_1 factor is not computable*, Bull. Symbolic Logic **30** (2024), no. 2, 181–198, [doi:10.1017/bsl.2024.7](#), [arXiv:2006.05629](#).
- [Hu] M. Hull, *Small cancellation in acylindrically hyperbolic groups*, Groups Geom. Dyn. **10** (2016), no. 4, 1077–1119, [doi:10.4171/GGD/377](#).
- [JNVWY] Z. Ji, A. Natarajan, T. Vidick, J. Wright, and H. Yuen, $\text{MIP}^* = \text{RE}$, Commun. ACM **64** (2021), no. 11, 131–138, [doi:10.1145/3485628](#), [arXiv:2001.04383](#).
- [K–T] H. V. Khanh and V. H. Thanh, *Matrix generators for the unit groups of $L_K(1, d)$* , preprint, 2026, [arXiv:2607.10351](#).
- [Kul6] G. Kun, *On sofic approximations of property (T) groups*, preprint, 2016, [arXiv:1606.04471](#).
- [KT19] G. Kun and A. Thom, *Inapproximability of actions and Kazhdan’s property (T)*, preprint, 2019, [arXiv:1901.03963](#).
- [Kor] A. I. Korchagin, *The MF-property for countable discrete groups*, Math. Notes **102** (2017), no. 2, 198–211, [doi:10.1134/S0001434617070227](#), [arXiv:1704.06906](#).
- [Lev] W. G. Leavitt, *The module type of a ring*, Trans. Amer. Math. Soc. **103** (1962), 113–130, [doi:10.1090/S0002-9947-1962-0132764-X](#).
- [Mil] J. Milnor, *Introduction to algebraic K-theory*, Annals of Mathematics Studies, vol. 72, Princeton University Press, Princeton, 1971.
- [OAI] OpenAI, *Nonsofic groups exist*, in *Ten advances in mathematics and theoretical computer science*, Chapter 3, 78–95, 2026, <https://cdn.openai.com/pdf/ten-proofs-oai.pdf>.
- [Os] D. Osin, *Acylindrically hyperbolic groups*, Trans. Amer. Math. Soc. **368** (2016), no. 2, 851–888, [doi:10.1090/tran/6343](#).
- [Pre] R. Preusser, *On general linear groups over exchange rings*, Linear and Multilinear Algebra **70** (2022), no. 4, 705–713, [doi:10.1080/03081087.2020.1743636](#).
- [Sch] C. Schafhauser, *Finite dimensional approximations of certain amalgamated free products of groups*, Groups Geom. Dyn. **20** (2026), no. 2, 607–615, [doi:10.4171/GGD/826](#), [arXiv:2306.02498](#).