

## Modeling and $R_0$ estimation of the 2010 Dengue Hemorrhagic Fever Outbreak in Hermosillo Sonora.

S. DÍAZ-INFANTE\*

CONACYT-Universidad de Sonora, Departamento de Matemáticas, Universidad de Sonora  
Blvd. Rosales y Luis Encinas S/N, Col. Centro, Hermosillo, Sonora, C.P. 83000, México

J. A. MONTOYA LAOS †

Universidad de Sonora, Departamento de Matemáticas, Universidad de Sonora  
Blvd. Rosales y Luis Encinas S/N, Col. Centro, Hermosillo, Sonora, C.P. 83000, México

AND

D. OLMOS LICEAGA ‡

Universidad de Sonora, Departamento de Matemáticas, Universidad de Sonora  
Blvd. Rosales y Luis Encinas S/N, Col. Centro, Hermosillo, Sonora, C.P. 83000, México  
[May 5, 2020]

We model and estimate the basic reproductive number of Dengue and Dengue Hemorrhagic Fever outbreak of the 2010 from Hermosillo Sonora. Our results suggest that serotype DENV-2 of the Dengue virus and the cross infection risk enhancement hypothesis, could explain the incidence of Dengue Hemorrhagic Fever cases reported by Secretaria de Salud del Estado de Sonora.

**Keywords:** Differential Equations; Dengue Hemorrhagic Fever; Boot strap;

### 1. Introduction

State the objectives of the work and provide an adequate background, avoiding a detailed literature survey or a summary of the results.

- State of the art (literature review).
- Knowledge gap.
- Objective.
- Methodology.
- Results.
- Conclusions.

\*Corresponding author. Present address: Departamento de Matemáticas, Universidad de Sonora Blvd. Rosales y Luis Encinas S/N, Col. Centro, Hermosillo, Sonora, C.P. 83000, México. Email: saul.diazinfante@unison.mx

†Email: montoya@unison.mx

‡Email: daniel.olmos@unison.mx

## 2. Introduction.

**CONTRIBUTION** Our objective is to explain the dengue hemorrhagic outbreak that occurred in the city of Hermosillo, located in the state of Sonora in 2010.

**SEVERITY** Dengue Classic Fever (DCF) Dengue Hemorrhagic Fever (DHF) description.

**ADE HYPOTHESIS** Dengue Virus Serotype 1 (DENV-1), Dengue Virus Serotype 2 (DENV-2) reinfection as cause of hemorrhagic.

**BACKGROUND** Dengue fever is one of the diseases that has advanced in the world due to climate change. One of the main strategies to control dengue has been the control of mosquito population.

As mentioned by Ernst et. al. (?) in Nogales there is much more propagating conditions than in Hermosillo. However, due to temperature, Dengue has not strike more Nogales. Can we implement a second variable (related to the conditions that a mosquito can reproduce, how many eggs are layed, how many hatch into larvae?).

We focus our studies in the state of Sonora, located at the Northwestern part of Mexico. An important case was discussed in (?), in which two populations, Hermosillo and Nogales, where the subject of study.

Several authors have considered the effects of temperature on mosquito dynamics and there has been also different approaches to understand dengue dynamics. Nonlinear ODE systems have been proposed. For example, Esteva and Yang (cita) consider a model based on epidemiological data to understand the effects of temperature on the intensity of dengue outbreaks. In (Beck-Johnson 2013) (to read carefully),...they do similar as EstevaYang. On (Lambrechts 2011) and (Liu-Helmersson 2014), there are studies on different parameters involved in the vectorial capacity as a function of temperature and diurnal temperature range (DTR). In (Marinho2016.pdf) Marinho et al studied the effect of temperature of the life cycle of the mosquito.

However as discussed in (?), temperature also plays an important role in the survival of the adult female mosquito. In their study, they considered two different populations from northwest Mexico, Nogales and Hermosillo. In their work, it was found that larval and pupal abundance was greater in Nogales, but dengue cases, where much higher in the city of Hermosillo. The importance of temperature was that adult mosquitoes were able to survive longer in Hermosillo and therefore, being able to transmit the virus.

From this study, it is clear that not only temperature has played an important role in dengue presence in the medium but also, there is a socioeconomical factor that has to be taken into account. In case of having the ideal temperature for having infected mosquitoes, but there are no favourable socio-economical conditions for propagation, then a possible outbreak might not occur but only might be an endemic situation.

## 3. Material and methods

It was reported that in 2010 there was present only DENV-1 circulating Hermosillo (Tesis de Pablo. La referencia que menciona ya no se encuentra disponible por internet.)

Provide sufficient detail to allow the work to be reproduced, with details of supplier and catalogue number when appropriate. Methods already published should be indicated by a reference: only relevant modifications should be described.

### 3.1 Data

**DATA DESCRIPTION (PABLO)** Socioeconomic description: homogeneity of regarding initial grow geographic area (Ana-Lucia)

#### Model

Our work aims to explain the DHF cases reported in the Hermosillo's 2010 outbreak. Particularly, our aim is to provide evidence based on a mathematical model and statistical parameter estimation that a second strain (DENV-2) was also present in the city in 2010, and thus being responsible for the presence of DHF.

Thus, our formulation only considers the time of the epidemic and supposes that at this period, the DENV-2 serotype invades for the first time the city. We understand the DHF cases as a consequence of reinfection with the DENV-2 serotype. That is, a fraction of individuals with immunity to serotype DENV-1—acquired by past outbreaks—increments their susceptibility to the DENV-2, and consequently develops hemorrhagic with a certain probability.

#### Important Hypothesis (Daniel-Saúl)

**PERMANENT IMMUNITY** Accordingly to WHO ((1997 (accessed May, 2018))), infection of Dengue caused by a serotype DEN-i induces long-life immunity to reinfection with this strain. Then, we assume in our formulation that a susceptible individual previously infected with Dengue of serotype DEN-i only could acquire Dengue of a different serotype. We also suppose that a susceptible individual who never get infected before, could obtain Dengue by any of serotypes DENV-1, or DENV-2, but only for a single time and during the outbreak period.

**ADE HYPOTHESIS** The processes and factors that produce DHF are still unclear. There is evidence that reinfection with a different serotype enhances the probability of developing plasma vascular permeability—the Antibody Dependent infection Enhancement (ADE) hypothesis ((see, e.g. Halstead, 1992, p. 295)). But there also exist studies that report first infection DHF cases Streatfield et al. ((1993)). We consider in our formulation the ADE hypothesis, that is, only a fraction of the second reinfection with serotype DENV-2 develops vascular leaking.

**DENV-2 REPORT CIRCULATION IN HERMOSILLO** According to Vázquez-Pichardo et al. ((2011)) and Reyes-Castro et al. ((2017)), in year 2010 only DENV-1 was present in the state of Sonora. Based on this information it is not clear that reinfection with a second strain in 2010 is the main cause of DHF. However, contrary to these studies, one of the main hypothesis in this work is that DENV-2 was present in Sonora in 2010. Our hypothesis is based on previous experiences, just as mentioned in Gómez Dantés et al. ((2014)). In this work, where the authors analyze the dengue situation in Mexico for the period 2000 to 2011, argue that the increase of DHF in 2001 in Yucatan was linked to the introduction of the DENV-2 strain. Therefore, part of our hypothesis is that in 2010 there was an introduction of the DENV-2 into the Sonora state.

**ASYMPTOMATIC AND REPORTED CASES** We assume that the 95% (pendiente) of the DCF are asymptomatic, whereas for DHF all the cases are reported. Therefore, the class  $Y_{-1}^{[h]}$  accounts for all the DHF cases, whereas a fraction  $p = 0.05$  of the sum  $I_1 + I_2 + Y_{-1}^{[c]}$  represent the confirmed cases of DCF.

de los por-  
asintomáticos,  
ho.int/en/news-  
ets/detail/dengue-  
engue says that  
asymptomatic

df

HOMOGENEITY ABOUT THE EARLY OUTBREAK STAGE Define the infection forces as

$$\begin{aligned} A_{I_1} &= \frac{\beta_M b}{N_H} I_1, & A_{I_2} &= \frac{\beta_M b}{N_H} I_2, \\ A_{Y_{-1}^{[h]}} &= \frac{\beta_M b}{N_H} Y_{-1}^{[h]}, & A_{Y_{-1}^{[c]}} &= \frac{\beta_M b}{N_H} Y_{-1}^{[c]}, \\ B_{M_1} &= \frac{\beta_H b}{N_H} M_1, & B_{M_2} &= \frac{\beta_H b}{N_H} M_2. \end{aligned} \quad (3.1)$$

VECTOR TRANSMISSION DYNAMICS Define

$$A_{\bullet} := A_{I_1} + A_{I_2} + A_{Y_{-1}^{[h]}} + A_{Y_{-1}^{[c]}}$$

as the total human infection force, that is, the sum of all human contributions to the vector infection. Then we describe the mosquito disease dynamics by

$$\begin{aligned} \frac{dM_S}{dt} &= \Lambda_M - A_{\bullet} M_S - \mu_M M_S \\ \frac{dM_1}{dt} &= A_{I_1} M_S - \mu_M M_1 \\ \frac{dM_2}{dt} &= \left( A_{I_2} + A_{Y_{-1}^{[h]}} + A_{Y_{-1}^{[c]}} \right) M_S - \mu_M M_2 \end{aligned} \quad (3.2)$$

Here  $M_S$ , is the vector susceptible class and  $M_1, M_2$  respectively denotes the vector Infected classes with DENV-1 and DENV-2.

**HOST DISEASE DYNAMICS** Susceptible individuals ( $S$ ) become infected for the first time with DENV-1 or DENV-2 after a successful mosquito bite and move to classes  $I_1$  and  $I_2$ , respectively. From here, they remain in the infected class for  $1/\alpha_c$  time units, after which, move to a recovered class  $R_S$ . As we are interested in a one year dynamics, for the rest of the epidemic they become immune to any serotype. A second class of susceptible individuals  $S_{-1}$ , consist on those who acquired DENV-1 in previous years and in the current year are susceptible only to DENV-2. Such individuals become infected with DENV-2 when exposed to infected mosquitoes with that serotype. In OhAinle et al. ((2011)) and Sangkawibha et al. ((1984)) it was observed that a more severe version of dengue occurs (might occur?) when an individual acquires dengue for a second time, and this happens to be DENV-2. Based on this assumption, an individual from  $S_{-1}$  moves to  $Y_{-1}^{[c]}$  or  $Y_{-1}^{[h]}$ , if the infection leads to DF or DHF, respectively. Finally, these infected individuals move to the recovered class  $R_{S_{-1}}$  at rates  $\alpha_c$  and  $\alpha_h$ , respectively. For our model,  $\mu_H$  is the human death rate;  $b$  is the number of bites per week per mosquito and  $\beta_H$  is the effectiveness of the bite. From the current hypothesis our model is given by

$$\begin{aligned}
\frac{dS}{dt} &= \mu_H N_S - (B_{M_1} + B_{M_2})S - \mu_H S \\
\frac{dI_1}{dt} &= B_{M_1} S - (\alpha_c + \mu_H) I_1 \\
\frac{dI_2}{dt} &= B_{M_2} S - (\alpha_c + \mu_H) I_2 \\
\frac{dR_S}{dt} &= \alpha_c I_2 - \mu_H R_S \\
\frac{dS_{-1}}{dt} &= \mu_H N_{S_{-1}} - \sigma B_{M_2} S_{-1} - \mu_H S_{-1} \\
\frac{dY_{-1}^{[c]}}{dt} &= (1 - \theta) \sigma B_{M_2} S_{-1} - (\alpha_c + \mu_H) Y_{-1}^{[c]} \\
\frac{dY_{-1}^{[h]}}{dt} &= \theta \sigma B_{M_2} S_{-1} - (\alpha_h + \mu_H) Y_{-1}^{[h]} \\
\frac{dR_{S_{-1}}}{dt} &= \alpha_c Y_{-1}^{[c]} + \alpha_h Y_{-1}^{[h]} - \mu_H R
\end{aligned} \tag{3.3}$$

Here, we take  $N_H = N_S + N_{S_{-1}}$  as the total number of individuals. For our formulation  $N_H, N_S$  and  $N_{S_{-1}}$  remain constant.  $N_S$  is the total number of individuals that are involved in the first infection dynamics ( $N_S = S + I_1 + I_2 + R_S$ ). On the other hand  $N_{S_{-1}}$  is the total number of individuals involved in the reinfection dynamics ( $N_{S_{-1}} = S_{-1} + Y_{-1}^{[c]} + Y_{-1}^{[h]} + R_{S_{-1}}$ ). Also, the recovered individuals in both classes can be considered as a single recovered class  $R = R_S + R_{S_{-1}}$  as our dynamics are taken only for one year. Then, our equations become

$$\begin{aligned}
\frac{dS}{dt} &= \mu_H N_S - (B_{M_1} + B_{M_2})S - \mu_H S \\
\frac{dI_1}{dt} &= B_{M_1} S - (\alpha_c + \mu_H) I_1 \\
\frac{dI_2}{dt} &= B_{M_2} S - (\alpha_c + \mu_H) I_2 \\
\frac{dS_{-1}}{dt} &= \mu_H N_{S_{-1}} - \sigma B_{M_2} S_{-1} - \mu_H S_{-1} \\
\frac{dY_{-1}^{[c]}}{dt} &= (1 - \theta) \sigma B_{M_2} S_{-1} - (\alpha_c + \mu_H) Y_{-1}^{[c]} \\
\frac{dY_{-1}^{[h]}}{dt} &= \theta \sigma B_{M_2} S_{-1} - (\alpha_h + \mu_H) Y_{-1}^{[h]} \\
\frac{dR}{dt} &= \alpha_c (I_1 + I_2 + Y_{-1}^{[c]}) + \alpha_h Y_{-1}^{[h]} - \mu_H R
\end{aligned} \tag{3.4}$$

Fix  $\Lambda_c = N_c$ .

Include information about the 2 strains

**BASIC REPRODUCTIVE NUMBER** The disease free equilibrium results

$$FDE = \left( \frac{\Lambda_M}{\mu_M}, 0, 0, N_H - N_{S_{-1}}, 0, N_{S_{-1}}, 0, 0, 0 \right).$$

Symbol	Meaning
$M_S$	Number of susceptible mosquitoes.
$M_1, M_2$	Number of infected mosquitoes with virus serotype DENV-1 or DENV-2.
$S$	Susceptible host population which, never has acquired dengue.
$S_{-1}$	Susceptible host population which is immune to serotype 1.
$I_1, I_2$	First time infected host population by serotype 1 and 2, respectively.
$Y_{-1}^{[h]}, Y_{-1}^{[c]}$	Second time infected host population with serotype 2, with DHF and DCF, respectively.

Table 1. Meaning of variables. Here we omit the explicit dependence of time.

Using the next generation operator method reported as in Feng and Velasco-Hernández ((1997)), we obtain the basic reproductive number

$$\begin{aligned}
 \psi &:= \frac{\beta_M b N_M}{\mu_M N_H} \\
 R_{0c} &:= \sqrt{\psi \left( \frac{\beta_H b N_S}{(\alpha_c + \mu_H) N_H} + \frac{\beta_H b (1 - \theta) \sigma N_{S_{-1}}}{(\alpha_c + \mu_H) N_H} \right)} \\
 R_{0h} &:= \sqrt{\left( \frac{\beta_M b N_M}{\mu_M N_H} \right) \left( \frac{\sigma \theta N_{S_{-1}}}{(\alpha_h + \mu_H) N_H} \right)} \\
 \mathcal{R}_0 &:= \sqrt{R_{0c}^2 + R_{0h}^2}.
 \end{aligned} \tag{3.5}$$

In this equation,  $R_{0c}$  and  $R_{0h}$ , are the basic reproductive numbers for classical and hemorrhagic dengue cases, respectively. From here,  $R_0$  provides a measure of how DF and DHF infected people influence the presence of new dengue cases (Either DF or DHF).  $R_{0h}$  measures the new hemorrhagic cases that arise from one hemorrhagic infected individual in a population of  $N_{S_{-1}}$  susceptible to strain 2 individuals, meanwhile  $R_{0c}$  provides a measure of how many new individuals will obtain DC fever (DF?) from an individual that has or has not have acquired dengue previously (from an individual that has either DF or DHF).

Observe that this  $R_0$  differs in some way to the traditional  $R_0$  where two different serotypes are involved (Feng and Velasco-Hernández ((1997)) include citations of  $R_0$  for two serotypes). This follows from the idea that we are interested in classic and hemorrhagic cases rather than the predominance of a serotype.

Discuss above parameters units. Weeks or days

### 3.2 $R_0$ estimation (Montoya)

$$\frac{dZ}{dt} = p \left( I_1 + I_2 + Y_{-1}^{[c]} \right). \quad (3.6)$$

## 4. Results

Results should be clear and concise.

### MODELING RESULTS (SAÚL-DANIEL)

### DATA ANALYSIS (SAÚL-MONTOYA)

#### 4.1 $R_0$ estimation (Montoya)

We suppose that the number of cases of classic and hemorrhagic dengue are observed at time points  $t_1, \dots, t_n$ . Here we assume that these processes, denoted by  $X_t$  and  $Y_t$  respectively, follow a Poisson distribution with mean  $\lambda_h(t) = Y_{-1}^{[h]}$  and  $\lambda_c(t) = Z$ , where

$$\frac{dZ}{dt} = p \left( I_1 + I_2 + Y_{-1}^{[c]} \right). \quad (4.1)$$

In our case the vector of parameters of the ordinary differential equations model is  $\phi = (\phi_1, \phi_2)$ , where  $\phi_1 = (\beta_H, \beta_M)$  is regarded as unknown and  $\phi_2 = (\alpha_c, \alpha_h, b, \mu_H, \mu_M, \sigma, \theta, p)$  is known in advance. We write  $\lambda_h(t)$  and  $\lambda_c(t)$  as  $\lambda_h(t; \phi_1)$  and  $\lambda_c(t; \phi_1)$  to emphasize this fact.

We use the likelihood approach to estimate the vector parameter  $\psi_1$  based on the observed samples  $\vec{x} = (x_{t_1}, \dots, x_{t_n})$  and  $\vec{y} = (y_{t_1}, \dots, y_{t_n})$ . The resulting likelihood function is thus

$$L(\phi_1) = \prod_{i=1}^n \left\{ \frac{1}{x_{t_i}!} [\lambda_h(t_i; \phi_1)]^{x_{t_i}} \exp[-\lambda_h(t_i; \phi_1)] \frac{1}{y_{t_i}!} [\lambda_c(t_i; \phi_1)]^{y_{t_i}} \exp[-\lambda_c(t_i; \phi_1)] \right\}. \quad (4.2)$$

The maximum likelihood estimate (MLE) of  $\phi_1$  is that value of  $\phi_1$  that maximizes  $L(\phi_1)$  in (4.2). We denote the MLE of  $\phi_1$  as  $\hat{\phi}_1$ .

We now consider profile-likelihood inference based on (1) for estimating the parameters of interest ( $R_{01}$ ,  $R_{02}$ , and  $\mathcal{R}_0$ ). Here we assume without loss of generality that  $\phi_1 = (\beta_H, \beta_M)$  can be rewritten as  $\phi_1 = (\gamma, \eta)$ , where  $\gamma$  is a scalar parameter of interest and  $\eta$  is a scalar nuisance parameter. For example, we may only be interested in  $R_{01}$ . In this case, we can rewrite the parameter  $\beta_M$  as a function of the parameters  $R_{01}$  and  $\beta_H$ ,

$$\beta_M = C \frac{R_{01}}{\beta_H},$$

where

$$C = \left[ \left( \frac{N_H - N_{S-1}}{\alpha_c + \mu_H} + \frac{(1 - \theta)\sigma N_{S-1}}{\alpha_c + \mu_H} \right) \left( \frac{b^2 \Lambda_M}{\mu_M^2 N_H^2} \right) \right]^{-1}$$

Thus, we reparametrize the model in terms of  $\phi_1 = (\gamma, \eta) = (R_{01}, \beta_H)$ , where  $\gamma = R_{01}$  is the parameter of interest and  $\eta = \beta_H$  is the nuisance parameter.

The profile likelihood and its corresponding relative likelihood function of  $\gamma$ , standardized to be one at the maximum of the likelihood function, are

$$L_{\max}(\gamma) = \max_{\eta} L(\phi_1 = (\gamma, \eta)),$$

$$R_{\max}(\gamma) = \frac{L_{\max}(\gamma)}{\max_{\phi_1} L(\phi_1)},$$

where  $L(\cdot)$  is the likelihood function given in (4.2). In particular, the relative profile likelihood varies between 0 and 1 and ranks all possible  $\gamma$  values based only on the observed samples  $\vec{x} = (x_{t_1}, \dots, x_{t_n})$  and  $\vec{y} = (y_{t_1}, \dots, y_{t_n})$ . Thus, a graph of  $R_{\max}(\gamma)$  allows to distinguish plausible and implausible values for  $\gamma$ .

A level  $\omega$  profile likelihood region (commonly an interval) for  $\gamma$  is given by

$$\{\gamma : R_{\max}(\gamma) \geq \omega\},$$

where  $0 \leq \omega \leq 1$ . We can assign a confidence level to the profile likelihood region of  $\gamma$  considering the asymptotical behavior of the likelihood ratio statistic  $D = -2 \ln R_{\max}(\gamma_0)$ . This is an asymptotic pivotal quantity having a Chi-squared distribution with one degree of freedom. Thus, approximate confidence levels of 99%, 95% and 90% can be ascribed to profile likelihood regions at  $\omega = 0.036, 0.146$ , and  $0.25$ , respectively.

## 5. Results

Results should be clear and concise.

MODELING RESULTS (SAÚL-DANIEL)

DATA ANALYSIS (SAÚL-MONTOYA)

$R_0$  AND PARAMETER INFERENCE (MONTOYA)

## 6. Discussion

**STATEMENT OF PRINCIPAL FINDINGS** We estimate that  $R_{0c} > 1$  and  $R_{0h} < 1$ . Thus DHF cases, cannot sustain new DHF cases. Further,  $R_{0h} < 1$  imply an exponential decay on the number of infected individuals. However, a small DHF outbreak arises despite the value of  $R_{0h}$ , we deduce that this DHF dynamics is a consequence of the intensity of the outbreak of DF, given by serotype 2.

We formulate a model to describe the evolution of DCF and DHF cases in Hermosillo for the 2010 outbreak and evaluate if ADE hypothesis could explain the DHF dynamics. The central hypotheses of this manuscript were:

- strain DENV-II circulated in the 2010 Hermosillo
- Dengue outbreak antibody-Dependent Enhancement hypothesis could explain the DHF high incidence
- unconfirmed cases represent almost 95 % of the total incidence.



With these hypotheses, we obtain a model that describes the incidence dynamics of DHF cases of the 2010 Hemosillo outbreak. Our parameters estimation does not reject these hypothesis and provide statistical evidence that the concerning reproductive number is in the 95 % interval [1.81253, 2.134538].

**STRENGTHS AND WEAKNESSES OF THE STUDY** Our  $\mathcal{R}_0$  estimation is consistent with other outbreak reports [\*]. To the best of our knowledge, this is the first attempt to fit the two exponential growth curves of the incidence according to Dengue Disease severity. Furter, our reproductive numeber  $\mathcal{R}_0$  results be the geometric mean of two expresion that are closed related with parameters transmission according to severity.

However, our formulation only considers an outbreak of one season. Thus classical asymptotic analysis has mathematical consistency but lacks biological meaning. In this line, we are preparing a version that achieves both—proper data fitting and consistent asymptotic behavior. We believe that adding a class of immunity according to the particular strain would be an option.

We take ideas from Zheng and Nie ((2018)); Nuraini et al. ((2007)); Feng and Velasco-Hernández ((1997)) to modulate the propensity of acquire Dengue due to a given serotype. Our model alos describe DHF transmission with a certain probability. Thus, combining this idea with the hypothesis of unreported incidence Yap et al. ((2013)); Guzmán and Kouri ((2002)), we achieve proper data fitting via maximum likelihood estimation.

The cause of DHF is still unclear; for example, in ESTEVA and YANG ((2015)) reports that the concentration of the virus explains the severity of Dengue. In this direction, Gomez et al. provide a model that considers a severity stratification according to the mosquito load virulence.

Our contribution describes scenarios wherein past Dengue outbreaks detected a dominant strain. Then decision-makers could project new dengue strain invasion scenarios. Thus, we believe that early serotype identification in any Dengue outbreak would be essential to make epidemiological decisions.

In short, our contribution suggest that early serotype identification in any Dengue outbreak would be essential infromation to make epidemiological decisions.

## 7. Conclusions

The main conclusions of the study may be presented in a short Conclusions section, which may stand alone or form a subsection of a Discussion or Results and Discussion section.

In this work we have presented a mathematical model to understand the 2010 dengue outbreak that occurred in Hermosillo, Mexico. The model includes infected classes of classic and hemorrhagic versions of dengue in order to adjust the observed data. To our knowledge, there is no published work

In this work we have presented a mathematical model to understand the 2010 dengue outbreak that occurred in Hermosillo, Mexico. The model includes infected classes of classic and hemorrhagic versions of dengue in order to adjust the observed data. To our knowledge, there is no published work

## References

- A. Balmaseda, S. N. Hammond, Y. Tellez, L. Imhoff, Y. Rodriguez, S. I. Saborío, J. C. Mercado, L. Perez, E. Vide, E. Almanza, et al.  
High seroprevalence of antibodies against dengue virus in a prospective study of schoolchildren in managua, nicaragua.  
*Tropical medicine & international health*, 11(6):935–942, 2006.

C. Chastel.

Infections inapparentes chez l'homme : un cheval de troie pour l'introduction et la diffusion des arbovirus transmis par des moustiques dans les régions non endémiques ?

*Bulletin de la Société de pathologie exotique*, 104(3):213, Jun 2011.

ISSN 1961-9049.

doi:10.1007/s13149-011-0165-1.

URL <https://doi.org/10.1007/s13149-011-0165-1>.

15

C. Chastel.

Eventual role of asymptomatic cases of dengue for the introduction and spread of dengue viruses in non-endemic regions.

*Frontiers in Physiology*, 3:70, 2012.

ISSN 1664-042X.

doi:10.3389/fphys.2012.00070.

URL <https://www.frontiersin.org/article/10.3389/fphys.2012.00070>.

15

L. ESTEVA and H. M. YANG.

ASSESSING THE EFFECTS OF TEMPERATURE AND DENGUE VIRUS LOAD ON DENGUE TRANSMISSION.

*Journal of Biological Systems*, 23(04):1550027, dec 2015.

ISSN 0218-3390.

doi:10.1142/S0218339015500278.

URL <http://www.worldscientific.com/doi/abs/10.1142/S0218339015500278>.

9

Z. Feng and J. X. Velasco-Hernández.

Competitive exclusion in a vector-host model for the dengue fever.

*Journal of Mathematical Biology*, 35(5):523–544, apr 1997.

ISSN 0303-6812.

doi:10.1007/s002850050064.

URL <http://link.springer.com/10.1007/s002850050064>.

6, 9, 15

H. Gómez Dantés, J. A. Farfán-Ale, and E. Sarti.

Epidemiological Trends of Dengue Disease in Mexico (2000–2011): A Systematic Literature Search and Analysis.

*PLoS Neglected Tropical Diseases*, 11(8):1–13 (e3158), 2014.

3

M. G. Guzmán and G. Kouri.

Dengue: an update.

*The Lancet Infectious Diseases*, 2(1):33–42, 2002.

ISSN 1473-3099.

doi:10.1016/S1473-3099(01)00171-2.

URL <http://www.sciencedirect.com/science/article/pii/S1473309901001712>.

9

- S. Halstead.  
The xxth century dengue pandemic: need for surveillance and research.  
*World health statistics quarterly. Rapport trimestriel de statistiques sanitaires mondiales*, 45(2-3): 292—298, 1992.  
ISSN 0379-8070.  
URL <http://europepmc.org/abstract/MED/1462664>.  
3
- N. Nuraini, E. Soewono, and K. A. Sidarto.  
Mathematical model of Dengue disease transmission with severe DHF compartment.  
*Bull. Malays. Math. Sci. Soc. (2)*, 30(2):143–157, 2007.  
ISSN 0126-6705.  
9
- M. OhAinle, A. Balmaseda, A. R. Macalalad, Y. Tellez, Z. M. C., S. Saborío, A. Nuñez, N. J. Lennon, B. W. Birren, A. Gordon, M. R. Henn, and E. Harris.  
Dynamics of Dengue Disease Severity Determined by the Interplay Between Viral Genetics and Serotype-Specific Immunity.  
*Sci Transl Med.*, 114(3):1 –29, 2011.  
doi:10.1126/scitranslmed.3003084.  
4
- S. T. R. Pinho, C. P. Ferreira, L. Esteva, F. R. Barreto, V. C. Morato e Silva, and M. G. L. Teixeira.  
Modelling the dynamics of dengue real epidemics.  
*Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 368(1933):5679–5693, 2010.  
ISSN 1364-503X.  
doi:10.1098/rsta.2010.0278.  
URL <http://rsta.royalsocietypublishing.org/content/368/1933/5679>.  
15
- P. A. Reyes-Castro, R. B. Harris, H. E. Brown, G. L. Christopherson, and K. C. Ernst.  
Spatio-temporal and neighborhood characteristics of two dengue outbreaks in two arid cities of Mexico.  
*Acta Tropica*, (167):174–182, 2017.  
3
- N. Sangkawibha, S. Rojanasuphot, A. Sompop, S. Viriyapongse, V. Jatanasen, Sujarti abd Salitul, B. Phanthumachinda, and S. B. Halstead.  
Risk factors in dengue shock syndrome: A prospective epidemiologic study in Rayong, Thailand: I. The 1980 Outbreak .  
*American Journal of Epidemiology*, 5(120):653–669, 1984.  
doi:10.1093/oxfordjournals.aje.a113932.  
4
- R. Streatfield, G. Bielby, and D. Sinclair.  
A primary dengue 2 epidemic with spontaneous haemorrhagic manifestations.  
*The Lancet*, 342(8870):560 – 561, 1993.

ISSN 0140-6736.

doi:[https://doi.org/10.1016/0140-6736\(93\)91692-F](https://doi.org/10.1016/0140-6736(93)91692-F).

URL <http://www.sciencedirect.com/science/article/pii/014067369391692F>.

Originally published as Volume 2, Issue 8870.

3

M. Vázquez-Pichardo, C. Rosales-Jiménez, A. Núñez-León, P. Rivera-Osorio, S. De La Cruz-Hernández, A. Ruiz-López, S. González-Mateos, I. López-Martínez, J. C. Rodríguez-Martínez, H. López-Gatell, and C. Alpuche-Aranda.

Dengue serotypes in Mexico during 2009-2010.

*Bol. Med. Hosp. Infant. Mex.*, 2(68):94–100, 2011.

3

WHO.

Dengue haemorrhagic fever: diagnosis, treatment, prevention and control. 2nd edition. Geneva : World health organization.

<http://www.who.int/csr/resources/publications/dengue/Denguepublication/en/>,

1997 (accessed May, 2018).

3

H. M. YANG, M. L. G. MACORIS, K. C. GALVANI, M. T. M. ANDRIGHETTI, and D. M. V. WANDERLEY.

Assessing the effects of temperature on the population of *Aedes aegypti*, the vector of dengue.

*Epidemiology and Infection*, 137(08):1188, aug 2009.

ISSN 0950-2688.

doi:10.1017/S0950268809002040.

URL <http://www.journals.cambridge.org/abstract{ }S0950268809002040>.

15

G. Yap, C. Li, A. Mutalib, Y.-L. Lai, and L.-C. Ng.

High Rates of Inapparent Dengue in Older Adults in Singapore.

*The American Journal of Tropical Medicine and Hygiene*, 88(6):1065–1069, 2013.

ISSN 0002-9637.

doi:10.4269/ajtmh.12-0150.

URL <https://www.ajtmh.org/content/journals/10.4269/ajtmh.12-0150>.

9

T. R. Yasuno M.

A study of biting habitats of *Aedes aegypti* in Bangkok, Thailand.

*Bulletin of the World Health Organization*, 43(2):319–325, 1990.

ISSN 0042-9686.

15

T.-T. Zheng and L.-F. Nie.

Modelling the transmission dynamics of two-strain Dengue in the presence awareness and vector control.

*Journal of Theoretical Biology*, 443:82–91, 2018.

ISSN 0022-5193.

*REFERENCES*

13 of 17

doi:10.1016/j.jtbi.2018.01.017.

URL <http://www.sciencedirect.com/science/article/pii/S0022519318300316>.

9

disiase\_flow.pdf

Figure 1. Flow diagram of model (??)

Symbol	Meaning	Reference	Range	units
$M_S(0)$ $M_1(0)$ , $M_2(0)$	Initial number of susceptible and infected mosquitoes.			
$N_H$	Total Susceptible population	INEGI (see ??)	283 493	
$b$	Biting rate	Yasuno M ((1990))	[10.36 , 33.39]	meals/week
$\Lambda_M$	Vector birth rate		$\mu_M \cdot N_M$	week <sup>-1</sup>
$\mu_M$	vector mortality rate	YANG et al. ((2009))	[0.252, 0.763]	week <sup>-1</sup>
$\mu_H$	Human mortality	—	0.000 273 973	week <sup>-1</sup>
$\beta_H$	Human infection  probability by vectors	Feng and Velasco- Hernández ((1997))	(0, 0.05]	—
$\beta_M$	Vector infection  probability by humans	Feng and Velasco- Hernández ((1997))	(0, 0.05]	—
$\alpha_c$	Mean recover rate from Classic Dengue	Pinho et al. ((2010))	[0.581, 1.75]	week <sup>-1</sup>
$\alpha_h$	Mean recover rate from Hemorrhagic Dengue	Pinho et al. ((2010))	[0.581, 1.75]	week <sup>-1</sup>
$\sigma$	Susceptibility to serotype  DENV-2.	Feng and Velasco- Hernández ((1997))	(0, 5)	
$p$	Ratio of asymptomatic cases	Balmaseda et al. ((2006)), Chastel ((2011, 2012))	$[\frac{1}{60}, \frac{1}{30}]$	—
$\theta$	Probability of acquire DHF as second infection			

Table 2. Parameter description

Parameters (time in weeks) for figs. 2 and 3		
$\Lambda_M = 30702.6139006,$	$\Lambda_S = 10.2385934233,$	$\Lambda_{S_{-1}} = 1.13762149148,$
$\alpha_c = 0.686615937276,$	$\alpha_h = 1.41310092256,$	$b = 12.7122333418,$
$\beta_H = 0.0478488977733,$	$\beta_H = 0.0478488977733,$	$\beta_M = 0.0361065995648,$
$\mu_H = 0.000273,$	$\mu_M = 0.307170720093,$	
$\sigma = 1.806480946,$		
$\theta = 0.1887501857,$		
$p = 0.126295209216,$	$h = 0.000189285714286,$	
$S(0) = 35598.0,$	$I_1(0) = 1.0,$	$I_2(0) = 1.0,$
$M_S(0) = 120000,$	$M_1(0) = 10,$	$M_2(0) = 10,$
$S_{-1}(0) = 4400.0,$	$Y_{-1}^{[c]}(0) = 0.0,$	
$Y_{-1}^{[h]}(0) = 0.0,$	$z(0) = 0.252590418433,$	$Rec(0) = 0.0,$

Table 3. Parameters of numerical example




Figure 2. DF and DHF numerical solutions versus Dengue data from 2010 Hermosillo outbreak. Python code and data in <https://github.com/SaulDiazInfante/Two-strains-dengue-model-data-fitting/tree/master/StochasticSearchPySimplifiedModel>






Figure 3. Evolutions of each stage.