

Optimal constant piecewise vaccination policies for COVID-19

Gabriel A. Salcedo-Varela^a, Francisco Peñuñuri^b, D. González-Sánchez^c, Saúl Díaz-Infante^{c,*}

^a Departamento de Matemáticas, Universidad de Sonora, Blvd. Luis Encinas y Rosales S/N, Hermosillo, Sonora, México, C.P. 83000.

^b Facultad de Ingeniería, Universidad Autónoma de Yucatán, A.P. 150, Cordemex, Mérida, Yucatán, México.

^c CONACYT-Universidad de Sonora, Departamento de Matemáticas, Blvd. Luis Encinas y Rosales S/N, Hermosillo, Sonora, México, C.P. 83000.

Abstract

BACKGROUND.

FINDINGS.

IMPLICATIONS.

Keywords: COVID-19, Optimal Control, COVAX, Vaccination, WHO-SAGE, DALYs.

1. Introduction

Main contribution and its relevance.

Background.

Vaccine development.

Problem setup.

Litterature review.

Papaer structure.

2. Covid-19 spread dynamics

Uncontrolled dynamics. We split the the constant population N in a base SEIR structure with segregation infected classes according with manifestation of symptoms. We also postulate the extra state Y_{IS} to fit commulative symptomatic cases reported in the databases from Mexico city during the exponential grow phase. Our dynamics

reads

$$\begin{aligned} L' &= \theta\mu N^* - \epsilon\lambda L - \delta_L L - \mu L, \\ S' &= (1 - \theta)\mu N^* + \delta_L L + \delta_R R - (\lambda + \mu)S, \\ E' &= \lambda(\epsilon L + S) - (\kappa + \mu)E, \\ I'_S &= p\kappa E - (\gamma_S + \delta_H + \underbrace{\mu I_S}_{\text{SDIV}} + \mu)I_S, \\ I'_A &= (1 - p)\kappa E - (\gamma_A + \mu)I_A, \\ H' &= \delta_H I_S - (\gamma_H + \mu_H + \mu)H, \\ R' &= \gamma_S I_S + \gamma_A I_A + \gamma_H H - (\delta_R + \mu)R, \\ D' &= \underbrace{\mu I_S}_{\text{SDIV}} + \mu_H H, \end{aligned} \tag{1}$$

$$\frac{dY_{IS}}{dt} = p\kappa E,$$

$$\lambda := \frac{\beta_A I_A + \beta_S I_S}{N^*},$$

$$N^*(t) = L + S + E + I_S + I_A + H + R.$$

See table[*] for notation and values.

Hypothesis. We consider that susceptible individuals become infected when they are in contact with asymptomatic individuals and individuals with symptoms, we will propose that a proportion of asymptomatic individuals have a way to get relief and not die. A proportion of individuals infected with symptoms may die from the disease or may be relieved.

We callibrate parameters of our base dynamics in (1) via Multichain Montecarlo (MCMC). To this end, we assume that the comulative incidence of new infected symptomatic cases CI_S follows a Poisson distribution with mean $\lambda_t = IC_s(t)$. Further, following [] we postulate priors for

*Corresponding author

Email addresses: adrian.acuna@unison.mx (Gabriel A. Salcedo-Varela), francisco.pa@uady.mx (Francisco Peñuñuri), dgonzalezsa@conacyt.mx (D. González-Sánchez), saúl.diazinfante@unison.mx (Saúl Díaz-Infante)

Parameter	Description
μ	Death rate
β_S	Infection rate between susceptible and symptomatic infected
β_A	Infection rate between susceptible and asymptomatic infected
λ_V	Vaccination rate
δ_V^{-1}	Vaccine-induced immunity
ε	Vaccine efficacy
κ^{-1}	Average incubation time
p	New asymptomatic generation proportion
θ	Proportion of individuals under lockdown
γ_S^{-1}	Average time of symptomatic recovery
γ_A^{-1}	Recovery average time of asymptomatic individuals
γ_H^{-1}	Recovery average time by hospitalization
δ_R^{-1}	Natural immunity
δ_H	Infected symptomatic hospitalization rate

Table 1: Parameters definition of model in eq. (1).




Figure 1: Fit of diary new cases of Mexico city during exponential growth.

p and κ

$$\begin{aligned}
Y_t &\sim \text{Poisson}(\lambda_t), \\
\lambda_t &= \int_0^t p \delta_e E, \\
p &\sim \text{Uniform}(0.3, 0.8), \\
\kappa &\sim \text{Gamma}(10, 50).
\end{aligned} \tag{2}$$

Using the reproductive number definition of VanDenDrishe, we obtain

$$R_0 := \frac{N^*(\beta_S p \kappa + \beta_A \kappa (1 - p))}{(\mu - \kappa)(\gamma_S + \mu_{I_s} + \gamma_A + \mu) N^* \mu}$$

Figure [...] displays data of cumulative confirmed cases of COVID-19 of Mexico city, and the fit of our model in eqs. (1) and (2).

3. Imperfect-preventive Covid-19 vaccination

Preventive vaccines.

Efficacy and vaccine-induced immunity.

Actual vaccine stage development.

Vaccination reproductive number.

Vaccination rate λ_V estimate.

Feasibility regions according to efficacy and vaccination rate.

$$\begin{aligned}
L' &= \theta \mu N^* - (\epsilon \lambda + \delta_L + \mu) L \\
S' &= (1 - \theta) \mu N^* + \delta_L L + \delta_V V + \delta_R R \\
&\quad - (\lambda + \lambda_V + \mu) S \\
E' &= \lambda (\epsilon L + (1 - \epsilon) V + S) - (\kappa + \mu) E \\
I_S' &= p \kappa E - (\delta_H + \gamma_S + \mu_{I_S} + \mu) I_S \\
I_A' &= (1 - p) \kappa E - (\gamma_A + \mu) I_A \\
H' &= \delta_H I_S - (\gamma_H + \mu_H + \mu) H \\
R' &= \gamma_S I_S + \gamma_A I_A + \gamma_H H - (\delta_R + \mu) R \\
D' &= \mu_{I_S} I_S + \mu_H H \\
V' &= \lambda_V S - [(1 - \epsilon) \lambda + \delta_V + \mu] V
\end{aligned}$$

$$\begin{aligned}
\frac{dX_{vac}}{dt} &= (u_V(t) + \lambda_V) [L + S + E + I_A + R] \\
\frac{dY_{I_S}}{dt} &= p \kappa E \\
\lambda &:= \frac{\beta_A I_A + \beta_S I_S}{N^*}
\end{aligned}$$

$$\begin{aligned}
L(0) &= L_0, \quad S(0) = S_0, \quad E(0) = E_0, \quad I_S(0) = I_{S_0}, \\
I_A(0) &= I_{A_0}, \quad H(0) = H_0, \quad R(0) = R_0, \quad D(0) = D_0, \\
V(0) &= 0, \quad X_{vac}(0) = 0, \\
X_{vac}(T) &= x_{coverage}, \\
N^*(t) &= L + S + E + I_S + I_A + H + R + V
\end{aligned} \tag{3}$$

4. Vaccination reproductive number

5. Optimal controlled version

Controlled Model. Now we model vaccination, treatment and lockdown as an optimal control problem. According to dynamics in eq. (1), we modulate the vaccination rate by a time-dependent control signal $u_V(t)$. We add compartment X_{vac} to count all the vaccine applications of susceptible, exposed, asymptomatic and recovered individuals. This process is modeled by

$$X'(t) = (\lambda_V + u_V(t))(S + E + I_A + R) \tag{4}$$

and describes the number of applied vaccines at time t . Consider

$$x(t) := (L, S, E, I_S, I_A, H, R, D, V, X_{vac})^\top(t)$$

and control signal $u_V(\cdot)$. We quantify the cost and reward of a vaccine strategy policy via the penalization functional

$$J(u_V) := \int_0^T a_S I_S + a_D D + \frac{1}{2} c_V u_V^2 ds. \tag{5}$$

In other words, we assume in functional J that pandemic cost is proportional to the symptomatic and death reported cases and that a vaccination policy implies quadratic consumption of resources.

Further, since we aim to simulate vaccination policies at different coverage scenarios, we impose the vaccination counter state's final time condition $X(T)$

$$\begin{aligned}
x(T) &= (\cdot, \cdot, \cdot, \cdot, \cdot, \cdot, \cdot, \cdot, X_{vac}(T))^\top, \in \Omega \\
X_{vac}(T) &= x_{coverage}, \\
x_{coverage} &\in \{\text{Low}(0.2), \text{Mid}(0.5), \text{High}(0.8)\}.
\end{aligned} \tag{6}$$

Thus, given the time horizon T , we impose that the last fraction of vaccinated populations corresponds to 20%, 50% or 80%, and the rest of final states as free. We also impose the path constraint

$$\Phi(x, t) := \kappa I_S(t) \leq B, \quad \forall t \in [0, T], \tag{7}$$

to ensure that healthcare services will not be overloaded. Here κ denotes hospitalization rate, and B is the load capacity of a health system.

Given a fixed time horizon and vaccine efficiency, we estimate the constant vaccination rate as the solution of

$$x_{coverage} = 1 - \exp(-\lambda_V T). \tag{8}$$

That is, λ_v denotes the constant rate to cover a fraction $x_{coverage}$ in time horizon T . Thus, according to this vaccination rate, we postulate a policy u_v that modulates vaccination rate according to λ_V as a baseline. That is, optimal vaccination amplifies or attenuates the estimated baseline λ_V in a interval $[\lambda_v^{\min}, \lambda_v^{\max}]$ to optimize functional $J(\cdot)$ —minimizing symptomatic, death reported cases and optimizing resources.

Our objective is minimize the cost functional (5)—over an appropriated functional space—subject to the dynamics in eqs. (1) and (4), boundary conditions, and the path constrain in (7). That is, we search for vaccination policies $u_V(\cdot)$, which solve the following optimal control problem

(OCP).

$$\min_{u \in \mathcal{U}} J(u) := \int_0^T [(a_D \mu_s + a_H \delta_H) I_S(r) + a_{I_S} p \kappa E(r)] dr$$

s. t.

$$\begin{aligned} L' &= \theta \mu N^* - \epsilon \lambda L - u_L(t) L - \mu L \\ S' &= (1 - \theta) \mu N^* + u_L(t) L + \delta_v V + \delta_R R \\ &\quad - [\lambda + (\lambda_V + u_V(t)) + \mu] S \\ E' &= \lambda(\epsilon L + (1 - \epsilon) V + S) - (\kappa + \mu) E \\ I_S' &= p \kappa E - (\gamma_S + \mu_{I_S} + \delta_H + \mu) I_S \\ I_A' &= (1 - p) \kappa E - (\gamma_A + \mu) I_A \\ H' &= \delta_H I_S - (\gamma_H + \mu_H + \mu) H \\ R' &= \gamma_S I_S + \gamma_A I_A + \gamma_H H - (\delta_R + \mu) R \\ D' &= \mu_{I_S} I_S + \mu_H H \\ V' &= (\lambda_V + u_V(t)) S - [(1 - \epsilon) \lambda + \delta_V + \mu] V \end{aligned}$$

$$\begin{aligned} \frac{dX_{vac}}{dt} &= (u_V(t) + \lambda_V) [L + S + E + I_A + R] \\ \frac{dY_{I_S}}{dt} &= p \kappa E \\ \lambda &:= \frac{\beta_A I_A + \beta_S I_S}{N^*} \end{aligned}$$

$$\begin{aligned} L(0) &= L_0, \quad S(0) = S_0, \quad E(0) = E_0, \quad I_S(0) = I_{S_0}, \\ I_A(0) &= I_{A_0}, \quad H(0) = H_0, \quad R(0) = R_0, \quad D(0) = D_0, \\ V(0) &= 0, \quad X_{vac}(0) = 0, \quad u_V(\cdot) \in [u_{\min}, u_{\max}], \\ X_{vac}(T) &= x_{coverage}, \quad \kappa I_S(t) \leq B, \quad \forall t \in [0, T], \\ N^*(t) &= L + S + E + I_S + I_A + H + R + V \end{aligned} \tag{9}$$

6. Parameter calibration

Bayesian estimation.

7. Optimal control problem

8. Numerical Results

Appendix A. Appendix

Consider the following cost functional that we want to minimize

$$\int_0^T C(t, X(t), u(t)) dt \tag{A.1}$$

subject to the dynamics

$$\dot{X}(t) = f(t, X(t), u(t)), \quad 0 \leq t \leq T, \tag{A.2}$$

and the initial state $X(0) = x_0$. Let $t_0 < t_1 < \dots < t_n$, with $t_0 = 0$ and $t_n = T$, be a partition of the interval $[0, T]$. We consider *piecewise constant controls* \tilde{u} of the form

$$\tilde{u}(t) = a_j \quad t_j \leq t < t_{j+1} \tag{A.3}$$

for $j = 0, \dots, n-1$. ASSUMPTION 1. ASSUMPTION 2. By Assumption 1, the system

$$\dot{X}(t) = f(t, X(t), a_0), \quad X(0) = x_0, \quad 0 \leq t \leq t_1,$$

has a unique solution $\tilde{X}_0(t; x_0, a_0)$ which is continuous in (x_0, a_0) . Next, put $x_1 := \tilde{X}_0(t_1; x_0, a_0)$ and consider the system

$$\dot{X}(t) = f(t, X(t), a_1), \quad X(t_1) = x_1, \quad t_1 \leq t \leq t_2,$$

which, again by Assumption 1, has a unique solution $\tilde{X}_1(t; x_1, a_1)$ continuous in (x_1, a_1) . By following this procedure, we end up having a recursive solution

$$\tilde{X}_{n-1}(t; x_{n-1}, a_{n-1}), \quad x_{n-1} := \tilde{X}_{n-2}(t_{n-1}; x_{n-2}, a_{n-1}), \quad t_{n-1} \leq t \leq t_n$$

Thus, for a control \tilde{u} of the form (A.3) and the corresponding solution path \tilde{X} , we have

$$\int_0^T C(t, \tilde{X}(t), \tilde{u}(t)) dt = \sum_{j=0}^{n-1} \int_{t_j}^{t_{j+1}} C(t, \tilde{X}_j(t), a_j) dt.$$

Notice that each \tilde{X}_j is a continuous function of (a_0, \dots, a_j) and x_0 . Therefore, by Assumption 2, the mapping

$$(a_0, \dots, a_{n-1}) \mapsto \sum_{j=0}^{n-1} \int_{t_j}^{t_{j+1}} C(t, \tilde{X}_j(t), a_j) dt$$

is continuous.

References

- [1] Yinon M Bar-on, Ron Sender, Avi I Flamholz, Rob Phillips, and Ron Milo. A quantitative compendium of COVID-19 epidemiology. pages 1–51.
- [2] Bruno Buonomo. Effects of information-dependent vaccination behavior on coronavirus outbreak: insights from a SIRI model. *Ricerche di Matematica*, 2020.
- [3] Anastasia Chatzilela, Edwin van Leeuwen, Oliver Ratmann, Marc Baguelin, and Nikolaos Demiris. Contemporary statistical inference for infectious disease models using Stan. *Epidemics*, 29(February):100367, 2019.
- [4] Nicholas G Davies, Adam J Kucharski, Rosalind M Eggo, Amy Gimma, CMMID COVID-19 Working Group, and W. John Edmunds. The effect of non-pharmaceutical interventions on COVID-19 cases, deaths and demand for hospital services in the UK: a modelling study. *medRxiv*, page 2020.04.01.20049908, 2020.
- [5] Laura Di Domenico, Giulia Pullano, Chiara E Sabbatini, Pierre-Yves Boëlle, and Vittoria Colizza. Currently under screening at medRxiv Expected impact of lockdown in Île-de-France and possible exit strategies. 2020.
- [6] Wandu Ding and Suzanne Lenhart. Introduction to optimal control for discrete time models with an application to disease modeling. In *Modeling paradigms and analysis of disease transmission models*, volume 75, pages 109–119. Amer. Math. Soc., Providence, RI, oct 2010.
- [7] Ramses Djidjou-Demasse, Yannis Michalakis, Marc Choisy, Micea T. Sofonea, and Samuel Alizon. Optimal COVID-19 epidemic control until vaccine deployment. *medRxiv*, page 2020.04.02.20049189, 2020.

Parameter	Median	Reference
q_r, ϵ	0.4, 0.3, 0.1	this study
β_S	$q_r \times 8.690\,483 \times 10^{-1}$	this study
β_A	$q_r \times 7.738\,431 \times 10^{-1}$	this study
κ	0.196\,078\,43	*
p	0.1213	*
θ	0.2,	this study
δ_L	0.04	postulated
δ_H	0.2	*
δ_V	0.002\,739\,726\,027\,397\,260\,3	$\delta_V^{-1} = 2$ years CanSinoBIO
δ_R	0.005\,555\,56	$\delta_R^{-1} \approx 180$ days
μ	$3.913\,894 \times 10^{-5}$	**
μ_{I_S}	0.0	
μ_H	0.016\,32	[FENG]
γ_S	0.092\,506\,94	*
γ_A	0.167\,504\,19	*
γ_H	$5.079\,869 \times 10^{-1}$	*
λ_V	0.000\,611\,35	
ε	0.7, 0.80, 0.9, 0.95	[PRESS RELESASES]
N	26\,446\,435	**
L_0	0.266\,260\,097\,021\,127\,96	
S_0	0.463\,606\,046\,009\,872	
E_0	0.000\,670\,33	*
I_{S_0}	9.283×10^{-5}	* * *
I_{A_0}	0.001\,209\,86	*
H_0	$1.341\,579\,69 \times 10^{-4}$	**
R_0	$2.661\,259\,39 \times 10^{-1}$	
D_0	0.001\,900\,74	**
X_{vac}^0	0.0	
V_0	0.0	
$Y_{I_S}^0$	0.122\,581\,64	
B	0.000\,359\,216\,658\,124\,242\,5	9500 beds/ N
a_{I_S}	0.002\,012\,775\,543\,825\,648\,6	DALY def
a_H	0.001\,411\,888\,738\,103\,725, or $a_H(x) := 0.001\,411\,888\,738\,103\,725 \log(\frac{1}{B-\kappa I_S})$	DALY def [Jo 2020]
a_D	7.25	DALY def

- [8] Neil M Ferguson, Daniel Laydon, Gemma Nedjati-gilani, Nat-suko Imai, Kylie Ainslie, Marc Baguelin, Sangeeta Bha-tia, Adhiratha Boonyasiri, Zulma Cucunubá, Gina Cuomo-dannenburg, Amy Dighe, Han Fu, Katy Gaythorpe, Will Green, Arran Hamlet, Wes Hinsley, Lucy C Okell, Sabine Van, Hay-ley Thompson, Robert Verity, Erik Volz, Haowei Wang, Yuan-rong Wang, Patrick G T Walker, Caroline Walters, Peter Win-skill, Charles Whittaker, Christl A Donnelly, Steven Riley, and Azra C Ghani. Impact of non-pharmaceutical interventions (NPIs) to reduce COVID- 19 mortality and healthcare demand. (March), 2020.
- [9] M. H. A. Biswas, L. T. Paiva, and MdR de Pinho. A SEIR model for control of infectious diseases with constraints. *Mathematical Biosciences and Engineering*, 11(4):761–784, 2014.
- [10] IHME COVID-19 health service utilization forecasting Team and Christopher JL Murray. Forecasting COVID-19 impact on hospital bed-days, ICU-days, ventilator-days and deaths by US state in the next 4 months. *medRxiv*, page 2020.03.27.20043752, mar 2020.
- [11] Gang Huang, Yasuhiro Takeuchi, Wanbiao Ma, and Daijun Wei. Global Stability for Delay SIR and SEIR Epidemic Models with Nonlinear Incidence Rate. *Bulletin of Mathematical Biology*, 72(5):1192–1207, 2010.
- [12] Jiwei Jia, Jian Ding, Siyu Liu, Guidong Liao, Jingzhi Li, B. E.N. Duan, Guoqing Wang, and R. A.N. Zhang. Modeling the control of COVID-19: Impact of policy interventions and meteorological factors. *Electronic Journal of Differential Equations*, 2020:1–21, 2020.
- [13] Abdelilah Kaddar, Abdelhadi Abta, and Hamad Talibi Alaoui. A comparison of delayed SIR and SEIR epidemic models. *Non-linear Analysis: Modelling and Control*, 16(2):181–190, apr 2011.
- [14] Qian Li, Biao Tang, Nicola Luigi Bragazzi, Yanni Xiao, and Jianhong Wu. Modeling the impact of mass influenza vaccina-tion and public health interventions on COVID-19 epidemics with limited detection capability. *Mathematical Biosciences*, 325(May):108378, jul 2020.
- [15] Gustavo Barbosa Libotte, Fran Sérgio Lobato, Gustavo Mendes Platt, and Antônio José da Silva Neto. Determination of an Op-timal Control Strategy for Vaccine Administration in COVID-19 Pandemic Treatment. (November 2019), apr 2020.
- [16] Manotosh Mandal, Soovoojeet Jana, Swapan Kumar Nandi, Anupam Khatua, Sayani Adak, and T.K. Kar. A model based study on the dynamics of COVID-19: Prediction and control. *Chaos, Solitons & Fractals*, 136:109889, jul 2020.
- [17] Van Kinh Nguyen and Esteban A. Hernandez-Vargas. Parame-ter Estimation in Mathematical Models of Viral Infections Using R. volume 1836, pages 531–549. 2018.
- [18] Jorge Nocedal, Andreas Wächter, and Richard A. Waltz. Adap-tive Barrier Update Strategies for Nonlinear Interior Methods. *SIAM Journal on Optimization*, 19(4):1674–1693, jan 2009.
- [19] Xinwei Wang, Haijun Peng, Boyang Shi, Dianheng Jiang, Sheng Zhang, and Biaosong Chen. Optimal vaccination strategy of a constrained time-varying SEIR epidemic model. *Communica-tions in Nonlinear Science and Numerical Simulation*, 67:37–48, 2019.