

# Optimal constant piecewise vaccination policies for COVID-19

Gabriel A. Salcedo-Varela<sup>a</sup>, Francisco Peñuñuri<sup>b</sup>, D. González-Sánchez<sup>c</sup>, Saúl  
Díaz-Infante<sup>c,\*</sup>

<sup>a</sup> Departamento de Matemáticas, Universidad de Sonora, Blvd. Luis Encinas y Rosales  
S/N, Hermosillo, Sonora, México, C.P. 83000.

<sup>b</sup> Facultad de Ingeniería, Universidad Autónoma de Yucatán, A.P. 150, Cordemex, Mérida,  
Yucatán, México.

<sup>c</sup> CONACYT-Universidad de Sonora, Departamento de Matemáticas, Blvd. Luis Encinas y  
Rosales S/N, Hermosillo, Sonora, México, C.P. 83000.

---

## Abstract

BACKGROUND.

FINDINGS.

IMPLICATIONS.

*Keywords:* COVID-19, Optimal Control, COVAX, Vaccination, WHO-SAGE,  
DALYs.

---

## 1. Introduction

*Main contribution and its relevance.*

*Background.*

*Vaccine development.*

*Problem setup.*

*Litterature review.*

*Papaer structure.*

## 2. Covid-19 spread dynamics

*Uncontrolled dynamics.* We split the the constant population  $N$  in a base SEIR structure with segregation infected classes according with manifestation of symptoms. We also postulate the extra state  $Y_{I_S}$  to fit commulative symptomatic

---

\*Corresponding author

*Email addresses:* `adrian.acuna@unison.mx` (Gabriel A. Salcedo-Varela),  
`francisco.pa@uady.mx` (Francisco Peñuñuri), `dgonzalezsa@conacyt.mx` (D.  
González-Sánchez), `saul.diazinfante@unison.mx` (Saúl Díaz-Infante)

cases reported in the databases from Mexico city during the exponential grow phase. Our dynamics reads

$$\begin{aligned}
L' &= \theta\mu N^* - \epsilon\lambda L - \delta_L L - \mu L, \\
S' &= (1 - \theta)\mu N^* + \delta_L L + \delta_R R - (\lambda + \mu)S, \\
E' &= \lambda(\epsilon L + S) - (\kappa + \mu)E, \\
I'_S &= p\kappa E - (\gamma_S + \delta_H + \underline{\mu_{I_S}}^{\text{SDIV}} + \mu)I_S, \\
I'_A &= (1 - p)\kappa E - (\gamma_A + \mu)I_A, \\
H' &= \delta_H I_S - (\gamma_H + \mu_H + \mu)H, \\
R' &= \gamma_S I_S + \gamma_A I_A + \gamma_H H - (\delta_R + \mu)R, \\
D' &= \underline{\mu_{I_S}}^{\text{SDIV}} I_S + \mu_H H, \\
\frac{dY_{I_S}}{dt} &= p\kappa E, \\
\lambda &:= \frac{\beta_A I_A + \beta_S I_S}{N^*}, \\
N^*(t) &= L + S + E + I_S + I_A + H + R.
\end{aligned} \tag{1}$$

See Table 1 for notation and references values.




Figure 1: Cumulative new symptomatic and confirmed COVID19 reported cases from Ciudad de Mexico between March, 10, to March 30 of 2020.

27

*Hypothesis.* We consider that susceptible individuals become infected when they are in contact with asymptomatic individuals and individuals with symptoms,

Parameter	Description
$\mu$	Death rate
$\beta_S$	Infection rate between susceptible and symptomatic infected
$\beta_A$	Infection rate between susceptible and asymptomatic infected
$\lambda_V$	Vaccination rate
$\delta_V^{-1}$	Vaccine-induced immunity
$\varepsilon$	Vaccine efficacy
$\kappa^{-1}$	Average incubation time
$p$	New asymptomatic generation proportion
$\theta$	Proportion of individuals under lockdown
$\gamma_S^{-1}$	Average time of symptomatic recovery
$\gamma_A^{-1}$	Recovery average time of asymptomatic individuals
$\gamma_H^{-1}$	Recovery average time by hospitalization
$\delta_R^{-1}$	Natural immunity
$\delta_H$	Infected symptomatic hospitalization rate

Table 1: Parameters definition of model in Equation (1).

we will propose that a proportion of asymptomatic individuals have a way to get relief and not die. A proportion of individuals infected with symptoms may die from the disease or may be relieved.

We calibrate parameters of our base dynamics in (1) via Multichain Metropolis (MCMC). To this end, we assume that the cumulative incidence of new infected symptomatic cases  $CI_S$  follows a Poisson distribution with mean  $\lambda_t = IC_s(t)$ . Further, following [] we postulate priors for  $p$  and  $\kappa$

$$\begin{aligned}
Y_t &\sim \text{Poisson}(\lambda_t), \\
\lambda_t &= \int_0^t p \delta_e E, \\
p &\sim \text{Uniform}(0.3, 0.8), \\
\kappa &\sim \text{Gamma}(10, 50).
\end{aligned} \tag{2}$$

Using the reproductive number definition of VanDenDrishe, we obtain

$$R_0 := \frac{N^*(\beta_S p \kappa + \beta_A \kappa (1 - p))}{(\mu - \kappa)(\gamma_S + \mu_{I_s} + \gamma_A + \mu) N^* \mu}$$




Figure 2: Fit of diary new cases of Mexico city during exponential growth.

Figure 2 displays data of coumulative confirmed cases of COVID-19 of Mexico city, and the fitt of our model in Equations (1) and (2).

### 3. Imperfect-preventive Covid-19 vaccination

*Preventive vaccines.*

*Efficacy and vaccine-induced immunity.*

*Actual vaccine stage development.*

*Vaccination reproductive number.*

*Vaccination rate  $\lambda_V$  estimate.*

*Feasibility regions according to efficacy and vaccination rate.*

$$\begin{aligned}
L' &= \theta \mu N^* - (\epsilon \lambda + \delta_L + \mu) L \\
S' &= (1 - \theta) \mu N^* + \delta_L L + \delta_V V + \delta_R R \\
&\quad - (\lambda + \lambda_V + \mu) S \\
E' &= \lambda (\epsilon L + (1 - \epsilon) V + S) - (\kappa + \mu) E \\
I'_S &= p \kappa E - (\delta_H + \gamma_S + \mu_{I_S} + \mu) I_S \\
I'_A &= (1 - p) \kappa E - (\gamma_A + \mu) I_A \\
H' &= \delta_H I_S - (\gamma_H + \mu_H + \mu) H \\
R' &= \gamma_S I_S + \gamma_A I_A + \gamma_H H - (\delta_R + \mu) R \\
D' &= \mu_{I_S} I_S + \mu_H H \\
V' &= \lambda_V S - [(1 - \epsilon) \lambda + \delta_V + \mu] V
\end{aligned}$$

$$\begin{aligned}
\frac{dX_{vac}}{dt} &= (u_V(t) + \lambda_V) [S + E + I_A + R] \\
\frac{dY_{I_S}}{dt} &= p \kappa E \\
\lambda &:= \frac{\beta_A I_A + \beta_S I_S}{N^*}
\end{aligned} \tag{3}$$

$$\begin{aligned}
L(0) &= L_0, \quad S(0) = S_0, \quad E(0) = E_0, \\
I_S(0) &= I_{S_0}, \quad I_A(0) = I_{A_0}, \quad H(0) = H_0, \\
R(0) &= R_0, \quad D(0) = D_0, \\
V(0) &= 0, \quad X_{vac}(0) = 0, \\
X_{vac}(T) &= x_{coverage}, \\
N^*(t) &= L + S + E + I_S + I_A + H + R + V.
\end{aligned}$$

#### 4. Parameter calibration

*Bayesian estimation.*

#### 5. Vaccination reproductive number

*$R_0$  definition.*

*No vaccine reproductive number.*

*Vaccine reproductive number.*

Parameter	Median	Reference
$q_r, \epsilon$	0.4, 0.3, 0.1	this study
$\beta_S$	$q_r \times 8.690\,483 \times 10^{-1}$	this study
$\beta_A$	$q_r \times 7.738\,431 \times 10^{-1}$	this study
$\kappa$	0.196\,078\,43	*
$p$	0.1213	*
$\theta$	0.2,	this study
$\delta_L$	0.04	postulated
$\delta_H$	0.2	*
$\delta_V$	0.002\,739\,726\,027\,397\,260\,3	$\delta_V^{-1} = 2$ years CanSinoBIO
$\delta_R$	0.005\,555\,56	$\delta_R^{-1} \approx 180$ days
$\mu$	$3.913\,894 \times 10^{-5}$	**
$\mu_{I_S}$	0.0	
$\mu_H$	0.016\,32	[FENG]
$\gamma_S$	0.092\,506\,94	*
$\gamma_A$	0.167\,504\,19	*
$\gamma_H$	$5.079\,869 \times 10^{-1}$	*
$\lambda_V$	0.000\,611\,35	
$\varepsilon$	0.7, 0.80, 0.9, 0.95	[PRESS RELESASES]
$N$	26\,446\,435	**
$L_0$	0.266\,260\,097\,021\,127\,96	
$S_0$	0.463\,606\,046\,009\,872	
$E_0$	0.000\,670\,33	*
$I_{S_0}$	$9.283 \times 10^{-5}$	** *
$I_{A_0}$	0.001\,209\,86	*
$H_0$	$1.341\,579\,69 \times 10^{-4}$	**
$R_0$	$2.661\,259\,39 \times 10^{-1}$	
$D_0$	0.001\,900\,74	**
$X_{vac}^0$	0.0	
$V_0$	0.0	
$Y_{I_S}^0$	0.122\,581\,64	
$B$	0.000\,359\,216\,658\,124\,242\,5	9500 beds/ $N$
$a_{I_S}$	0.002\,012\,775\,543\,825\,648\,6	DALY def
$a_H$	0.001\,411\,888\,738\,103\,725, or $a_H(x) := 0.001\,411\,888\,738\,103\,725 \log(\frac{1}{B-\kappa I_S})$	DALY def [Jo 2020]
$a_D$	7.25	DALY def

51 *Efficacy, coverage and vaccination rate.*

52 Here Gabriel's R not calculations.<sup>SDIV</sup>

$$\frac{\kappa (\epsilon \mu p \theta \beta_A - \epsilon \mu p \theta \beta_s + \epsilon p \theta \beta_A \delta_H)}{\gamma_A \mu I_S \gamma_A \mu I_S} \quad (4)$$

[SDIV 1]  
Here contour  
plots figure  
as function  
of efficacy  
and vaccina-  
tion rate




Figure 3: R not contour plot as function of efficacy and vaccination rate

53

## 54 6. Optimal controlled version

55 *Controlled Model.* Now we model vaccination, treatment and lockdown as a  
56 optimal control problem. According to dynamics in Equation (1), we modu-  
57 late the vaccination rate with a time-dependent control signal  $u_V(t)$ . We add  
58 compartment  $X_{vac}$  to count all the vaccine applications of susceptible, exposed,  
59 asymptomatic and recovered individuals. This process is modeled by

$$X'(t) = (\lambda_V + u_V(t))(S + E + I_A + R) \quad (5)$$

and describes the number of applied vaccines at time  $t$ . Consider

$$x(t) := (L, S, E, I_S, I_A, H, R, D, V, X_{vac})^\top(t)$$

60 and control signal  $u_v(\cdot)$ . We quantify the cost and reward of a vaccine strategy  
61 policy via the penalization functional

$$J(u_V) := \int_0^T a_S I_S + a_d D + \frac{1}{2} c_V u_v^2 ds. \quad (6)$$

62 In other words, we assume in functional  $J$  that pandemic cost is proportional to  
63 the symptomatic and death reported cases and that a vaccination policy implies  
64 quadratic consumption of resources.

Further, since we aim to simulate vaccination policies at different coverage scenarios, we impose the vaccination counter state's final time condition  $X(T)$

$$\begin{aligned} x(T) &= (\cdot, \cdot, \cdot, \cdot, X_{vac}(T))^T, \in \Omega \\ X_{vac}(T) &= x_{coverage}, \\ x_{coverage} &\in \{\text{Low}(0.2), \text{Mid}(0.5), \text{High}(0.8)\}. \end{aligned} \tag{7}$$

Thus, given the time horizon  $T$ , we impose that the last fraction of vaccinated populations corresponds to 20%, 50% or 80%, and the rest of final states as free. We also impose the path constraint

$$\Phi(x, t) := \kappa I_S(t) \leq B, \quad \forall t \in [0, T], \tag{8}$$

to ensure that healthcare services will not be overloaded. Here  $\kappa$  denotes hospitalization rate, and  $B$  is the load capacity of a health system.

Given a fixed time horizon and vaccine efficiency, we estimate the constant vaccination rate as the solution of

$$x_{coverage} = 1 - \exp(-\lambda_V T). \tag{9}$$

That is,  $\lambda_v$  denotes the constant rate to cover a fraction  $x_{coverage}$  in time horizon  $T$ . Thus, according to this vaccination rate, we postulate a policy  $u_v$  that modulates vaccination rate according to  $\lambda_V$  as a baseline. That is, optimal vaccination amplifies or attenuates the estimated baseline  $\lambda_V$  in a interval  $[\lambda_v^{\min}, \lambda_v^{\max}]$  to optimize functional  $J(\cdot)$ —minimizing symptomatic, death reported cases and optimizing resources.

Our objective is minimize the cost functional (6)—over an appropriated functional space—subject to the dynamics in equations (1) and (5), boundary conditions, and the path constrain in (8). That is, we search for vaccination policies



83  $u_V(\cdot)$ , which solve the following optimal control problem (OCP).

$$\begin{aligned}
\min_{u \in \mathcal{U}} J(u) &:= \int_0^T [(a_D \mu_s + a_H \delta_H) I_S(r) + a_{I_S} p \kappa E(r)] dr \\
\text{s. t.} \\
L' &= \theta \mu N^* - \epsilon \lambda L - u_L(t) L - \mu L \\
S' &= (1 - \theta) \mu N^* + u_L(t) L + \delta_v V + \delta_R R \\
&\quad - [\lambda + (\lambda_V + u_V(t)) + \mu] S \\
E' &= \lambda(\epsilon L + (1 - \epsilon) V + S) - (\kappa + \mu) E \\
I_S' &= p \kappa E - (\gamma_S + \mu_{I_S} + \delta_H + \mu) I_S \\
I_A' &= (1 - p) \kappa E - (\gamma_A + \mu) I_A \\
H' &= \delta_H I_S - (\gamma_H + \mu_H + \mu) H \\
R' &= \gamma_S I_S + \gamma_A I_A + \gamma_H H - (\delta_R + \mu) R \\
D' &= \mu_{I_S} I_S + \mu_H H \\
V' &= (\lambda_V + u_V(t)) S - [(1 - \epsilon) \lambda + \delta_V + \mu] V
\end{aligned} \tag{10}$$

$$\begin{aligned}
\frac{dX_{vac}}{dt} &= (u_V(t) + \lambda_V) [L + S + E + I_A + R] \\
\frac{dY_{I_S}}{dt} &= p \kappa E \\
\lambda &:= \frac{\beta_A I_A + \beta_S I_S}{N^*}
\end{aligned}$$

$$\begin{aligned}
L(0) &= L_0, \quad S(0) = S_0, \quad E(0) = E_0, \quad I_S(0) = I_{S_0}, \\
I_A(0) &= I_{A_0}, \quad H(0) = H_0, \quad R(0) = R_0, \quad D(0) = D_0, \\
V(0) &= 0, \quad X_{vac}(0) = 0, \quad u_V(\cdot) \in [u_{\min}, u^{\max}], \\
X_{vac}(T) &= x_{coverage}, \quad \kappa I_S(t) \leq B, \quad \forall t \in [0, T], \\
N^*(t) &= L + S + E + I_S + I_A + H + R + V
\end{aligned}$$

## 84 7. Numerical Results

### 85 Changes (compact)

86 **Author: anonymous**

87 No changes.

88 **Author: SDIV**

89 Added ..... 1

90 Deleted ..... 2

91 Commented ..... 1

92

93 **Appendix A. Appendix**

94 Consider the following cost functional that we want to minimize

$$\int_0^T C(t, X(t), u(t)) dt \quad (\text{A.1})$$

95 subject to the dynamics

$$\dot{X}(t) = f(t, X(t), u(t)), \quad 0 \leq t \leq T, \quad (\text{A.2})$$

96 and the initial state  $X(0) = x_0$ . Let  $t_0 < t_1 < \dots < t_n$ , with  $t_0 = 0$  and  $t_n = T$ ,  
 97 be a partition of the interval  $[0, T]$ . We consider *piecewise constant controls*  $\tilde{u}$   
 98 of the form

$$\tilde{u}(t) = a_j \quad t_j \leq t < t_{j+1} \quad (\text{A.3})$$

99 for  $j = 0, \dots, n-1$ .

100 ASSUMPTION 1.

101 ASSUMPTION 2.

102 By Assumption 1, the system

$$\dot{X}(t) = f(t, X(t), a_0), \quad X(0) = x_0, \quad 0 \leq t \leq t_1,$$

103 has a unique solution  $\tilde{X}_0(t; x_0, a_0)$  which is continuous in  $(x_0, a_0)$ . Next, put  
 104  $x_1 := \tilde{X}_0(t_1; x_0, a_0)$  and consider the system

$$\dot{X}(t) = f(t, X(t), a_1), \quad X(t_1) = x_1, \quad t_1 \leq t \leq t_2,$$

which, again by Assumption 1, has a unique solution  $\tilde{X}_1(t; x_1, a_1)$  continuous in  
 $(x_1, a_1)$ . By following this procedure, we end up having a recursive solution

$$\begin{aligned} & \tilde{X}_{n-1}(t; x_{n-1}, a_{n-1}), \\ & x_{n-1} := \tilde{X}_{n-2}(t_{n-1}; x_{n-2}, a_{n-1}), \quad t_{n-1} \leq t \leq T. \end{aligned}$$

105 Thus, for a control  $\tilde{u}$  of the form (A.3) and the corresponding solution path  $\tilde{X}$ ,  
 106 we have

$$\int_0^T C(t, \tilde{X}(t), \tilde{u}(t)) dt = \sum_{j=0}^{n-1} \int_{t_j}^{t_{j+1}} C(t, \tilde{X}_j(t), a_j) dt.$$

107 Notice that each  $\tilde{X}_j$  is a continuous function of  $(a_0, \dots, a_j)$  and  $x_0$ . Therefore,  
 108 by Assumption 2, the mapping

$$(a_0, \dots, a_{n-1}) \mapsto \sum_{j=0}^{n-1} \int_{t_j}^{t_{j+1}} C(t, \tilde{X}_j(t), a_j) dt$$

109 is continuous.

## 110 References

- 111 [1] Yinon M Bar-on, Ron Sender, Avi I Flamholz, Rob Phillips, and Ron Milo.  
112 A quantitative compendium of COVID-19 epidemiology. pages 1–51.
- 113 [2] Bruno Buonomo. Effects of information-dependent vaccination behavior on  
114 coronavirus outbreak: insights from a SIRI model. *Ricerche di Matematica*,  
115 2020.
- 116 [3] Anastasia Chatzilena, Edwin van Leeuwen, Oliver Ratmann, Marc  
117 Baguelin, and Nikolaos Demiris. Contemporary statistical inference for in-  
118 fectious disease models using Stan. *Epidemics*, 29(February):100367, 2019.
- 119 [4] Nicholas G Davies, Adam J Kucharski, Rosalind M Eggo, Amy Gimma,  
120 CMMID COVID-19 Working Group, and W. John Edmunds. The effect  
121 of non-pharmaceutical interventions on COVID-19 cases, deaths and de-  
122 mand for hospital services in the UK: a modelling study. *medRxiv*, page  
123 2020.04.01.20049908, 2020.
- 124 [5] Laura Di Domenico, Giulia Pullano, Chiara E Sabbatini, Pierre-Yves  
125 Boëlle, and Vittoria Colizza. Currently under screening at medRxiv Ex-  
126 pected impact of lockdown in Île-de-France and possible exit strategies.  
127 2020.
- 128 [6] Wandi Ding and Suzanne Lenhart. Introduction to optimal control for  
129 discrete time models with an application to disease modeling. In *Modeling*  
130 *paradigms and analysis of disease transmission models*, volume 75, pages  
131 109–119. Amer. Math. Soc., Providence, RI, oct 2010.
- 132 [7] Ramses Djidjou-Demasse, Yannis Michalakis, Marc Choisy, Micea T. So-  
133 fonea, and Samuel Alizon. Optimal COVID-19 epidemic control until vac-  
134 cine deployment. *medRxiv*, page 2020.04.02.20049189, 2020.
- 135 [8] Neil M Ferguson, Daniel Laydon, Gemma Nedjati-gilani, Natsuko Imai,  
136 Kylie Ainslie, Marc Baguelin, Sangeeta Bhatia, Adhiratha Boonyasiri,  
137 Zulma Cucunubá, Gina Cuomo-dannenburg, Amy Dighe, Han Fu, Katy  
138 Gaythorpe, Will Green, Arran Hamlet, Wes Hinsley, Lucy C Okell, Sabine  
139 Van, Hayley Thompson, Robert Verity, Erik Volz, Haowei Wang, Yuan-  
140 rong Wang, Patrick G T Walker, Caroline Walters, Peter Winskill, Charles  
141 Whittaker, Christl A Donnelly, Steven Riley, and Azra C Ghani. Impact of  
142 non-pharmaceutical interventions ( NPIs ) to reduce COVID- 19 mortality  
143 and healthcare demand. (March), 2020.
- 144 [9] M. H. A. Biswas, L. T. Paiva, and MdR de Pinho. A SEIR model for  
145 control of infectious diseases with constraints. *Mathematical Biosciences*  
146 *and Engineering*, 11(4):761–784, 2014.
- 147 [10] IHME COVID-19 health service utilization forecasting Team and Christo-  
148 pher JL Murray. Forecasting COVID-19 impact on hospital bed-days,

- 149 ICU-days, ventilator-days and deaths by US state in the next 4 months.  
150 *medRxiv*, page 2020.03.27.20043752, mar 2020.
- 151 [11] Gang Huang, Yasuhiro Takeuchi, Wanbiao Ma, and Daijun Wei. Global  
152 Stability for Delay SIR and SEIR Epidemic Models with Nonlinear Inci-  
153 dence Rate. *Bulletin of Mathematical Biology*, 72(5):1192–1207, 2010.
- 154 [12] Jiwei Jia, Jian Ding, Siyu Liu, Guidong Liao, Jingzhi Li, B. E.N. Duan,  
155 Guoqing Wang, and R. A.N. Zhang. Modeling the control of COVID-  
156 19: Impact of policy interventions and meteorological factors. *Electronic*  
157 *Journal of Differential Equations*, 2020:1–21, 2020.
- 158 [13] Abdelilah Kaddar, Abdelhadi Abta, and Hamad Talibi Alaoui. A com-  
159 parison of delayed SIR and SEIR epidemic models. *Nonlinear Analysis:*  
160 *Modelling and Control*, 16(2):181–190, apr 2011.
- 161 [14] Qian Li, Biao Tang, Nicola Luigi Bragazzi, Yanni Xiao, and Jianhong Wu.  
162 Modeling the impact of mass influenza vaccination and public health inter-  
163 ventions on COVID-19 epidemics with limited detection capability. *Math-*  
164 *ematical Biosciences*, 325(May):108378, jul 2020.
- 165 [15] Gustavo Barbosa Libotte, Fran Sérgio Lobato, Gustavo Mendes Platt, and  
166 Antônio José da Silva Neto. Determination of an Optimal Control Strategy  
167 for Vaccine Administration in COVID-19 Pandemic Treatment. (November  
168 2019), apr 2020.
- 169 [16] Manotosh Mandal, Soovoojeet Jana, Swapan Kumar Nandi, Anupam  
170 Khatua, Sayani Adak, and T.K. Kar. A model based study on the dy-  
171 namics of COVID-19: Prediction and control. *Chaos, Solitons & Fractals*,  
172 136:109889, jul 2020.
- 173 [17] Van Kinh Nguyen and Esteban A. Hernandez-Vargas. Parameter Estima-  
174 tion in Mathematical Models of Viral Infections Using R. volume 1836,  
175 pages 531–549. 2018.
- 176 [18] Jorge Nocedal, Andreas Wächter, and Richard A. Waltz. Adaptive Bar-  
177 rier Update Strategies for Nonlinear Interior Methods. *SIAM Journal on*  
178 *Optimization*, 19(4):1674–1693, jan 2009.
- 179 [19] Xinwei Wang, Haijun Peng, Boyang Shi, Dianheng Jiang, Sheng Zhang,  
180 and Biaosong Chen. Optimal vaccination strategy of a constrained time-  
181 varying SEIR epidemic model. *Communications in Nonlinear Science and*  
182 *Numerical Simulation*, 67:37–48, 2019.