

Optimal constant piecewise vaccination policies for COVID-19

Gabriel A. Salcedo-Varela^a, Francisco Peñuñuri^b, D. González-Sánchez^c, Saúl Díaz-Infante^{c,*}

^a Departamento de Matemáticas, Universidad de Sonora, Blvd. Luis Encinas y Rosales S/N, Hermosillo, Sonora, México, C.P. 83000.

^b Facultad de Ingeniería, Universidad Autónoma de Yucatán, A.P. 150, Cordemex, Mérida, Yucatán, México.

^c CONACYT-Universidad de Sonora, Departamento de Matemáticas, Blvd. Luis Encinas y Rosales S/N, Hermosillo, Sonora, México, C.P. 83000.

Abstract

BACKGROUND.

FINDINGS.

IMPLICATIONS.

Keywords: COVID-19, Optimal Control, COVAX, Vaccination, WHO-SAGE, DALYs.

1. Introduction

Main contribution and its relevance.

Background.

Vaccine development.

Problem setup.

Litterature review.

Papaer structure.

2. Covid-19 spread dynamics

Uncontrolled dynamics. We split a given population of size N in the base SEIR structure with segregation infected classes according to the manifestation of symptoms. Let $L, S, E, I_S, I_A, H, R, D$ respectively denote the class of an individual according to its current state, namely

Lockdown (L) All individuals that has with null mobility and that remains under isolation

*Corresponding author

Email addresses: adrian.acuna@unison.mx (Gabriel A. Salcedo-Varela), francisco.paquady.mx (Francisco Peñuñuri), dgonzalezsa@conacyt.mx (D. González-Sánchez), saul.diazinfante@unison.mx (Saúl Díaz-Infante)

Preprint submitted to Annual Reviews in Control December 20, 2020

28 **Suceptible** (S) Individual under risk

29 **Exposed** (E) Population fraction that host SARS-CoV-2 but cannot infect

30 **Infected-Symptomatic** (I_S) Population infected fraction with symptoms and
31 reported as confirmed case

32 **Infected-Asymptomatic** (I_A) Infected individual whit transitory or null symp-
33 toms and unreported

34 **Hospitalized** (H) Infected population that requires hospitalization or inten-
35 sive care.

36 **Recover or removed** (R) Population that recovers from infection and devel-
37 ops partial immunity

38 **Death** (D) Population fraction that death by COVID-19

39 To fit data of cumulative reported symptomatic cases, we postulated the counter
40 state Y_{I_S} and made the following hypothesis.

41 **Hypothesis 1.** (H-1) We suppose that at least 30 %of the population is un-
42 der lock-down and that eventually a fraction of this class move to the
43 susceptible compartment at rate δ_L .

44 (H-2) Force infection is defined as the probablity of acquire COVID-19 given
45 the contact with a symptomatic or asympotomatic individual. Thus we
46 normalize under live population N^*

47 (H-3) Susceptible individuals become exposed but not infectious When they are
48 in contact with asymptomatic or symptomatic individuals. Thus β_S, β_A
49 denote probability of infectious given the contact with a symptomatic or
50 asymptomatic infectious individuals.

51 (H-4) After a period of latency of $1/\kappa = 5.1$ days, an exposed individual became
52 infected. Being p the probability of develop symptoms and $(1 - p)$ the
53 probability of became infectious but asymptomatic. Thus $p\kappa E$ denotes
54 the event of become infectious and develop symptoms given that the
55 individual has been exposed

56 (H-5) Asymptomatic individuals not die or get in a Hospital

57 (H-6) A fraction μ_H of symptomatic individuals die by COVID-19 without hos-
58 pitalization

59 ,

Thus we formulate the following Ordinary Differential Equation (ODE)

$$\begin{aligned}
L' &= \theta\mu N^* - \epsilon\lambda L - \delta_L L - \mu L, \\
S' &= (1 - \theta)\mu N^* + \delta_L L + \delta_R R - (\lambda + \mu)S, \\
E' &= \lambda(\epsilon L + S) - (\kappa + \mu)E, \\
I_S' &= p\kappa E - (\gamma_S + \delta_H + \underline{\mu_{I_S}}^{\text{SDIV}} + \mu)I_S, \\
I_A' &= (1 - p)\kappa E - (\gamma_A + \mu)I_A, \\
H' &= \delta_H I_S - (\gamma_H + \mu_H + \mu)H, \\
R' &= \gamma_S I_S + \gamma_A I_A + \gamma_H H - (\delta_R + \mu)R, \\
D' &= \underline{\mu_{I_S}}^{\text{SDIV}} + \mu_H H, \\
\frac{dY_{I_S}}{dt} &= p\kappa E, \\
\lambda &:= \frac{\beta_A I_A + \beta_S I_S}{N^*}, \\
N^*(t) &= L + S + E + I_S + I_A + H + R.
\end{aligned} \tag{1}$$

See Table 1 for notation and references values.




Figure 1: Cummulative new symptomatic and confirmed COVID19 reported cases from Ciudad de Mexico between March, 10, to March 30 of 2020.

We callibrate parameters of our base dynamics in (1) via Multichain Monte-carlo (MCMC). To this end, we assume that the cumulative incidence of new infected symptomatic cases CI_S follows a Poisson distribution with mean

Parameter	Description
μ	Death rate
β_S	Infection rate between susceptible and symptomatic infected
β_A	Infection rate between susceptible and asymptomatic infected
λ_V	Vaccination rate
δ_V^{-1}	Vaccine-induced immunity
ε	Vaccine efficacy
κ^{-1}	Average incubation time
p	New asymptomatic generation proportion
θ	Proportion of individuals under lockdown
γ_S^{-1}	Average time of symptomatic recovery
γ_A^{-1}	Recovery average time of asymptomatic individuals
γ_H^{-1}	Recovery average time by hospitalization
δ_R^{-1}	Natural immunity
δ_H	Infected symptomatic hospitalization rate

Table 1: Parameters definition of model in Equation (1).

$\lambda_t = IC_s(t)$. Further, following [] we postulate priors for p and κ

$$\begin{aligned}
Y_t &\sim \text{Poisson}(\lambda_t), \\
\lambda_t &= \int_0^t p \delta_e E, \\
p &\sim \text{Uniform}(0.3, 0.8), \\
\kappa &\sim \text{Gamma}(10, 50).
\end{aligned} \tag{2}$$

Using the reproductive number definition of VanDenDrishe, we obtain

$$R_0 := \frac{N^*(\beta_S p \kappa + \beta_A \kappa (1 - p))}{(\mu - \kappa)(\gamma_S + \mu_{I_s} + \gamma_A + \mu) N^* \mu}$$

Figure 2 displays data of coumulative confirmed cases of COVID-19 of Mexico city, and the fitt of our model in Equations (1) and (2).

3. Imperfect-preventive Covid-19 vaccination

Preventive vaccines.




Figure 2: Fit of diary new cases of Mexico city during exponential growth.

70 *Efficacy and vaccine-induced immunity.*

71 *Actual vaccine stage development.*

72 *Vaccination reproductive number.*

73 *Vaccination rate λ_V estimate.*

Feasibility regions according to efficacy and vaccination rate.

$$\begin{aligned}
L' &= \theta \mu N^* - (\epsilon \lambda + \delta_L + \mu) L \\
S' &= (1 - \theta) \mu N^* + \delta_L L + \delta_V V + \delta_R R \\
&\quad - (\lambda + \lambda_V + \mu) S \\
E' &= \lambda (\epsilon L + (1 - \epsilon) V + S) - (\kappa + \mu) E \\
I'_S &= p \kappa E - (\delta_H + \gamma_S + \mu_{I_S} + \mu) I_S \\
I'_A &= (1 - p) \kappa E - (\gamma_A + \mu) I_A \\
H' &= \delta_H I_S - (\gamma_H + \mu_H + \mu) H \\
R' &= \gamma_S I_S + \gamma_A I_A + \gamma_H H - (\delta_R + \mu) R \\
D' &= \mu_{I_S} I_S + \mu_H H \\
V' &= \lambda_V S - [(1 - \epsilon) \lambda + \delta_V + \mu] V
\end{aligned}$$

$$\begin{aligned}
\frac{dX_{vac}}{dt} &= (u_V(t) + \lambda_V) [S + E + I_A + R] \\
\frac{dY_{I_S}}{dt} &= p \kappa E \\
\lambda &:= \frac{\beta_A I_A + \beta_S I_S}{N^*}
\end{aligned} \tag{3}$$

$$\begin{aligned}
L(0) &= L_0, \quad S(0) = S_0, \quad E(0) = E_0, \\
I_S(0) &= I_{S_0}, \quad I_A(0) = I_{A_0}, \quad H(0) = H_0, \\
R(0) &= R_0, \quad D(0) = D_0, \\
V(0) &= 0, \quad X_{vac}(0) = 0, \\
X_{vac}(T) &= x_{coverage}, \\
N^*(t) &= L + S + E + I_S + I_A + H + R + V.
\end{aligned}$$

74 4. Parameter calibration

75 *Bayesian estimation.*

76 5. Vaccination reproductive number

77 *R₀ definition.*

78 *No vaccine reproductive number.*

79 *Vaccine reproductive number.*

Parameter	Median	Reference
q_r, ϵ	0.4, 0.3, 0.1	this study
β_S	$q_r \times 8.690\,483 \times 10^{-1}$	this study
β_A	$q_r \times 7.738\,431 \times 10^{-1}$	this study
κ	0.196\,078\,43	*
p	0.1213	*
θ	0.2,	this study
δ_L	0.04	postulated
δ_H	0.2	*
δ_V	0.002\,739\,726\,027\,397\,260\,3	$\delta_V^{-1} = 2$ years CanSinoBIO
δ_R	0.005\,555\,56	$\delta_R^{-1} \approx 180$ days
μ	$3.913\,894 \times 10^{-5}$	**
μ_{I_S}	0.0	
μ_H	0.016\,32	[FENG]
γ_S	0.092\,506\,94	*
γ_A	0.167\,504\,19	*
γ_H	$5.079\,869 \times 10^{-1}$	*
λ_V	0.000\,611\,35	
ε	0.7, 0.80, 0.9, 0.95	[PRESS RELESASES]
N	26\,446\,435	**
L_0	0.266\,260\,097\,021\,127\,96	
S_0	0.463\,606\,046\,009\,872	
E_0	0.000\,670\,33	*
I_{S_0}	9.283×10^{-5}	** *
I_{A_0}	0.001\,209\,86	*
H_0	$1.341\,579\,69 \times 10^{-4}$	**
R_0	$2.661\,259\,39 \times 10^{-1}$	
D_0	0.001\,900\,74	**
X_{vac}^0	0.0	
V_0	0.0	
$Y_{I_S}^0$	0.122\,581\,64	
B	0.000\,359\,216\,658\,124\,242\,5	9500 beds/ N
a_{I_S}	0.002\,012\,775\,543\,825\,648\,6	DALY def
a_H	0.001\,411\,888\,738\,103\,725, or $a_H(x) := 0.001\,411\,888\,738\,103\,725 \log(\frac{1}{B-\kappa I_S})$	DALY def [Jo 2020]
a_D	7.25	DALY def

80 *Efficacy, coverage and vaccination rate.*

81 Here Gabriel's R not calculations.^{SDIV}

$$\frac{\kappa (\epsilon \mu p \theta \beta_A - \epsilon \mu p \theta \beta_s + \epsilon p \theta \beta_A \delta_H)}{\gamma_A \mu I_S \gamma_A \mu I_S} \quad (4)$$

[SDIV 1]
Here contour
plots figure
as function
of efficacy
and vaccina-
tion rate




Figure 3: R not contour plot as function of efficacy and vaccination rate

82

83 6. Optimal controlled version

84 *Controlled Model.* Now we model vaccination, treatment and lockdown as a
85 optimal control problem. According to dynamics in Equation (1), we modu-
86 late the vaccination rate with a time-dependent control signal $u_V(t)$. We add
87 compartment X_{vac} to count all the vaccine applications of susceptible, exposed,
88 asymptomatic and recovered individuals. This process is modeled by

$$X'(t) = (\lambda_V + u_V(t))(S + E + I_A + R) \quad (5)$$

and describes the number of applied vaccines at time t . Consider

$$x(t) := (L, S, E, I_S, I_A, H, R, D, V, X_{vac})^\top(t)$$

89 and control signal $u_v(\cdot)$. We quantify the cost and reward of a vaccine strategy
90 policy via the penalization functional

$$J(u_L, u_V) := \int_0^T a_S I_S + a_d D + \frac{1}{2} (c_L u_L^2 + c_V u_V^2) ds. \quad (6)$$

91 In other words, we assume in functional J that pandemic cost is proportional to
92 the symptomatic and death reported cases and that a vaccination policy implies
93 quadratic consumption of resources.

Further, since we aim to simulate vaccination policies at different coverage scenarios, we impose the vaccination counter state's final time condition $X(T)$

$$\begin{aligned} x(T) &= (\cdot, \cdot, \cdot, \cdot, X_{vac}(T))^\top, \in \Omega \\ X_{vac}(T) &= x_{coverage}, \\ x_{coverage} &\in \{\text{Low}(0.2), \text{Mid}(0.5), \text{High}(0.8)\}. \end{aligned} \tag{7}$$

Thus, given the time horizon T , we impose that the last fraction of vaccinated populations corresponds to 20%, 50% or 80%, and the rest of final states as free. We also impose the path constraint

$$\Phi(x, t) := \kappa I_S(t) \leq B, \quad \forall t \in [0, T], \tag{8}$$

to ensure that healthcare services will not be overloaded. Here κ denotes hospitalization rate, and B is the load capacity of a health system.

Given a fixed time horizon and vaccine efficiency, we estimate the constant vaccination rate as the solution of

$$x_{coverage} = 1 - \exp(-\lambda_V T). \tag{9}$$

That is, λ_v denotes the constant rate to cover a fraction $x_{coverage}$ in time horizon T . Thus, according to this vaccination rate, we postulate a policy u_v that modulates vaccination rate according to λ_V as a baseline. That is, optimal vaccination amplifies or attenuates the estimated baseline λ_V in a interval $[\lambda_v^{\min}, \lambda_v^{\max}]$ to optimize functional $J(\cdot)$ —minimizing symptomatic, death reported cases and optimizing resources.

Our objective is minimize the cost functional (6)—over an appropriated functional space—subject to the dynamics in equations (1) and (5), boundary conditions, and the path constrain in (8). That is, we search for vaccination policies

112 $u_V(\cdot)$, which solve the following optimal control problem (OCP).

$$\begin{aligned}
\min_{u \in \mathcal{U}} J(u) &:= \int_0^T [(a_D \mu_s + a_H \delta_H) I_S(r) + a_{I_S} p \kappa E(r)] dr \\
\text{s. t.} \\
L' &= \theta \mu N^* - \epsilon \lambda L - u_L(t) L - \mu L \\
S' &= (1 - \theta) \mu N^* + u_L(t) L + \delta_v V + \delta_R R \\
&\quad - [\lambda + (\lambda_V + u_V(t)) + \mu] S \\
E' &= \lambda(\epsilon L + (1 - \epsilon) V + S) - (\kappa + \mu) E \\
I_S' &= p \kappa E - (\gamma_S + \mu_{I_S} + \delta_H + \mu) I_S \\
I_A' &= (1 - p) \kappa E - (\gamma_A + \mu) I_A \\
H' &= \delta_H I_S - (\gamma_H + \mu_H + \mu) H \\
R' &= \gamma_S I_S + \gamma_A I_A + \gamma_H H - (\delta_R + \mu) R \\
D' &= \mu_{I_S} I_S + \mu_H H \\
V' &= (\lambda_V + u_V(t)) S - [(1 - \epsilon) \lambda + \delta_V + \mu] V
\end{aligned} \tag{10}$$

$$\begin{aligned}
\frac{dX_{vac}}{dt} &= (u_V(t) + \lambda_V) [L + S + E + I_A + R] \\
\frac{dY_{I_S}}{dt} &= p \kappa E \\
\lambda &:= \frac{\beta_A I_A + \beta_S I_S}{N^*}
\end{aligned}$$

$$\begin{aligned}
L(0) &= L_0, \quad S(0) = S_0, \quad E(0) = E_0, \quad I_S(0) = I_{S_0}, \\
I_A(0) &= I_{A_0}, \quad H(0) = H_0, \quad R(0) = R_0, \quad D(0) = D_0, \\
V(0) &= 0, \quad X_{vac}(0) = 0, \quad u_V(\cdot) \in [u_{\min}, u^{\max}], \\
X_{vac}(T) &= x_{coverage}, \quad \kappa I_S(t) \leq B, \quad \forall t \in [0, T], \\
N^*(t) &= L + S + E + I_S + I_A + H + R + V
\end{aligned}$$

113 7. Numerical Results

114 Changes (compact)

115 **Author: anonymous**

116 No changes.

117 **Author: SDIV**

118 Added 1

119 Deleted 2

120 Commented 1

121

122 **Appendix A. Appendix**

123 Consider the following cost functional that we want to minimize

$$\int_0^T C(t, X(t), u(t)) dt \quad (\text{A.1})$$

124 subject to the dynamics

$$\dot{X}(t) = f(t, X(t), u(t)), \quad 0 \leq t \leq T, \quad (\text{A.2})$$

125 and the initial state $X(0) = x_0$. Let $t_0 < t_1 < \dots < t_n$, with $t_0 = 0$ and $t_n = T$,
 126 be a partition of the interval $[0, T]$. We consider *piecewise constant controls* \tilde{u}
 127 of the form

$$\tilde{u}(t) = a_j \quad t_j \leq t < t_{j+1} \quad (\text{A.3})$$

128 for $j = 0, \dots, n-1$.

129 ASSUMPTION 1.

130 ASSUMPTION 2.

131 By Assumption 1, the system

$$\dot{X}(t) = f(t, X(t), a_0), \quad X(0) = x_0, \quad 0 \leq t \leq t_1,$$

132 has a unique solution $\tilde{X}_0(t; x_0, a_0)$ which is continuous in (x_0, a_0) . Next, put
 133 $x_1 := \tilde{X}_0(t_1; x_0, a_0)$ and consider the system

$$\dot{X}(t) = f(t, X(t), a_1), \quad X(t_1) = x_1, \quad t_1 \leq t \leq t_2,$$

which, again by Assumption 1, has a unique solution $\tilde{X}_1(t; x_1, a_1)$ continuous in
 (x_1, a_1) . By following this procedure, we end up having a recursive solution

$$\begin{aligned} & \tilde{X}_{n-1}(t; x_{n-1}, a_{n-1}), \\ & x_{n-1} := \tilde{X}_{n-2}(t_{n-1}; x_{n-2}, a_{n-1}), \quad t_{n-1} \leq t \leq T. \end{aligned}$$

134 Thus, for a control \tilde{u} of the form (A.3) and the corresponding solution path \tilde{X} ,
 135 we have

$$\int_0^T C(t, \tilde{X}(t), \tilde{u}(t)) dt = \sum_{j=0}^{n-1} \int_{t_j}^{t_{j+1}} C(t, \tilde{X}_j(t), a_j) dt.$$

136 Notice that each \tilde{X}_j is a continuous function of (a_0, \dots, a_j) and x_0 . Therefore,
 137 by Assumption 2, the mapping

$$(a_0, \dots, a_{n-1}) \mapsto \sum_{j=0}^{n-1} \int_{t_j}^{t_{j+1}} C(t, \tilde{X}_j(t), a_j) dt$$

138 is continuous.

139 References

- 140 [1] Yinon M Bar-on, Ron Sender, Avi I Flamholz, Rob Phillips, and Ron Milo.
141 A quantitative compendium of COVID-19 epidemiology. pages 1–51.
- 142 [2] Bruno Buonomo. Effects of information-dependent vaccination behavior on
143 coronavirus outbreak: insights from a SIRI model. *Ricerche di Matematica*,
144 2020.
- 145 [3] Anastasia Chatzilena, Edwin van Leeuwen, Oliver Ratmann, Marc
146 Baguelin, and Nikolaos Demiris. Contemporary statistical inference for in-
147 fectious disease models using Stan. *Epidemics*, 29(February):100367, 2019.
- 148 [4] Nicholas G Davies, Adam J Kucharski, Rosalind M Eggo, Amy Gimma,
149 CMMID COVID-19 Working Group, and W. John Edmunds. The effect
150 of non-pharmaceutical interventions on COVID-19 cases, deaths and de-
151 mand for hospital services in the UK: a modelling study. *medRxiv*, page
152 2020.04.01.20049908, 2020.
- 153 [5] Laura Di Domenico, Giulia Pullano, Chiara E Sabbatini, Pierre-Yves
154 Boëlle, and Vittoria Colizza. Currently under screening at medRxiv Ex-
155 pected impact of lockdown in Île-de-France and possible exit strategies.
156 2020.
- 157 [6] Wandi Ding and Suzanne Lenhart. Introduction to optimal control for
158 discrete time models with an application to disease modeling. In *Modeling*
159 *paradigms and analysis of disease transmission models*, volume 75, pages
160 109–119. Amer. Math. Soc., Providence, RI, oct 2010.
- 161 [7] Ramses Djidjou-Demasse, Yannis Michalakis, Marc Choisy, Micea T. So-
162 fonea, and Samuel Alizon. Optimal COVID-19 epidemic control until vac-
163 cine deployment. *medRxiv*, page 2020.04.02.20049189, 2020.
- 164 [8] Neil M Ferguson, Daniel Laydon, Gemma Nedjati-gilani, Natsuko Imai,
165 Kylie Ainslie, Marc Baguelin, Sangeeta Bhatia, Adhiratha Boonyasiri,
166 Zulma Cucunubá, Gina Cuomo-dannenburg, Amy Dighe, Han Fu, Katy
167 Gaythorpe, Will Green, Arran Hamlet, Wes Hinsley, Lucy C Okell, Sabine
168 Van, Hayley Thompson, Robert Verity, Erik Volz, Haowei Wang, Yuan-
169 rong Wang, Patrick G T Walker, Caroline Walters, Peter Winskill, Charles
170 Whittaker, Christl A Donnelly, Steven Riley, and Azra C Ghani. Impact of
171 non-pharmaceutical interventions (NPIs) to reduce COVID- 19 mortality
172 and healthcare demand. (March), 2020.
- 173 [9] M. H. A. Biswas, L. T. Paiva, and MdR de Pinho. A SEIR model for
174 control of infectious diseases with constraints. *Mathematical Biosciences*
175 *and Engineering*, 11(4):761–784, 2014.
- 176 [10] IHME COVID-19 health service utilization forecasting Team and Christo-
177 pher JL Murray. Forecasting COVID-19 impact on hospital bed-days,

- 178 ICU-days, ventilator-days and deaths by US state in the next 4 months.
179 *medRxiv*, page 2020.03.27.20043752, mar 2020.
- 180 [11] Gang Huang, Yasuhiro Takeuchi, Wanbiao Ma, and Daijun Wei. Global
181 Stability for Delay SIR and SEIR Epidemic Models with Nonlinear Inci-
182 dence Rate. *Bulletin of Mathematical Biology*, 72(5):1192–1207, 2010.
- 183 [12] Jiwei Jia, Jian Ding, Siyu Liu, Guidong Liao, Jingzhi Li, B. E.N. Duan,
184 Guoqing Wang, and R. A.N. Zhang. Modeling the control of COVID-
185 19: Impact of policy interventions and meteorological factors. *Electronic*
186 *Journal of Differential Equations*, 2020:1–21, 2020.
- 187 [13] Abdelilah Kaddar, Abdelhadi Abta, and Hamad Talibi Alaoui. A com-
188 parison of delayed SIR and SEIR epidemic models. *Nonlinear Analysis:*
189 *Modelling and Control*, 16(2):181–190, apr 2011.
- 190 [14] Qian Li, Biao Tang, Nicola Luigi Bragazzi, Yanni Xiao, and Jianhong Wu.
191 Modeling the impact of mass influenza vaccination and public health inter-
192 ventions on COVID-19 epidemics with limited detection capability. *Math-*
193 *ematical Biosciences*, 325(May):108378, jul 2020.
- 194 [15] Gustavo Barbosa Libotte, Fran Sérgio Lobato, Gustavo Mendes Platt, and
195 Antônio José da Silva Neto. Determination of an Optimal Control Strategy
196 for Vaccine Administration in COVID-19 Pandemic Treatment. (November
197 2019), apr 2020.
- 198 [16] Manotosh Mandal, Soovoojeet Jana, Swapan Kumar Nandi, Anupam
199 Khatua, Sayani Adak, and T.K. Kar. A model based study on the dy-
200 namics of COVID-19: Prediction and control. *Chaos, Solitons & Fractals*,
201 136:109889, jul 2020.
- 202 [17] Van Kinh Nguyen and Esteban A. Hernandez-Vargas. Parameter Estima-
203 tion in Mathematical Models of Viral Infections Using R. volume 1836,
204 pages 531–549. 2018.
- 205 [18] Jorge Nocedal, Andreas Wächter, and Richard A. Waltz. Adaptive Bar-
206 rier Update Strategies for Nonlinear Interior Methods. *SIAM Journal on*
207 *Optimization*, 19(4):1674–1693, jan 2009.
- 208 [19] Xinwei Wang, Haijun Peng, Boyang Shi, Dianheng Jiang, Sheng Zhang,
209 and Biaosong Chen. Optimal vaccination strategy of a constrained time-
210 varying SEIR epidemic model. *Communications in Nonlinear Science and*
211 *Numerical Simulation*, 67:37–48, 2019.