

An stochastic HIV Epidemic Model Driven with fBM Death Uncertainty

Saúl Díaz-Infante ^{a,*}, Francisco Delgado-Vences ^b, Arely ¹

^a CONACYT-Universidad de Sonora, Departamento de Matemáticas, Boulevard Luis Encinas y Rosales S/N, Col. Centro, Hermosillo, Sonora, México.

^b CONACYT-UNAM, Instituto de Matemáticas, Sede Oaxaca, México.

Abstract

Keywords: fBM, stochastic differential equation.

1. Introduction

2. Model formulation

Here we consider the model reported in [5] as block base to design our stochastic model. The deterministic dynamics obeys the following equation:

$$\begin{aligned}\frac{dS}{dt} &= \mu K - c\beta(I + bJ)S - \mu S, \\ \frac{dI}{dt} &= c\beta(I + bJ)S - (\mu + k_1)I + \alpha J, \\ \frac{dJ}{dt} &= k_1 I - (\mu + k_2 + \alpha)J, \\ \frac{dA}{dt} &= k_2 J - (\mu + \mu_d)A.\end{aligned}\tag{1}$$

Our aim is to quantify uncertainty by a fractional Ornstein–Uhlenbeck process. To this end, we fit a mean reverting fractional process to the disease-related AIDS rate μ_d . That is, we estimate the reverting parameters $\gamma_d, \bar{\mu}_d, \sigma_d$ and the Hurst parameter H of the fractional stochastic process

$$\frac{d\mu_d(t)}{dt} = \gamma_d (\bar{\mu}_d - \mu_d(t)) dt + \sigma_d dB_t^H.\tag{2}$$

Here B_t^H denotes a fractional Brownian motion with Hurst parameter $H \in [\frac{1}{2}, 1)$.

3. Hurst parameter estimation

4. Existence of a unique positive invariant solution

5. Numerical results

6. Discussion

- [1] B. Boufoussi, S. Hajji, Neutral stochastic functional differential equations driven by a fractional Brownian motion in a Hilbert space, Statistics & Probability Letters 82 (2012) 1549–1558. URL: <http://dx.doi.org/10.1016/j.spl.2012.04.013><http://linkinghub.elsevier.com/retrieve/pii/S0167715212001563>. doi:10.1016/j.spl.2012.04.013.

*Corresponding author

Email addresses: saul.diazinfante@unison.mx (Saúl Díaz-Infante), delgado@im.unam.mx (Francisco Delgado-Vences), delgado@im.unam.mx (Arely)

- [2] D. T. Nguyen, Mackey–Glass equation driven by fractional Brownian motion, *Physica A: Statistical Mechanics and its Applications* 391 (2012) 5465–5472. URL: <http://dx.doi.org/10.1016/j.physa.2012.06.013><http://linkinghub.elsevier.com/retrieve/pii/S0378437112005109>. doi:10.1016/j.physa.2012.06.013.
- [3] Y. Ma, Q. Zhang, M. Ye, Mean-square dissipativity of numerical methods for a class of resource-competition models with fractional Brownian motion, *Systems Science & Control Engineering* 5 (2017) 268–277. URL: <https://doi.org/10.1080/21642583.2017.1333469><https://www.tandfonline.com/doi/full/10.1080/21642583.2017.1333469>. doi:10.1080/21642583.2017.1333469.
- [4] Y. Mishura, G. Shevchenko, The rate of convergence for Euler approximations of solutions of stochastic differential equations driven by fractional Brownian motion, *Stochastics* 80 (2008) 489–511. doi:10.1080/17442500802024892.
- [5] L. Cai, X. Li, M. Ghosh, B. Guo, Stability analysis of an HIV/AIDS epidemic model with treatment, *Journal of Computational and Applied Mathematics* 229 (2009) 313–323. URL: <https://www.sciencedirect.com/science/article/pii/S0377042708005815><http://linkinghub.elsevier.com/retrieve/pii/S0377042708005815>. doi:10.1016/j.cam.2008.10.067.
- [6] M. J. Garrido-Atienza, P. E. Kloeden, A. Neuenkirch, Discretization of Stationary Solutions of Stochastic Systems Driven by Fractional Brownian Motion, *Applied Mathematics and Optimization* 60 (2009) 151–172. URL: <http://link.springer.com/10.1007/s00245-008-9062-9>. doi:10.1007/s00245-008-9062-9.
- [7] J. Traebert, E. Traebert, F. Schuelter-Trevisol, J. J. Cortez Escalante, I. J. C. Schneider, The burden of AIDS: a time series analysis of thirty-five years of the epidemic in Brazil, *AIDS Care* 0 (2018) 1–8. URL: <https://doi.org/10.1080/09540121.2018.1456642><https://www.tandfonline.com/doi/full/10.1080/09540121.2018.1456642>. doi:10.1080/09540121.2018.1456642.