

# A stochastic model of mortality rate with memory

## A stochastic model of mortality rate with memory

**Arely Ornelas · Francisco  
Delgado-Vences · Saul Diaz-Infante**

Received: March 12, 2020/ Accepted: date

**Abstract** An accurate estimation of mortality rates is essential to make decisions. For example, the ensures companies, investment projects, among others, projects its operations according to estimations based on these rates. However, the vast number of variables and its intricate relation implies a challenge for the estimation of these rates.

We assume the following hypothesis: the mortality rate has a strong relationship with its owns past. In this line, we propose a stochastic model with long-term memory that describes mortality. Then, using data from Italy, we provide statistical evidence via Hurst parameter estimation that does not reject our hypothesis. Further, we extract a subset of data to evaluate its forecasting performance, and we observe a good estimation.

To the best of our knowledge, our contribution is the first attempt that includes long term memory in the formulation of a model to describe mortality rates. Our results suggest that the hypothesis of an imperfect correlation intensity across generations would be more realistic and that sex is an important variable to consider in next formulations.

---

Francisco Delgado-Vences  
Conacyt-Universidad Nacional Autónoma de México.  
Instituto de Matemáticas, Oaxaca, México  
E-mail: delgado@im.unam.mx  
ORCID:

Arely Ornelas  
Conacyt-Instituto Politecnico Nacional-CICIMAR,  
La Paz, México  
E-mail: arely.ornelas@conacyt.mx  
ORCID:

Saul Diaz-Infante  
Conacyt-Universidad de Sonora  
Departamento de Matemáticas,  
Hermosillo, Sonoran México E-mail: sdinfante@conacyt.mx  
ORCID: 0000-0001-9559-1293

## 1 Introduction

Future planning in the demographic, economic, and actuarial areas is crucial. For instance, proper planning in social programs, government budgets, cost of insurance, and others depends on the forecast. However, constant changes in technology, lifestyle, migration, to name a few, make predicting a demanding task. Mortality impacts directly in cash and therefore need a reliable future projection.

Previous work has only focused on deterministic or stochastic models without memory effects. [Pitacco et al.(2009)Pitacco, Denuit, Haberman and Olivieri] review the first mortality tables and models. [Milevsky and Promislow(2001)] report a linear SDE driven by Brownian motion that describes the mortality hazard rate. In [Giacometti et al.(2011)Giacometti, Ortobelli and Bertocchi], the authors extend the Milevesky model to an SDE with time-dependent diffusion and study a type of autoregressive model for the logarithm of the hazard rate. [Jevtić et al.(2013)Jevtić, Luciano and Vigna] formulate a cohort-based model with imperfect correlation across generations and use data from UK to estimate parameters.

The present paper aims to show statistical evidence that mortality rates follow a stochastic process with long-range dependence (LRD), that is, a stochastic process with memory. According to [Prakasa Rao(2010)], a stochastic process or time series is LRD, if it has persistence behavior—below we give a formal definition.

We base our stochastic formulation in the fractional Brownian Motion (fBM) [Mandelbrot and Van Ness(1968)]. Since fBM is a generalization of the standard Brownian Motion (BM) that still satisfies self-similarity and is LRD, fBM results to be a natural noise model to describe LRD behavior.

The main idea is to extend the stochastic model reported by Milevesky and Promislow to a model with LRD and verify its performance to fitting and forecasting with real data.

We obtain statistical evidence—via Hurst parameter estimation—that Italy mortality rate data is LRD. Our model captures women’s mortality rate dynamics. But, the results suggest that stratification by gender would be a direction for future formulations.

Add relevant use and implications of our results

After this brief introduction, Section 2 reviews the fBM and presents the fractional Ornstein-Uhlenbeck (fOU) process. Section 3 outlines the applied method for parameter estimation. In Section 4, we implement our formulation to fitting data of mortality from Italy. Further, in this section, we also run forecasting to a subset of the data and evaluate its efficiency. Finally, we conclude in Section 6 with relevant implications and perspectives.

## 2 Fractional Brownian motion

We consider the Gaussian process  $\{B_t^H, t \geq 0\}$ , with  $H \in (0, 1)$ , and with zero-mean and covariance function given by

$$R_H(t, s) := \mathbb{E}(B_s^H B_t^H) = \frac{1}{2}(t^{2H} + s^{2H} - |t - s|^{2H}). \quad (1)$$

This stochastic process is called a *fractional Brownian motion* (fBm) and was introduced by Kolmogoroff [Kolmogoroff(1940)] and studied by Mandelbrot and Van Ness in [Mandelbrot and Van Ness(1968)]. The parameter  $H$  is called Hurst index because of the statistical analysis developed by the climatologist Hurst [Hurst(1951)]. The fBm is a generalization of Brownian motion without independent increments, in fact it is a continuous-time Gaussian process.

fBm process has the properties of self-similarity and stationary increments. Its sample-paths are almost nowhere differentiable but almost-all Hölder continuous for any order strictly less than  $H$ . That is, for each trajectory, there exists a finite constant  $C$  such that for every  $\epsilon > 0$

$$\mathbb{E}(|B_t^H - B_s^H|) \leq C|t - s|^{H-\epsilon}.$$

For  $H = \frac{1}{2}$  the covariance of fBM can be written as  $R_{1/2}(t, s) = \min(s, t)$ , thus process  $B_t^{1/2}$ , is equivalent to the standard Brownian motion. However, if  $H \neq \frac{1}{2}$ , then this increments are *not* independent. Let  $X_n = B_n^H - B_{n-1}^H$ ,  $n \geq 1$ . Then  $\{X_n, n \geq 1\}$  is a Gaussian stationary sequence with unit variance and covariance function if

$$\begin{aligned} \rho_H(n) &= \frac{1}{2} \left( (n+1)^{2H} + (n-1)^{2H} - (2n)^{2H} \right) \\ &\approx H(2H-1)n^{2H-2} \rightarrow 0, \quad \text{when } n \rightarrow \infty. \end{aligned}$$

Therefore,

- if  $H > \frac{1}{2}$ , then  $\rho_H(n) > 0$  for  $n$  large enough and  $\sum_{n=1}^{\infty} \rho_H(n) = \infty$ . In this case, we say that process  $X_n$  is persistent with positive correlation and that  $X_n$  has the *long-range dependence* property
- if  $H < \frac{1}{2}$ , then  $\rho_H(n) < 0$  for  $n$  large enough and  $\sum_{n=1}^{\infty} \rho_H(n) < \infty$ . This is an anti-persistent process with negative correlation.

For further information on fBM see [Prakasa Rao(2010), Nualart(2006), Mishura(2008)].

### 2.1 Fractional Ornstein-Uhlenbeck process (fOU)

The fOU is an SDE driven by a fractional Brownian motion. The same model was used in [Milevsky and Promislow(2001)] or [Giacometti et al.(2011)Giacometti, Ortobelli and Bertocchi].

As we mentioned before, the survival probability  $S(t, T)$  of an individual aged  $x$  in the period  $[t, T]$ , is given in the Equation (??) and  $h_x(t)$  is the stochastic force of mortality or hazard rate given by the Equation (??).

We will assume that  $Y_t$  is an stochastic process that satisfies the SDE:

$$dY_t^H = -\lambda Y_t^H dt + \sigma dB_t^H, \quad (2)$$

where  $B_t^H$  is a fBM with Hurst parameter  $1/2 \leq H < 1$ ,  $Y_0 = 0$ , and  $\sigma, \lambda > 0$ . This SDE is the fractional Ornstein-Uhlenbeck process.

There are substantial differences in trying to solve Equation (2) with respect to the method use in [Milevsky and Promislow(2001)] or [Giacometti et al.(2011)Giacometti, Ortobelli and Bertocchi]. We now discuss some of these differences. First we interpret the SDE (2) as

$$Y_t^H = -\lambda \int_0^t Y_s^H ds + \sigma B_t^H. \quad (3)$$

The equation above lacks of a stochastic integral because we consider the case with additive noise. Nevertheless, we can consider the general case with multiplicative noise, in that case we define a stochastic integral with respect to fractional Brownian motion as a pathwise Riemann–Stieltjes integral, see, e.g. [Young(1936)] for the original definition and [Dudley and Norvaiša(2011)] for advanced results.

Coming back to Equation (3), [Cheridito et al.(2003)Cheridito, Kawaguchi and Maejima] introduced the fractional Ornstein-Uhlenbeck process (fOU) and they showed that the process

$$Y_t^H = \sigma \int_0^t e^{-\lambda(t-u)} dB_u^H, \quad (4)$$

is the unique almost surely continuous-path process which solves (3) (see also Theorem 1.24 in [Prakasa Rao(2010)]). The integral in Equation (4) is a pathwise Riemann–Stieltjes integral. The fOU process is neither Markovian nor a semimartingale for  $H \in (1/2, 1)$  (see [Dudley and Norvaiša(2011)]) but remains Gaussian and ergodic.

Moreover, when  $H \in (1/2, 1)$ ,  $Y_t$  even presents the long-range dependence property (see Cheridito [Cheridito et al.(2003)Cheridito, Kawaguchi and Maejima] or [Prakasa Rao(2010)]).

The variance of the fOU process  $Y_t$  is given by the following expression (see[Zeng et al.(2012)Zeng, Chen and Yang]):

$$Var(Y_t) = \sigma^2 2H e^{-2\lambda t} \int_0^t s^{2H-1} e^{2\lambda s} ds. \quad (5)$$

Notice that when  $H = 1/2$  we get

$$Var(Y_t) = \frac{\sigma^2}{2\lambda} (1 - e^{-2\lambda t}), \quad (6)$$

which is the variance of the standard Ornstein-Uhlenbeck process (see for instance [Mikosch(1998)] page 143).

If we consider the constant  $\alpha_1 = T^{-H}$  and the Equation (5), the expression for the variance of  $\alpha_1 Y_t$  is given by

$$\begin{aligned} Var(\alpha_1 Y_t) &= \alpha_1^2 Var(Y_t) = \alpha_1^2 \sigma^2 2H \int_0^t s^{2H-1} e^{-2\lambda(t-s)} ds \\ &\leq \alpha_1^2 \sigma^2 2H \int_0^t s^{2H-1} ds = \alpha_1^2 \sigma^2 2H \frac{s^{2H}}{2H} \Big|_{s=0}^t \\ &= \alpha_1^2 \sigma^2 t^{2H} = \sigma^2 (t/T)^{2H}, \end{aligned} \quad (7)$$

then  $Var(\alpha_1 Y_t) \leq \sigma^2$  since  $0 \leq t \leq T$ , this implies that the variance of  $\alpha_1 Y_t$  is bounded by a constant that does not depend on time. We will use  $\alpha_1$  to control the variance of the process  $Y_t$ .

### 3 Estimation of the parameters

In this section we will describe a methodology to estimate the parameters. We will use some R-libraries in order to estimate the parameters.

We need to estimate  $\alpha_0, \alpha_1$  as well as  $\sigma, \lambda$  for the SDE model that we described in the previous section. Furthermore, The Hurst parameter ( $H$ ) involved in the driven fractional Brownian motion will be also estimated, however, this estimation is highly complicated. To solve this problem we will use the empirical evidence that the Hurst value in the equation (2) is preserved, this means that the value of the Hurst parameters  $H$ , in the equation (2), for the fBm  $B_t^H$  and the one for the fractional Gaussian noise  $Y_t^H$  are the same. Observe that  $\alpha_1 = T^{-H}$  will be calculated using the estimated Hurst parameter.

#### 3.1 Estimation of the parameter $\alpha_0$ .

The model is given by the equation (??). In order to estimate  $\alpha_0$  we will assume that  $\alpha_1 = 1$ . Moreover,  $h_0$  will be fixed to the initial value of the observed rates  $h(t)$  Taking ln we obtain

$$\ln h(t) = \ln h_0 + \alpha_0 t + Y_t. \quad (8)$$

One simple method to estimate the parameter  $\alpha_0$  is by minimizing the sum of the square errors. Let  $S$  be given by

$$S := \sum_{t_{initial}}^{t_{final}} \left( \ln h(t) - \ln h_0 - \widehat{\alpha_0} t \right)^2.$$

Taking derivative of  $S$  with respect to  $\widehat{\alpha}_0$  we get

$$\frac{\partial S}{\partial \widehat{\alpha}_0} = -2 \sum_{t_{initial}}^{t_{final}} \left( \ln h(t) - \ln h_0 - \widehat{\alpha}_0 t \right) t = 0,$$

and from this equation we obtain  $\widehat{\alpha}_0$ :

$$\widehat{\alpha}_0 = \frac{\sum_t t \ln h(t) - \ln h(0) \sum_t t}{\sum_t t^2}. \quad (9)$$

Once we have estimated  $\alpha_0$  we proceed to estimate the Hurst parameter,  $\sigma$  and  $\lambda$ .

### 3.2 Relation between the Hurst parameter and the H-index in the FOU

In this subsection we will discuss the procedure we have used to estimate the parameter  $H$ .

We will use the following empirical fact. Suppose that a fOU process is driving with a fBM with a given Hurst parameter  $H_0$ . Yerlikaya-Okzurt et al ([Yerlikaya-Özkurt et al.(2014)Yerlikaya-Özkurt, Vardar-Acar, Yolcu-Okur and Weber]) have shown a relationship between the Hurst parameter  $H$  of the fractional Brownian motion and the Hurst parameter of the fractional Gaussian noise given by an SDE. In fact, they have found statistical evidence that the fOU should have the same value  $H_0$  that the fBM (see table 1 Yerlikaya-Okzurt et al). Therefore, at least empirically, the value of  $H$  is the same. Then, it is possible to choose the same value of the parameter  $H$  for both processes. A formal proof of this fact, up to our knowledge, is missed.

The subsequent sections are devoted to present several methods to estimate the Hurst parameter for the fBM.

### 3.3 Estimation of the self-similarity index $H$ for the fBM

The last subsection allows us to estimate the parameter  $H$  in one simple way. According to equation (8), the residuals are given by the expression

$$\hat{Y}_t = \ln h(t) - \ln h_0 - \widehat{\alpha}_0 t.$$

$\hat{Y}_t$  is a fractional Gaussian noise, so that we can use it to estimate  $H$ . Afterwards we will use  $\hat{H}$  to approximate the Hurst parameters of the fractional Brownian motion  $B_t^H$ . For this purpose we will review some methods to estimate the parameter  $H$ .

### 3.3.1 *R over S Analysis*

Following [Weron(2002)]. The analysis begins dividing a time series  $\{Z_i\}$  of length  $L$  into  $d$  subseries of length  $n$  and denote it by  $\{Z_{i,m}\}$ ,  $m = 1, \dots, d$ . Then, for each subseries  $\{Z_{i,m}\}$ ,  $m = 1, \dots, d$ :

1. Find the mean  $E_m$  and standard deviation  $S_m$ .
2. Normalize the data  $Z_{i,m}$  by subtracting the sample mean  $X_{i,m} = Z_{i,m} - E_m$  for  $i = 1, \dots, n$ .
3. Create a cumulative time series  $Y_{i,m} = \sum_{j=1}^i X_{j,m}$  for  $i = 1, \dots, n$ .
4. Find the range  $R_m = \max\{Y_{1,m}, \dots, Y_{n,m}\} - \min\{Y_{1,m}, \dots, Y_{n,m}\}$ ;
5. Rescale the range  $R_m/S_m$ .
6. Calculate the mean value of the rescaled range for all subseries of length  $n$

$$(R/S)_n = \frac{1}{d} \sum_{m=1}^d R_m/S_m.$$

It can be shown (see [Weron(2002)]) that the  $R/S$  statistic asymptotically follows the relation:

$$(R/S)_n \sim cn^H,$$

where  $c$  is a constant. Thus, the value of  $H$  can be obtained by running a simple linear regression over a sample of increasing time horizons

$$\log(R/S)_n = \log c + H \log n.$$

Equivalently, we can plot the  $(R/S)_n$  statistics against  $n$  on a double-logarithmic paper. If the returns process is white noise then the plot is roughly a straight line with slope 0.5. If the process is persistent then the slope  $H$  is greater than 0.5; if it is anti-persistent then the slope  $H$  is less than 0.5. The “significance” level of the estimated parameter  $H$  is usually chosen to be one over the square root of sample length, i.e. the standard deviation of a Gaussian white noise.

A major drawback of the  $R/S$  analysis is that no asymptotic distribution theory has been derived for the Hurst parameter  $H$ . The only results known are for the rescaled (but not by standard deviation) range  $R_m$  itself, see [Lo(1991)].

### 3.3.2 *Method of rescaled range analysis $R/S$*

Following [Prakasa Rao(2010)], chapter 9. This method was suggested by Hurst (1951). The series  $\{X_j, 1 \leq j \leq N-2\}$  is divided into  $K$  nonoverlapping blocks such that each block contains  $M$  elements where  $M$  is the integer part of  $N/K$ . Let  $t_i = M(i-1)$ , where  $t_i = M(i-1)$  is the starting point of the  $i$ th block for  $i = 1, \dots, K$ . Define

$$R(t_i, r) = \max[W(t_1, 1), \dots, W(t_i, r)] - \min[W(t_1, 1), \dots, W(t_i, r)],$$

where  $r$  takes values in natural number whenever  $r$  satisfy the inequality  $t_i + r \leq N$ . Moreover,  $W(t_i, k)$  is set as

$$W(t_i, k) = \sum_{j=0}^{k-1} X_{t_i+j} - k \left( \frac{1}{r} \sum_{j=0}^{r-1} X_{t_i+j} \right), \quad k = 1, \dots, r.$$

Note that  $R(t_i, r) \geq 0$  since  $W(t_i, r) = 0$  and the quantity  $R(t_i, r)$  can be computed only when  $t_i + r \leq N$ . Define

$$S^2(t_i, r) = \frac{1}{r} \sum_{j=0}^{r-1} X_{t_i+j}^2 - \left( \frac{1}{r} \sum_{j=0}^{r-1} X_{t_i+j} \right)^2.$$

The ratio  $R(t_i, r)/S(t_i, r)$  is called the rescaled adjusted range. It is computed for a number of values of  $r$  that makes sense according to the definition. Observe that, for each value of  $r$ , we obtain a number of  $R/S$  samples. The number of samples decrease as  $r$  increases. However, the resulting samples are not independent. It is believed that the  $R/S$ -statistic is proportional to  $r^H$  as  $r \rightarrow \infty$  for the fractional Gaussian noise. Assuming this property, it is possible to regress  $\log(R/S)$  against  $\log(r)$  to obtain an estimator for  $H$ .

### 3.3.3 FDWhittle Estimator

Following [Park et al.(2011)] Park, Hernández-Campos, Le, Marron, Park, Pipiras, Smith, Smith, Trovero and Zhu]. The Local Whittle Estimator (LWE) is a semiparametric Hurst parameter estimator based on the periodogram. It assumes that the spectral density  $f(\omega)$  of the process can be approximated by the function

$$f_{c,H}(\omega) = c\omega^{1-2H}, \quad (10)$$

for frequencies  $\omega$  in a neighborhood of the origin,  $c$  is a constant. The periodogram of a time series  $\{X_t, 1 \leq t \leq N\}$  is defined by

$$I_N(\omega) = \frac{1}{2\pi N} \left| \sum_{t=1}^N X_t e^{i\omega t} \right|^2,$$

where  $i = \sqrt{-1}$ . Usually, it is evaluated at the Fourier Frequencies  $\omega_{j,N} = \frac{2\pi j}{N}$ ,  $0 \leq j \leq [N/2]$ . Note that the periodogram is the norm of the Discrete Fourier transform of the time series (see Section 6.1.2 in [Priestley(1981)] for instance).

The LWE of the Hurst parameter,  $\hat{H}_{LWE}(m)$  is implicitly defined by minimizing

$$\sum_{j=1}^m \log f_{c,H}(\omega_j, N) + \frac{I_N(\omega_j, N)}{f_{c,H}(\omega_j, N)},$$

with respect to  $c$  and  $H$ , with  $f_{c,H}$  defined in (10).



### 3.4 Estimation of $\sigma$ and $\lambda$

There are several methods to estimate parameters  $\sigma$  and  $\lambda$ . For instance see [Prakasa Rao(2010)] or the references in [Neuenkirch and Tindel(2014)] or in [Kubilius and Mishura(2012)]. In the following section we will do a brief review of some of these methods.

#### 3.4.1 Estimation $\sigma$ .

Brouste and Iacus [Brouste and Iacus(2013)] proposed some consistent and asymptotically Gaussian estimators for the parameters  $\sigma, \lambda$  and  $H$  of the discretely observed fractional Ornstein-Uhlenbeck process solution expressed in the stochastic differential equation. There is a restriction on the estimation of the drift  $\lambda$ : the results are valid only in the case when  $1/2 < H < 3/4$ .

The key point of this method of estimation is that the Hurst exponent  $H$  and the diffusion coefficient  $\sigma$  can be estimated without estimating  $\lambda$ . We will use this method to estimate the parameters  $\sigma$  and  $\lambda$ . Notice that  $H$  was already estimated.

Let  $\mathbf{a} = (a_0, \dots, a_K)$  be a discrete filter of order  $L \geq 1$  and length  $K + 1$ ,  $K \in \mathbb{N}$  and we require  $L \leq K$ , i.e.

$$\sum_{k=0}^K a_k k^j = 0 \quad \text{for } 0 \leq j \leq L-1 \quad \text{and} \quad \sum_{k=0}^K a_k k^L \neq 0.$$

Let it be normalized

$$\sum_{k=0}^K (-1)^{1-k} a_k = 1.$$

We will also consider a *dilated* filter  $\mathbf{a}^{(2)}$  associated to  $\mathbf{a}$ . For  $0 \leq k \leq K$  we define

$$a_k^{(2)} = \begin{cases} a_{k'}, & \text{if } k = 2k' \\ 0, & \text{otherwise.} \end{cases}$$

Since  $\sum_{k=0}^{2K} a_k^2 k^j = 2^j \sum_{k=0}^K a_k k^j$  then the filter  $\mathbf{a}^{(2)}$  has the same order than  $\mathbf{a}$ .

Let  $Y^T = (Y_t : 0 \leq t \leq T)$  be the sample path of the solution of (4). A discretization of  $Y^T$  is

$$(X_n := Y_{n\Delta_N}, n = 0, \dots, N), \quad N \in \mathbb{N},$$

where  $\Delta_N = T/N$  and  $N$  is the number of observations of  $Y_t$ . We denote by

$$V_{N,\mathbf{a}} := \sum_{i=0}^{N-K} \left( \sum_{k=0}^K a_k X_{i+k} \right)^2,$$

the *generalized quadratic variation* associated to the filter  $\mathbf{a}$  (see for instance [Istas and Lang(1997)]). Then, define the following estimators for  $H$  and  $\sigma$ .

$$\hat{H}_N := \frac{1}{2} \log_2 \left( \frac{V_{N,\mathbf{a}^2}}{V_{N,\mathbf{a}}} \right), \quad (11)$$

$$\hat{\sigma}_N := \left( -2 \frac{V_{N,\mathbf{a}}}{\sum_{k,l} a_k a_l |k-l|^{2\hat{H}_N} \Delta_N^{2\hat{H}_N}} \right)^{1/2}. \quad (12)$$

Brouste and Iacus (see Th. 1 in [Brouste and Iacus(2013)]) have shown the next result.

**Theorem 1** *Let  $\mathbf{a}$  be a filter of order  $L \geq 2$ . Then, both estimators  $\hat{H}_N$  and  $\hat{\sigma}_N$  are strongly consistent, i.e.*

$$(\hat{H}_N, \hat{\sigma}_N) \xrightarrow{\text{a.s.}} (H, \sigma) \quad \text{as } N \rightarrow +\infty.$$

Moreover, we have asymptotic normality property: for all  $H \in (0, 1)$ ,

$$\begin{aligned} \sqrt{N}(\hat{H}_N - H) &\xrightarrow{\mathcal{L}} N(0, \Gamma_1(\mathbf{a}, \sigma, H)) \quad \text{as } N \rightarrow +\infty, \\ \frac{\sqrt{N}}{\log N}(\hat{\sigma}_N - \sigma) &\xrightarrow{\mathcal{L}} N(0, \Gamma_2(\mathbf{a}, \sigma, H)) \quad \text{as } N \rightarrow +\infty, \end{aligned}$$

where  $\Gamma_1$  and  $\Gamma_2$  are symmetric positive definite matrices depending on  $\sigma, H$  and the filter  $\mathbf{a}$ .  $\square$

With this result is possible to obtain an estimator for the parameters  $\sigma$ .

We will use the following two filters:

- Classical filter. Let  $K > 0$  and define

$$a_k := \frac{(-1)^{1-k}}{2^k} \binom{K}{k} = \frac{(-1)^{1-k}}{2^k} \frac{K!}{k!(K-k)!} \quad \text{for } 0 \leq k \leq K.$$

- Daubechies filters (see [Daubechies(1992)] for the original definition). We set  $K = 4$  and define the Daubechies filter by

$$\frac{1}{\sqrt{2}}(0.48296291314453, -0.8365163037378, 0.22414386804201, 0.12940952255126).$$

### 3.4.2 Estimation of the drift parameter $\lambda$ when both $H$ and $\sigma$ are known

Hu and Nualart, [Hu and Nualart(2010)] have shown that

$$\lim_{t \rightarrow \infty} \text{Var}(Y_t) = \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t Y_t^2 dt = \frac{\sigma^2 \Gamma(2H+1)}{2\lambda^{2H}} := \mu_2.$$

This equation gives a  $\lambda$  estimator, namely

$$\hat{\lambda}_N = \left( \frac{2\hat{\mu}_{2,N}}{\hat{\sigma}_N^2 \Gamma(2\hat{H}_N + 1)} \right)^{-\frac{1}{2\hat{H}_N}}, \quad (13)$$

where  $\hat{\mu}_{2,N}$  is the empirical moment of order 2, i.e

$$\hat{\mu}_{2,N} = \frac{1}{N} \sum_{n=1}^N X_N^2.$$

Set  $T_N = N\Delta_N$ . We have the next result.

**Theorem 2** *Let  $H \in (\frac{1}{2}, \frac{3}{4})$  and a mesh satisfying the condition  $N\Delta_N^p \rightarrow 0$ ,  $p > 1$ , and  $\Delta_N(\log N)^2 \rightarrow 0$  as  $N \rightarrow +\infty$ . Then, as  $N \rightarrow +\infty$ ,*

$$\hat{\lambda}_N \xrightarrow{\text{a.s.}} \lambda,$$

and

$$\sqrt{T_N}(\hat{\lambda}_N - \lambda) \xrightarrow{\mathcal{L}} N(0, \Gamma_3(\sigma, H)),$$

where  $\Gamma_3(\sigma, H) = \lambda \left( \frac{\sigma H}{2H} \right)^2$  and

$$\sigma_H^2 = (4H + 1) \left( 1 + \frac{\Gamma(1 - 4H)\Gamma(4H - 1)}{\Gamma(2 - 2H)\Gamma(2H)} \right).$$

□

For the proof see Theorem 2 in [Brouste and Iacus(2013)].

#### 4 Results with the fractional Ornstein-Uhlenbeck model

In this section we present the estimated mortality rates with the use of the model described in section 1. We obtain the data from the website of Human Mortality Database for the Italian population between 1950 to 2004.

In first place, we present the estimation of the  $H$  parameter. In second place, we present the results on simulated mortality rates using equations (12)-(13) to estimate the parameters  $\sigma, \lambda$ . The parameter  $\alpha_0$  has been fixed with the use of equation (9).

We run 10000 simulations of the mortality rates from ages 0 to age 90. To do that, we have simulate a fractional Brownian motion  $B_t^H$  and using equation (??) we have estimated the mortality rate. To run the fBm simulations we have used the function **fbm** which is includes in the R library **somebm**. We also include a 95.5% confidence interval.

We present the results for women and men in sections 4.2 and 4.3, respectively.


Estimations and predictions were performed using R Ver. 3.2.3 (R Core Team, 2015), and specialized packages Fractal (Time Series Modeling and Analysis Version 2.0-1, 2016), Pracma (Practical Numerical Math Functions 2.0.7, 2017) and somebm (some Brownian motions simulation function Version 0.1, 2016).

#### 4.1 Hurst estimation

For the Hurst parameter estimate we have used three R routines: FDWhittle, RoverS and hurstexp. The two first routines are from the fractal library, while the latter is from the pracma library.

The former routine estimate the Hurst parameter by Whittle's method as was described in the subsection 3.3. RoverS routine estimate  $\hat{H}$  by rescaled range (R/S) method. The hurstexp routine estimate  $\hat{H}$  using R/S analysis.

Finally, figures 1 and 2 show the estimated Hurst parameter for women and men separately.




**Fig. 1** Estimated Hurst parameter using R-routines.

With the rescaled range  $R/S$  and  $R/S$  methods we obtain a consistent estimator for the Hurst parameter in the sense that they do not present dramatic changes through the time. Moreover, the  $H$  estimated with these two methods take values in the interval  $(0.57, 0.80)$  approximately. This tells us that the data has the long memory property as was mentioned in section 2. Same results are obtained for the men and women.

Notice that the estimated parameters for  $H$  using Whittle method have high variation through the time, in opposition to those obtained with the other two methods so that the estimated Hurst parameters using this method do not perform well in the simulations.

The high variation on the Hurst estimated values could be explained because Whittle method uses the periodogram to estimate  $H$  while the other two methods uses the raw data. The two approaches using Whittle likelihood and raw data are very different and hence they give very different estimates of  $H$ . Since rescaled range  $R/S$  and  $R/S$  methods have estimated very similar  $H$ , we decide to use the Hurst coefficients obtained with the method of  $R/S$  to perform the mortality rates simulations.



**Fig. 2** Estimated Hurst parameter using R-routines.

change to  
eps format

#### 4.2 Results for women

We present the results for 10000 path simulations of the mortality rates for ages: 0, 5, 25, 50, 60, 70, 80, 90. We graph the historical mortality rate, the mean of all simulations and the 95% confidence interval. See figures 3 and 4.

In general, for all ages, the model is well fitted, in particular, after the 80's. Nevertheless, there are some time periods where the model is not so good as we want to. For instance, the model underestimates the mortality rate for women age 0 during the period of 50s to 80s and for women age 50, 60, 80 and 90 during 60s to 80s approximately. Moreover, it also overestimates the mortality rate for women age 25 during the period of 50s to 80s.


For older ages (see figure 4) we observe that for ages 60 and 70 the estimation is well fitted trough the years. We notice that predicted rates are not so far away and that the historical rates are inside the confidence interval. For the very oldest ages the estimation is not so good as for earlier ages.

The main difference is in 50's when the absolute number of living persons arriving to those ages were small so that the variability of the estimates is larger.

All these suggest that a better model could include a short and a long-term memory process, so that the model could help us to control the short-term variations in a better way.

### 4.3 Results for men


As in the case for women, we present results for 10000 simulations of the mortality rates for ages: 0, 5, 25, 50, 60, 70, 80, 90. We graph the historical rate mortality, the mean of all simulations and the 95.5% confidence interval. See figures 5 and 6.



**Fig. 5** Simulations for the rate mortality with the fOU model: ages 0, 5, 25, 50 and  $N = 10000$ . ····· IC at 95% - - - simulation mean — Data mortality rates

As in the case for women, the proposed model for men is well fitted. We observe an increase in the rates mortality at ages between 25 to 35, this caused a overestimation in the first 35 years and latter a underestimation of the mortality rates. As was mentioned before, if we include in the model a short-term process, we believe the model could be better fitted. The main example of a short-term process to try is a  $AR(p)$  with  $p \leq 2$  or 3.





**Fig. 6** Simulations for the rate mortality with the fOU model, ages 60, 70, 80, 90 and  $N = 10000$ . ···· IC at 95% - - - simulation mean — Data mortality rates


For older ages, we observed that the data is irregular, so it is necessary to use a more complex model to fit this data.

## 5 Forecast


When we use our model to forecast and compare with the real data between the years 2005 to 2014. In general, from our results we observe that the forecast for women are good for almost all ages we tested. For men the variability of the

results is strong and the behavior of the forecast is in general not good as those for women; for instance for ages smaller than 10 the results are quite similar than for women, even for ages between 10 and 45 it is possible to consider the results just good. However, for ages greater than 50, the results are not good, in fact, for older ages the results are bad: the model overestimate the mortality rates.


We present the results in the figures 7 for women at ages 0, 25, 50, 90. For men we present more ages to illustrate that the results are bad as the ages increases. see figures 5 and 9 at ages 0, 25, 50, 90.



**Fig. 7** Forecast for the rate mortality with the fOU model. Women at ages 0, 25, 50, 90.  
⋯ IC at 95%    - - simulation mean    — Data mortality rates.



**Fig. 8** Forecast for the rate mortality with the fOU model. Men at ages 0, 10, 25, 40.  
 ..... IC at 95%    - - - simulation mean    / Data mortality rates.



**Fig. 9** Forecast for the rate mortality with the fOU model. Men at ages 50, 60, 70, 90.  
 ..... IC at 95%    - - - simulation mean    — Data mortality rates.

## 6 Conclusions

We have applied our proposed model to the Italian mortality rates with a geometric-type fractional Ornstein-Uhlenbeck process. Our main hypothesis was that, for a fixed age, the mortality rates changes through the time slowly, so that a stochastic differential equations that captures the long-range dependence could be a good model. We have used a stochastic differential equation

with a fractional Brownian motion as a driven noise with  $H \in (0.5, 1)$  in order to satisfy the long-range dependence property. With the data we have fixed the Hurst coefficient and we have confirmed our hypothesis since we have found that the estimated Hurst is in  $(0.58, 0.8)$ .

Notice that we have consider a more general model that the one used in [Giacometti et al.(2011)Giacometti, Ortobelli and Bertocchi]. This is because we have included the possibility that the Hurst parameter could be equal to  $1/2$ , which is the case when the fractional Brownian motion becomes a standard Brownian motion. Therefore, when  $H = 1/2$  we recover the Giacometti, Ortobelli and Bertocchi model.

The model is, specially for women, well behaved. For men at some ages we found some shortcomings that suggest the use of more terms in order to improve the model. The long-range dependence model proposed in this paper is good enough to reproduce the mortality rates. If we add some extra terms to make it more flexible to reproduce the cases where the mortality rates have more variations then it will generate a more accurate model. We are starting to work on this extension of the model. Moreover, a multiplicative noise model will be the subject of a future research.




## References

- Brouste and Iacus(2013). Brouste, A. and Iacus, S. M. (2013) Parameter estimation for the discretely observed fractional Ornstein-Uhlenbeck process and the Yuima R package. *Comput. Statist.*, **28**, 1529–1547. URL: <https://doi.org/10.1007/s00180-012-0365-6>.
- Cheridito et al.(2003)Cheridito, Kawaguchi and Maejima. Cheridito, P., Kawaguchi, H. and Maejima, M. (2003) Fractional Ornstein-Uhlenbeck processes. *Electron. J. Probab.*, **8**, no. 3, 14. URL: <https://doi.org/10.1214/EJP.v8-125>.
- Daubechies(1992). Daubechies, I. (1992) *Ten lectures on wavelets*, vol. 61 of *CBMS-NSF Regional Conference Series in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA. URL: <https://doi.org/10.1137/1.9781611970104>.
- Dudley and Norvaiša(2011). Dudley, R. M. and Norvaiša, R. (2011) *Concrete functional calculus*. Springer Monographs in Mathematics. Springer, New York. URL: <https://doi.org/10.1007/978-1-4419-6950-7>.
- Giacometti et al.(2011)Giacometti, Ortobelli and Bertocchi. Giacometti, R., Ortobelli, S. and Bertocchi, M. (2011) A stochastic model for mortality rate on Italian data. *J. Optim. Theory Appl.*, **149**, 216–228. URL: <https://doi.org/10.1007/s10957-010-9771-5>.
- Hu and Nualart(2010). Hu, Y. and Nualart, D. (2010) Parameter estimation for fractional Ornstein-Uhlenbeck processes. *Statist. Probab. Lett.*, **80**, 1030–1038. URL: <https://doi.org/10.1016/j.spl.2010.02.018>.
- Hurst(1951). Hurst, H. E. (1951) Long-term storage capacity of reservoirs. *Trans. Amer. Soc. Civil Eng.*, **116**, 770–799.
- Istas and Lang(1997). Istas, J. and Lang, G. (1997) Quadratic variations and estimation of the local Hölder index of a Gaussian process. *Ann. Inst. H. Poincaré Probab. Statist.*, **33**, 407–436. URL: [https://doi.org/10.1016/S0246-0203\(97\)80099-4](https://doi.org/10.1016/S0246-0203(97)80099-4).


- Jevtić et al.(2013) Jevtić, Luciano and Vigna. Jevtić, P., Luciano, E. and Vigna, E. (2013) Mortality surface by means of continuous time cohort models. *Insurance Math. Econom.*, **53**, 122–133. URL: <https://doi.org/10.1016/j.insmatheco.2013.04.005>.
- Kolmogoroff(1940). Kolmogoroff, A. N. (1940) Wienerische Spiralen und einige andere interessante Kurven im Hilbertschen Raum. *C. R. (Doklady) Acad. Sci. URSS (N.S.)*, **26**, 115–118.
- Kubilius and Mishura(2012). Kubilius, K. and Mishura, Y. (2012) The rate of convergence of Hurst index estimate for the stochastic differential equation. *Stochastic Process. Appl.*, **122**, 3718–3739. URL: <https://doi.org/10.1016/j.spa.2012.06.011>.
- Lo(1991). Lo, A. W. (1991) Long-term memory in stock market prices. *Econometrica*, **59**, 1279–1313.
- Mandelbrot and Van Ness(1968). Mandelbrot, B. B. and Van Ness, J. W. (1968) Fractional Brownian motions, fractional noises and applications. *SIAM Rev.*, **10**, 422–437. URL: <https://doi.org/10.1137/1010093>.
- Mikosch(1998). Mikosch, T. (1998) *Elementary stochastic calculus—with finance in view*, vol. 6 of *Advanced Series on Statistical Science & Applied Probability*. World Scientific Publishing Co., Inc., River Edge, NJ. URL: <https://doi.org/10.1142/9789812386335>.
- Milevsky and Promislow(2001). Milevsky, M. A. and Promislow, S. D. (2001) Mortality derivatives and the option to annuitise. vol. 29, 299–318. URL: [https://doi.org/10.1016/S0167-6687\(01\)00093-2](https://doi.org/10.1016/S0167-6687(01)00093-2). 4th IME Conference (Barcelona, 2000).
- Mishura(2008). Mishura, Y. S. (2008) *Stochastic calculus for fractional Brownian motion and related processes*, vol. 1929 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin. URL: <https://doi.org/10.1007/978-3-540-75873-0>.
- Neuenkirch and Tindel(2014). Neuenkirch, A. and Tindel, S. (2014) A least square-type procedure for parameter estimation in stochastic differential equations with additive fractional noise. *Stat. Inference Stoch. Process.*, **17**, 99–120. URL: <https://doi.org/10.1007/s11203-013-9084-z>.
- Nualart(2006). Nualart, D. (2006) Fractional Brownian motion: stochastic calculus and applications. In *International Congress of Mathematicians. Vol. III*, 1541–1562. Eur. Math. Soc., Zürich.
- Park et al.(2011) Park, Hernández-Campos, Le, Marron, Park, Pipiras, Smith, Smith, Trovero and Zhu. Park, C., Hernández-Campos, F., Le, L., Marron, J. S., Park, J., Pipiras, V., Smith, F. D., Smith, R. L., Trovero, M. and Zhu, Z. (2011) Long-range dependence analysis of Internet traffic. *J. Appl. Stat.*, **38**, 1407–1433. URL: <https://doi.org/10.1080/02664763.2010.505949>.
- Pitacco et al.(2009) Pitacco, Denuit, Haberman and Olivieri. Pitacco, E., Denuit, M., Haberman, S. and Olivieri, A. (2009) *Modelling longevity dynamics for pensions and annuity business*. Oxford University Press. URL: <https://www.amazon.com/Modelling-Longevity-Dynamics-Pensions-Mathematics/dp/0199547270?SubscriptionId=AKIAIOBINVYXZQZ2U3A&tag=chimbori05-20&linkCode=xm2&camp=2025&creative=165953&creativeASIN=0199547270>.
- Prakasa Rao(2010). Prakasa Rao, B. L. S. (2010) *Statistical inference for fractional diffusion processes*. Wiley Series in Probability and Statistics. John Wiley & Sons, Ltd., Chichester.
- Priestley(1981). Priestley, M. B. (1981) *Spectral analysis and time series. Vol. 1*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London-New York. Univariate series, Probability and Mathematical Statistics.
- Weron(2002). Weron, R. (2002) Estimating long-range dependence: finite sample properties and confidence intervals. *Phys. A*, **312**, 285–299. URL: [https://doi.org/10.1016/S0378-4371\(02\)00961-5](https://doi.org/10.1016/S0378-4371(02)00961-5).
- Yerlikaya-Özkurt et al.(2014) Yerlikaya-Özkurt, Vardar-Acar, Yolcu-Okur and Weber. Yerlikaya-Özkurt, F., Vardar-Acar, C., Yolcu-Okur, Y. and Weber, G.-W. (2014) Estimation of the Hurst parameter for fractional Brownian motion using the CMARS method. *J. Comput. Appl. Math.*, **259**, 843–850. URL: <https://doi.org/10.1016/j.cam.2013.08.001>.
- Young(1936). Young, L. C. (1936) An inequality of the Hölder type, connected with Stieltjes integration. *Acta Math.*, **67**, 251–282. URL: <https://doi.org/10.1007/BF02401743>.

---

Zeng et al.(2012)Zeng, Chen and Yang. Zeng, C., Chen, Y. and Yang, Q. (2012) The fBm-driven Ornstein-Uhlenbeck process: probability density function and anomalous diffusion. *Fract. Calc. Appl. Anal.*, **15**, 479–492. URL: <https://doi.org/10.2478/s13540-012-0034-z>.

$\widehat{h(t)}$  for women at age 0 $\widehat{h(t)}$  for women at age 5 $\widehat{h(t)}$  for women at age 25 $\widehat{h(t)}$  for women at age 50





**Fig. 4** Simulations for the rate mortality with the fOU model: ages 60, 70, 80, 90 and  $N = 10000$ .