Recommended Baseline Bloodwork Before Starting a Peptide Protocol

Before beginning any peptide, GLP-1, longevity, recovery, or performance-enhancement protocol, we recommend obtaining baseline bloodwork. This allows you to monitor progress, identify potential deficiencies, and make informed decisions about your health.

Essential Bloodwork (Men & Women)

Panel	Tests
Complete Blood Count (CBC)	White Blood Cells (WBC); Red Blood Cells (RBC); Hemoglobin; Hematocrit; Platelets
Comprehensive Metabolic Panel (CMP)	Glucose; Kidney Function (Creatinine, BUN, eGFR); Liver Function (ALT, AST, GGT); Electrolytes (Sodium, Potassium, Chloride)
Lipid Panel	Total Cholesterol; LDL Cholesterol; HDL Cholesterol; Triglycerides
Blood Sugar & Metabolic Health	Fasting Glucose; Fasting Insulin; HbA1c
Inflammation Markers	High-Sensitivity C-Reactive Protein (hs-CRP); Homocysteine
Thyroid Function	TSH; Free T3; Free T4
Nutrient Status	Vitamin D (25-OH); Vitamin B12; Ferritin; Iron Panel

Men’s Hormone Panel

Hormones; Total Testosterone; Free Testosterone; SHBG; Estradiol (Sensitive); DHEA-S; PSA; IGF-1; Cortisol (AM)

Women’s Hormone Panel

Hormones; Estradiol; Progesterone; Testosterone; DHEA-S; FSH; LH; Cortisol (AM); For cycling women:; Blood should ideally be collected during days 19-21 of the menstrual cycle for progesterone assessment.

Advanced Longevity Testing (Optional)

For clients seeking deeper optimization:; ApoB; Lipoprotein(a); Fasting Insulin; Omega-3 Index; Uric Acid; Magnesium (RBC); Zinc; Selenium; Biological Age Testing; Continuous Glucose Monitoring (CGM)

When to Retest

Protocol	Retest
Fat Loss Protocols	(Retatrutide, Tirzepatide, MOTS-C); Baseline; 8-12 weeks; 6 months
Recovery & Longevity Protocols	(BPC-157, TB500, GHK-Cu, NAD+, SS-31); Baseline; 3-6 months
Growth Hormone Protocols	(CJC-1295, Ipamorelin, Sermorelin, Tesamorelin, IGF-1 LR3); Baseline; 8-12 weeks; Every 6 months thereafter

Important Disclaimer

Peptides are intended for research and educational purposes. This bloodwork guide is not medical advice and should not replace consultation with a qualified healthcare professional. Always discuss laboratory findings and treatment decisions with your physician.