Glutathione 1500mg

1500mg Vial

BAC Water Added	Concentration	300mg Dose syringe draw	600mg Dose syringe draw
1.5mL	1,000mg/mL	30 units (0.3mL)	60 units (0.6mL)
3mL Recommended	500mg/mL	60 units (0.6mL)	120 units (1.2mL) Over 1mL — needs a larger syringe or two draws
5mL	300mg/mL	100 units (1mL)	200 units (2mL) Over 1mL — needs a larger syringe or two draws

Units are the marks on a U-100 insulin syringe (100 units = 1mL): the amount drawn for one dose, not the number of doses in the vial.

Reconstitute: add **3 mL** of bacteriostatic water to the 1500 mg vial. Resulting concentration: **500 mg/mL**.

DOSE	SYRINGE UNITS	ML VOLUME	SCHEDULE
200 mg	40 units	0.4 mL	2-3x/week IM or SubQ Starting
400 mg	80 units	0.8 mL	2-3x/week IM or SubQ Mid-range
600 mg	120 units	1.2 mL	2-3x/week IM or SubQ Functional-medicine upper range

Math assumes U-100 insulin syringes (1 mL = 100 units). Verify your syringe matches before injecting. Round half-units to the nearest visible mark.

TIP:

Make sure to keep it in the fridge!