
PROTOCOL

NAD+ 200mg Protocol



01 Reconstitution

Vial Size: NAD+ 200mg

Add: 1mL bacteriostatic water

Concentration: 200mg/mL

Storage: Refrigerate after mixing.

Use a U100 insulin syringe.

*** Optimal Energy & Longevity**

Dose

50mg

Draw

25 units

Frequency

3 times weekly

02 Timing

Morning, preferably fasted

Cycle

4–8 weeks or ongoing

One vial provides

4 doses

Approximately 1.3 weeks at 3 injections per week

03 Recovery / Burnout Protocol

Dose

100mg

Draw

50 units

Frequency

3 times weekly

04 Timing

Morning, preferably fasted

Cycle

4–8 weeks

One vial provides

2 doses

Approximately 1 week at 3 injections per week

05 Unit Conversion Guide

Dose	Units
25mg	12.5 units
50mg	25 units
75mg	37.5 units
100mg	50 units
150mg	75 units
200mg	100 units

06 Timing

Morning administration is preferred.

Fasted administration is ideal.

Wait approximately 30 minutes before eating.

Stay well hydrated throughout the day.

07 Suggested Stack Options

Cellular Energy Stack

NAD+ 50mg (25 units) – 3x weekly

MOTS-c – 5mg Monday, Wednesday, Friday

Mitochondrial Support Stack

NAD+ 50–100mg – 3x weekly

SS-31 – 5–10mg daily

Performance & Recovery Stack

NAD+ 100mg (50 units) 3x weekly

MOTS-c 5mg 3x weekly

Creatine 5g daily

Electrolytes

High-protein diet

08 Weekly Schedule

Day	Morning
Monday	NAD+
Tuesday	—
Wednesday	NAD+
Thursday	—
Friday	NAD+

08 Weekly Schedule · continued

Day	Morning
Saturday	—
Sunday	—

09 Benefits

Supports cellular energy production

Promotes mitochondrial health

May improve mental clarity and focus

Supports healthy aging and longevity

Assists recovery from stress and burnout

Supports overall metabolic function

Educational purposes only. Individual responses vary, and protocols should be personalized when appropriate.