

COMP 580 – Assignment 4

Section 2: Adaptive Sampling and Random Sampling

Saurav Kumar Gupta (SG275)

December 2, 2025

Setup

We have a set of non-negative numbers

$$S = \{w_1, w_2, \dots, w_n\},$$

and we want to compute the sum

$$T = \sum_{i=1}^n w_i.$$

Assume that element w_i is sampled with probability p_i , where

$$p_i > 0, \quad \sum_{i=1}^n p_i = 1.$$

Sampling with these probabilities induces a distribution D on S . We form a sampled set SS by sampling according to these probabilities. The assignment defines the estimator

$$\hat{T} = \sum_{w_j \in SS} \frac{w_j}{p_j}.$$

We are asked to:

- Prove that \hat{T} is an unbiased estimator of T .
- Calculate $\text{Var}(\hat{T})$ in terms of the w_i 's and p_i 's.
- Comment on how to design the p_i 's to obtain smaller variance than random sampling with $p_i = 1/n$.

Let I_i be the indicator random variable of whether w_i is included in SS :

$$I_i = \begin{cases} 1, & \text{if } w_i \in SS, \\ 0, & \text{otherwise.} \end{cases}$$

Then $\mathbb{P}(I_i = 1) = p_i$ and

$$\hat{T} = \sum_{i=1}^n I_i \frac{w_i}{p_i}.$$

Using linearity of expectation,

$$\begin{aligned}\mathbb{E}[\hat{T}] &= \mathbb{E}\left[\sum_{i=1}^n I_i \frac{w_i}{p_i}\right] = \sum_{i=1}^n \mathbb{E}\left[I_i \frac{w_i}{p_i}\right] \\ &= \sum_{i=1}^n \frac{w_i}{p_i} \mathbb{E}[I_i] = \sum_{i=1}^n \frac{w_i}{p_i} p_i \\ &= \sum_{i=1}^n w_i = T.\end{aligned}$$

Thus

$$\boxed{\mathbb{E}[\hat{T}] = T,}$$

so \hat{T} is an unbiased estimator of T .

Write

$$\hat{T} = \sum_{i=1}^n Y_i, \quad \text{where} \quad Y_i := I_i \frac{w_i}{p_i}.$$

We assume that the sampling of different elements is independent, so that the I_i (and hence the Y_i) are independent.

2.1 Variance of a single term

The random variable Y_i takes value $\frac{w_i}{p_i}$ with probability p_i and 0 with probability $1 - p_i$. We already have $\mathbb{E}[Y_i] = w_i$. Also,

$$\mathbb{E}[Y_i^2] = \left(\frac{w_i}{p_i}\right)^2 \mathbb{P}(I_i = 1) + 0^2 \mathbb{P}(I_i = 0) = \left(\frac{w_i}{p_i}\right)^2 p_i = \frac{w_i^2}{p_i}.$$

Therefore

$$\begin{aligned}\text{Var}(Y_i) &= \mathbb{E}[Y_i^2] - (\mathbb{E}[Y_i])^2 \\ &= \frac{w_i^2}{p_i} - w_i^2 \\ &= w_i^2 \left(\frac{1}{p_i} - 1\right) = w_i^2 \frac{1 - p_i}{p_i}.\end{aligned}$$

2.2 Variance of the estimator

Since the Y_i are independent,

$$\begin{aligned}\text{Var}(\hat{T}) &= \text{Var}\left(\sum_{i=1}^n Y_i\right) = \sum_{i=1}^n \text{Var}(Y_i) \\ &= \sum_{i=1}^n w_i^2 \frac{1 - p_i}{p_i}.\end{aligned}$$

Equivalently,

$$\text{Var}(\hat{T}) = \sum_{i=1}^n \left(\frac{w_i^2}{p_i} - w_i^2\right) = \left(\sum_{i=1}^n \frac{w_i^2}{p_i}\right) - \sum_{i=1}^n w_i^2.$$

3. Choice of p_i and comparison to uniform sampling

3.1 Uniform random sampling

Under uniform random sampling, $p_i = 1/n$ for all i . Plugging into the variance expression,

$$\begin{aligned}\text{Var}_{\text{uniform}}(\hat{T}) &= \sum_{i=1}^n w_i^2 \frac{1 - \frac{1}{n}}{\frac{1}{n}} \\ &= \sum_{i=1}^n w_i^2 (n-1) \\ &= (n-1) \sum_{i=1}^n w_i^2.\end{aligned}$$

3.2 Designing p_i for smaller variance

From the general variance formula,

$$\text{Var}(\hat{T}) = \sum_{i=1}^n w_i^2 \frac{1 - p_i}{p_i} = \left(\sum_{i=1}^n \frac{w_i^2}{p_i} \right) - \sum_{i=1}^n w_i^2,$$

we see that for fixed w_i the term that depends on the sampling design is

$$F(p_1, \dots, p_n) = \sum_{i=1}^n \frac{w_i^2}{p_i},$$

subject to

$$p_i > 0, \quad \sum_{i=1}^n p_i = 1.$$

To minimize the variance of \hat{T} , we minimize F under these constraints. Using a Lagrange multiplier λ ,

$$\mathcal{L}(p_1, \dots, p_n, \lambda) = \sum_{i=1}^n \frac{w_i^2}{p_i} + \lambda \left(\sum_{i=1}^n p_i - 1 \right).$$

For each i ,

$$\frac{\partial \mathcal{L}}{\partial p_i} = -\frac{w_i^2}{p_i^2} + \lambda = 0 \quad \implies \quad p_i^2 = \frac{w_i^2}{\lambda} \quad \implies \quad p_i \propto w_i.$$

Thus, to obtain smaller variance than uniform sampling, we should choose sampling probabilities p_i so that larger w_i have larger p_i . In particular, the choice

$$p_i = \frac{w_i}{\sum_{j=1}^n w_j}$$

makes F (and hence $\text{Var}(\hat{T})$) as small as possible under the given constraints, and therefore gives better (smaller) variance than uniform random sampling when the w_i are not all equal.