



FOREST VERIFICATION REPORT

ANALYZED ZONE INFORMATION

Latitude	-1.4537	Longitude	30.0693
Zone size (km)	10	Analysis date	2026-08-26

61

FOREST INTEGRITY SCORE / 100 Degraded Forest

DETAILED RESULTS

Indicator	Value	Interpretation
Current vegetation (NDVI)	0.592	Good
NDVI Trend 3 years (Sentinel-2)	-0.024	Stable
10-year trend (Landsat 8/9)	+0.272	Improving
40-year trend (Landsat 1984)	+0.315	Improving
Forest Status (FAO)	Forest recovery	Recuperation
Annual NDVI rate	+0.79% / year	Amelioration
NDVI change 5 years	-0.043	Stable
AI Classification	Temperate Forest	Broadleaf/mixed forest, moderate climate
Sentinel-1 Radar (VV)	-8.065 dB	Dense forest (radar)
Forest cover	94.8%	Excellent
Fire pixels detected (FIRMS/year)	549 pixels	~137 events estimated
Overall trend	stable	
Satellites used	- ESA Sentinel-2 - NASA FIRMS - ESA Sentinel-1 - NASA Landsat 8/9 - NASA Landsat 4/5	
Deforestation Alert	CAUTION — Light forest degradation	
Area change (3 years)	-353.8 ha	-3.61%
Annual rate (area/yr)	-117.9 ha/yr	Loss
Area change (10 years)	+9012.9 ha	+2019.92%
Annual rate 10yr (area)	+901.3 ha/yr	Recovery

SCORE BREAKDOWN

Component	Points	Maximum
Current vegetation (NDVI)	18	25
NDVI trend (3 years)	8	25
Forest cover	25	25
Sentinel-1 Radar (VV)	10	15
Fire score (0=many fires, 10=none)	0	10
TOTAL	61	100

FIRE DETECTION METHODOLOGY: Fire pixels are detected by NASA FIRMS using VIIRS (375m) and MODIS (1km) thermal anomaly sensors. Raw pixel count represents thermal detections over 12 months in the analysis zone. Estimated fire events are calculated using NASA clustering methodology (~4 pixels per fire event average). This is a statistical approximation. For precise fire event mapping, consult NASA FIRMS directly at firms.modaps.eosdis.nasa.gov

This report was automatically generated by CanopySat using real satellite data. contact@canopysat.org - canopysat.org