

# pH determines the energetic efficiency of the cyanobacterial CO<sub>2</sub> concentrating mechanism: mathematical supplement

Niall Mangan, Avi Flamholz, Rachel Hood, Ron Milo & David Savage

## 1 Equations when RuBisCO is saturated

The analytic solution for the CO<sub>2</sub> and HCO<sub>3</sub><sup>-</sup> concentration in the carboxysome when RuBisCO is saturated is:

$$C_{carboxysome} = \frac{N}{M} - \frac{R_c^3 V_{max} P}{3MD} \quad (1)$$

$$H_{carboxysome} = K_{eq}(pH) C_{carboxysome} \quad (2)$$

where,

$$N = (j_c + k_m^{eff}(pH_{out})) H_{out} ((k_m^C + \alpha) G^C + \frac{D}{R_b^2}) + k_m^C C_{out} (k_m^{eff} G^H + \alpha G^C + \frac{D}{R_b^2}) \quad (3)$$

$$M = K_{eq} * k_m^{eff} \left( (\alpha + k_m^C) G^C + \frac{D}{R_b^2} \right) + k_m^C \left( k_m^{eff} G^H + \frac{D}{R_b^2} \right) + \alpha k_m^{eff} G^H \quad (4)$$

$$P = ((\alpha + k_m^C) G^C + \frac{D}{R_b^2}) (k_m^{eff} G^H + \frac{D}{R_b^2}) \quad (5)$$

$$G^C = \frac{D}{R_c^2 k_c^C} + \frac{1}{R_c} - \frac{1}{R_b} \quad (6)$$

$$G^H = \frac{D}{R_c^2 k_c^H} + \frac{1}{R_c} - \frac{1}{R_b} \quad (7)$$

The derivation of this equation can be found in the supplementary material of (Mangan & Brenner, eLife 2014). Here we have made a few modifications: (1) kept track of the carboxysome permeability to CO<sub>2</sub>,  $k_c^C$ , and HCO<sub>3</sub><sup>-</sup>,  $k_c^H$ , independently, (2) substituted the pH dependent equilibrium constant for the carbonic anhydrase reaction,  $K_{eq}(ph) = \frac{V_{ca} K_{ba}}{V_{ba} K_{ca}}$ , (3) written the CO<sub>2</sub> → HCO<sub>3</sub><sup>-</sup> reaction with  $\alpha$  as the linear reaction rate (in Mangan 2014 the linear rate was  $\alpha/K_\alpha$ ), (4) we have replaced the membrane permeability to HCO<sub>3</sub><sup>-</sup> with the effective membrane permeability to the bicarbonate pool, and designated when is dependent on the external pH,  $k_m^{eff}(pH_{out})$ . This term only appears once in

equation 3 for  $N$ . For all other  $k_m^{eff} = k_m^{eff}(pH_{in})$  values it is dependent on the pH inside the cell, so we have dropped indicating the pH dependence to simplify the formulas.

## 2 Analysis of membrane permeability effects

### 2.1 Cell permeability compared to diffusive velocities

Examining equation (6) we note that for large carboxysome permeability  $1/R_c$  will be the dominant term, and for smaller carboxysome permeability values the first term will be larger and dominate. Therefore  $G^C \geq 1/R_c$ . Studying the equations (3-7) we note that the terms  $((\alpha + k_m^C)G^C + \frac{D}{R_b^2})$  appears repeatedly. We use the following argument:

$$(\alpha + k_m^C)G^C \geq (\alpha + k_m^C)/R_c \gg D/R_b^2, \quad (8)$$

$$\text{if } (\alpha + k_m^C) \gg DR_c/R_b^2 \quad (9)$$

For even a small 20 nm diameter ( $R_c = 10^{-6}$  cm) carboxysome this will hold as  $k_m^C \approx 0.3$  cm/s and  $DR_c/R_b^2 = 4 \times 10^{-3}$  cm/s from the values in Table S1. So the membrane permeability to CO<sub>2</sub> could be an order of magnitude too high in our model and this would still be a reasonable assumption. Therefore we will substitute

$$(\alpha + k_m^C)G^C + D/R_b^2 \approx (\alpha + k_m^C)G^C. \quad (10)$$

Inserting this into equations (1-5) we get

$$\begin{aligned} C_{carboxysome} = & \frac{(j_c + k_m^{eff}(pH_{out}))H_{out}(k_m^C + \alpha)G^C + k_m^C C_{out}(k_m^{eff}G^H + \alpha G^C + \frac{D}{R_b^2})}{K_{eq}k_m^{eff}(\alpha + k_m^C)G^C + k_m^C \left(k_m^{eff}G^H + \frac{D}{R_b^2}\right) + \alpha k_m^{eff}G^H} \\ & - \frac{R_c^3 V_{max}(\alpha + k_m^C)G^C(k_m^{eff}G^H + \frac{D}{R_b^2})/(3D)}{K_{eq}k_m^{eff}(\alpha + k_m^C)G^C + k_m^C \left(k_m^{eff}G^H + \frac{D}{R_b^2}\right) + \alpha k_m^{eff}G^H}. \quad (11) \end{aligned}$$

We can divide through by  $(k_m^C + \alpha)$  to obtain:

$$\begin{aligned} C_{carboxysome} = & \frac{(j_c + k_m^{eff}(pH_{out}))H_{out}G^C + \frac{k_m^C}{(k_m^C + \alpha)}C_{out}(k_m^{eff}G^H + \alpha G^C + \frac{D}{R_b^2})}{K_{eq}k_m^{eff}G^C + \frac{k_m^C}{(k_m^C + \alpha)} \left(k_m^{eff}G^H + \frac{D}{R_b^2}\right) + \frac{\alpha}{(k_m^C + \alpha)}k_m^{eff}G^H} \\ & - \frac{R_c^3 V_{max}G^C(k_m^{eff}G^H + \frac{D}{R_b^2})/(3D)}{K_{eq}k_m^{eff}G^C + \frac{k_m^C}{(k_m^C + \alpha)} \left(k_m^{eff}G^H + \frac{D}{R_b^2}\right) + \frac{\alpha}{(k_m^C + \alpha)}k_m^{eff}G^H}. \quad (12) \end{aligned}$$

We now want to examine the remaining terms in the membrane permeability to CO<sub>2</sub>,  $k_m^C$ .

## 2.2 Membrane permeability to CO<sub>2</sub> has little effect.

There are two parameter groupings in equation (12) containing  $k_m^C$ :

$$\frac{k_m^C}{k_m^C + \alpha} \quad (13)$$

$$\frac{\alpha}{k_m^C + \alpha} \quad (14)$$

Therefore if  $k_m^C > \alpha$  or  $\text{CO}_2 \rightarrow \text{HCO}_3^-$  conversion is negligible the first term (13) reduces to 1, and the second reduces to  $1/k_m^C$ . We will return to the case where this conversion is not negligible later.

With these two simplifications we obtain:

$$C_{\text{carboxysome}} = \frac{(j_c + k_m^{\text{eff}}(pH_{\text{out}}))H_{\text{out}}G^C + C_{\text{out}}(k_m^{\text{eff}}G^H + \alpha G^C + \frac{D}{R_b^2})}{K_{eq}k_m^{\text{eff}}G^C + \left(k_m^{\text{eff}}G^H + \frac{D}{R_b^2}\right) + \frac{1}{k_m^C}k_m^{\text{eff}}G^H} - \frac{R_c^3 V_{\text{max}}G^C(k_m^{\text{eff}}G^H + \frac{D}{R_b^2})/(3D)}{K_{eq}k_m^{\text{eff}}G^C + \left(k_m^{\text{eff}}G^H + \frac{D}{R_b^2}\right) + \frac{1}{k_m^C}k_m^{\text{eff}}G^H}. \quad (15)$$

Examining equation (15), note that the only appearance of the membrane permeability to CO<sub>2</sub> is now in the denominator which we can rewrite as  $k_m^{\text{eff}}(G^C K_{eq} + \frac{G^H}{k_m^C}) + \left(k_m^{\text{eff}}G^H + \frac{D}{R_b^2}\right)$ . Using this equation, we can write a strong bound on when the membrane permeability will effect the function of the CCM.

We find  $k_m^C$  has no significant effect when  $K_{eq}G^C > \frac{G^H}{k_m^C}$  or  $k_m^C > \frac{G^H}{G^C K_{eq}}$ . If we assume that the carboxysome permeability to CO<sub>2</sub> will always be smaller than or equal to the permeability to HCO<sub>3</sub><sup>-</sup> ( $k_c^C \geq k_c^H$ ) then  $G^H \geq G^C$  and  $\frac{G^H}{G^C} \leq 1$ , so  $k_m^C$  will be negligible as long as  $k_m^C > 1/K_{eq}$ . For pH > 6.6,  $1/K_{eq} > 0.3$  and therefore the assumed value of  $k_m^C = 0.3$  will be negligible. However, if the cell operated in a lower pH regime and the membrane permeability was substantially lower to CO<sub>2</sub> it would begin to effect the CO<sub>2</sub> concentration.

Thus far we have made a series of observations about the size of terms compared to the membrane permeability to CO<sub>2</sub> and found that when  $(\alpha + k_m^C) \gg DR_c/R_b^2$ ,  $k_m^C > \alpha$  and  $k_m^C > \frac{G^H}{G^C K_{eq}} \approx 1/K_{eq}$  the CO<sub>2</sub> concentration in the carboxysome reduces to

$$C_{\text{carboxysome}} = \frac{(j_c + k_m^{\text{eff}}(pH_{\text{out}}))H_{\text{out}}G^C + C_{\text{out}}(k_m^{\text{eff}}G^H + \alpha G^C + \frac{D}{R_b^2})}{k_m^{\text{eff}}(G^C K_{eq} + G^H) + \frac{D}{R_b^2}} - \frac{R_c^3 V_{\text{max}}G^C(k_m^{\text{eff}}G^H + \frac{D}{R_b^2})/(3D)}{k_m^{\text{eff}}(G^C K_{eq} + G^H) + \frac{D}{R_b^2}}. \quad (16)$$

We can make a similar argument taking the equation for the CO<sub>2</sub> concen-

tration at the cell membrane:

$$C_{cytosol}(r = R_b) = \frac{k_m^C C_{out} - (\alpha + k_m^C) C_{carboxysome}}{(\alpha + k_m^C) G^C + D/R_b^2} G^C + C_{carboxysome} \approx C_{out} \quad (17)$$

This means that the  $\text{CO}_2$  leakage term will be negligible since the cytosolic  $\text{CO}_2$  concentration will be approximately equal to the external  $\text{CO}_2$  concentration. The  $\text{HCO}_3^-$  transport required to sustain a given internal inorganic carbon pool will then be:

$$\begin{aligned} j_c H_{out} &= \left( \frac{R_c^3}{3R_b^2} V_{max} - k_m^C (C_{out} - C_{cytosol}) - k_m^{eff} H_{out} + k_m^{eff} H_{cytosol} \right) \\ &= \left( \frac{R_c^3}{3R_b^2} V_{max} - k_m^{eff} H_{out} + k_m^{eff} H_{cytosol} \right) \end{aligned} \quad (18)$$

We can calculate  $H_{carboxysome} = K_{eq} C_{carboxysome}$  from equation (17), and is therefore also independent of  $k_m^C$ . In previous work we showed that

$$H_{cytosol} = \frac{(j_c + k_m^{eff}(pH_{out}))H_{out} + \frac{\alpha}{K_a} C_{cytosol}(r = R_b) - k_m^{eff} H_{carboxysome}}{k_m^{eff} G^H + \frac{D}{R_b^2}} G^H \quad (19)$$

We have now shown that all the terms in  $H_{cyto}$  are negligibly dependent on the membrane permeability to  $\text{CO}_2$ . Therefore, the  $\text{HCO}_3^-$  transport level require to satisfy equation (18) is independent of the membrane permeability to  $\text{CO}_2$ . This observation is consistent with the low flux of  $\text{CO}_2$  leakage in main text Figure 2.

### 2.3 Without facilitated $\text{CO}_2$ uptake external $\text{CO}_2$ has little effect

Unless conversion from  $\text{CO}_2$  to  $\text{HCO}_3^-$  is large we note that the second  $C_{out}$  term in equation(15) is negligible for the regimes we study. We will revisit  $\text{CO}_2$  uptake and recycling later. Comparing this term against the first term in the numerator, again allows us to put a quantitative description on when this regime holds. Additionally we find that when the transport of  $\text{HCO}_3^-$  is significant ( $j_c > k_m^{eff}(pH_{out})$ ) we arrive at

$$C_{carboxysome} = \frac{j_c H_{out} G^C - R_c^3 V_{max} G^C (k_m^{eff} G^H + \frac{D}{R_b^2}) / (3D)}{k_m^{eff} (G^C K_{eq} + G^H) + \frac{D}{R_b^2}} \quad (20)$$

$$H_{carboxysome} = K_{eq} C_{carboxysome} \quad (21)$$

### 3 Effect of Carboxysome permeability

Recalling the equation for  $G^C = \frac{D}{R_c^2 k_c^C} + \frac{1}{R_c} - \frac{1}{R_b}$ , we can see that the carboxysome permeability to  $\text{CO}_2$  will only matter if  $\frac{D}{R_c^2 k_c^C} > \frac{1}{R_c}$ . In other words the carboxysome permeability to  $\text{CO}_2$ ,  $k_c^C$ , begins to effectively trap  $\text{CO}_2$  in the carboxysome when  $k_c^C < \frac{D}{R_c} \approx 2 \text{ cm/s}$  for our base case of a 100 nm carboxysome ( $R_c = 50 \text{ nm}$ ). Similarly  $G^H \approx \frac{D}{R_c^2 k_c^H}$  when  $k_c^H < \frac{D}{R_c}$ . As common thinking is that  $k_c^H \geq k_c^C$ ,  $k_c^H < \frac{D}{R_c}$  may not always hold when  $k_c^C < \frac{D}{R_c}$ .

#### 3.0.1 Different carboxysome permeability for $\text{HCO}_3^-$

An existing hypothesis in the CCM literature is that the carboxysome has differential permeability and is more permeable to  $\text{HCO}_3^-$  and less permeable to  $\text{CO}_2$ . Intuitively this would allow more  $\text{HCO}_3^-$  into the carboxysome and trap more  $\text{CO}_2$ , thereby accumulating more inorganic carbon in the form of  $\text{CO}_2$ . We use our model to test whether differential carboxysome permeability enables higher carboxysomal  $\text{CO}_2$  concentration for the same level of  $\text{HCO}_3^-$  transport. In the Figure S12 we show the  $k_c$  vs  $j_c$  phase space where we have plotted the carboxysome permeability to  $\text{CO}_2$ ,  $k_c^C$ , on the y-axis. We plot different ratios (1, 10, 100, 1000) between  $k_c^C$  and the carboxysome permeability to  $\text{HCO}_3^-$ ,  $k_c^H = \text{ratio} \times k_c^C$ .

Examining Figure S12, we see that making the carboxysome more permeable to  $\text{HCO}_3^-$  does not improve the function of the CCM as drastically as one might assume. The "turn on" of  $\text{CO}_2$  accumulation with decreasing permeability is unaffected by changes to  $k_c^H$ , and depends only on the permeability to  $\text{CO}_2$ ,  $k_c^C$ . The "turn off" of accumulation for lower carboxysome permeabilities is greatly affected by the permeability of the carboxysome to  $\text{HCO}_3^-$ ,  $k_c^H$ . These two effects are exactly what we previously discussed as defining the carboxysome permeability optimum.

As we start at the top of the y-axis and decrease the carboxysome permeability the following occurs: At high permeability not enough  $\text{CO}_2$  is trapped, but  $\text{HCO}_3^-$  enters readily. As we moved to lower permeabilities  $\text{CO}_2$  begins to be trapped, but there is a window where  $\text{HCO}_3^-$  still enters enough to supply the system. Eventually the carboxysome begins to restrict  $\text{HCO}_3^-$  entry. If the carboxysome is more permeable to  $\text{HCO}_3^-$  than to  $\text{CO}_2$  then the window where  $\text{CO}_2$  trapping is effective without restricting  $\text{HCO}_3^-$  entry broadens. The width of this window (on the y-axis) will also depend strongly on how much of the  $\text{CO}_2$  is being fixed.

The "turn off" of the optimum, caused by not allowing enough  $\text{HCO}_3^-$  into the carboxysome, does slightly increase the amount of transport required to saturate RuBisCO at the carboxysome optimum. The reduction in transport required, and therefore CCM cost is around 5% when going from a  $k_c^C$  to  $k_c^H$  ratio of 1 to 1000.

## 4 Effect of membrane permeability to $\text{H}_2\text{CO}_3$

The sensitivity of the cost to our assumption for the value of the membrane permeability to  $\text{H}_2\text{CO}_3$  can be determined from the equation derived previously. If we are in a regime where  $\text{CO}_2$  leakage is negligible, as is the regime presented in the main paper, the second line holds.

$$\begin{aligned} j_c H_{out} &= \left( \frac{R_c^3}{3R_b^2} V_{max} - k_m^C (C_{out} - C_{cytosol}) - k_m^{eff}(pH_{out})H_{out} + k_m^{eff}(pH_{in})H_{cytosol} \right) \\ &= \left( \frac{R_c^3}{3R_b^2} V_{max} - k_m^{eff}(pH_{out})H_{out} + k_m^{eff}(pH_{in})H_{cytosol} \right) \quad (22) \end{aligned}$$

In this equation  $k_m^{eff} = k_m^{H_2CO_3} \times 10^{(pK_1 - pH)}$ . Therefore, the leakage of  $H_{total}$  out of the cell will depend linearly on what we assume for  $k_m^{H_2CO_3}$ . This linear dependence is passed on to the active  $\text{HCO}_3^-$  transport required to replenish the leaked inorganic carbon, and therefore onto the CCM cost. In Figure S14 you can see this effect, where going from  $k_m^{H_2CO_3} = 3 \times 10^{-2}$  to  $k_m^{H_2CO_3} = 3 \times 10^{-3}$  (an order of magnitude change), decreases the active  $\text{HCO}_3^-$  transport needed by an order of magnitude. Decreasing to  $k_m^{H_2CO_3} = 3 \times 10^{-4}$  is a little less than an order of magnitude, indicating that the linear dependence breaks down and  $\text{CO}_2$  leakage would become important for that value. There is also an order of magnitude change in the optimal carboxysome permeability from  $10^{-4}$  to  $10^{-5}$  across the 2 order of magnitude change in  $k_m^{H_2CO_3}$  we are checking.