

# COMP0009 Logic slides 3. Propositional Tableau.

October 12, 2023

## Proof by Contradiction

- ▶ To prove a formula  $\phi$ ,

## Proof by Contradiction

- ▶ To prove a formula  $\phi$ , first
- ▶ Assume  $\neg\phi$ ,

## Proof by Contradiction

- ▶ To prove a formula  $\phi$ , first
- ▶ Assume  $\neg\phi$ , then
- ▶ Deduce a contradiction  $q \wedge \neg q$ .

## Proof by Contradiction

- ▶ To prove a formula  $\phi$ , first
- ▶ Assume  $\neg\phi$ , then
- ▶ Deduce a contradiction  $q \wedge \neg q$ . Hence,
- ▶ Conclude  $\phi$

## Propositional Tableaux

- ▶ Given the formula  $\phi$ , a tableau can tell us whether it is **satisfiable or not**

## Propositional Tableaux

- ▶ Given the formula  $\phi$ , a tableau can tell us whether it is **satisfiable or not**
- ▶ It does so by **decomposing** the formulae according to certain rules and growing a labelled tree to the point that only literals are left

## Propositional Tableaux

- ▶ Given the formula  $\phi$ , a tableau can tell us whether it is **satisfiable or not**
- ▶ It does so by **decomposing** the formulae according to certain rules and growing a labelled tree to the point that only literals are left
- ▶ A branch is closed if it has one node labelled by a proposition and another node labelled by its negation.



## Propositional Tableaux

- ▶ Given the formula  $\phi$ , a tableau can tell us whether it is **satisfiable or not**
- ▶ It does so by **decomposing** the formulae according to certain rules and growing a labelled tree to the point that only literals are left
- ▶ A branch is closed if it has one node labelled by a proposition and another node labelled by its negation.
- ▶ A tableau is closed if every branch is closed.

## Propositional Tableaux


- ▶ Given the formula  $\phi$ , a tableau can tell us whether it is **satisfiable or not**
- ▶ It does so by **decomposing** the formulae according to certain rules and growing a labelled tree to the point that only literals are left
- ▶ A branch is closed if it has one node labelled by a proposition and another node labelled by its negation.
- ▶ A tableau is closed if every branch is closed.
- ▶ If the tableau closes then  $\phi$  is **unsatisfiable**

## Propositional Tableaux

- ▶ Given the formula  $\phi$ , a tableau can tell us whether it is **satisfiable or not**
- ▶ It does so by **decomposing** the formulae according to certain rules and growing a labelled tree to the point that only literals are left
- ▶ A branch is closed if it has one node labelled by a proposition and another node labelled by its negation.
- ▶ A tableau is closed if every branch is closed.
- ▶ If the tableau closes then  $\phi$  is **unsatisfiable**
- ▶ If the tableau never closes then  $\phi$  is **satisfiable**


## The idea

To test whether  $((\phi \wedge \neg\phi) \vee (\psi \wedge \neg\psi))$  is satisfiable



## The idea

To test whether  $((\phi \wedge \neg\phi) \vee (\psi \wedge \neg\psi))$  is satisfiable



Both branches close  $\Rightarrow$  not satisfiable.

## Tableaux

- Formally, a tableau  $T$  is a type of binary tree where every node is labelled by a formula.

## Tableaux

- ▶ Formally, a tableau  $T$  is a type of binary tree where every node is labelled by a formula.
- ▶ At first,  $T$  has only one node, containing the formula  $\phi$  whose satisfiability we are investigating.

## Tableaux

- ▶ Formally, a tableau  $T$  is a type of binary tree where every node is labelled by a formula.
- ▶ At first,  $T$  has only one node, containing the formula  $\phi$  whose satisfiability we are investigating.
- ▶ Every formula in the tableau, except literals, gets “expanded” and ticked.
- ▶ If a branch of  $T$  contains  $p$  and  $\neg p$  then it is **closed**. Otherwise it is **open**.
- ▶ If every branch of  $T$  is closed then  $T$  is **closed**. Otherwise it is **open**.



# Tableau Expansion Rules


$\alpha$  formulas

$$(\phi \wedge \psi)$$

Observe that  $(\phi \wedge \psi)$  is true if and only if  $\phi$  is true and  $\psi$  is true.

# Tableau Expansion Rules

$\alpha$  formulas



Observe that  $(\phi \wedge \psi)$  is true if and only if  $\phi$  is true and  $\psi$  is true.

**Means that nodes corresponding to  $\phi$  and  $\psi$  are added at every leaf of  $T$  below the current node  $(\phi \wedge \psi)$ .**

# Tableau Expansion Rules

$\alpha$  formulas

Three other kinds of  $\alpha$  formulas.

$$\begin{array}{c} \neg\neg\phi \\ | \\ \phi \end{array}$$

# Tableau Expansion Rules

$\alpha$  formulas

Three other kinds of  $\alpha$  formulas.

$$\begin{array}{c} \neg\neg\phi \\ | \\ \phi \end{array}$$

$$\begin{array}{c} \neg(\phi \vee \psi) \quad \checkmark \\ | \\ \neg\phi \\ | \\ \neg\psi \end{array}$$

**Observe that**  $\neg(\phi \vee \psi) \equiv \neg\phi \wedge \neg\psi$

# Tableau Expansion Rules

$\alpha$  formulas

Three other kinds of  $\alpha$  formulas.

$$\begin{array}{c} \neg\neg\phi \\ | \\ \phi \end{array}$$

$$\begin{array}{c} \neg(\phi \vee \psi) \quad \checkmark \\ | \\ \neg\phi \\ | \\ \neg\psi \end{array}$$

$$\begin{array}{c} \neg(\phi \rightarrow \psi) \quad \checkmark \\ | \\ \phi \\ | \\ \neg\psi \end{array}$$

**Observe that**  $\neg(\phi \vee \psi) \equiv \neg\phi \wedge \neg\psi$

**Observe that**  $\neg(\phi \rightarrow \psi) \equiv \neg(\neg\phi \vee \psi) \equiv \phi \wedge \neg\psi$

# Tableau Expansion Rules


$\beta$  formulas

$$(\phi \vee \psi)$$

For this case, at each leaf of the tableau below the current node, we create two new leaves, one labelled  $\phi$  the other labelled  $\psi$ .

# Tableau Expansion Rules

$\beta$  formulas



For this case, at each leaf of the tableau below the current node, we create two new leaves, one labelled  $\phi$  the other labelled  $\psi$ .

Observe that  $(\phi \vee \psi)$  is true if and only if  $\phi$  is true or  $\psi$  is true.

# Tableau Expansion Rules

$\beta$  formulas

There are two other kinds of  $\beta$  formulas.

$$\begin{array}{c} \neg(\phi \wedge \psi) \quad \checkmark \\ \swarrow \quad \searrow \\ \neg\phi \quad \neg\psi \end{array}$$

**Observe that**  $\neg(\phi \wedge \psi) \equiv \neg\phi \vee \neg\psi$



# Tableau Expansion Rules

$\beta$  formulas

There are two other kinds of  $\beta$  formulas.



**Observe that**  $\neg(\phi \wedge \psi) \equiv \neg\phi \vee \neg\psi$

**Observe that**  $(\phi \rightarrow \psi) \equiv \neg\phi \vee \psi$

## Tableaux

- ▶ A tableau is **complete** if every node is either ticked (already expanded) or a literal.
- ▶ If  $\phi$  is at the root of a **complete open** tableau, then  $\phi$  is **satisfiable**.

## Propositional Tableaux

**Remember:** In a proof system we are interested in validity, not just satisfiability. Fortunately, we know that:

$\phi$  is satisfiable  $\iff \neg\phi$  is not valid

$\phi$  is valid  $\iff \neg\phi$  is not satisfiable

## Propositional Tableaux

**Remember:** In a proof system we are interested in validity, not just satisfiability. Fortunately, we know that:

$\phi$  is satisfiable  $\iff \neg\phi$  is not valid

$\phi$  is valid  $\iff \neg\phi$  is not satisfiable

**Solution:** In order to test whether a formula  $\phi$  is valid we see whether the tableau for  $\neg\phi$  closes

**Example 1: Is formula  $((p \wedge q) \rightarrow r) \rightarrow ((p \rightarrow r) \vee (q \rightarrow r))$  valid?**

---

$$(1) \neg(((p \wedge q) \rightarrow r) \rightarrow ((p \rightarrow r) \vee (q \rightarrow r)))$$

**Example 1: Is formula  $((p \wedge q) \rightarrow r) \rightarrow ((p \rightarrow r) \vee (q \rightarrow r))$  valid?**


---

$$(1) \neg(((p \wedge q) \rightarrow r) \rightarrow ((p \rightarrow r) \vee (q \rightarrow r))) \quad \checkmark$$


$$(2) ((p \wedge q) \rightarrow r)$$

$$(3) \neg((p \rightarrow r) \vee (q \rightarrow r))$$

**Example 1: Is formula  $((p \wedge q) \rightarrow r) \rightarrow ((p \rightarrow r) \vee (q \rightarrow r))$  valid?**




**Example 1: Is formula  $((p \wedge q) \rightarrow r) \rightarrow ((p \rightarrow r) \vee (q \rightarrow r))$  valid?**







**Example 1: Is formula  $((p \wedge q) \rightarrow r) \rightarrow ((p \rightarrow r) \vee (q \rightarrow r))$  valid?**




**Example 1: Is formula  $((p \wedge q) \rightarrow r) \rightarrow ((p \rightarrow r) \vee (q \rightarrow r))$  valid?**




**Example 1: Is formula  $((p \wedge q) \rightarrow r) \rightarrow ((p \rightarrow r) \vee (q \rightarrow r))$  valid?**



**Example 1: Is formula  $((p \wedge q) \rightarrow r) \rightarrow ((p \rightarrow r) \vee (q \rightarrow r))$  valid?**



**Example 1: Is formula  $((p \wedge q) \rightarrow r) \rightarrow ((p \rightarrow r) \vee (q \rightarrow r))$  valid?**



Closed tableau — **root formula is unsatisfiable, original formula is valid.**

**Example 2: Is formula  $((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r)$  valid?**

$$(1) \neg(((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r))$$

**Example 2: Is formula  $((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r)$  valid?**


---

$$(1) \neg(((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r)) \quad \checkmark$$

$$(2) ((p \rightarrow r) \vee (q \rightarrow r))$$

$$(3) \neg((p \vee q) \rightarrow r)$$

**Example 2: Is formula  $((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r)$  valid?**





**Example 2:** Is formula  $((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r)$  valid?

$$(1) \neg(((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r)) \quad \checkmark$$
 $\alpha(1)$ 
$$(2) ((p \rightarrow r) \vee (q \rightarrow r)) \quad \checkmark$$
$$(3) \neg((p \vee q) \rightarrow r) \quad \checkmark$$
$$(4) \ (p \rightarrow r)$$
 $\beta(2)$ 
$$(5) \ (\overline{q} \rightarrow r)$$
 $\alpha(3)$  $\alpha(3)$ 
$$(6) (p \vee q)$$
$$(8) (p \vee q)$$


1

1


$$(7) \neg r$$

(9)  $\neg r$


**Example 2: Is formula  $((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r)$  valid?**




**Example 2: Is formula  $((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r)$  valid?**




**Example 2: Is formula  $((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r)$  valid?**



**Example 2: Is formula  $((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r)$  valid?**



**Example 2: Is formula  $((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r)$  valid?**



Two open branches — **root formula is satisfiable, original formula is not valid.**

## Disjunctive Normal Form (DNF)

A **DNF** formula is a disjunctions of clauses where each clause is a conjunction of literals. E.g.

$$(p \wedge \neg q \wedge r) \vee (p \wedge q) \vee (\neg p \wedge \neg q \wedge \neg r))$$

Also

$$p \vee \neg q \vee (p \wedge q)$$

Every propositional formula has an equivalent formula in DNF. A **CNF** formula is a conjunction of clauses where each clause is disjunction of literals. E.g.

$$(p \vee q) \wedge (\neg p \vee \neg r) \wedge q$$

## Converting a formula to DNF

1. By tableau
2. By truth table
3. By logical equivalences (De Morgan's laws, distribution laws etc., later)



## Using tableau to find DNF equivalent


- ▶ Place  $\phi$  at root of new tableau.
- ▶ Expand until tableau  $T$  is completed (no nodes left to expand).
- ▶ For each open branch  $\Theta$  of  $T$  let

$$C_{\Theta} = \bigwedge \{\text{literals in } \Theta\}$$

- ▶ Then

$$\phi \equiv \bigvee_{\text{open branches } \Theta} C_{\Theta}$$

**Example 2: Is formula  $((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r)$  valid?**



The **DNF** for  $\neg(((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r))$  is  $(\neg r \wedge \neg p \wedge q) \vee (\neg r \wedge \neg q \wedge p)$

## DNF by truth table

$$\neg(((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r))$$

$p$	$q$	$r$	$\neg(((p \rightarrow r) \vee (q \rightarrow r)) \rightarrow ((p \vee q) \rightarrow r))$	
0	0	0	0	
0	0	1	0	
0	1	0	1	$(\neg p \wedge q \wedge \neg r)$
0	1	1	0	$\vee$
1	0	0	1	$(p \wedge \neg q \wedge \neg r)$
1	0	1	0	
1	1	1	0	

## Problem

- ▶ How do you convert a propositional formula into an equivalent CNF?
- ▶ E.g.

$$((p \wedge \neg q) \vee (q \wedge \neg r)) \equiv ??$$