
```
%Kyle O'Connor
%January 19, 2016
%Lecutre 2
%
clear
close all
home
a = [1 \ 2 \ -5; \ 3 \ 1 \ 4; \ 1 \ 2 \ -1]
b = [7 \ 8 \ 9; 6 \ -5 \ 4; 3 \ 2 \ -1]
c = [1 \ 2 \ 3; \ 4 \ 5 \ 6]
d = [1 \ 2 \ 3; \ 4 \ 5 \ 6]
e = [1 \ 2 \ -5; \ 3 \ 1 \ 4; \ 4 \ 3 \ -1]
a =
                 -5
     1
           2
     3
            1
                  4
     1
            2
                  -1
b =
     7
           8
                  9
     6
           -5
                  4
            2
     3
                  -1
c =
     1
           2
                  3
            5
d =
     1
            2
                   3
            5
e =
     1
            2
                  -5
     3
             1
                  4
             3
                  -1
```

1

```
clear
close all
home
x = 0:10
y1 = (-8-2*x)/3;
y2 = 9 - (3*x);
plot(x,y1,'r.-',x,y2,'b.-')
a = [2 \ 3; \ 3 \ 1]
b = [-8; 9]
det(a)
inv(a)
inv(a)*b
x =
    0 1 2 3 4 5 6 7 8 9 10
a =
      3
    2
    3
b =
   -8
    9
ans =
   -7
ans =
  -0.1429 0.4286
  0.4286 -0.2857
ans =
   5
   -6
```

2



Published with MATLAB® R2015a