

Designing a Long-Term Effectiveness Model of Next-Generation LLINs Using R Software

Koissi Savi (Ph.D.)

Harvard School of Medicine

January 17, 2024

- 1 Introduction to Long-Term Effectiveness Models
- 2 Setting Up the Environment
- 3 Exploratory Data Analysis
- 4 Building Cox Proportional Hazards Model

- 1 Introduction to Long-Term Effectiveness Models
- 2 Setting Up the Environment
- 3 Exploratory Data Analysis
- 4 Building Cox Proportional Hazards Model

Overview of LLINs

- In malaria-endemic regions LLINs represent the most widely used malaria prevention and control intervention, especially in Africa where LLINs and other vector control interventions are predicted to have averted 1.7 billion malaria cases and 10.6 million malaria deaths between 2000 and 2020 (WHO 2021).

Overview of LLINs

- In malaria-endemic regions LLINs represent the most widely used malaria prevention and control intervention, especially in Africa where LLINs and other vector control interventions are predicted to have averted 1.7 billion malaria cases and 10.6 million malaria deaths between 2000 and 2020 (WHO 2021).
- The materials commonly used for the nets vary from polyester to composites of high-density polyethylene and low-density polyethylene with a maximum lifespan of only around three years.

Overview of LLINs

- In malaria-endemic regions LLINs represent the most widely used malaria prevention and control intervention, especially in Africa where LLINs and other vector control interventions are predicted to have averted 1.7 billion malaria cases and 10.6 million malaria deaths between 2000 and 2020 (WHO 2021).
- The materials commonly used for the nets vary from polyester to composites of high-density polyethylene and low-density polyethylene with a maximum lifespan of only around three years.
- Many LLINs are still coated using technology that involves an insecticidal resin that binds to the surface of the net, which has proven to be effective (Kilian et al., 2008), while others are incorporated with the insecticide during fiber formation (Bhatt et al., 2015)


Overview of LLINs

- In malaria-endemic regions LLINs represent the most widely used malaria prevention and control intervention, especially in Africa where LLINs and other vector control interventions are predicted to have averted 1.7 billion malaria cases and 10.6 million malaria deaths between 2000 and 2020 (WHO 2021).
- The materials commonly used for the nets vary from polyester to composites of high-density polyethylene and low-density polyethylene with a maximum lifespan of only around three years.
- Many LLINs are still coated using technology that involves an insecticidal resin that binds to the surface of the net, which has proven to be effective (Kilian et al., 2008), while others are incorporated with the insecticide during fiber formation (Bhatt et al., 2015)


Components of a Long-Term Effectiveness Models

Pyrethroid-treated Nets (Olyset™)


Components of a Long-Term Effectiveness Models




Components of a Long-Term Effectiveness Models




Components of a Long-Term Effectiveness Models



Components of a Long-Term Effectiveness Models



Components of a Long-Term Effectiveness Models



Additional LLINs Brake et al. 2022

Summary of commercially available LLINs

Name	Material	Insecticide	Dose (g/kg)	Denier	Reference
World Health Organization qualified/prequalified					
Olyset Net	Polyethylene	Permethrin	20.0	150	Tarimo & Cosmas (2018); WHO (2020d)
Olyset Plus	Polyethylene	Permethrin	20.0	150	Oumbouke et al. (2019); WHO (2020d)
		PBO	10.0		
Veeralin	Polyethylene	Alpha-cypermethrin	6.0	130	Oumbouke et al. (2019); WHO (2020d)
		Piperonyl butoxide	2.2		
MAGnet	HDPE	Alpha-cypermethrin	5.8	150	Oumbouke et al. (2019)
Permanet 2.0	Polyester	Deltamethrin	1.8	75	Willis et al. (2013); WHO (2020d)
			1.4	100	
Interceptor	Polyester	Alpha-cypermethrin	6.7	75	Lissenden (2020); WHO (2020d)
Interceptor G2	Polyester	Alpha-cypermethrin	3.2	75	Bayili et al. (2017); Lissenden (2020); WHO (2020d)
			2.4	100	
		Chlorfenapyr	6.4	75	
			4.8	100	
Royal Sentry	HDPE	Alpha-cypermethrin	5.8	150	Lissenden (2020); WHO (2020d)
Royal Sentry 2.0	HDPE	Alpha-cypermethrin	5.8	120	Lissenden (2020); WHO (2020d)
Royal Guard	HDPE	Alpha-cypermethrin	5.5	120 or 150	Lissenden (2020); WHO (2020d)
		Pyriproxyfen	5.0		
Permanet 3.0	Polyester	Deltamethrin	Roof: 4.0	75 or 100	Tungu et al. (2010); Lissenden (2020); WHO (2020d)
			Side: 2.8	75	
			Side: 2.1	100	
		Piperonyl butoxide	Roof: 25	75 or 100	
Duranet	HDPE	Alpha-cypermethrin	5.8	150	Lissenden (2020); WHO (2020d)
Duranet Plus	HDPE	Alpha-cypermethrin	6.0	150	Lissenden (2020); WHO (2020d)
	LDPE	Piperonyl butoxide	2.2		
Miranet	HDPE	Alpha-cypermethrin	4.5	135	Lissenden (2020); WHO (2020d)
Yahe	Polyester	Deltamethrin	2.3	50	Lissenden (2020); WHO (2020d)
			1.85	75	
			1.4	100	
Safenet	Polyester	Alpha-cypermethrin	6.7	75	Lissenden (2020); WHO (2020d)
			5.0	100	
Yorkool	Polyester	Deltamethrin	1.8	75	Ketoh et al. (2018); Lissenden (2020)
			1.4	100	
Panda Net 2.0	Polyethylene	Deltamethrin	1.8	75	UNICEF (2020); WHO (2020d)
			1.4	100	
Tsara	Polyethylene	Deltamethrin	2.5	120	Lissenden (2020)
Tsara Boost	Polyethylene	Deltamethrin	3.0	130	Kasinathan et al. (2019); Lissenden (2020)
		Piperonyl butoxide	11.0		
Tsara Soft	Polyester	Deltamethrin	2.7	75	Lissenden (2020); WHO (2020d)
			2.0	100	
				150	
Reliefnet Reverte	Polyethylene	Deltamethrin	1.8	120	WHO (2021)
Tsara Plus	Polyester	Deltamethrin	Roof: 3.0; Sides: 2.5	Roof: 130; Sides: 100	Lissenden (2020); WHO (2020d)
		Piperonyl butoxide	Roof: 11.0		
Not approved					

Durability of LLINs

- The main factors for chemical durability are wash resistance, indicated by the retention index, and diffusion rate of the insecticide in the polymer matrix

Durability of LLINs

- The main factors for chemical durability are wash resistance, indicated by the retention index, and diffusion rate of the insecticide in the polymer matrix
- Retention index is given by

$$r = n \sqrt{\frac{t_n}{t_0}} \quad (1)$$

where t_n is the active ingredient content (g/kg) after n washes, and t_0 is the active ingredient content (g/kg) after 0 washes, and n the number of washes

Durability of LLINs

- The main factors for chemical durability are wash resistance, indicated by the retention index, and diffusion rate of the insecticide in the polymer matrix
- Retention index is given by

$$r = n \sqrt{\frac{t_n}{t_0}} \quad (1)$$

where t_n is the active ingredient content (g/kg) after n washes, and t_0 is the active ingredient content (g/kg) after 0 washes, and n the number of washes

- The diffusion rate of an insecticide in a polymer matrix is typically described by Fick's Law of Diffusion. Fick's Law is a fundamental equation governing diffusion, and it can be expressed as follows:

$$J = -D \frac{dC}{dx} \quad (2)$$

where J is the diffusion flux (the amount of substance crossing a unit area per unit time), D is the diffusion coefficient (a property of the material), dC is the concentration gradient of the substance, and dx is the distance over which the diffusion occurs.

Durability of LLINs

- The main factors for chemical durability are wash resistance, indicated by the retention index, and diffusion rate of the insecticide in the polymer matrix
- Retention index is given by

$$r = n\sqrt{\frac{t_n}{t_0}} \quad (1)$$

where t_n is the active ingredient content (g/kg) after n washes, and t_0 is the active ingredient content (g/kg) after 0 washes, and n the number of washes

- The diffusion rate of an insecticide in a polymer matrix is typically described by Fick's Law of Diffusion. Fick's Law is a fundamental equation governing diffusion, and it can be expressed as follows:

$$J = -D \frac{dC}{dx} \quad (2)$$

where J is the diffusion flux (the amount of substance crossing a unit area per unit time), D is the diffusion coefficient (a property of the material), dC is the concentration gradient of the substance, and dx is the distance over which the diffusion occurs.

- Physical durability of the nets is analyzed by the size of holes and tears found in the net after a monitored period of use by the proportional hole index.

Durability of LLINs

- The main factors for chemical durability are wash resistance, indicated by the retention index, and diffusion rate of the insecticide in the polymer matrix
- Retention index is given by

$$r = n\sqrt{\frac{t_n}{t_0}} \quad (1)$$

where t_n is the active ingredient content (g/kg) after n washes, and t_0 is the active ingredient content (g/kg) after 0 washes, and n the number of washes

- The diffusion rate of an insecticide in a polymer matrix is typically described by Fick's Law of Diffusion. Fick's Law is a fundamental equation governing diffusion, and it can be expressed as follows:

$$J = -D \frac{dC}{dx} \quad (2)$$

where J is the diffusion flux (the amount of substance crossing a unit area per unit time), D is the diffusion coefficient (a property of the material), dC is the concentration gradient of the substance, and dx is the distance over which the diffusion occurs.

- Physical durability of the nets is analyzed by the size of holes and tears found in the net after a monitored period of use by the proportional hole index.

Challenges and Opportunities

- LLINs are a cost-effective strategy due to the reduced cost of manufacturing and the cost of delivery models (Arroz et al., 2019).

Challenges and Opportunities

- LLINs are a cost-effective strategy due to the reduced cost of manufacturing and the cost of delivery models (Arroz et al., 2019).
- Distribution costs can be reduced by improving the overall product so that it remains effective and useable for longer periods.

Challenges and Opportunities

- LLINs are a cost-effective strategy due to the reduced cost of manufacturing and the cost of delivery models (Arroz et al., 2019).
- Distribution costs can be reduced by improving the overall product so that it remains effective and useable for longer periods.
- LLINs are produced with inexpensive material leading to the rapid loss of the physical integrity of nets and thereby generating additional costs to combat pyrethroid resistance.

Challenges and Opportunities

- LLINs are a cost-effective strategy due to the reduced cost of manufacturing and the cost of delivery models (Arroz et al., 2019).
- Distribution costs can be reduced by improving the overall product so that it remains effective and useable for longer periods.
- LLINs are produced with inexpensive material leading to the rapid loss of the physical integrity of nets and thereby generating additional costs to combat pyrethroid resistance.
- It was estimated that roughly 2.3 billion long-lasting insecticidal nets (LLINs) are disposed of in 2022 being part of a potential source of 1600 million tonnes of plastic waste generated every year (WHO 2021). Nevertheless, the World Health Organization (WHO) has recommended the proper ways of disposing or handling the old LLINs such as;

Challenges and Opportunities

- LLINs are a cost-effective strategy due to the reduced cost of manufacturing and the cost of delivery models (Arroz et al., 2019).
- Distribution costs can be reduced by improving the overall product so that it remains effective and useable for longer periods.
- LLINs are produced with inexpensive material leading to the rapid loss of the physical integrity of nets and thereby generating additional costs to combat pyrethroid resistance.
- It was estimated that roughly 2.3 billion long-lasting insecticidal nets (LLINs) are disposed of in 2022 being part of a potential source of 1600 million tonnes of plastic waste generated every year (WHO 2021). Nevertheless, the World Health Organization (WHO) has recommended the proper ways of disposing or handling the old LLINs such as;
 - 1 continue to be used even if they have holes until they are replaced with new LLINs

Challenges and Opportunities

- LLINs are a cost-effective strategy due to the reduced cost of manufacturing and the cost of delivery models (Arroz et al., 2019).
- Distribution costs can be reduced by improving the overall product so that it remains effective and useable for longer periods.
- LLINs are produced with inexpensive material leading to the rapid loss of the physical integrity of nets and thereby generating additional costs to combat pyrethroid resistance.
- It was estimated that roughly 2.3 billion long-lasting insecticidal nets (LLINs) are disposed of in 2022 being part of a potential source of 1600 million tonnes of plastic waste generated every year (WHO 2021). Nevertheless, the World Health Organization (WHO) has recommended the proper ways of disposing or handling the old LLINs such as;
 - 1 continue to be used even if they have holes until they are replaced with new LLINs
 - 2 not to dispose of the old LLINs in water bodies,

Challenges and Opportunities

- LLINs are a cost-effective strategy due to the reduced cost of manufacturing and the cost of delivery models (Arroz et al., 2019).
- Distribution costs can be reduced by improving the overall product so that it remains effective and useable for longer periods.
- LLINs are produced with inexpensive material leading to the rapid loss of the physical integrity of nets and thereby generating additional costs to combat pyrethroid resistance.
- It was estimated that roughly 2.3 billion long-lasting insecticidal nets (LLINs) are disposed of in 2022 being part of a potential source of 1600 million tonnes of plastic waste generated every year (WHO 2021). Nevertheless, the World Health Organization (WHO) has recommended the proper ways of disposing or handling the old LLINs such as;
 - 1 continue to be used even if they have holes until they are replaced with new LLINs
 - 2 not to dispose of the old LLINs in water bodies,
 - 3 collection of the old LLINs for disposal by National Malaria programs,

Challenges and Opportunities

- LLINs are a cost-effective strategy due to the reduced cost of manufacturing and the cost of delivery models (Arroz et al., 2019).
- Distribution costs can be reduced by improving the overall product so that it remains effective and useable for longer periods.
- LLINs are produced with inexpensive material leading to the rapid loss of the physical integrity of nets and thereby generating additional costs to combat pyrethroid resistance.
- It was estimated that roughly 2.3 billion long-lasting insecticidal nets (LLINs) are disposed of in 2022 being part of a potential source of 1600 million tonnes of plastic waste generated every year (WHO 2021). Nevertheless, the World Health Organization (WHO) has recommended the proper ways of disposing or handling the old LLINs such as;
 - 1 continue to be used even if they have holes until they are replaced with new LLINs
 - 2 not to dispose of the old LLINs in water bodies,
 - 3 collection of the old LLINs for disposal by National Malaria programs,
 - 4 incineration of old LLINs

Challenges and Opportunities

- LLINs are a cost-effective strategy due to the reduced cost of manufacturing and the cost of delivery models (Arroz et al., 2019).
- Distribution costs can be reduced by improving the overall product so that it remains effective and useable for longer periods.
- LLINs are produced with inexpensive material leading to the rapid loss of the physical integrity of nets and thereby generating additional costs to combat pyrethroid resistance.
- It was estimated that roughly 2.3 billion long-lasting insecticidal nets (LLINs) are disposed of in 2022 being part of a potential source of 1600 million tonnes of plastic waste generated every year (WHO 2021). Nevertheless, the World Health Organization (WHO) has recommended the proper ways of disposing or handling the old LLINs such as;
 - 1 continue to be used even if they have holes until they are replaced with new LLINs
 - 2 not to dispose of the old LLINs in water bodies,
 - 3 collection of the old LLINs for disposal by National Malaria programs,
 - 4 incineration of old LLINs
 - 5 formulating guidelines, policies, and regulations by the NMCPs in collaboration with national environmental authorities

Challenges and Opportunities

- LLINs are a cost-effective strategy due to the reduced cost of manufacturing and the cost of delivery models (Arroz et al., 2019).
- Distribution costs can be reduced by improving the overall product so that it remains effective and useable for longer periods.
- LLINs are produced with inexpensive material leading to the rapid loss of the physical integrity of nets and thereby generating additional costs to combat pyrethroid resistance.
- It was estimated that roughly 2.3 billion long-lasting insecticidal nets (LLINs) are disposed of in 2022 being part of a potential source of 1600 million tonnes of plastic waste generated every year (WHO 2021). Nevertheless, the World Health Organization (WHO) has recommended the proper ways of disposing or handling the old LLINs such as;
 - 1 continue to be used even if they have holes until they are replaced with new LLINs
 - 2 not to dispose of the old LLINs in water bodies,
 - 3 collection of the old LLINs for disposal by National Malaria programs,
 - 4 incineration of old LLINs
 - 5 formulating guidelines, policies, and regulations by the NMCPs in collaboration with national environmental authorities

Long-Term Effectiveness Models of LLITNs

- Long-term effectiveness models for LLINs aim to assess and predict the continued efficacy of these nets over an extended period. Here's an outline of key components that could be included in long-term effectiveness models of LLINs:

Long-Term Effectiveness Models of LLITNs

- Long-term effectiveness models for LLINs aim to assess and predict the continued efficacy of these nets over an extended period. Here's an outline of key components that could be included in long-term

effectiveness models of LLINs:

- 1 Net Longevity i.e. Assess the physical durability and lifespan of the LLIN material. It is possible to consider factors such as wear and tear, exposure to environmental conditions, and user behaviors that may impact the net's longevity.

Long-Term Effectiveness Models of LLITNs

- Long-term effectiveness models for LLINs aim to assess and predict the continued efficacy of these nets over an extended period. Here's an outline of key components that could be included in long-term effectiveness models of LLINs:
 - 1 Net Longevity i.e. Assess the physical durability and lifespan of the LLIN material. It is possible to consider factors such as wear and tear, exposure to environmental conditions, and user behaviors that may impact the net's longevity.
 - 2 Insecticide Persistence i.e. Model the decay or loss of insecticidal properties over time.

Long-Term Effectiveness Models of LLITNs

- Long-term effectiveness models for LLINs aim to assess and predict the continued efficacy of these nets over an extended period. Here's an outline of key components that could be included in long-term

effectiveness models of LLINs:

- 1 Net Longevity i.e. Assess the physical durability and lifespan of the LLIN material. It is possible to consider factors such as wear and tear, exposure to environmental conditions, and user behaviors that may impact the net's longevity.
- 2 Insecticide Persistence i.e. Model the decay or loss of insecticidal properties over time.
- 3 Entomological factor i.e. Incorporate mosquito behavior and population dynamics where it is possible to assess changes in mosquito species, resistance patterns, and feeding habits over the LLIN's lifespan.

Long-Term Effectiveness Models of LLITNs

- Long-term effectiveness models for LLINs aim to assess and predict the continued efficacy of these nets over an extended period. Here's an outline of key components that could be included in long-term

effectiveness models of LLINs:

- 1 Net Longevity i.e. Assess the physical durability and lifespan of the LLIN material. It is possible to consider factors such as wear and tear, exposure to environmental conditions, and user behaviors that may impact the net's longevity.
- 2 Insecticide Persistence i.e. Model the decay or loss of insecticidal properties over time.
- 3 Entomological factor i.e. Incorporate mosquito behavior and population dynamics where it is possible to assess changes in mosquito species, resistance patterns, and feeding habits over the LLIN's lifespan.
- 4 Disease Epidemiology i.e. Integrate malaria incidence data to correlate LLIN usage with disease prevention (Savi et al. 2024 (Bonus)).

Long-Term Effectiveness Models of LLITNs

- Long-term effectiveness models for LLINs aim to assess and predict the continued efficacy of these nets over an extended period. Here's an outline of key components that could be included in long-term

effectiveness models of LLINs:

- 1 Net Longevity i.e. Assess the physical durability and lifespan of the LLIN material. It is possible to consider factors such as wear and tear, exposure to environmental conditions, and user behaviors that may impact the net's longevity.
- 2 Insecticide Persistence i.e. Model the decay or loss of insecticidal properties over time.
- 3 Entomological factor i.e. Incorporate mosquito behavior and population dynamics where it is possible to assess changes in mosquito species, resistance patterns, and feeding habits over the LLIN's lifespan.
- 4 Disease Epidemiology i.e. Integrate malaria incidence data to correlate LLIN usage with disease prevention (Savi et al. 2024 (Bonus)).
- 5 Explore spatial heterogeneity in LLIN effectiveness within a given region (Savi et al. 2024 (Bonus)).

Long-Term Effectiveness Models of LLITNs

- Long-term effectiveness models for LLINs aim to assess and predict the continued efficacy of these nets over an extended period. Here's an outline of key components that could be included in long-term

effectiveness models of LLINs:

- 1 Net Longevity i.e. Assess the physical durability and lifespan of the LLIN material. It is possible to consider factors such as wear and tear, exposure to environmental conditions, and user behaviors that may impact the net's longevity.
- 2 Insecticide Persistence i.e. Model the decay or loss of insecticidal properties over time.
- 3 Entomological factor i.e. Incorporate mosquito behavior and population dynamics where it is possible to assess changes in mosquito species, resistance patterns, and feeding habits over the LLIN's lifespan.
- 4 Disease Epidemiology i.e. Integrate malaria incidence data to correlate LLIN usage with disease prevention (Savi et al. 2024 (Bonus)).
- 5 Explore spatial heterogeneity in LLIN effectiveness within a given region (Savi et al. 2024 (Bonus)).

Understanding Entomological Factors in LLIN Long-Term Effectiveness Models

- The presence of resistance in malaria vectors to the insecticides used in the LLINs makes it necessary not only to monitor the bio-efficacy of the nets with resistant mosquitoes but also to update the resistance status in endemic areas to obtain better results from control strategies.

Understanding Entomological Factors in LLIN Long-Term Effectiveness Models

- The presence of resistance in malaria vectors to the insecticides used in the LLINs makes it necessary not only to monitor the bio-efficacy of the nets with resistant mosquitoes but also to update the resistance status in endemic areas to obtain better results from control strategies.
- Assess the durability and bio-efficacy of the active ingredients of LLINs in the community;

Understanding Entomological Factors in LLIN Long-Term Effectiveness Models

- The presence of resistance in malaria vectors to the insecticides used in the LLINs makes it necessary not only to monitor the bio-efficacy of the nets with resistant mosquitoes but also to update the resistance status in endemic areas to obtain better results from control strategies.
- Assess the durability and bio-efficacy of the active ingredients of LLINs in the community;
- Examine whether entomological outcomes from adapted experimental hut trials can predict epidemiological and transmission outcomes.

Understanding Entomological Factors in LLIN Long-Term Effectiveness Models

- The presence of resistance in malaria vectors to the insecticides used in the LLINs makes it necessary not only to monitor the bio-efficacy of the nets with resistant mosquitoes but also to update the resistance status in endemic areas to obtain better results from control strategies.
- Assess the durability and bio-efficacy of the active ingredients of LLINs in the community;
- Examine whether entomological outcomes from adapted experimental hut trials can predict epidemiological and transmission outcomes.

Key Entomological Concepts

Some mosquito behaviors are relevant to LLIN effectiveness

Key Entomological Concepts

Some mosquito behaviors are relevant to LLIN effectiveness

- **Biting habits** refer to the fact that mosquitoes are most active from dusk to dawn. However, depending on the species, mosquitoes may also bite during the day. For example, *Aedes* mosquitoes feed during the day, while *Culex* mosquitoes feed mostly at night.

Key Entomological Concepts

Some mosquito behaviors are relevant to LLIN effectiveness

- **Biting habits** refer to the fact that mosquitoes are most active from dusk to dawn. However, depending on the species, mosquitoes may also bite during the day. For example, *Aedes* mosquitoes feed during the day, while *Culex* mosquitoes feed mostly at night.
- **Resting patterns** refer to the fact that Some *Anopheles* mosquitoes feed indoors, while others feed outdoors. Some *Anopheles* mosquitoes prefer to rest indoors after feeding, while others prefer to rest outdoors.

Key Entomological Concepts

Some mosquito behaviors are relevant to LLIN effectiveness

- **Biting habits** refer to the fact that mosquitoes are most active from dusk to dawn. However, depending on the species, mosquitoes may also bite during the day. For example, *Aedes* mosquitoes feed during the day, while *Culex* mosquitoes feed mostly at night.
- **Resting patterns** refer to the fact that Some *Anopheles* mosquitoes feed indoors, while others feed outdoors. Some *Anopheles* mosquitoes prefer to rest indoors after feeding, while others prefer to rest outdoors.
- **Feeding preferences** refer to the fact that some mosquitoes feed on plant nectar and other fluids from plants. Female mosquitoes also feed on blood for the development of their ovaries and eggs. When searching for blood hosts, some mosquitoes may have preferential behavior for certain species.

Key Entomological Concepts

Some mosquito behaviors are relevant to LLIN effectiveness

- **Biting habits** refer to the fact that mosquitoes are most active from dusk to dawn. However, depending on the species, mosquitoes may also bite during the day. For example, *Aedes* mosquitoes feed during the day, while *Culex* mosquitoes feed mostly at night.
- **Resting patterns** refer to the fact that Some *Anopheles* mosquitoes feed indoors, while others feed outdoors. Some *Anopheles* mosquitoes prefer to rest indoors after feeding, while others prefer to rest outdoors.
- **Feeding preferences** refer to the fact that some mosquitoes feed on plant nectar and other fluids from plants. Female mosquitoes also feed on blood for the development of their ovaries and eggs. When searching for blood hosts, some mosquitoes may have preferential behavior for certain species.
- **Anthrophagy**: species that feed on humans mainly.

Key Entomological Concepts

Some mosquito behaviors are relevant to LLIN effectiveness

- **Biting habits** refer to the fact that mosquitoes are most active from dusk to dawn. However, depending on the species, mosquitoes may also bite during the day. For example, *Aedes* mosquitoes feed during the day, while *Culex* mosquitoes feed mostly at night.
- **Resting patterns** refer to the fact that Some *Anopheles* mosquitoes feed indoors, while others feed outdoors. Some *Anopheles* mosquitoes prefer to rest indoors after feeding, while others prefer to rest outdoors.
- **Feeding preferences** refer to the fact that some mosquitoes feed on plant nectar and other fluids from plants. Female mosquitoes also feed on blood for the development of their ovaries and eggs. When searching for blood hosts, some mosquitoes may have preferential behavior for certain species.
- **Anthrophagy**: species that feed on humans mainly.
- **Contact irritant**: a chemical that stimulates mosquitoes to move away from the source after physical contact occurs.

Key Entomological Concepts

Some mosquito behaviors are relevant to LLIN effectiveness

- **Biting habits** refer to the fact that mosquitoes are most active from dusk to dawn. However, depending on the species, mosquitoes may also bite during the day. For example, *Aedes* mosquitoes feed during the day, while *Culex* mosquitoes feed mostly at night.
- **Resting patterns** refer to the fact that Some *Anopheles* mosquitoes feed indoors, while others feed outdoors. Some *Anopheles* mosquitoes prefer to rest indoors after feeding, while others prefer to rest outdoors.
- **Feeding preferences** refer to the fact that some mosquitoes feed on plant nectar and other fluids from plants. Female mosquitoes also feed on blood for the development of their ovaries and eggs. When searching for blood hosts, some mosquitoes may have preferential behavior for certain species.
- **Anthrophagy**: species that feed on humans mainly.
- **Contact irritant**: a chemical that stimulates mosquitoes to move away from the source after physical contact occurs.
- **Endophagy**: species that have a preference to feed indoors.

Key Entomological Concepts

Some mosquito behaviors are relevant to LLIN effectiveness

- **Biting habits** refer to the fact that mosquitoes are most active from dusk to dawn. However, depending on the species, mosquitoes may also bite during the day. For example, *Aedes* mosquitoes feed during the day, while *Culex* mosquitoes feed mostly at night.
- **Resting patterns** refer to the fact that Some *Anopheles* mosquitoes feed indoors, while others feed outdoors. Some *Anopheles* mosquitoes prefer to rest indoors after feeding, while others prefer to rest outdoors.
- **Feeding preferences** refer to the fact that some mosquitoes feed on plant nectar and other fluids from plants. Female mosquitoes also feed on blood for the development of their ovaries and eggs. When searching for blood hosts, some mosquitoes may have preferential behavior for certain species.
- **Anthrophagy**: species that feed on humans mainly.
- **Contact irritant**: a chemical that stimulates mosquitoes to move away from the source after physical contact occurs.
- **Endophagy**: species that have a preference to feed indoors.
- **Endophily**: an inherent tendency to rest indoors after feeding (mosquitoes may feed indoors or outdoors).

Key Entomological Concepts

Some mosquito behaviors are relevant to LLIN effectiveness

- **Biting habits** refer to the fact that mosquitoes are most active from dusk to dawn. However, depending on the species, mosquitoes may also bite during the day. For example, *Aedes* mosquitoes feed during the day, while *Culex* mosquitoes feed mostly at night.
- **Resting patterns** refer to the fact that Some *Anopheles* mosquitoes feed indoors, while others feed outdoors. Some *Anopheles* mosquitoes prefer to rest indoors after feeding, while others prefer to rest outdoors.
- **Feeding preferences** refer to the fact that some mosquitoes feed on plant nectar and other fluids from plants. Female mosquitoes also feed on blood for the development of their ovaries and eggs. When searching for blood hosts, some mosquitoes may have preferential behavior for certain species.
- **Anthrophagy**: species that feed on humans mainly.
- **Contact irritant**: a chemical that stimulates mosquitoes to move away from the source after physical contact occurs.
- **Endophagy**: species that have a preference to feed indoors.
- **Endophily**: an inherent tendency to rest indoors after feeding (mosquitoes may feed indoors or outdoors).
- **Exophagy**: species that have a preference to feed outdoors mainly.

Key Entomological Concepts

Some mosquito behaviors are relevant to LLIN effectiveness

- **Biting habits** refer to the fact that mosquitoes are most active from dusk to dawn. However, depending on the species, mosquitoes may also bite during the day. For example, *Aedes* mosquitoes feed during the day, while *Culex* mosquitoes feed mostly at night.
- **Resting patterns** refer to the fact that Some *Anopheles* mosquitoes feed indoors, while others feed outdoors. Some *Anopheles* mosquitoes prefer to rest indoors after feeding, while others prefer to rest outdoors.
- **Feeding preferences** refer to the fact that some mosquitoes feed on plant nectar and other fluids from plants. Female mosquitoes also feed on blood for the development of their ovaries and eggs. When searching for blood hosts, some mosquitoes may have preferential behavior for certain species.
- **Anthrophagy**: species that feed on humans mainly.
- **Contact irritant**: a chemical that stimulates mosquitoes to move away from the source after physical contact occurs.
- **Endophagy**: species that have a preference to feed indoors.
- **Endophily**: an inherent tendency to rest indoors after feeding (mosquitoes may feed indoors or outdoors).
- **Exophagy**: species that have a preference to feed outdoors mainly.
- **Exophily**: species that have a preference to rest outdoors mainly.

Key Entomological Concepts

Some mosquito behaviors are relevant to LLIN effectiveness

- **Biting habits** refer to the fact that mosquitoes are most active from dusk to dawn. However, depending on the species, mosquitoes may also bite during the day. For example, *Aedes* mosquitoes feed during the day, while *Culex* mosquitoes feed mostly at night.
- **Resting patterns** refer to the fact that Some *Anopheles* mosquitoes feed indoors, while others feed outdoors. Some *Anopheles* mosquitoes prefer to rest indoors after feeding, while others prefer to rest outdoors.
- **Feeding preferences** refer to the fact that some mosquitoes feed on plant nectar and other fluids from plants. Female mosquitoes also feed on blood for the development of their ovaries and eggs. When searching for blood hosts, some mosquitoes may have preferential behavior for certain species.
- **Anthrophagy**: species that feed on humans mainly.
- **Contact irritant**: a chemical that stimulates mosquitoes to move away from the source after physical contact occurs.
- **Endophagy**: species that have a preference to feed indoors.
- **Endophily**: an inherent tendency to rest indoors after feeding (mosquitoes may feed indoors or outdoors).
- **Exophagy**: species that have a preference to feed outdoors mainly.
- **Exophily**: species that have a preference to rest outdoors mainly.

Key Entomological concepts - 2

- **Spatial repellent/deterrent:** a chemical that stimulates mosquitoes to move away from the source without the need for physical contact.

Key Entomological concepts - 2

- **Spatial repellent/deterrent:** a chemical that stimulates mosquitoes to move away from the source without the need for physical contact.
- **Toxic chemical action:** knockdown or death of mosquitoes after physical contact with the chemical.

Key Entomological concepts - 2

- **Spatial repellent/deterrent:** a chemical that stimulates mosquitoes to move away from the source without the need for physical contact.
- **Toxic chemical action:** knockdown or death of mosquitoes after physical contact with the chemical.
- **Vectorial capacity (VC):** the total number of infectious mosquito bites on humans that will arise from a single infected person on a single day. VC is a measure used in epidemiology to assess the potential for a vector population to transmit a pathogen to humans. It is often associated with the transmission of vector-borne diseases such as malaria. The basic formula for calculating vectorial capacity is:

$$VC = \frac{ma^2 p^n}{-ln(p)} \quad (3)$$

where m : The number of female mosquitoes per person. It quantifies the density of mosquitoes in the population; a : The human biting rate, representing the average number of bites a mosquito takes on a human per day; p : The daily survival rate of mosquitoes. It denotes the probability that a mosquito survives each day; n : The extrinsic incubation period, referring to the time it takes for the pathogen to develop inside the mosquito before it can be transmitted.

Key Entomological concepts - 2

- **Spatial repellent/deterrent:** a chemical that stimulates mosquitoes to move away from the source without the need for physical contact.
- **Toxic chemical action:** knockdown or death of mosquitoes after physical contact with the chemical.
- **Vectorial capacity (VC):** the total number of infectious mosquito bites on humans that will arise from a single infected person on a single day. VC is a measure used in epidemiology to assess the potential for a vector population to transmit a pathogen to humans. It is often associated with the transmission of vector-borne diseases such as malaria. The basic formula for calculating vectorial capacity is:

$$VC = \frac{ma^2 p^n}{-ln(p)} \quad (3)$$

where m : The number of female mosquitoes per person. It quantifies the density of mosquitoes in the population; a : The human biting rate, representing the average number of bites a mosquito takes on a human per day; p : The daily survival rate of mosquitoes. It denotes the probability that a mosquito survives each day; n : The extrinsic incubation period, referring to the time it takes for the pathogen to develop inside the mosquito before it can be transmitted.

- **Zoophagy:** species that feed on animals mainly.

Key Entomological concepts - 2

- **Spatial repellent/deterrent:** a chemical that stimulates mosquitoes to move away from the source without the need for physical contact.
- **Toxic chemical action:** knockdown or death of mosquitoes after physical contact with the chemical.
- **Vectorial capacity (VC):** the total number of infectious mosquito bites on humans that will arise from a single infected person on a single day. VC is a measure used in epidemiology to assess the potential for a vector population to transmit a pathogen to humans. It is often associated with the transmission of vector-borne diseases such as malaria. The basic formula for calculating vectorial capacity is:

$$VC = \frac{ma^2 p^n}{-ln(p)} \quad (3)$$

where m : The number of female mosquitoes per person. It quantifies the density of mosquitoes in the population; a : The human biting rate, representing the average number of bites a mosquito takes on a human per day; p : The daily survival rate of mosquitoes. It denotes the probability that a mosquito survives each day; n : The extrinsic incubation period, referring to the time it takes for the pathogen to develop inside the mosquito before it can be transmitted.

- **Zoophagy:** species that feed on animals mainly.

Key Entomological concept - End

- **Entomological Inoculation Rate (EIR)**: is a measure used in epidemiology to estimate the average number of infectious mosquito bites that a person receives in a specific period, typically per year. It is a critical parameter in understanding the transmission dynamics of vector-borne diseases, such as malaria. The formula for calculating EIR is as follows:

$$EIR = \frac{a \times m \times b \times p}{s} \quad (4)$$

where a : The human biting rate, indicating the average number of mosquito bites on a person per night; m : The sporozoite rate, which is the proportion of mosquitoes in the population that are infected with the pathogen and capable of transmitting it; b : The probability that a mosquito bite results in a transmission event (probability of transmission per bite); p : The daily survival rate of mosquitoes, representing the probability that a mosquito survives each day; s : The duration of the sporogonic cycle, which is the time it takes for the pathogen to develop and become infective inside the mosquito.

- 1 Introduction to Long-Term Effectiveness Models
- 2 Setting Up the Environment**
- 3 Exploratory Data Analysis
- 4 Building Cox Proportional Hazards Model

Initiation to R

- Gentle initiation to R
- What else can R be used for?
- Let's get our hands dirty

Crunching Numbers

- **Step 1:** Load a dataset

Crunching Numbers

- **Step 1:** Load a dataset
`data(mtcars)`
- **Step 2:** View the dataset.

Crunching Numbers

- **Step 1:** Load a dataset
`data(mtcars)`
- **Step 2:** View the dataset.
`head(mtcars)`
- **Step 3:** Create a simple plot.

Crunching Numbers

- **Step 1:** Load a dataset

```
data(mtcars)
```

- **Step 2:** View the dataset.

```
head(mtcars)
```

- **Step 3:** Create a simple plot.

```
plot(mtcars$mpg, mtcars$hp, main = "Miles per Gallon vs Horsepower", xlab = "Miles per Gallon",
```


Computation of VC and EIR

- Hands -on

1 Introduction to Long-Term Effectiveness Models

2 Setting Up the Environment

3 Exploratory Data Analysis

Experiment overview

Hypothesis testing

Prediction

4 Building Cox Proportional Hazards Model

1 Introduction to Long-Term Effectiveness Models

2 Setting Up the Environment

3 Exploratory Data Analysis

Experiment overview

Hypothesis testing

Prediction


4 Building Cox Proportional Hazards Model

Data stewardship

- Dr. Konkon Alphonse

Data stewardship


- Dr. Konkon Alphonse



- Dr. Zougbedji David

Data stewardship

• Dr. Konkon Alphonse



• Dr. Zougbedji David



Application using real-world data

- The study aims to investigate the effectiveness of various malaria interventions and treatments in reducing mortality rates among affected populations.

Application using real-world data

- The study aims to investigate the effectiveness of various malaria interventions and treatments in reducing mortality rates among affected populations.
- **Objective:** The primary objective of this study is to evaluate the impact of different treatments on mortality rates associated with malaria. We will specifically focus on understanding how distinct interventions, represented by various treatments, influence the outcomes of individuals over time.

Application using real-world data

- The study aims to investigate the effectiveness of various malaria interventions and treatments in reducing mortality rates among affected populations.
- **Objective:** The primary objective of this study is to evaluate the impact of different treatments on mortality rates associated with malaria. We will specifically focus on understanding how distinct interventions, represented by various treatments, influence the outcomes of individuals over time.
- **Data Source:** Our analysis draws upon a comprehensive dataset encompassing variables such as treatment types, mortality rates, and other relevant factors. The dataset spans a specified period and includes information from diverse regions, providing a nuanced understanding of the effectiveness of interventions.

Application using real-world data

- The study aims to investigate the effectiveness of various malaria interventions and treatments in reducing mortality rates among affected populations.
- **Objective:** The primary objective of this study is to evaluate the impact of different treatments on mortality rates associated with malaria. We will specifically focus on understanding how distinct interventions, represented by various treatments, influence the outcomes of individuals over time.
- **Data Source:** Our analysis draws upon a comprehensive dataset encompassing variables such as treatment types, mortality rates, and other relevant factors. The dataset spans a specified period and includes information from diverse regions, providing a nuanced understanding of the effectiveness of interventions.
- Detailed analyses are accessible here

Application using real-world data

- The study aims to investigate the effectiveness of various malaria interventions and treatments in reducing mortality rates among affected populations.
- **Objective:** The primary objective of this study is to evaluate the impact of different treatments on mortality rates associated with malaria. We will specifically focus on understanding how distinct interventions, represented by various treatments, influence the outcomes of individuals over time.
- **Data Source:** Our analysis draws upon a comprehensive dataset encompassing variables such as treatment types, mortality rates, and other relevant factors. The dataset spans a specified period and includes information from diverse regions, providing a nuanced understanding of the effectiveness of interventions.
- Detailed analyses are accessible here

1 Introduction to Long-Term Effectiveness Models

2 Setting Up the Environment

3 Exploratory Data Analysis

Experiment overview

Hypothesis testing

Prediction

4 Building Cox Proportional Hazards Model

Efficacy testing

- H_0 : The treated nets significantly increase the number of dead mosquitoes compared to untreated nets.

Efficacy testing

- H0: The treated nets significantly increase the number of dead mosquitoes compared to untreated nets.
- Traditionally ANOVA is used to analyze such a dataset but often basic assumptions (normal population distribution; same variance; independent) are not met.

Efficacy testing

- H_0 : The treated nets significantly increase the number of dead mosquitoes compared to untreated nets.
- Traditionally ANOVA is used to analyze such a dataset but often basic assumptions (normal population distribution; same variance; independent) are not met.
- Using a generalized linear model (GLM) with a Poisson or negative binomial family would be appropriate. The Poisson distribution is often used for counting data, but if there's overdispersion (variance greater than the mean), a negative binomial distribution may be more suitable.

Efficacy testing

- H_0 : The treated nets significantly increase the number of dead mosquitoes compared to untreated nets.
- Traditionally ANOVA is used to analyze such a dataset but often basic assumptions (normal population distribution; same variance; independent) are not met.
- Using a generalized linear model (GLM) with a Poisson or negative binomial family would be appropriate. The Poisson distribution is often used for counting data, but if there's overdispersion (variance greater than the mean), a negative binomial distribution may be more suitable.
- Nonetheless, we will access the most fitted testing for models of the Poisson family namely Poisson regression, Negative Binomial regression, and Zero-Inflated Poisson or Negative Binomial models.

Efficacy testing

- H_0 : The treated nets significantly increase the number of dead mosquitoes compared to untreated nets.
- Traditionally ANOVA is used to analyze such a dataset but often basic assumptions (normal population distribution; same variance; independent) are not met.
- Using a generalized linear model (GLM) with a Poisson or negative binomial family would be appropriate. The Poisson distribution is often used for counting data, but if there's overdispersion (variance greater than the mean), a negative binomial distribution may be more suitable.
- Nonetheless, we will access the most fitted testing for models of the Poisson family namely Poisson regression, Negative Binomial regression, and Zero-Inflated Poisson or Negative Binomial models.
- Further details starting from line 123

Efficacy testing

- H_0 : The treated nets significantly increase the number of dead mosquitoes compared to untreated nets.
- Traditionally ANOVA is used to analyze such a dataset but often basic assumptions (normal population distribution; same variance; independent) are not met.
- Using a generalized linear model (GLM) with a Poisson or negative binomial family would be appropriate. The Poisson distribution is often used for counting data, but if there's overdispersion (variance greater than the mean), a negative binomial distribution may be more suitable.
- Nonetheless, we will access the most fitted testing for models of the Poisson family namely Poisson regression, Negative Binomial regression, and Zero-Inflated Poisson or Negative Binomial models.
- Further details starting from line 123

1 Introduction to Long-Term Effectiveness Models

2 Setting Up the Environment

3 Exploratory Data Analysis

Experiment overview

Hypothesis testing

Prediction

4 Building Cox Proportional Hazards Model

Prediction

- Here we investigate determinants of mortality
- Detail is available here

- 1 Introduction to Long-Term Effectiveness Models
- 2 Setting Up the Environment
- 3 Exploratory Data Analysis
- 4 Building Cox Proportional Hazards Model**

Concept

- The Cox Proportional Hazards model helps you understand how different factors influence the "hazard" of the event happening over time.

Concept

- The Cox Proportional Hazards model helps you understand how different factors influence the "hazard" of the event happening over time.
- **Hazard:** In survival analysis, "hazard" refers to the risk of an event occurring at any given time. It's like the probability of an event happening in a small time interval.

Concept

- The Cox Proportional Hazards model helps you understand how different factors influence the "hazard" of the event happening over time.
- **Hazard:** In survival analysis, "hazard" refers to the risk of an event occurring at any given time. It's like the probability of an event happening in a small time interval.
- **Proportional Hazards:** The key idea of the Cox model is that the hazard of an event for any individual is a constant multiple of the hazard for any other individual, and this constant is the same over time. In other words, the ratio of hazards between two individuals remains constant over time.

Concept

- The Cox Proportional Hazards model helps you understand how different factors influence the "hazard" of the event happening over time.
- **Hazard:** In survival analysis, "hazard" refers to the risk of an event occurring at any given time. It's like the probability of an event happening in a small time interval.
- **Proportional Hazards:** The key idea of the Cox model is that the hazard of an event for any individual is a constant multiple of the hazard for any other individual, and this constant is the same over time. In other words, the ratio of hazards between two individuals remains constant over time.
- **Covariates:** The model considers covariates or factors that might affect the hazard. These could be things like different treatments, ages, or any other relevant characteristic of the individuals being studied.

Concept

- The Cox Proportional Hazards model helps you understand how different factors influence the "hazard" of the event happening over time.
- **Hazard:** In survival analysis, "hazard" refers to the risk of an event occurring at any given time. It's like the probability of an event happening in a small time interval.
- **Proportional Hazards:** The key idea of the Cox model is that the hazard of an event for any individual is a constant multiple of the hazard for any other individual, and this constant is the same over time. In other words, the ratio of hazards between two individuals remains constant over time.
- **Covariates:** The model considers covariates or factors that might affect the hazard. These could be things like different treatments, ages, or any other relevant characteristic of the individuals being studied.
- **Regression Analysis:** The Cox model uses a form of regression analysis to estimate how each covariate influences the hazard. It provides hazard ratios, which tell you how much more or less likely an event is to occur based on changes in the covariates.

Concept

- The Cox Proportional Hazards model helps you understand how different factors influence the "hazard" of the event happening over time.
- **Hazard:** In survival analysis, "hazard" refers to the risk of an event occurring at any given time. It's like the probability of an event happening in a small time interval.
- **Proportional Hazards:** The key idea of the Cox model is that the hazard of an event for any individual is a constant multiple of the hazard for any other individual, and this constant is the same over time. In other words, the ratio of hazards between two individuals remains constant over time.
- **Covariates:** The model considers covariates or factors that might affect the hazard. These could be things like different treatments, ages, or any other relevant characteristic of the individuals being studied.
- **Regression Analysis:** The Cox model uses a form of regression analysis to estimate how each covariate influences the hazard. It provides hazard ratios, which tell you how much more or less likely an event is to occur based on changes in the covariates.
- Example

Thank You