

Introduction to Biophysics II: Quantum physics & biology

Lecture XX: Circular dichroism

Circular polarization
Circular dichroism (CD)
Molecular symmetry, handedness, relevance to biology
Magnetic transition dipole
Absorption of right and left circularly polarized light, rotational strength
- Interaction with magnetic and electric transition dipoles
- Excitonic circular dichroism
Making circularly polarized light
CD spectrometer design

Reading:
Modern optical spectroscopy, chapter 10 (the 'green' book)
Excitons, van Amerongen et al.

by S.Savikhin, Purdue University

1

Circularly polarized light

Left-handed (counter-clockwise)

Right-handed (clockwise)

+: left
-: right

$$\vec{E}^{\pm}(t, z) = E_0 \left[\cos(\omega t - kz) \hat{x} \pm \sin(\omega t - kz) \hat{y} \right]$$

$$\omega = \frac{2\pi c}{\lambda} \quad k = \frac{2\pi}{\lambda}$$

Linear polarization:

$$\vec{E}(t, z) = E_0 \cos(\omega t - kz) \hat{x}$$

$$\vec{B}(t, z) = B_0 \cos(\omega t - kz) \hat{y}$$

$$B_0 = E_0 / c$$

by S.Savikhin, Purdue University

2

Circularly polarized light

Linear: $\vec{E}_{lin}(t, z) = E_0 \cos(\omega t - kz) \hat{x}$ $\omega = \frac{2\pi c}{\lambda}$ $k = \frac{2\pi}{\lambda}$

+ : Left $\vec{E}^+(t, z) = E_0 [\cos(\omega t - kz) \hat{x} \pm \sin(\omega t - kz) \hat{y}]$
 - : Right

In this representation, the energy density (intensity) of circularly polarized light is twice that of linearly polarized light:

$$u_{lin} = \frac{1}{2} \epsilon_0 E_0^2 \quad u_{circ} = \epsilon_0 E_0^2$$

Often, an alternative equation is used where the energy density of circularly polarized light is the same as that of linearly polarized light:

$$\vec{E}^\pm(t, z) = \frac{1}{\sqrt{2}} E_0 [\cos(\omega t - kz) \hat{x} \pm \sin(\omega t - kz) \hat{y}] \Rightarrow u_{circ} = \frac{1}{2} \epsilon_0 E_0^2$$

Alternatively, using complex space (math simplifies):

$$\begin{cases} \vec{E}^\pm = E_0 (\hat{x} \pm i\hat{y}) e^{-i\omega t} \\ \vec{B}^\pm = \frac{1}{c} \hat{z} \times \vec{E}^\pm = \mp \frac{i}{c} \vec{E}^\pm \end{cases} \quad \text{OR:} \quad \begin{cases} \vec{E}^\pm = \frac{1}{\sqrt{2}} E_0 (\hat{x} \pm i\hat{y}) e^{-i\omega t} \\ \vec{B}^\pm = \frac{1}{c} \hat{z} \times \vec{E}^\pm = \mp \frac{i}{c} \vec{E}^\pm \end{cases}$$

Note: linear polarization: $\vec{E}_{lin} = \frac{1}{\sqrt{2}} (\vec{E}^+ + \vec{E}^-) = E_0 \hat{x} e^{i(kz - \omega t)}$

by S.Savikhin, Purdue University

3

Circular dichroism (CD)

Circular Dichroism: Absorption of circularly polarized light depends on the handedness of light, L (left) or R (right)

$$A_L \equiv A^+ = A + \Delta A$$

A: Conventional absorption

$$A_R \equiv A^- = A - \Delta A$$

ΔA : Change due to circular polarization

CD in optical density units: $\Delta A_{CD} = A_L - A_R = 2\Delta A$ It depends on the wavelength

Molar extinction CD: $\Delta \epsilon_{CD}(\lambda) = \epsilon_L(\lambda) - \epsilon_R(\lambda) = \Delta A_{CD} / (cl)$ c – molar concentration
 l – optical thickness

CD measured as ellipticity in degrees:

Recall linear polarization: $\vec{E}_{lin} = \frac{1}{\sqrt{2}} (\vec{E}^+ + \vec{E}^-) = E_0 \hat{x} e^{i(kz - \omega t)}$

- Linear polarization is an equal superposition of L and R-polarized light

- The sample absorbs one more than the other one

→ After the sample polarization is elliptical

$$\begin{cases} E_x(z, t) = E_0 \cos \theta \cos(kz - \omega t) \\ E_y(z, t) = -E_0 \sin \theta \cos(kz - \omega t) \end{cases}$$

Ellipticity θ :

For small θ (and ΔA)

Ellipticity: ↑

$$\tan \theta = \frac{E_R - E_L}{E_R + E_L} \approx \theta [\text{rad}] \approx \ln(10) \frac{\Delta A_{CD}}{4}$$

Unit conversion



$$\theta [\text{mdeg}] \approx 33 \times 10^3 \Delta A_{CD}$$

by S.Savikhin, Purdue University


4

The origin of CD (chirality)

Two enantiomers of a generic amino acid that is chiral (handedness)



Alpha helix



Most amino acids have two enantiomers:
Left-handed: denoted S (Sinistral) or L (Left)
Right-handed: denoted D (Dextral) or R (Right)

Natural amino acids are mostly of L-type; D-type is rare.

Alpha helices in proteins, as well as DNA, are almost always right-handed

Circular Dichroism: Absorption of circularly polarized light depends on the handedness of light and molecules

by S.Savikhin, Purdue University



5

Examples of Circular Dichroism Spectra

The difference in absorption of left and right circularly polarized light

$$\Delta \epsilon_{CD}(\lambda) = \ln(10) \frac{\Delta A_{CD}}{4} / (cl)$$

$$\Delta A_{CD} = A_L - A_R$$




<https://www.bif.mit.edu/new-page-1-1>

CD spectra of proteins with defined secondary structure (left) and CD spectra of representative proteins with varying conformations (right).
Greenfield, N. *Nature Protocols*, p. 2876, v. 1, 2007.

by S.Savikhin, Purdue University



6

Example: using CD signals to study proteins



PrP: A major prion protein in the nervous system
It is crucial for neuron health and has been linked to synaptic formation and function.

Prion diseases occur PrP^C transforms into an abnormal, pathogenic isoform PrP^{Sc}, forming infectious amyloid fibrils that cause neurodegeneration, such as Creutzfeldt-Jakob disease (CJD) in humans and scrapie in animals.



by S.Savikhin, Purdue University

7

Accounting for the magnetic dipole moment

The intrinsic CD of a molecule is the result of the interaction of a molecule with both the electric and magnetic fields of a light electromagnetic wave

Absorption of light, earlier lecture: interaction with light $\tilde{H}'(t) = -\mathbf{E}(t) \cdot \tilde{\mu}$

Light also has a magnetic field component $B_0 = \frac{E_0}{c}$

The *electric* field interacts with the *electric* transition dipole
The *magnetic* field interacts with the *magnetic* transition dipole

$$\tilde{H}'(t) = -\mathbf{E}(t) \cdot \tilde{\mu} - \mathbf{B}(t) \cdot \tilde{\mathbf{m}}$$

↑ Magnetic moment operator

A magnetic moment stems from the orbital motion of charges and spins


by S.Savikhin, Purdue University

8


Magnetic (dipole) moment

$$\tilde{H}'(t) = -\mathbf{E}(t) \cdot \tilde{\mu} - \mathbf{B}(t) \cdot \tilde{\mathbf{m}}$$

Magnetic field of a molecule due to orbital motion of electrons (Bohr model):



(a) Current-carrying loop



(b) Hydrogen atom

$$I = \frac{e}{T} = \frac{ev}{2\pi r}$$

$$A = \pi r^2$$

$$m = IA = \frac{ev}{2\pi r} \pi r^2 = \frac{evr}{2}$$

$$\text{SI: } \tilde{\mathbf{m}} = \frac{-e}{2} (\mathbf{v} \times \mathbf{r}) = \frac{-e}{2m_e} (\mathbf{p} \times \mathbf{r})$$

$$\text{CGS: } \tilde{\mathbf{m}} = \frac{-e}{2m_e c} (\mathbf{p} \times \mathbf{r})$$

Magnetic moment: μ
determines the torque an object
experiences in an external magnetic field

Magnetic moment operator (CGS):

$$\tilde{\mathbf{m}} = \frac{e}{2m_e} (\mathbf{r} \times \mathbf{p} + g_e \mathbf{S})$$

Angular momentum: $\tilde{\mathbf{L}} = \mathbf{r} \times \mathbf{p}$

Electron *g-factor* = 2.00232

Electron spin: negligible contribution
to transition dipole
→ ignore

Linear momentum operator: $\tilde{\mathbf{p}} = -i\hbar \nabla$

by S.Savikhin, Purdue University

9

Magnetic moment-driven transitions

$$\tilde{H}'(t) = -\mathbf{E}(t) \cdot \tilde{\mu} - \mathbf{B}(t) \cdot \tilde{\mathbf{m}}$$

Transition between states *a* and *b*: need matrix elements:

$$\tilde{H}'_{ba} = -\vec{E} \cdot \langle \psi_b | \tilde{\mu} | \psi_a \rangle - \vec{B} \cdot \langle \psi_b | \tilde{\mathbf{m}} | \psi_a \rangle$$

	Electric	Magnetic
Transition matrix element $\tilde{H}'_{ba} =$	$-\vec{E} \cdot \langle \psi_b \tilde{\mu} \psi_a \rangle$	$-\vec{B} \cdot \langle \psi_b \tilde{\mathbf{m}} \psi_a \rangle$
Transition probability	$\propto \vec{E}_0 \cdot \tilde{\mu}_{ba} ^2$	$\propto \vec{B} \cdot \tilde{\mathbf{m}}_{ba} ^2$
Transition dipole moment	$\tilde{\mu}_{ba} \equiv \langle \psi_b \tilde{\mu} \psi_a \rangle$	$\tilde{\mathbf{m}}_{ba} \equiv \langle \psi_b \tilde{\mathbf{m}} \psi_a \rangle$

See lectures 3-4

By analogy

Note: typically, $|\tilde{\mathbf{m}}_{ba}| \ll |\tilde{\mu}_{ba}|$

The absorption strength driven by the magnetic component of light is about 2-4 orders of magnitude weaker than that driven by the electric component

by S.Savikhin, Purdue University

10

Magnetic transition dipole moment

Consider the transition from ψ_a to ψ_b

$$\tilde{m} = \frac{e}{2m_e} \vec{r} \times (-i\hbar \tilde{\nabla})$$

Find: $\vec{m}_{ba} \equiv \langle \psi_b | \tilde{m} | \psi_a \rangle$

$$\vec{m}_{ba} = \langle \psi_b | \tilde{m} | \psi_a \rangle = \left\langle \psi_b \left| \frac{e}{2m_e} \vec{r} \times (-i\hbar \tilde{\nabla}) \right| \psi_a \right\rangle$$

Magnetic transition dipole moment

$$\vec{m}_{ba} = -i \frac{e\hbar}{2m_e} \langle \psi_b | \vec{r} \times \tilde{\nabla} | \psi_a \rangle$$

Magnetic dipole moment is imaginary!

Note: SI units, CGS: divide by the speed of light c

Or: $\vec{m}_{ba} = -im_B \langle \psi_b | \vec{r} \times \tilde{\nabla} | \psi_a \rangle$ Bohr magneton: $m_B \equiv \frac{e\hbar}{2m_e}$

Transition probability: $\propto |\vec{B} \cdot \vec{m}_{ba}|^2$ m_B - magnetic dipole moment of the lowest-energy orbital of the hydrogen atom

HOMO (ψ_h) to LUMO (ψ_l) with 1 electron on HOME orbital:

$$\vec{m}_{ba} = -i \frac{e\hbar}{2m_e} \langle \psi_l | \vec{r} \times \tilde{\nabla} | \psi_h \rangle$$

by S.Savikhin, Purdue University

11

Magnetic transition moment, HOMO → LUMO

Consider the transition from HOMO (ψ_h) to LUMO (ψ_l) with 2 electrons on HOMO

Ground state $\Psi_a = \psi_h(1)\psi_h(2)$

Excited state $\Psi_b = 2^{-1/2} [\psi_h(1)\psi_l(2) + \psi_h(2)\psi_l(1)]$

Since electrons are indistinguishable, use the Slater-type orbital for the excited state

Find: $\vec{m}_{ba} \equiv \langle \Psi_b | \vec{m} | \Psi_a \rangle$ $\vec{m} = \vec{m}_1 + \vec{m}_2$ (2 electrons)

$$\vec{m}_{ba} = 2^{-1/2} \langle \psi_h(1)\psi_l(2) + \psi_h(2)\psi_l(1) | \vec{m}_1 + \vec{m}_2 | \psi_h(1)\psi_h(2) \rangle$$

$$\vec{m} = \frac{e}{2m_e} \vec{r} \times (-i\hbar \tilde{\nabla})$$

$$\vec{m}_{ba} = 2^{-1/2} \left[\langle \psi_h(1)\psi_l(2) | \vec{m}_1 + \vec{m}_2 | \psi_h(1)\psi_h(2) \rangle + \langle \psi_h(2)\psi_l(1) | \vec{m}_1 + \vec{m}_2 | \psi_h(1)\psi_h(2) \rangle \right]$$

$$\begin{aligned} \text{BLUE} &= \langle \psi_h(1)\psi_l(2) | \vec{m}_1 | \psi_h(1)\psi_h(2) \rangle + \langle \psi_h(1)\psi_l(2) | \vec{m}_2 | \psi_h(1)\psi_h(2) \rangle \\ &= \langle \psi_h(1) | \vec{m}_1 | \psi_h(1) \rangle \langle \psi_l(2) | \psi_h(2) \rangle + \langle \psi_h(1) | \psi_h(1) \rangle \langle \psi_l(2) | \vec{m}_2 | \psi_h(2) \rangle \\ &= \langle \psi_l(2) | \vec{m}_2 | \psi_h(2) \rangle \quad \leftarrow =0 \quad \leftarrow =1 \\ \text{RED} &= \langle \psi_l(1) | \vec{m}_1 | \psi_h(1) \rangle \end{aligned}$$

Blue and red are equal:
Only integration variable changes

$$\vec{m}_{ba} = 2^{-1/2} \cdot 2 \langle \psi_l | \vec{m} | \psi_h \rangle = 2^{1/2} \left\langle \psi_l \left| \frac{e}{2m_e} \vec{r} \times (-i\hbar \tilde{\nabla}) \right| \psi_h \right\rangle = -i \frac{\sqrt{2}e\hbar}{2m_e} \langle \psi_l | \vec{r} \times \tilde{\nabla} | \psi_h \rangle$$

$$\vec{m}_{ba} = -i \frac{\sqrt{2}e\hbar}{2m_e} \langle \psi_l | \vec{r} \times \tilde{\nabla} | \psi_h \rangle$$

Transition strength $|m_{ba}|^2$ is twice that of HOME → LUMO with 1 electron

by S.Savikhin, Purdue University

12

Magnetic transition dipole moment

CAN SKIP

$$\vec{m}_{ba} = -i \frac{\sqrt{2}\hbar e}{2m_e} \langle \psi_l | \vec{r} \times \vec{\nabla} | \psi_h \rangle$$

Consider a molecule with an orbital constructed as a linear combination of $2p$ orbitals



$$\text{HOMO: } \psi_h = \sum_t C_t^h p_t$$

$$\text{LUMO: } \psi_l = \sum_s C_s^l p_s$$

$$\vec{m}_{ba} = -i \frac{\sqrt{2}\hbar e}{2m_e} \left\langle \sum_s C_s^l p_s \left| \vec{r} \times \vec{\nabla} \right| \sum_t C_t^h p_t \right\rangle$$

$$\vec{m}_{ba} = -i \frac{\sqrt{2}\hbar e}{2m_e} \sum_s \sum_t C_s^{l*} C_t^h \langle p_s | \vec{r} \times \vec{\nabla} | p_t \rangle$$

by S.Savikhin, Purdue University

13

Magnetic transition dipole moment

CAN SKIP

$$\vec{m}_{ba} = -i \frac{\sqrt{2}\hbar e}{2m_e} \sum_s \sum_t C_s^{l*} C_t^h \langle p_s | \vec{r} \times \vec{\nabla} | p_t \rangle$$

We will now split the vector $\vec{r}$ into the sum of vector $\vec{R}_t$ to the center of atom t and the position of electron $\vec{r}_t$ with respect to that: $\vec{r} \rightarrow \vec{R}_t + \vec{r}_t$

$$\vec{m}_{ba} = -i \frac{\sqrt{2}\hbar e}{2m_e} \sum_s \sum_t C_s^{l*} C_t^h \langle p_s | (\vec{r}_t + \vec{R}_t) \times \vec{\nabla} | p_t \rangle$$

$$\vec{m}_{ba} = -i \frac{\sqrt{2}\hbar e}{2m_e} \sum_s \sum_t C_s^{l*} C_t^h \left[\langle p_s | \vec{r}_t \times \vec{\nabla} | p_t \rangle + \langle p_s | \vec{R}_t \times \vec{\nabla} | p_t \rangle \right]$$

$$\vec{m}_{ba} = -i \frac{\sqrt{2}\hbar e}{2m_e} \sum_s \sum_t C_s^{l*} C_t^h \left[\langle p_s | \vec{r}_t \times \vec{\nabla} | p_t \rangle + \vec{R}_t \times \langle p_s | \vec{\nabla} | p_t \rangle \right]$$

by S.Savikhin, Purdue University

14

Magnetic transition dipole moment

CAN SKIP

$$\vec{m}_{ba} = -i \frac{\sqrt{2}\hbar e}{2m_e} \sum_s \sum_t C_s^* C_t \left[\langle p_s | \vec{r}_i \times \vec{\nabla} | p_t \rangle + \vec{R}_i \times \langle p_s | \vec{\nabla} | p_t \rangle \right]$$

positions of electrons with respect to the atom

The overlap between the 2p orbitals of the same atom is the main contributor

Relative positions of atoms

For orbitals on the same atom, $R_i=0$

For orbitals on different atoms, the overlap integral is small

→ Ignore this term

The first term cross product:

$$\langle p_s | \vec{r}_i \times \vec{\nabla} | p_t \rangle = \left\langle p_s \left| y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} \right| p_t \right\rangle \hat{x} - \left\langle p_s \left| x \frac{\partial}{\partial z} - z \frac{\partial}{\partial x} \right| p_t \right\rangle \hat{y} + \left\langle p_s \left| x \frac{\partial}{\partial y} - y \frac{\partial}{\partial x} \right| p_t \right\rangle \hat{z}$$

Consider x-component

$$\left\langle p_y \left| y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} \right| p_z \right\rangle = \left\langle p_y \left| y \frac{\partial p_z}{\partial z} \right\rangle - \left\langle p_y \left| z \frac{\partial p_z}{\partial y} \right\rangle$$

by S.Savikhin, Purdue University

15

Magnetic transition dipole moment

CAN SKIP

Let's investigate the first term (x-component) when

$p_s : 2p_y \text{ orbital}$
 $p_t : 2p_z \text{ orbital}$

$$\left\langle p_y \left| y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} \right| p_z \right\rangle = \left\langle p_y \left| y \frac{\partial p_z}{\partial z} \right\rangle - \left\langle p_y \left| z \frac{\partial p_z}{\partial y} \right\rangle$$

See the comprehensive discussion in the "green" book
But D and E graphs in the book are swapped!

X-component is non-zero

Can show that for these orbitals, the y and z components are zero

→ **Magnetic moment is oriented along the x-axis for these orbitals**

by S.Savikhin, Purdue University

16

Magnetic transition dipole moment

CAN SKIP

$$\vec{m}_{ba} = -i \frac{\sqrt{2}\hbar e}{2m_e} \sum_s \sum_t C_s^* C_t^h \left[\langle p_s | \vec{r}_t \times \vec{\nabla} | p_t \rangle + \vec{R}_t \times \langle p_s | \vec{\nabla} | p_t \rangle \right]$$

$$\langle p_s | \vec{r}_t \times \vec{\nabla} | p_t \rangle = \langle p_s | y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} | p_t \rangle \hat{x} - \langle p_s | x \frac{\partial}{\partial z} - z \frac{\partial}{\partial x} | p_t \rangle \hat{y} + \langle p_s | x \frac{\partial}{\partial y} - y \frac{\partial}{\partial x} | p_t \rangle \hat{z}$$

For p_y and p_z orbitals on the same atom, $\vec{m}_{ba}$ is oriented along the x-axis.

If the p_y and p_z are on different atoms, the orientation of $\vec{m}_{ba}$ is the same, but with a much lower amplitude due to poor overlap of the wavefunctions

by S.Savikhin, Purdue University



17

Magnetic transition moment: example

CAN SKIP

Excitation of a **carbonyl group** involves $2p_y$ and $2p_z$ orbitals:

Transition $n \rightarrow \pi^*$, from the nonbonding orbital of the oxygen atom to π^* molecular orbital constructed of oxygen and carbon $2p$ orbitals.



$$p_{y(O)} \longrightarrow 2^{-1/2} (p_{z(C)} - p_{z(O)})$$

Electric transition moment:

$$\vec{\mu}_{ba} = 2^{-1/2} \langle (p_{z(C)} - p_{z(O)}) | \vec{\mu} | p_{y(O)} \rangle = 2^{-1/2} \langle p_{z(C)} | \vec{\mu} | p_{y(O)} \rangle - 2^{-1/2} \langle -p_{z(O)} | \vec{\mu} | p_{y(O)} \rangle$$

Magnetic transition moment:

$$\vec{m}_{ba} = 2^{-1/2} \langle (p_{z(C)} - p_{z(O)}) | \vec{m} | p_{y(O)} \rangle \text{ Is not zero (as shown in prev. slide)}$$

The extinction coefficient is small, <100 (for electric transitions in dyes, it is >10,000)

by S.Savikhin, Purdue University

18

Magnetic transition moment: properties

$$\vec{m}_{ba} = -i \frac{e\hbar}{2m_e} \langle \psi_b | \vec{r} \times \vec{\nabla} | \psi_a \rangle$$

$$\text{Bohr magneton: } m_B \equiv \frac{e\hbar}{2m_e}$$

The magnetic transition moment is weak

In most molecules, absorption due to the magnetic transition dipole moment is 2-4 orders of magnitude weaker than absorption due to the electric transition dipole moment.

Right-hand rule

The direction of $\mathbf{m}$ is determined by the "rotation" of charge density during the transition

For planar molecules, $\mathbf{m}$ points out of the plane (perpendicular)

For flat (planar) molecules (such as chlorophyll), the magnetic transition dipole moment typically involves "circular" movement of electrons within the plane of the molecule (such as $\pi \rightarrow \pi^*$ transitions).

Symmetry:

In some planar molecules, transitions that are electrically forbidden ($\mu=0$) can be magnetically allowed, if the transition involves a change in orbital angular momentum that mimics a current loop

by S.Savikhin, Purdue University


19

Magnetic vs electric transition dipole absorption

Magnetic transition

Formaldehyde (CH_2O):

Its $n \rightarrow \pi^*$ transition, the transition is strictly electric-dipole forbidden by symmetry. It becomes observable primarily through magnetic dipole transitions




Formaldehyde absorption spectrum in the UV range, re-plotted from the data of Meller et al [30]. Peaks are not fully resolved at the resolution of this spectrum (0.01 nm).

https://www.researchgate.net/figure/Formaldehyde-absorption-spectrum-in-the-UV-range-re-plotted-from-the-data-of-Meller-et_fig1_287158085

Electric transition

Chlorophylls & carotenoids



https://www.researchgate.net/figure/upper-Differences-in-absorption-spectra-of-chlorophyll-a-chlorophyll-b-and-b-carotene_fig1_40799750

by S.Savikhin, Purdue University

20

Interaction with circularly polarized light

Given the right combination of μ and m , absorption of R and L polarization is different $\rightarrow$ circular dichroism

Recall lecture 3: transition probability

Given Hamiltonian $\tilde{H} = \tilde{H}_0 + \tilde{H}'$

For harmonic interaction $\tilde{H}' = \tilde{H}'(\vec{r}) e^{i\omega t}$

Transition probability: $P_{a \rightarrow b} \approx \frac{|H'_{ba}(\vec{r})|^2}{\hbar^2} \frac{\sin^2[(\omega_{ba} - \omega)t/2]}{(\omega_{ba} - \omega)^2}$

Have: $\tilde{H}'(t) = -\vec{E}(t) \cdot \tilde{\mu} - \vec{B}(t) \cdot \tilde{m}$

$$\left\{ \begin{array}{l} \tilde{E}^{\pm} = E_0 (\hat{x} \pm i\hat{y}) e^{i(kz - \omega t)} \\ \tilde{B}^{\pm} = \frac{1}{c} \hat{z} \times \tilde{E}^{\pm} = \mp \frac{i}{c} \tilde{E}^{\pm} \end{array} \right. \xrightarrow[\text{Time-independent part}]{\text{Circular polarization}} \left\{ \begin{array}{l} \tilde{E}^{\pm}(\vec{r}) = E_0 (\hat{x} \pm i\hat{y}) e^{ikz} \\ \tilde{B}^{\pm}(\vec{r}) = \mp \frac{i}{c} \tilde{E}^{\pm}(\vec{r}) \end{array} \right.$$

$$\tilde{H}'_{ba} \equiv \langle \psi_b | \tilde{H}'(\vec{r}) | \psi_a \rangle = \langle \psi_b | \tilde{E}(\vec{r}) \cdot \tilde{\mu} | \psi_a \rangle + \langle \psi_b | \tilde{B}(\vec{r}) \cdot \tilde{m} | \psi_a \rangle$$

$$= \tilde{E}(\vec{r}) \cdot \tilde{\mu}_{ba} + \tilde{B}(\vec{r}) \cdot \tilde{m}_{ba}$$

by S.Savikhin, Purdue University

21

Interaction with circularly polarized light

$$H'_{ba} = \tilde{E}(\vec{r}) \cdot \tilde{\mu}_{ba} + \tilde{B}(\vec{r}) \cdot \tilde{m}_{ba} \quad \left\{ \begin{array}{l} \tilde{E}^{\pm}(\vec{r}) = E_0 (\hat{x} \pm i\hat{y}) e^{ikz} \\ \tilde{B}^{\pm}(\vec{r}) = \mp \frac{i}{c} \tilde{E}^{\pm}(\vec{r}) \end{array} \right. \quad \begin{array}{l} + = \text{left} \\ - = \text{right} \end{array}$$

Without loss of generality: molecule is at $z=0$ (molecule is small):

$$H'_{ba} = E_0 \left[(\hat{x} \pm i\hat{y}) \cdot \tilde{\mu}_{ba} \mp \frac{i}{c} (\hat{x} \pm i\hat{y}) \cdot \tilde{m}_{ba} \right] \quad \text{Important: keep a close attention to sign changes, + to - and } \pm \text{ to } \mp$$

Transition probability is proportional to the square:

$$|H_{ba}|^2 = E_0^2 \left[(\hat{x} \mp i\hat{y}) \cdot \tilde{\mu}_{ba}^* \pm \frac{i}{c} (\hat{x} \mp i\hat{y}) \cdot \tilde{m}_{ba}^* \right] \left[(\hat{x} \pm i\hat{y}) \cdot \tilde{\mu}_{ba} \mp \frac{i}{c} (\hat{x} \pm i\hat{y}) \cdot \tilde{m}_{ba} \right]$$

Notes:

$\tilde{\mu} \equiv \tilde{\mu}_{ba} = \tilde{\mu}_{ba}^*$: the electric transition dipole is a real number (lectures 3-4)

$\tilde{m} \equiv \tilde{m}_{ba} = -\tilde{m}_{ba}^*$: m is purely imaginary

$$|H_{ba}|^2 = E_0^2 \left[(\hat{x} \mp i\hat{y}) \cdot \tilde{\mu}_{ba} \mp \frac{i}{c} (\hat{x} \mp i\hat{y}) \cdot \tilde{m}_{ba} \right] \left[(\hat{x} \pm i\hat{y}) \cdot \tilde{\mu}_{ba} \mp \frac{i}{c} (\hat{x} \pm i\hat{y}) \cdot \tilde{m}_{ba} \right]$$

$$|H_{ba}|^2 = E_0^2 \left[(\mu_x \mp i\mu_y) \mp \frac{i}{c} (m_x \mp im_y) \right] \left[(\mu_x \pm i\mu_y) \mp \frac{i}{c} (m_x \pm im_y) \right]$$

by S.Savikhin, Purdue University

22

Interaction with circularly polarized light

$$|H_{ba}|^2 = E_0^2 \left[(\mu_x \mp i\mu_y) \mp \frac{i}{c} (m_x \mp im_y) \right] \left[(\mu_x \pm i\mu_y) \mp \frac{i}{c} (m_x \pm im_y) \right]$$

Expand four terms:

$$\begin{aligned} (\mu_x \mp i\mu_y)(\mu_x \pm i\mu_y) &= \mu_x^2 + \mu_y^2 & \mp \frac{i}{c} (\mu_x \mp i\mu_y)(m_x \pm im_y) &= \mp \frac{i}{c} (\mu_x m_x \pm i\mu_x m_y \mp i\mu_y m_x + \mu_y m_y) \\ -\frac{1}{c^2} (m_x \mp im_y)(m_x \pm im_y) &= -\frac{1}{c^2} [m_x^2 + m_y^2] & \mp \frac{i}{c} (m_x \mp im_y)(\mu_x \pm i\mu_y) &= \mp \frac{i}{c} (\mu_x m_x \mp i\mu_x m_y \pm i\mu_y m_x + \mu_y m_y) \end{aligned}$$

$$|\tilde{H}'_{ba}|^2 = E_0^2 \left([\mu_x^2 + \mu_y^2] \mp \frac{2i}{c} [\mu_x m_x + \mu_y m_y] + \frac{1}{c^2} [m_x^2 + m_y^2] \right)$$

Assume the transition moments are in the xy-plane
(largest interaction):

$$|\tilde{H}'_{ba}|^2 = E_0^2 |\vec{\mu}|^2 \mp 2iE_0 B_0 \vec{\mu} \cdot \vec{m} + B_0^2 |\vec{m}|^2$$

$$\begin{cases} \vec{E}^\pm = 2^{-1/2} E_0 (\hat{x} \pm i\hat{y}) e^{-i\omega t} \\ \vec{B}^\pm = \frac{1}{c} \hat{z} \times \vec{E}^\pm = \mp \frac{i}{c} \vec{E}^\pm \end{cases}$$

E and B are in
the xy-plane

by S.Savikhin, Purdue University

23

Interaction with circularly polarized light

$$|\tilde{H}'_{ba}|^2 = E_0^2 |\vec{\mu}|^2 \mp 2iE_0 B_0 \vec{\mu} \cdot \vec{m} + B_0^2 |\vec{m}|^2$$

$$\begin{aligned} E_0^2 |\vec{\mu}|^2 & \text{ Absorption due to the electric transition dipole moment, strength } D_E = |\vec{\mu}|^2 & \text{ For linear: } |\vec{E} \cdot \vec{\mu}|^2 \\ B_0^2 |\vec{m}|^2 & \text{ Absorption due to the magnetic transition dipole moment, strength } D_B = |\vec{m}|^2 & |\vec{B} \cdot \vec{m}|^2 \\ \mp 2iE_0 B_0 \vec{\mu} \cdot \vec{m} & \text{ Absorption due to the magnetic and electric transition dipole moments } D_{L,R} = \mp 2 \text{Im}(\vec{\mu} \cdot \vec{m}) \end{aligned}$$

$$\text{CD difference } \Delta A_{CD} \propto |\tilde{H}'_{ba}|_+^2 - |\tilde{H}'_{ba}|_-^2 = E_0 B_0 [-4 \text{Im}(\vec{\mu}_{ba} \cdot \vec{m}_{ba})]$$

Define Rotational Strength: $\mathcal{R} \equiv -\text{Im}(\vec{\mu}_{ba} \cdot \vec{m}_{ba}) = \frac{D_L - D_R}{4}$ (Rosenfeld equation, 1928)
R is a real number

$$D_L - D_R = 4\mathcal{R}$$

The coefficient 4 stems from Rosenfeld's definition (apparently?)

by S.Savikhin, Purdue University

24

Interaction with circularly polarized light

	Maximum in optimal orientation	Averaging over all possible orientations		Strength
		For circular	For linear	
$ \vec{E} \cdot \vec{\mu} ^2 =$	$E_0^2 D_E$	$\frac{2}{3} E_0^2 D_E$	$\frac{1}{3} E_0^2 D_E$	$D_E \equiv \vec{\mu} ^2$
$ \vec{B} \cdot \vec{m} ^2 =$	$B_0^2 D_B$	$\frac{2}{3} B_0^2 D_B$	$\frac{1}{3} B_0^2 D_B$	$D_B \equiv \vec{m} ^2$
CD difference	$E_0 B_0 (4\mathcal{R})$	$\frac{2}{3} E_0 B_0 (4\mathcal{R})$		$\mathcal{R} = -\text{Im}(\vec{\mu}_{ba} \cdot \vec{m}_{ba})$

↑ 2/3 vs 1/3 ↑

Note: for the same E_0 , the energy density of circularly polarized light is twice as large as for linearly polarized light, hence the 2x difference in averages for circular and linear polarizations.

If average absorption is expressed as a function of light energy density, i.e., use $1/\sqrt{2}$ coefficient in the expression for light fields, then the angular averaging coefficient is the same 1/3 for circular and linear polarizations!

by S.Savikhin, Purdue University

25

Rotational strength from experiment

Rotational strength: $\mathcal{R} = \frac{D_L - D_R}{4}$ (transition dipole strengths, quantum)

One can find D and R from experimental measurements of molar extinction spectra

$$\text{D from absorption: } D = \frac{3hc \cdot 10^3 \ln(10)}{8\pi^3 N_A} \int \frac{\epsilon(\tilde{\nu})}{\tilde{\nu}} d\tilde{\nu}$$

$$\text{R from absorption difference: } \mathcal{R} = \frac{3hc \cdot 10^3 \ln(10)}{32\pi^3 N_A} \int \frac{\Delta\epsilon(\tilde{\nu})}{\tilde{\nu}} d\tilde{\nu}$$

If you have a CD spectrum, you can estimate the rotational strength of a specific band using the *Gaussian approximation*:

$$\mathcal{R} \approx 1.23 \times 10^{-42} \cdot \frac{\Delta\epsilon_{\max} \cdot \Delta\tilde{\nu}}{\tilde{\nu}_{\max}}$$

- $\Delta\epsilon_{\max}$ is the peak molar circular dichroism.
- $\tilde{\nu}_{\max}$ is the wavenumber at the peak.
- $\Delta\tilde{\nu}$ is the half-width of the band at 1/e of its maximum height.

$$\text{Also: } \frac{\mathcal{R}}{D} = \frac{\Delta A_{CD}}{4A} = -\frac{\text{Im}(\vec{\mu}_{ba} \cdot \vec{m}_{ba})}{\mu_{ba}^2} = -\frac{|m_{ba}|}{|\mu_{ba}|} \cos \alpha_{m\mu}$$

by S.Savikhin, Purdue University

26

Intrinsic CD: planar molecules

Planar molecules have no intrinsic CD

For flat (planar) molecules (such as chlorophyll), the magnetic transition dipole moment typically involves "circular" movement of electrons within the plane of the molecule (such as $\pi \rightarrow \pi^*$ transitions). Thus, the magnetic transition moment points perpendicular to the molecular plane, while the electric transition dipole is in the plane of the molecule $\rightarrow$ there is no intrinsic CD ($\vec{\mu} \cdot \vec{m} = 0$).


by S.Savikhin, Purdue University

27

Oligomers: excitonic CD

Handwaving explanation

Example: a strongly coupled dimer



$$H\Psi_i = E_i\Psi_i$$

$$H = \begin{pmatrix} E & V \\ V & E \end{pmatrix}$$



Delocalized states

$$\Psi_1^* \sim |10\rangle + |01\rangle$$



$$\vec{\eta}_1 \Rightarrow \vec{\mu}_1 + \vec{\mu}_2$$

$$\Psi_2^* \sim |10\rangle - |01\rangle$$

$$\vec{\eta}_2 \Rightarrow \vec{\mu}_1 - \vec{\mu}_2$$



The electric field direction is different at the two locations



For in-phase transition: $\vec{\eta}_1 \Rightarrow \vec{\mu}_1 + \vec{\mu}_2$

Right circularly polarized light is absorbed better

For out-of-phase transition: $\vec{\eta}_2 \Rightarrow \vec{\mu}_1 - \vec{\mu}_2$

Left circularly polarized light is absorbed better

CD stems from the spatial distribution of coupled transition dipole moments

The main CD effect stems from electric moments; magnetic moments are weak

by S.Savikhin, Purdue University

28

Oligomers: excitonic CD

$$P_{a \rightarrow b} \approx \frac{|H'_{ba}(\vec{r})|^2}{\hbar^2} \frac{\sin^2[(\omega_{ba} - \omega)t/2]}{(\omega_0 - \omega)^2} \quad (\text{Lecture 3})$$

$$\tilde{H}'_{ba} \equiv \tilde{H}'_{ba}(\vec{r}) = \langle \psi_b | \tilde{H}'(\vec{r}) | \psi_a \rangle$$

Dipole operator (lecture 4)
Ground state $\psi_a(\vec{r}_i)$

$$\text{Interaction with light: } \tilde{H}'_{ba} = \langle \psi_b | \vec{E}^\pm(\vec{r}) \cdot \hat{\mu} | \psi_a \rangle$$

$$\text{Excited excitonic (delocalized) state } n: \psi_n(\vec{r}) = \sum_i c_{ni} \phi_i(\vec{r}_i) \quad (\text{Lecture 11})$$

Excited state

$$\tilde{H}'_{na} = \vec{E}^\pm(\vec{r}) \cdot \left\langle \sum_i c_{ni} \phi_i | \hat{\mu} | \psi_a \right\rangle = \sum_i c_{ni} \vec{E}^\pm(\vec{r}) \cdot \langle \phi_i | \hat{\mu} | \psi_a \rangle = \sum_i c_{ni} \vec{E}^\pm(\vec{r}) \cdot \vec{\mu}_i$$

Excited states of individual molecules (basis)

$$\text{Circularly polarized light propagating along the z-axis: } \vec{E}^\pm(\vec{r}) = \frac{1}{\sqrt{2}} E_0 (\hat{x} \pm i\hat{y}) e^{ikz}$$

$$\tilde{H}'_{na} = \sum_i c_{ni} \frac{E_0}{\sqrt{2}} (\hat{x} \pm i\hat{y}) e^{ikz_i} \cdot \vec{\mu}_i$$

Not index, the square root of -1
z-component of the chromophore's position

by S.Savikhin, Purdue University

29

Oligomers: excitonic CD

$$\tilde{H}'_{na} = \sum_i c_{ni} \frac{E_0}{\sqrt{2}} (\hat{x} \pm i\hat{y}) e^{ikz_i} \cdot \vec{\mu}_i$$

$$\text{Transition strength is proportional to: } |\tilde{H}'_{na}|^2 = \tilde{H}'_{na}{}^* \tilde{H}'_{na}$$

$$|\tilde{H}'_{na}|^2 = \frac{1}{2} |E_0|^2 \left[\sum_i c_{ni}^* (\hat{x} \mp i\hat{y}) \cdot \vec{\mu}_i e^{-ikz_i} \right] \left[\sum_j c_{nj} (\hat{x} \pm i\hat{y}) \cdot \vec{\mu}_j e^{ikz_j} \right]$$

$$= \frac{1}{2} |E_0|^2 \sum_i \sum_j c_{ni}^* c_{nj} [(\hat{x} \mp i\hat{y}) \cdot \vec{\mu}_i] [(\hat{x} \pm i\hat{y}) \cdot \vec{\mu}_j] e^{ik(z_j - z_i)}$$

$$= \frac{1}{2} |E_0|^2 \sum_i \sum_j c_{ni}^* c_{nj} [(\mu_{ix} \mp i\mu_{iy})(\mu_{jx} \pm i\mu_{jy})] e^{-ikz_{ij}}$$

$$= \frac{1}{2} |E_0|^2 \sum_i \sum_k c_{ni}^* c_{nj} [\mu_{ix}\mu_{jx} \pm i\mu_{ix}\mu_{jy} \mp i\mu_{iy}\mu_{jx} + \mu_{iy}\mu_{jy}] e^{-ikz_{ij}}$$

by S.Savikhin, Purdue University

30

Oligomers: excitonic CD

$$\left| \tilde{H}'_{na} \right|^2 = \frac{1}{2} |E_0|^2 \sum_i \sum_k c_{ni}^* c_{kj} \left[\mu_{ix} \mu_{jx} \pm i \mu_{ix} \mu_{jy} \mp i \mu_{iy} \mu_{jx} + \mu_{iy} \mu_{jy} \right] e^{-ikz_{ij}}$$

Cancel in difference

Circular dichroism: $\Delta A_{CD} = A_L - A_R \propto \Delta \left(\left| \tilde{H}'_{na} \right|^2 \right)_{CD} = \left| \tilde{H}'_{na} \right|_+^2 - \left| \tilde{H}'_{na} \right|_-^2$

$$\Delta \left(\left| \tilde{H}'_{na} \right|^2 \right)_{CD} = |E_0|^2 \sum_i \sum_k c_{ni}^* c_{kj} \left[i \left(\vec{\mu}_{ix} \vec{\mu}_{jy} - \vec{\mu}_{iy} \vec{\mu}_{jx} \right) \right] e^{-ikz_{ij}}$$

↑ z-component of cross-product $\vec{\mu}_i \times \vec{\mu}_j$

$$\Delta \left(\left| \tilde{H}'_{na} \right|^2 \right)_{CD} = |E_0|^2 \sum_i \sum_j c_{ni}^* c_{nj} \left[i \hat{z} \cdot \left(\vec{\mu}_i \times \vec{\mu}_j \right) \right] e^{-ikz_{ij}}$$

$e^{-ikz_{ij}} = \cos(kz_{ij}) - i \sin(kz_{ij})$

The distance between interacting pigments is short, $z_{ij} \sim 10 \text{ \AA}$,
while the wavelength of light is $\lambda \sim 6000 \text{ \AA} \rightarrow kz_{ij} \sim 1/100$.

For small angles $\sin(x) \sim x$, $\cos(x) \sim 1$:

$$e^{-ikz_{ij}} \approx 1 - ikz_{ij}$$

$$k = \frac{2\pi}{\lambda}$$

by S.Savikhin, Purdue University

31

Oligomers: excitonic CD

$$\Delta \left(\left| \tilde{H}'_{na} \right|^2 \right)_{CD} = |E_0|^2 \sum_i \sum_j c_{ni}^* c_{nj} \left[i \hat{z} \cdot \left(\vec{\mu}_i \times \vec{\mu}_j \right) \right] e^{-ikz_{ij}} \quad e^{-ikz_{ij}} \approx 1 - ikz_{ij}$$

$$\Delta \left(\left| \tilde{H}'_{na} \right|^2 \right)_{CD} \approx |E_0|^2 \sum_i \sum_j c_{ni}^* c_{nj} \left[i \hat{z} \cdot \left(\vec{\mu}_i \times \vec{\mu}_j \right) \right] + |E_0|^2 \sum_i \sum_j c_{ni}^* c_{nj} \left[\hat{z} \cdot \left(\vec{\mu}_i \times \vec{\mu}_j \right) \right] kz_{ij}$$

Notes: $\left(\vec{\mu}_i \times \vec{\mu}_j \right) = - \left(\vec{\mu}_j \times \vec{\mu}_i \right)$ and $z_{ji} \equiv -z_{ij}$

Sums always have terms with ij and ji;

The first sum is odd with respect to ij to ji substitution, $\rightarrow$ it will be 0:

$$\Delta \left(\left| \tilde{H}'_{na} \right|^2 \right)_{CD} = |E_0|^2 \sum_i \sum_j c_{ni}^* c_{nj} \left[\hat{z} \cdot \left(\vec{\mu}_i \times \vec{\mu}_j \right) \right] kz_{ij}$$

$k = \frac{2\pi}{\lambda}$

$$\Delta \left(\left| \tilde{H}'_{na} \right|^2 \right)_{CD} = \frac{2\pi}{\lambda} |E_0|^2 \sum_i \sum_j c_{ni}^* c_{nj} \left[\hat{z} \cdot \left(\vec{\mu}_i \times \vec{\mu}_j \right) \right] \hat{z} \cdot \vec{r}_{ij}$$

Average over all possible directions of z: $\overline{(\hat{z} \cdot \vec{A})(\hat{z} \cdot \vec{B})} = \frac{1}{3} \vec{A} \cdot \vec{B}$

$$\overline{\Delta \left(\left| \tilde{H}'_{na} \right|^2 \right)_{CD}} = |E_0|^2 \frac{2\pi}{3\lambda} \sum_i \sum_j c_{ni}^* c_{nj} \vec{r}_{ij} \cdot \left(\vec{\mu}_i \times \vec{\mu}_j \right)$$

by S.Savikhin, Purdue University

32

Oligomers: excitonic CD

$$\overline{\Delta \left(\left| \tilde{H}_{na} \right|^2 \right)_{CD}} = |E_0|^2 \frac{2\pi}{3\lambda} \sum_i \sum_j c_{ni}^* c_{nj} \vec{r}_{ij} \cdot (\vec{\mu}_i \times \vec{\mu}_j) \quad z_{ij} \equiv z_i - z_j = \hat{z} \cdot \vec{r}_{ij}$$

Direction from j to i

This is the difference in apparent transition dipole strength for left (D_L) and right (D_R) circularly polarized light absorption averaged over all directions. The rotational strength is defined as

$$\mathcal{R} = \frac{D_L - D_R}{4}$$

And the averaging coefficient (1/3) is dropped (as earlier for intrinsic CD)*:

Rotational strength

$$\mathcal{R}_n = \frac{\pi}{2\lambda} \sum_{i,j} c_{ni} c_{nj} \left[\vec{r}_{ij} \cdot (\vec{\mu}_i \times \vec{\mu}_j) \right]$$

*Here, we used $2^{-1/2}$ coefficient in the expression for the amplitude E_0 , so the averaging coefficient was 1/3 and not 2/3

by S.Savikhin, Purdue University

33

Oligomers: excitonic CD

Rotational strength for excitonic transition n :

$$\mathcal{R}_n = \frac{\pi}{2\lambda} \sum_{i,j} c_{ni} c_{nj} \left[\vec{r}_{ij} \cdot (\vec{\mu}_i \times \vec{\mu}_j) \right] \quad \vec{r}_{ij} \equiv \vec{r}_i - \vec{r}_j$$

Vector from j to i
if $i=j$, $r_{ij} = 0$

Transition dipole strength for excitonic transition:

$$D_n = \left| \sum_i c_{ni} \vec{\mu}_i \right|^2 = \sum_{i,j} c_{ni} c_{nj} \vec{\mu}_i \cdot \vec{\mu}_j$$

To average over all possible directions x1/3


Notes on oligomeric CD:

- Must superimpose the effects of intrinsic molecular $\mathcal{R}$ and excitonic $\mathcal{R}$
- For a flat molecule (like chlorophyll), there is no intrinsic CD
- We ignored CD due to couplings between magnetic dipole moments and magnetic-electric dipole moments for different molecules (they are typically negligible compared to excitonic CD)
- The actual ΔA_{CD} for a single oligomer oriented in a specific way could be positive or negative, even if the (averaged) $\mathcal{R}$ is zero!
- The excitonic CD spectrum is conservative, i.e., integrated intensity is zero

by S.Savikhin, Purdue University

34


Excitonic CD: homodimer example



Delocalized states

$$\mathcal{R} = \frac{\pi}{2\lambda} \sum_{i,j} c_{ki} c_{kj} [\vec{r}_{ij} \cdot (\vec{\mu}_i \times \vec{\mu}_j)]$$

Note: if $i=j$, $r_{ij} = 0$




$$\Psi_+ = \frac{1}{\sqrt{2}} \psi_1 + \frac{1}{\sqrt{2}} \psi_2 \quad \vec{\mu}_+ = \frac{1}{\sqrt{2}} \vec{\mu}_1 + \frac{1}{\sqrt{2}} \vec{\mu}_2$$

$$\Psi_- = \frac{1}{\sqrt{2}} \psi_1 - \frac{1}{\sqrt{2}} \psi_2 \quad \vec{\mu}_- = \frac{1}{\sqrt{2}} \vec{\mu}_1 - \frac{1}{\sqrt{2}} \vec{\mu}_2$$

$$\mathcal{R}_{\pm} = \frac{\pi}{2\lambda_{\pm}} \left[\pm \frac{1}{2} \vec{r}_{12} \cdot (\vec{\mu}_1 \times \vec{\mu}_2) \pm \frac{1}{2} \vec{r}_{21} \cdot (\vec{\mu}_2 \times \vec{\mu}_1) \right]$$

$$\vec{r}_{21} = -\vec{r}_{12} \quad \vec{\mu}_2 \times \vec{\mu}_1 = -\vec{\mu}_1 \times \vec{\mu}_2$$

$$\mathcal{R}_{\pm} = \pm \frac{\pi}{2\lambda_{\pm}} \vec{r}_{12} \cdot (\vec{\mu}_1 \times \vec{\mu}_2) \quad \vec{r}_{12} \equiv \vec{r}_1 - \vec{r}_2$$



While the absorption strengths of each band could be different, the CD signals have the same magnitudes but opposite signs


→ CD spectra can distinguish both bands even if one is 'dark.'

→ Since bands are positive and negative, one can distinguish overlapping bands

by S.Savikhin, Purdue University

35

Excitonic CD: the Fenna Matthews Olson complex



$$3. \text{ Calculate: } D_e = \left| \sum_{n,m=1}^N c_{en} \vec{\mu}_n \right|^2 = \sum_{n,m=1}^N c_{en} c_{em}^* \vec{\mu}_n \cdot \vec{\mu}_m$$

$$R_e = \sum_{n,m=1}^N c_{en} c_{em}^* \vec{r}_{nm} \cdot (\vec{\mu}_n \times \vec{\mu}_m)$$

T= 77K


1. Hamiltonian in the basis of pigment's wavefunctions

$$H = \begin{bmatrix} 12420 & -106 & 8 & -5 & 6 & -8 & -4 \\ -106 & 12560 & 28 & 6 & 2 & 13 & 1 \\ 8 & 28 & 12140 & -62 & -1 & -9 & 17 \\ -5 & 6 & -62 & 12335 & -70 & -19 & -57 \\ 6 & 2 & -1 & -70 & 12460 & 40 & -2 \\ -8 & 13 & -9 & -19 & 40 & 12500 & 32 \\ -4 & 1 & 17 & -57 & -2 & 32 & 12400 \end{bmatrix} \quad \Psi_n = \begin{bmatrix} c_{n1} \\ c_{n2} \\ \vdots \\ c_{n7} \end{bmatrix}$$

2. Solve TISE: $\mathbf{H}\Psi_n = E_n\Psi_n$

Obtain (by $\mathbf{H}$ diagonalization):
 E_n – exciton transition energies (eigenvalues)

c_{ni} – expansion coefficients (eigenvectors)



by S.Savikhin, Purdue University

36

Making circularly polarized light

How to make circularly polarized light?

$$\begin{aligned}\vec{E}^{\pm}(t, z) &= E_0 \left[\hat{x} \cos(\omega t - kz) \pm \hat{y} \sin(\omega t - kz) \right] \\ &= E_0 \left[\hat{x} \cos(\omega t - kz) + \hat{y} \cos(\omega t - kz + \theta) \right]\end{aligned}$$

$$k = \frac{2\pi}{\lambda}$$

$\theta = \pm\pi/2$ - Phase shift for one axis results in circular polarization
(shift of π will rotate polarization by 90 degrees)

$\theta = \pm\pi/2$ corresponds to optical path delay by quarter wavelength, $\lambda/4$


$$\vec{E}^{\pm}(t, z) = E_0 \left[\hat{x} \cos(\omega t - kz) + \hat{y} \cos(\omega t - k[z + n_{\theta}\lambda]) \right]$$

n_{θ} - delay in wavelengths

Wave-plate: refraction index is different for light polarized along orthogonal axes

Quarter-wave plate: difference in optical path is $\lambda/4$ ($+n\lambda$), $n_{\theta} = 1/4$. By rotating $\lambda/4$ -plate by 90 degrees, one can switch between L and R polarizations

Half-wave plate: can rotate polarization by any angle up to 90 degrees




by S.Savikhin, Purdue University


37

Measuring circular dichroism

CD spectrometer with L/R handedness modulation



PEM, Photo-Elastic Modulator:
Soundwave modulates the density of glass and the refractive index



Reference $\sin(\omega t)$ Signal $S(t)$

Lock-in amplifier

Detects the signal at a certain frequency

Fourier trick: $\Delta T_{CD} = a_{\omega} = \int S(t) \sin(\omega t) dt$

Compressing glass along one axis changes the refractive index (and speed of light)


Typically, the standing acoustic wave frequency is ~ 50 kHz, so light alternates between R and L at this frequency.

by S.Savikhin, Purdue University

38

CD spectrometer

Push-button, no brains needed to operate



ZEMOS CD spectrometer

by S.Savikhin, Purdue University

39

EXTRA

by S.Savikhin, Purdue University

40

Interaction with circularly polarized light

Generated by Gemini AI

$$\tilde{H}'(t) = -\mathbf{E}(t) \cdot \tilde{\boldsymbol{\mu}} - \mathbf{B}(t) \cdot \tilde{\mathbf{m}}$$

Transition probability is: $W_{ba} \propto |\tilde{H}'_{ba}|^2$

Note: assume that the light frequency is in resonance with the transition

$$\tilde{H}'_{ba} = \langle \psi_b | -\vec{E} \cdot \tilde{\boldsymbol{\mu}} - \vec{B} \cdot \tilde{\mathbf{m}} | \psi_a \rangle = -\vec{E} \langle \psi_b | \tilde{\boldsymbol{\mu}} | \psi_a \rangle - \vec{B} \langle \psi_b | \tilde{\mathbf{m}} | \psi_a \rangle = -\vec{E} \cdot \tilde{\boldsymbol{\mu}}_{ba} - \vec{B} \cdot \tilde{\mathbf{m}}_{ba}$$

Circularly polarized light:
$$\begin{cases} \vec{E}^{\pm} = E_0 (\hat{x} \pm i\hat{y}) e^{i(kz - \omega t)} \\ \vec{B}^{\pm} = \frac{1}{c} \hat{z} \times \vec{E}^{\pm} = \mp \frac{i}{c} \vec{E}^{\pm} \end{cases}$$

+ = left
- = right

Circular dichroism: $\Delta A_{CD} = W_{ba}^{+} - W_{ba}^{-}$

$$|\tilde{H}'_{ba}^{\pm}|^2 = \left| -\vec{E}^{\pm} \cdot \tilde{\boldsymbol{\mu}}_{ba} \pm \frac{i}{c} \vec{E}^{\pm} \cdot \tilde{\mathbf{m}}_{ba} \right|^2$$

$$|\tilde{H}'_{ba}^{\pm}|^2 = E_0^2 \left(|\tilde{\boldsymbol{\mu}}_{ba}|^2 \pm \frac{2i}{c} \tilde{\boldsymbol{\mu}}_{ba} \cdot \tilde{\mathbf{m}}_{ba} + \frac{1}{c^2} |\tilde{\mathbf{m}}_{ba}|^2 \right)$$

The first and last terms are the same for the right and left polarizations:

$$R = \text{Im}(\tilde{\boldsymbol{\mu}}_{ba} \cdot \tilde{\mathbf{m}}_{ba})$$

BUT RIGHT ANSWER

by S.Savikhin, Purdue University

41

Gemini: rotational strength in radians vs absorption difference of left and right polarized light, equation

- $\Delta\epsilon$: $\epsilon_L - \epsilon_R$ (Molar circular dichroism, L/mol-cm).
- λ : Wavelength (nm).

Key Relationships

- **Absorption Difference (ΔA)**: The direct experimental measurement is $\Delta A = A_L - A_R$.
- **Ellipticity (θ) and Radians**: If measured as ellipticity θ in radians, it is related to absorbance difference by $\tan \theta = \frac{E_R - E_L}{E_R + E_L} \approx \Delta A \frac{\ln(10)}{4}$.
- **Small Angle Approximation**: Since ΔA is small, the ellipticity in radians is often approximated directly from the absorbance difference:

$$\Delta A \approx \frac{4\theta_{\text{rad}}}{\ln(10)}$$

Wikipedia

Summary Table of Units

Quantity	Common Unit	Relationship to R
Rotational Strength (R)	10^{-40} cgs (Debye $\cdot$ BM)	Fundamental parameter
CD ($\Delta\epsilon$)	$L \cdot \text{mol}^{-1} \cdot \text{cm}^{-1}$	$\int \Delta\epsilon dv$
Ellipticity (θ)	Millidegrees (m°) or Radians	$\Delta A \cdot 32.982 \approx m^\circ$

Key Takeaway: The rotational strength is the integrated value of the difference in absorption ($\epsilon_L - \epsilon_R$) scaled by a factor that includes fundamental physical constants, allowing us to relate light-matter interaction to molecular geometry.

by S.Savikhin, Purdue University

42

1. The "1/4" Rule is for Dipole Strengths (D)

The factor of 1/4 applies when you are looking at the **Dipole Strength** (the transition's intensity), which has units of D^2 (Debye squared):

$$R = \frac{D_L - D_R}{4}$$

Here, D_L and D_R are the dipole strengths for left and right polarized light, respectively.

2. The Relationship to Molar Circular Dichroism ($\Delta\epsilon$)

Rotational strength R is related to the difference in molar absorption ($\Delta\epsilon = \epsilon_L - \epsilon_R$), but it requires an **integral over the entire absorption band** because R represents the total "area" of the transition, not a single point value.

The standard conversion is:

$$R \propto \int \frac{\epsilon_L(\nu) - \epsilon_R(\nu)}{\nu} d\nu$$

by S.Savikhin, Purdue University

43

1. Fundamental Integral Relationship

The most common form of this relationship uses **molar circular dichroism** ($\Delta\epsilon = \epsilon_L - \epsilon_R$) and the wavenumber ($\tilde{\nu}$):

$$R = \frac{3hc \cdot 10^3 \ln(10)}{32\pi^3 N_A} \int \frac{\Delta\epsilon(\tilde{\nu})}{\tilde{\nu}} d\tilde{\nu}$$

In standard cgs units (where R is in 10^{-40} erg·cm³ or Debye-Bohr magnetons), this simplifies to:

$$R \approx 2.296 \times 10^{-39} \int \frac{\Delta\epsilon(\tilde{\nu})}{\tilde{\nu}} d\tilde{\nu}$$

2. Gaussian Approximation for a Single Band

If the absorption difference band follows a Gaussian distribution, you can convert the rotational strength directly from the peak properties of the band without a complex integral:

$$R \approx 1.23 \times 10^{-42} \cdot \frac{\Delta\epsilon_{\max} \cdot \Delta\tilde{\nu}}{\tilde{\nu}_{\max}}$$

Summary of Units

- R : Rotational Strength (units of 10^{-40} erg·cm³ or 1 Debye-Bohr magneton $\approx 0.9273 \times 10^{-38}$ erg·cm³).
- $\Delta\epsilon$: Molar circular dichroism (M^{-1} cm⁻¹).
- $\tilde{\nu}$: Wavenumber (cm⁻¹). 

by S.Savikhin, Purdue University

44

The relationship between the **dipole strength** (D) and the **molar absorption coefficient** (ϵ) is an integral relationship, just like the one for rotational strength.

In simple terms, D is the **total area** under the absorption curve, while ϵ is the **height** of the curve at a specific frequency.

1. The Fundamental Equation

The dipole strength of an electronic transition is proportional to the integral of the molar absorption coefficient (ϵ) over the absorption band:

$$D = \frac{3hc \cdot 10^3 \ln(10)}{8\pi^3 N_A} \int \frac{\epsilon(\tilde{\nu})}{\tilde{\nu}} d\tilde{\nu}$$

In standard units (where D is in **Debye²** and $\tilde{\nu}$ is in **cm⁻¹**), this simplifies to the commonly used numerical constant:

$$D \approx 9.184 \times 10^{-3} \int \frac{\epsilon(\tilde{\nu})}{\tilde{\nu}} d\tilde{\nu}$$

2. Gaussian Approximation

If the absorption peak looks like a standard bell curve (Gaussian), you can estimate D without doing a complex integral:

$$D \approx 9.184 \times 10^{-3} \frac{\epsilon_{\max} \cdot \Delta\tilde{\nu}_{1/2}}{\tilde{\nu}_{\max}}$$

by S.Savikhin, Purdue University

45

Directional Properties

- **Orbital Magnetic Moment:** For a pure p_z orbital ($m_l = 0$), the orbital magnetic moment along the z -axis is **zero**. This is because the electron has no net angular momentum about the z -axis in this state.
- **Transition Magnetic Dipole Moment:** If a transition occurs (e.g., from a p_x to a p_y orbital), the transition magnetic moment vector is typically directed **perpendicular** to the plane containing the transition's orbital movement.
- **Transition Electric Dipole Moment:** For a common electronic transition like $s \rightarrow p_z$, the transition dipole moment is directed strictly along the **z -axis**.  Reddit +5

by S.Savikhin, Purdue University

46

In a flat (planar) molecule, the **transition magnetic moment ($\vec{m}$)** typically points **perpendicular to the molecular plane**. This directionality is tied to the way charge "rotates" during a transition, similar to how current in a flat loop of wire generates a magnetic field normal to the loop's surface. [Physics LibreTexts +1](#)

Directional Characteristics

- **Perpendicular Orientation:** For transitions that involve "circular" movement of electrons within the plane (such as $\pi \rightarrow \pi^*$ or certain $n \rightarrow \pi^*$ transitions), the resulting magnetic moment is directed along the axis normal to the molecular plane.
- **Magnetic vs. Electric Dipole:** While the **transition electric dipole moment ($\vec{\mu}$)** usually lies **in-plane** for $\pi \rightarrow \pi^*$ transitions, the magnetic counterpart $\vec{m}$ is out-of-plane.
- **Chirality and Optical Activity:** In achiral flat molecules like formaldehyde or benzene, the direction of $\vec{m}$ is often zero or arbitrarily defined unless symmetry is broken. In chiral systems or supramolecular stacks, a strong out-of-plane $\vec{m}$ is a key indicator of **circular dichroism** and magnetic-dipole-allowedness. [National Institutes of Health \(.gov\) +4](#)

by S.Savikhin, Purdue University

47

Key Physical Principles

1. **Right-Hand Rule:** The direction of $\vec{m}$ is determined by the "rotation" of charge density during the transition; using your right hand, fingers curling with the charge rotation will point the thumb in the direction of the magnetic moment.
2. **Symmetry Constraints:** In many planar molecules, transitions that are **electrically forbidden** (where $\vec{\mu} = 0$) can be **magnetically allowed** (where $\vec{m} \neq 0$), specifically if the transition involves a change in orbital angular momentum that mimics a current loop.
3. **Magnitude:** Typical values for strong magnetic transitions in flat aromatic systems are around **0.3 to 0.4 Bohr magnetons (μ_B)**. [National Institutes of Health \(.gov\) +4](#)


by S.Savikhin, Purdue University

48

Formaldehyde (CH₂O):

In its $n \rightarrow \pi^*$ transition, the transition is strictly electric-dipole forbidden by symmetry. It becomes observable primarily through magnetic dipole transitions or by "stealing" intensity via molecular vibrations (vibronic coupling).

Convert to extinction:
 $\epsilon \approx \sigma \times (2.615 \times 10^{20})$



Formaldehyde absorption spectrum in the UV range, re-plotted from the data of Meller et al [30]. Peaks are not fully resolved at the resolution of this spectrum (0.01 nm).

by S.Savikhin, Purdue University

49

Magnetic transition dipole moment

Consider the transition from HOMO (ψ_h) to LUMO (ψ_l)

$$\tilde{m}_{ba} \equiv \langle \psi_b | \tilde{m} | \psi_a \rangle$$

Ground state $\Psi_a = \psi_h(1)\psi_h(2)$

Excited state $\Psi_b = 2^{-1/2}\psi_h(1)\psi_l(2) + 2^{-1/2}\psi_h(2)\psi_l(1)$ Since electrons are indistinguishable, use the Slater-type function for the excited state

Find: $\tilde{m}_{ba} \equiv \langle \psi_b | \tilde{m} | \psi_a \rangle$

$$\tilde{m} = \frac{e}{2m_e c} \vec{r} \times \vec{p}$$

$$\tilde{m}_{ba} = \frac{1}{\sqrt{2}} \left\langle \psi_h(1)\psi_l(2) + \psi_h(2)\psi_l(1) \left| \frac{e}{2m_e c} (\vec{r} \times \{-i\hbar \vec{\nabla}\}) \right| \psi_h(1)\psi_h(2) \right\rangle$$

$$\tilde{m}_{ba} = -i\hbar \frac{1}{\sqrt{2}} \frac{e}{2m_e c} \left[\langle \psi_h(1)\psi_l(2) | \vec{r} \times \vec{\nabla} | \psi_h(1)\psi_h(2) \rangle + \langle \psi_h(2)\psi_l(1) | \vec{r} \times \vec{\nabla} | \psi_h(1)\psi_h(2) \rangle \right]$$

Acts only on transition $h \rightarrow l$

$$\tilde{m}_{ba} = -i\hbar \frac{1}{\sqrt{2}} \frac{e}{2m_e c} \left[\langle \psi_h(1) | \psi_h(1) \rangle \langle \psi_l(2) | \vec{r} \times \vec{\nabla} | \psi_h(2) \rangle + \langle \psi_h(2) | \psi_h(2) \rangle \langle \psi_l(1) | \vec{r} \times \vec{\nabla} | \psi_h(1) \rangle \right]$$

$$\tilde{m}_{ba} = -i\hbar \frac{1}{\sqrt{2}} \frac{e}{2m_e c} \left[\langle \psi_l(2) | \vec{r} \times \vec{\nabla} | \psi_h(2) \rangle + \langle \psi_l(1) | \vec{r} \times \vec{\nabla} | \psi_h(1) \rangle \right]$$

Terms are equal.
The only difference is the integration variable (1 or 2)

$$\tilde{m}_{ba} = -i \frac{\sqrt{2}\hbar e}{2m_e c} \langle \psi_l | \vec{r} \times \vec{\nabla} | \psi_h \rangle$$

Magnetic dipole moment is imaginary!

Note: CGS units, SI: erase speed of light c

by S.Savikhin, Purdue University

50