

MEDAGENTIX AI CLINICAL PORTAL

AUTOMATED MEDICAL DIAGNOSTIC PRESCRIPTION SHEET

Date: 2026-06-04 23:17

Facility: MedAgentix Virtual Care Unit

PATIENT PROFILE		VITALS RECORDED	
Age:	N/A	Heart Rate:	110 bpm
Gender:	N/A	Oxygen Sat. (SpO2):	91%
Vital Alerts:	Oxygen Level=92 (Alert)	Blood Pressure (BP):	165/100 mmHg
		Body Temp:	99.2 °F

PRIMARY DIAGNOSIS: Heart attack

Severity: Critical

ML Model Confidence: 96.4% (HIGH)

Consensus Agreement: 67% Agent Voting

Diagnostic Source Path: Differential Agent


CLINICAL REASONING & RELEVANT PATHWAYS

High confidence (96.4%) from differential agent. Disease 'Heart attack' predicted with strong agreement.

Differential Candidates Assessed: Bronchial Asthma (Score: 0.002)

AI EXPLAINABILITY & BIOMARKER REASONING (SHAP)

Below is the Shapley additive explanations (SHAP) feature importance chart. It shows which patient biomarkers and symptoms contributed to the model's primary diagnostic decision. **Blue bars (positive)** indicate features that supported this diagnosis, while **red bars (negative)** indicate factors that pushed against it.



Rx - PRESCRIBED MEDICATIONS

Medication Name	Dosage	Route	Timing & Frequency
Atorvastatin (Lipitor)	40-80 mg once daily	Oral	Once Daily

Aspirin	325 mg stat, then 75-100 mg daily	Oral (chew immediately)	Once Daily
Clopidogrel (Plavix)	75 mg once daily	Oral	Once Daily
Nitroglycerin (GTN)	0.4 mg sublingual as needed	Sublingual	Once Daily
Metoprolol (Lopressor)	25-50 mg twice daily	Oral	Once Daily

RECOMMENDED CLINICAL DIAGNOSTIC LAB TESTS

• ECG (Electrocardiogram) • Troponin-I / Troponin-T • Chest X-Ray • Echocardiography • CBC (Complete Blood Count)

CRITICAL CLINICAL ALERTS & TREATMENT PLAN

- Alert 1: [!] Geriatric patient -- consider reduced dosage and potential drug interactions. Monitor renal and hepatic function.
- Alert 2: [!] Atorvastatin (Lipitor): Contraindicated in Active liver disease, pregnancy. Verify patient history before administration.
- Alert 3: [!] Aspirin: Contraindicated in Active bleeding, aspirin allergy. Verify patient history before administration.
- Alert 4: [!] Clopidogrel (Plavix): Contraindicated in Active pathological bleeding. Verify patient history before administration.
- Alert 5: [!] Nitroglycerin (GTN): Contraindicated in Severe hypotension, use of PDE5 inhibitors. Verify patient history before administration.
- Alert 6: [!] Metoprolol (Lopressor): Contraindicated in Severe bradycardia, heart block. Verify patient history before administration.
- Alert 7: [!!] Severe/Critical condition -- hospitalization may be required. Ensure immediate medical supervision.

Care Plan Instruction: Emergency intervention. ICU admission may be required.

Clinical Follow-Up: Immediate specialist intervention required.

Signature: _____
Attending Virtual Physician, MD

Authorized Verification Seal:
[MedAgentix Autonomous System Validated]

■ *DISCLAIMER: This is an AI-generated diagnostic assessment for informational purposes only. It does NOT constitute medical advice. Always consult a qualified healthcare professional.*