



FRIEDRICH-ALEXANDER  
UNIVERSITÄT  
ERLANGEN-NÜRNBERG



## Lectures on Quantum Theory

---

Course delivered in 2015 by

**Dr Frederic P. Schuller**

*Friedrich-Alexander-Universität Erlangen-Nürnberg,  
Institut für Theoretische Physik III*

*E-mail:* [fps@aei.mpg.de](mailto:fps@aei.mpg.de)

Notes taken by Simon Rea & Richie Dadhley & Edited by Jonah Herr

*E-mail:* [s.rea.hw@gmail.com](mailto:s.rea.hw@gmail.com)

*E-mail:* [richie@dadhley.com](mailto:richie@dadhley.com)

*E-mail:* [jonah.herr@tutanota.de](mailto:jonah.herr@tutanota.de)

Last updated on April 16, 2019

## ACKNOWLEDGMENTS

This set of lecture notes accompanies Frederic Schuller's course on Quantum Theory, taught in the summer of 2015 at the Friedrich-Alexander-Universität Erlangen-Nürnberg as part of the Elite Graduate Programme.

The entire course is hosted on YouTube at the following address:

[www.youtube.com/playlist?list=PLPH7f\\_7ZlzxQVx5jRjbfRGEzWY\\_upS5K6](http://www.youtube.com/playlist?list=PLPH7f_7ZlzxQVx5jRjbfRGEzWY_upS5K6)

These lecture notes are not endorsed by Dr. Schuller or the University.

While I have tried to correct typos and errors made during the lectures (some helpfully pointed out by YouTube commenters), I have also taken the liberty to add and/or modify some of the material at various points in the notes. Any errors that may result from this are, of course, mine.

If you have any comments regarding these notes, feel free to get in touch. Visit my blog for the most up to date version of these notes

<http://mathswithphysics.blogspot.com>

My gratitude goes to Dr. Schuller for lecturing this course and for making it available on YouTube.

*Simon Rea*

I have picked up the notes from where Simon left off (Lecture 8). I have also been through the first lectures and added small details that I thought helpful. I have tried to stay consistent with Simon's inclusion of additional valuable material throughout the remainder of the course. As with Simon's comment above, any mistakes made because of this are, of course, mine.

I have also made the up-to-date version of the notes available on my blog site

<https://richie291.wixsite.com/theoreticalphysics>

I would like to extend a message of thanks to Simon for providing these notes (and the notes for Dr. Schuller's other courses) online, I have personally found them very useful.

I would equally like to thank Jonah Herr for looking over these notes and editing them to make them more readable. This is done both by including more information (such as the table in section 3.4) as well as in more subtle ways (such as introducing a diamond symbol to indicate the end of a definition).

I would also like to show my gratitude to Dr. Schuller for putting his courses on YouTube, I have found them both very informative and interesting, a credit to his brilliant teaching ability.

*Richie Dadhley*

---

## CONTENTS

---

<b>Introduction</b>	<b>1</b>
<b>1 Axioms of quantum mechanics</b>	<b>2</b>
1.1 Desiderata	2
1.2 The axioms of quantum mechanics	5
<b>2 Banach spaces</b>	<b>10</b>
2.1 Generalities on Banach spaces	10
2.2 Bounded linear operators	11
2.3 Extension of bounded linear operators	19
<b>3 Separable Hilbert Spaces</b>	<b>23</b>
3.1 Relationship between norms and inner products	23
3.2 Hamel versus Schauder	31
3.3 Separable Hilbert spaces	33
3.4 Unitary maps	35
<b>4 Projectors, bras and kets</b>	<b>39</b>
4.1 Projectors	39
4.2 Closed linear subspaces	40
4.3 Orthogonal projections	41
4.4 Riesz representation theorem, bras and kets	44
<b>5 Measure theory</b>	<b>48</b>
5.1 General measure spaces and basic results	48
5.2 Borel $\sigma$ -algebras	54
5.3 Lebesgue measure on $\mathbb{R}^d$	56
5.4 Measurable maps	58
5.5 Push-forward of a measure	59
<b>6 Integration of measurable functions</b>	<b>60</b>
6.1 Characteristic and simple functions	60
6.2 Integration of non-negative measurable simple functions	62
6.3 Integration of non-negative measurable functions	64
6.4 Lebesgue integrable functions	70
6.5 The function spaces $L^p(M, \Sigma, \mu)$	74

<b>7</b>	<b>Self-adjoint and essentially self-adjoint operators</b>	<b>79</b>
7.1	Adjoint operators	79
7.2	The adjoint of a symmetric operator	81
7.3	Closability, closure, closedness	82
7.4	Essentially self-adjoint operators	84
7.5	Criteria for self-adjointness and essential self-adjointness	85
<b>8</b>	<b>Spectra and perturbation theory</b>	<b>86</b>
8.1	Resolvent map and spectrum	86
8.2	The spectrum of a self-adjoint operator	87
8.3	Perturbation theory for point spectra of self-adjoint operators	88
<b>9</b>	<b>Case study: momentum operator</b>	<b>95</b>
9.1	The Momentum Operator	95
9.2	Absolutely Continuous Functions and Sobolev Spaces	95
9.3	Momentum Operator on a Compact Interval	97
9.4	Momentum Operator on a Circle	101
<b>10</b>	<b>Inverse spectral theorem</b>	<b>103</b>
10.1	Projection-valued measures	104
10.2	Real and Complex Valued Borel Measures Induced by a PVM	106
10.3	Integration With Respect to a PVM	106
10.4	The Inverse Spectral Theorem	113
<b>11</b>	<b>Spectral theorem</b>	<b>114</b>
11.1	Measurable Function Applied To A Spectrally Decomposable Self Adjoint Operator	114
11.2	Reconstruct PVM From a Spectrally Decomposable, Self Adjoint Operator	115
11.3	Construction Of PVM From A Self Adjoint Operator	118
11.4	Commuting Operators	119
<b>12</b>	<b>Stone's Theorem and Construction of Observables</b>	<b>122</b>
12.1	One Parameter Groups and Their Generators	122
12.2	Stone's Theorem	124
12.3	Domains of Essential Self Adjointness ("Cores")	127
12.4	Position, Momentum and Angular Momentum	127
12.5	Schwartz Space $S(\mathbb{R}^d)$	129
<b>13</b>	<b>Spin</b>	<b>131</b>
13.1	General Spin	133
13.2	Derivation of Pure Point Spectrum	133
13.3	Pure Spin- $j$ Systems	139

<b>14 Composite Systems</b>	<b>141</b>
14.1 Tensor Product of Hilbert Spaces	142
14.2 Practical Rules for Tensor Products of Vectors	147
14.3 Tensor Product Between Operators	147
14.4 Symmetric and Antisymmetric Tensor Products	148
14.5 Collapse of Notation	150
14.6 Entanglement	150
<b>15 Total Spin of Composite System</b>	<b>152</b>
15.1 Total Spin	152
15.2 Eigenbasis For The Composite System in Terms of Simultaneous Eigenvec- tors of $A^2 \otimes \mathbb{1}$ , $\mathbb{1} \otimes B^2$ , $A_3 \otimes \mathbb{1}$ and $\mathbb{1} \otimes B^3$	154
15.3 Conversion to Eigenbasis in Terms of $J^2$ , $J_3$ , $A^2 \otimes \mathbb{1}$ , $\mathbb{1} \otimes B^2$	154
15.4 Value of $m$	155
15.5 Clebsch-Gordan Coefficients for Maximal $j$	156
15.6 Clebsch-Gordan Coefficients For Lower Than Max $j$	157
15.7 Total Spin of Composite System	158
<b>16 Quantum Harmonic Oscillator</b>	<b>160</b>
16.1 What People Say	160
16.2 The Quantum Harmonic Oscillator	161
16.3 The Energy Spectrum	162
<b>17 Measurements of Observables</b>	<b>167</b>
17.1 Spectral Decomposition of $H$ For The Quantum Harmonic Oscillator	167
17.2 Probability to Measure a Certain Value	169
17.3 Measurements of Pure States	170
17.4 Preparation of Pure States	173
17.5 Mixed States	174
<b>18 The Fourier Operator</b>	<b>176</b>
18.1 The Fourier Operator on Schwartz Space	176
18.2 Inverse of Fourier Operator	179
18.3 Extension of $\mathfrak{F}$ to $L^2(\mathbb{R}^d)$	181
18.4 Convolutions	182
<b>19 The Schrodinger Operator</b>	<b>185</b>
19.1 Domain of Self Adjointness of $H_{\text{free}}$	185
19.2 Spectrum of $H_{\text{free}}$	186
19.3 Time Evolution	187
19.4 Physical Interpretation (for the Patient Observer)	188


<b>20 Periodic Potentials</b>	<b>190</b>
20.1 Basics of Rigged Hilbert Space	191
20.2 Fundamental Solutions and Fundamental Matrix	192
20.3 Translation Operator	193
20.4 Application to Our Quantum Problem	195
20.5 Energy Bands	196
20.6 Quantitative Calculation (Outline)	196
<b>21 Relativistic Quantum Mechanics</b>	<b>197</b>
21.1 Heuristic Derivation of the Schrödinger Equation	197
21.2 Heuristic Derivation of the Relativistic Schrödinger Equation (Klein-Gordon Equation)	198
21.3 Dirac Equation	199
21.4 The Deep Route of the Problem	200
21.5 Feynman Diagrams	201
<b>Further readings</b>	<b>203</b>

---

## INTRODUCTION

Quantum mechanics has a reputation for being a difficult subject, and it really deserves that reputation. It is, indeed, very difficult. This is partly due to the fact that, unlike classical mechanics or electromagnetism, it is very different from what we feel the world is. But the fault is on us. The world does *not* behave in the way that we feel it should from our everyday experience. Of course, the reason why classical mechanics works so well for modelling stones, rockets and planets is that the masses involved are much larger than those of, say, elementary particles, while the speeds are much slower than the speed of light. However, even the stone that one throws doesn't follow a trajectory governed by Newton's axioms. In fact, it doesn't follow a trajectory at all. The very idea of a point particle following a trajectory turns out to be entirely wrong. So don't worry if your classical mechanics course didn't go well. It's all wrong anyway!

We know from the double slit experiment that the reality is more complicated. The result of the experiment can be interpreted as the electron going through both slits and neither slit at the same time, and in fact taking every possible path. The experiment has been replicated with objects much larger than an electron<sup>1</sup>, and in principle it would work even if we used a whale (which is not a fish!).



---

<sup>1</sup>Eibenberger et al., Matter-wave interference with particles selected from a molecular library with masses exceeding 10000 amu, <https://arxiv.org/abs/1310.8343>

## 1 Axioms of quantum mechanics

People discovered what was wrong with classical mechanics bit by bit and, consequently, the historical development of quantum mechanics was highly “non-linear”. Rather than following this development, we will afford the luxury of having a well-working theory of quantum mechanics, and we will present it from the foundations up. We begin by writing down a list things we would like to have.

## 1.1 Desiderata<sup>2</sup>

A working theory of quantum mechanics would need to account for the following.

- (a) *Measurements of observables, unlike in classical mechanics, don't just range over an interval  $I \subseteq \mathbb{R}$ .*

Recall that in classical mechanics an observable is a map  $F: \Gamma \rightarrow \mathbb{R}$ , where  $\Gamma$  is the phase space of the system, typically given by the cotangent space  $T^*Q$  of some configuration manifold  $Q$ . The map is taken to be at least continuous with respect to the standard topology on  $\mathbb{R}$  and an appropriate topology on  $\Gamma$ , and hence if  $\Gamma$  is connected, we have  $F(\Gamma) = I \subseteq \mathbb{R}$ .

Consider, for instance, the two-body problem. We have a potential  $V(r) = -\frac{1}{r}$  and, assuming that the angular momentum  $L$  is non-zero, the energy observable (or Hamiltonian)  $H$  satisfies  $H(\Gamma) = [E_{\min}, \infty) \subset \mathbb{R}$ .

However, measurements of the spectrum of the hydrogen atom give the following values for the energies (in electronvolts) assumed by the electron

$$\{-13.6 \times \frac{1}{n^2} \mid n \in \mathbb{N}^+\} \cup (0, \infty).$$

Hence, we need to turn to new mathematics in which we can define a notion of observable that allows for a spectrum of measurement results for a quantum observable  $A$  of the form

$$\sigma(A) = \text{discrete part} \cup \text{continuous part}.$$

An example would be the energies of the hydrogen atom

Figure 1 shows a schematic representation of the energy spectrum of a hydrogen atom. The horizontal axis represents energy in eV. A vertical line is drawn at -13.6 eV. To the left of this line, there are several discrete energy levels represented by vertical tick marks. To the right of this line, there is a shaded gray region representing a continuous spectrum. The axis is labeled with -13.6 eV and 0 eV.

Note that one of the parts may actually be empty. For instance, as we will later show, the simple quantum harmonic oscillator has the following energy spectrum

$$\sigma(H) = \begin{array}{c} \text{---} \frac{1}{2}\hbar\omega \hspace{1.5cm} (\frac{1}{2} + n)\hbar\omega \text{---} \end{array}$$

while the spectrum of the position operator  $Q$  is  $\sigma(Q) = \mathbb{R}$ .

<sup>2</sup>Educated term for “wishlist”.




Also, the continuous part need not be connected, as is the case with spectrum of the Hamiltonian an electron in a periodic potential

$$\sigma(H) = \text{---} \boxed{\phantom{000}} \text{---} \boxed{\phantom{000}} \text{---} \boxed{\phantom{000}} \text{---} \rightarrow$$

It turns out that self-adjoint linear maps on a complex Hilbert space provide a suitable formalism to describe the observables of quantum mechanics.

- (b) *An irreducible impact that each measurement has on the state of a quantum system.*


The crucial example demonstrating this is the Stern-Gerlach experiment, which consists in the following. Silver atoms are heated up in an oven and sent against a screen with a hole. The atoms passing through the hole are then subjected to an inhomogeneous magnetic field, which deflects them according to the component of their angular momentum in the direction of the field. Finally, a screen detects the various deflections.




Since the angular momentum distribution of the silver atoms coming from the oven is random, we would expect an even distribution of values of the component along the direction of the magnetic field to be recorded on the final screen, as in  $S$ . However, the impact pattern actually detected is that on the  $W\tau\varphi$  screen. In fact, 50% of the incoming atoms impact at the top and we say that their angular momentum component is  $\uparrow$ , and the other 50% hit the bottom region, and we say that their angular momentum component is  $\downarrow$ . This is another instance of our earlier point: there seem to be only two possible values for the component of angular momentum along the direction of the magnetic field, i.e. the spectrum is discrete. Hence, this is not particularly surprising at this point.

Let us now consider successive iterations of this experiment. Introduce some system of cartesian coordinates  $(x, y, z)$  and let  $\text{SG}(x)$  and  $\text{SG}(z)$  denotes a Stern-Gerlach apparatus whose magnetic field points in the  $x$  and  $z$ -direction, respectively.

Suppose that we sent the atoms through a first  $\text{SG}(z)$  apparatus, and then we use the  $z^\uparrow$ -output as the input of a second  $\text{SG}(z)$  apparatus.




The second  $\text{SG}(z)$  apparatus finds no  $z^\downarrow$ -atoms. This is not surprising since, intuitively, we “filtered out” all the  $z^\downarrow$ -atoms with the first apparatus. Suppose now that we feed the  $z^\uparrow$  output of a  $\text{SG}(z)$  apparatus into a  $\text{SG}(x)$  apparatus.



Experimentally, we find that about half of the atoms are detected in the state  $x^\uparrow$  and half in the state  $x^\downarrow$ . This is, again, not surprising since we only filtered out the  $z^\downarrow$  atoms, and hence we can interpret this result as saying that the  $x^\uparrow$ ,  $x^\downarrow$  states are independent from the  $z^\uparrow$ ,  $z^\downarrow$ .

If our ideas of “filtering states out” is correct, then feeding the  $x^\uparrow$ -output of the previous set-up to another  $\text{SG}(z)$  apparatus should clearly produce a 100%  $z^\uparrow$ -output, since we already filtered out all the  $z^\downarrow$  ones in the previous step.



Surprisingly, the output is again 50-50. The idea behind this result is the following. The  $\text{SG}(z)$  apparatus left the atoms in a state such that a repeated measurement with the  $\text{SG}(z)$  apparatus would give the same result, and similarly for the  $\text{SG}(x)$

apparatus. However, the measurement of the  $\text{SG}(x)$  apparatus somehow altered the state of the atoms in such a way as to “reset” them with respect to a measurement by the  $\text{SG}(z)$  apparatus. For more details on the Stern-Gerlach experiment and further conclusions one can draw from its results, you should consult the book *Modern Quantum Mechanics* by J. J. Sakurai. The conclusion that we are interested in here is that measurements can alter the state of a system.

- (c) *Even if the state  $\rho$  of a quantum system is completely known, the only prediction one can make for the measurement of some observable  $A$  is the probability that the measured value, which is an element of the spectrum  $\sigma(A)$ , lies within a Borel-measurable subset  $E \subseteq \mathbb{R}$ , denoted by  $\mu_\rho^A(E)$ .*

In particular, one cannot predict which concrete outcome an isolated measurement will produce. This is even more annoying given that the precise impact that a measurement has on the state of the system (see previous point) depends on the observed outcome of the measurement.

A suitable theory that accommodates all known experimental facts has been developed between 1900 and 1927 on the physics side by, among others, Schrödinger, Heisenberg and Dirac, and on the mathematical side almost single-handedly by von Neumann who invented a massive proportion of a field known today as functional analysis.

## 1.2 The axioms of quantum mechanics

We will now present the axioms of quantum mechanics by using notions and terminology that will be defined later in the course. In this sense, this section constitutes a preview of the next few lectures.

**Axiom 1** (Quantum systems and states). *To every quantum system there is associated a separable complex Hilbert space  $(\mathcal{H}, +, \cdot, \langle \cdot | \cdot \rangle)$ . The states of the system are all positive, trace-class linear maps  $\rho: \mathcal{H} \rightarrow \mathcal{H}$  for which  $\text{Tr } \rho = 1$ .*

Remark 1.1. Throughout the quantum mechanics literature, it is stated that the unit, or normalised, elements  $\psi \in \mathcal{H}$  (that is,  $\langle \psi | \psi \rangle = 1$ ) are the states of the quantum system. This is not correct.

States can be pure or mixed. A state  $\rho: \mathcal{H} \rightarrow \mathcal{H}$  is called *pure* if

$$\exists \psi \in \mathcal{H} : \forall \alpha \in \mathcal{H} : \rho(\alpha) = \frac{\langle \psi | \alpha \rangle}{\langle \psi | \psi \rangle} \psi.$$

Thus, we can associate to each pure state  $\rho$  an element  $\psi \in \mathcal{H}$ . However, this correspondence is not one-to-one. Even if we restrict to pure states and impose the normalisation condition, there can be many  $\psi \in \mathcal{H}$  representing the same pure state  $\rho$ .

Therefore, it is wrong to say that the states of the quantum system are the normalised elements of the Hilbert space, since they do not represent all the states of the system, and do not even represent uniquely the states that they do represent.

The terms used in Axiom 1 are defined as follows.

**Definition.** A *complex Hilbert space* is a tuple  $(\mathcal{H}, +, \cdot, \langle \cdot | \cdot \rangle)$  where

- $\mathcal{H}$  is a set
- $+$  is a map  $+: \mathcal{H} \times \mathcal{H} \rightarrow \mathcal{H}$
- $\cdot$  is a map  $\cdot: \mathbb{C} \times \mathcal{H} \rightarrow \mathcal{H}$  (typically suppressed in the notation)

such that the triple  $(\mathcal{H}, +, \cdot)$  is a vector space over  $\mathbb{C}$ , and

- $\langle \cdot | \cdot \rangle$  is a *sesqui-linear<sup>3</sup> inner product*, i.e. a map  $\langle \cdot | \cdot \rangle: \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{C}$  satisfying
    - (i)  $\langle \varphi | \psi \rangle = \overline{\langle \psi | \varphi \rangle}$  (conjugate symmetry/Hermitian property)
    - (ii)  $\langle \varphi | z\psi_1 + \psi_2 \rangle = z\langle \varphi | \psi_1 \rangle + \langle \varphi | \psi_2 \rangle$  (linearity in the second argument)
    - (iii)  $\langle \psi | \psi \rangle \geq 0$  and  $\langle \psi | \psi \rangle = 0 \Leftrightarrow \psi = 0_{\mathcal{H}}$  (positive-definiteness)
- for all  $\varphi, \psi_1, \psi_2 \in \mathcal{H}$  and  $z \in \mathbb{C}$ ,

and moreover

- $\mathcal{H}$  is a *complete metric space* with respect to the metric induced by the norm induced in turn by the sesqui-linear map  $\langle \cdot | \cdot \rangle$ . Explicitly, for every sequence  $\phi: \mathbb{N} \rightarrow \mathcal{H}$  that satisfies the *Cauchy property*, namely

$$\forall \varepsilon > 0 : \exists N \in \mathbb{N} : \forall n, m \geq N : \|\phi_n - \phi_m\| < \varepsilon,$$

where  $\phi_n := \phi(n)$  and  $\|\psi\| := \sqrt{\langle \psi | \psi \rangle}$ , then the sequence converges in  $\mathcal{H}$ , i.e.

$$\exists \varphi \in \mathcal{H} : \forall \varepsilon > 0 : \exists N \in \mathbb{N} : \forall n \geq N : \|\varphi - \phi_n\| < \varepsilon.$$

◇

Note that the  $\mathbb{C}$ -vector space  $(\mathcal{H}, +, \cdot)$  need not be finite-dimensional and, in fact, we will mostly work with infinite-dimensional Hilbert spaces.

**Definition.** A map  $A: \mathcal{D}_A \rightarrow \mathcal{H}$ , where the subspace  $\mathcal{D}_A \subseteq \mathcal{H}$  is called the *domain* of  $A$ , is a *linear map* if

$$\forall \varphi, \psi \in \mathcal{D}_A : \forall z \in \mathbb{C} : A(z\varphi + \psi) = zA(\varphi) + A(\psi).$$

◇

From now on, if there is no risk of confusion, we will write  $A\varphi := A(\varphi)$  in order to spare some brackets. We will be particularly interested in special types of linear map.

---

<sup>3</sup>sesqui is Latin for “one and a half”.

**Definition.** A linear map  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  is *densely defined* if  $\mathcal{D}_A$  is *dense* in  $\mathcal{H}$ , i.e.

$$\forall \psi \in \mathcal{H} : \forall \varepsilon > 0 : \exists \alpha \in \mathcal{D}_A : \|\alpha - \psi\| < \varepsilon.$$

◇

**Definition.** A linear map  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  is said to be *positive* if

$$\forall \psi \in \mathcal{D}_A : \langle \psi | A \psi \rangle \geq 0.$$

◇

**Definition.** A linear map  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  is said to be of *trace-class* if  $\mathcal{D}_A = \mathcal{H}$  and, for any orthonormal basis  $\{e_n\}$  of  $\mathcal{H}$ , the sum/series

$$\sum_n \langle e_n | A e_n \rangle < \infty.$$

◇

If  $A: \mathcal{H} \rightarrow \mathcal{H}$  is of trace-class, one can show that the value of  $\sum_n \langle e_n | A e_n \rangle$  does not depend on the choice of orthonormal basis  $\{e_n\}$ .

**Definition.** Let  $A: \mathcal{H} \rightarrow \mathcal{H}$  be of trace-class. Then the *trace* of  $A$  is

$$\text{Tr } A := \sum_n \langle e_n | A e_n \rangle$$

where  $\{e_n\}$  is any orthonormal basis of  $\mathcal{H}$ .

◇

**Axiom 2** (Observables). *The observables of a quantum system are the self-adjoint linear maps  $A: \mathcal{D}_A \rightarrow \mathcal{H}$ .*

While the notion of a self-adjoint map is easy to define in finite-dimensional spaces, it is much more subtle for infinite-dimensional spaces.

**Definition.** A densely defined linear map  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  is said to be of *self-adjoint* if it coincides with its adjoint map  $A^*: \mathcal{D}_{A^*} \rightarrow \mathcal{H}$ , that is

- $\mathcal{D}_A = \mathcal{D}_{A^*}$
- $\forall \varphi \in \mathcal{D}_A : A \varphi = A^* \varphi.$

◇

**Definition.** The *adjoint map*  $A^*: \mathcal{D}_{A^*} \rightarrow \mathcal{H}$  of a linear map  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  is defined by

- $\mathcal{D}_{A^*} := \{\psi \in \mathcal{H} \mid \forall \alpha \in \mathcal{D}_A : \exists \eta \in \mathcal{H} : \langle \psi | A \alpha \rangle = \langle \eta | \alpha \rangle\}$
- $A^* \psi := \eta.$

◇

We will later show that the adjoint map is well-defined, i.e. for each  $\alpha \in \mathcal{D}_A$  and  $\psi \in \mathcal{H}$  there exists at most one  $\eta \in \mathcal{H}$  such that  $\langle \psi | A \alpha \rangle = \langle \eta | \alpha \rangle$ .

*Remark 1.2.* If we defined  $\mathcal{D}_{A^*}$  by requiring that  $\eta \in \mathcal{D}_A$ , we would obtain a notion of self-adjointness which has undesirable properties. In particular, the spectrum (to be defined later) of a self-adjoint operator would not be guaranteed to be a subset of  $\mathbb{R}$ .

**Axiom 3** (Measurement). *The probability that a measurement of an observable  $A$  on a system that is in the state  $\rho$  yields a result in the Borel set  $E \subseteq \mathbb{R}$  is given by*

$$\mu_\rho^A(E) := \text{Tr}(P_A(E) \circ \rho)$$

*where the map  $P_A: \text{Borel}(\mathbb{R}) \rightarrow \mathcal{L}(\mathcal{H})$ , from the Borel-measurable subsets of  $\mathbb{R}$  to the Banach space of bounded linear maps on  $\mathcal{H}$ , is the unique projection-valued measure that is associated with the self-adjoint map  $A$  according to the spectral theorem.*

We will later see that the composition of a bounded map with a trace-class map is again of trace-class, so that  $\text{Tr}(P_A(E) \circ \rho)$  is well-defined. For completeness, the spectral theorem states that for any self-adjoint map  $A$  there exists a projection-valued measure  $P_A$  such that  $A$  can be represented in terms of the Lebesgue-Stieltjes integral as

$$A = \int_{\mathbb{R}} \lambda \, dP_A(\lambda).$$

This is the infinite-dimensional analogue of the diagonalisation theorem for symmetric or Hermitian matrices on finite-dimensional vector spaces, and it is the theorem in which the first half of the course will find its climax.

**Axiom 4** (Unitary dynamics). *In a time interval  $(t_1, t_2) \subseteq \mathbb{R}$  in which no measurement occurs, the state  $\rho$  at time  $t_1$ , denoted  $\rho(t_1)$ , is related to the state  $\rho$  at time  $t_2$ , denoted  $\rho(t_2)$ , by*

$$\rho(t_2) = U(t_2 - t_1) \rho(t_1) U^{-1}(t_2 - t_1)$$

*with the unitary evolution operator  $U$  defined as*

$$U(t) := \exp\left(-\frac{i}{\hbar} H t\right),$$

*where  $H$  is the energy observable and, for any observable  $A$  and  $f: \mathbb{R} \rightarrow \mathbb{C}$ , we define*

$$f(A) := \int_{\mathbb{R}} f(\lambda) \, dP_A(\lambda).$$

Note that, as was the case for the previous axiom, the spectral theorem is crucial since it is needed to define the unitary evolution operator.

**Axiom 5** (Projective dynamics). *The state  $\rho_{\text{after}}$  of a quantum system immediately following the measurement of an observable  $A$  is*

$$\rho_{\text{after}} := \frac{P_A(E) \circ \rho_{\text{before}} \circ P_A(E)}{\text{Tr}(P_A(E) \circ \rho_{\text{before}} \circ P_A(E))}$$

*where  $\rho_{\text{before}}$  is the state immediately preceding the measurement and  $E \subseteq \mathbb{R}$  is the smallest Borel set in which the actual outcome of the measurement happened to lie.*

## 2 Banach spaces

Hilbert spaces are a special type of a more general class of spaces known as Banach spaces. We are interested in Banach spaces not just for the sake generality, but also because they naturally appear in Hilbert space theory. For instance, the space of bounded linear maps on a Hilbert space is not itself a Hilbert space, but only a Banach space.

### 2.1 Generalities on Banach spaces

We begin with some basis notions from metric space theory.

**Definition.** A *metric space* is a pair  $(X, d)$ , where  $X$  is a set and  $d$  is a *metric* on  $X$ , that is, a map  $d: X \times X \rightarrow \mathbb{R}$  satisfying

$$(i) \quad d(x, x) \geq 0 \quad \text{(non-negativity)}$$

$$(ii) \quad d(x, y) = 0 \Leftrightarrow x = y \quad \text{(identity of indiscernibles)}$$

$$(iii) \quad d(x, y) = d(y, x) \quad \text{(symmetry)}$$

$$(iv) \quad d(x, z) \leq d(x, y) + d(y, z) \quad \text{(triangle inequality)}$$

for all  $x, y, z \in X$ .  $\diamond$

**Definition.** A sequence  $\{x_n\}_{n \in \mathbb{N}}$  in a metric space  $(X, d)$  is said to *converge* to an element  $x \in X$ , written  $\lim_{n \rightarrow \infty} x_n = x$ , if

$$\forall \varepsilon > 0 : \exists N \in \mathbb{N} : \forall n \geq N : d(x_n, x) < \varepsilon.$$

$\diamond$

A sequence in a metric space can converge to at most one element.

**Definition.** A *Cauchy sequence* in a metric space  $(X, d)$  is a sequence  $\{x_n\}_{n \in \mathbb{N}}$  such that

$$\forall \varepsilon > 0 : \exists N \in \mathbb{N} : \forall n, m \geq N : d(x_n, x_m) < \varepsilon.$$

$\diamond$

Any convergent sequence is clearly a Cauchy sequence.



**Definition.** A metric space  $(X, d)$  is said to be *complete* if every Cauchy sequence converges to some  $x \in X$ .  $\diamond$

A natural metric on a vector space is that induced by a norm. Note that every convergent sequence is a Cauchy sequence, the converse does not hold in general.

**Definition.** A *normed space* is a (complex) vector space  $(V, +, \cdot)$  equipped with a *norm*, that is, a map  $\|\cdot\|: V \rightarrow \mathbb{R}$  satisfying

$$(i) \quad \|f\| \geq 0 \quad (\text{non-negativity})$$

$$(ii) \quad \|f\| = 0 \Leftrightarrow f = 0 \quad (\text{definiteness})$$

$$(iii) \quad \|z \cdot f\| = |z| \|f\| \quad (\text{homogeneity/scalability})$$

$$(iv) \quad \|f + g\| \leq \|f\| + \|g\| \quad (\text{triangle inequality/sub-additivity})$$

for all  $f, g \in V$  and all  $z \in \mathbb{C}$ .  $\diamond$

Once we have a norm  $\|\cdot\|$  on  $V$ , we can define a metric  $d$  on  $V$  by

$$d(f, g) := \|f - g\|.$$

Then we say that the normed space  $(V, \|\cdot\|)$  is *complete* if the metric space  $(V, d)$ , where  $d$  is the metric induced by  $\|\cdot\|$ , is complete. Note that we will usually suppress inessential information in the notation, for example writing  $(V, \|\cdot\|)$  instead of  $(V, +, \cdot, \|\cdot\|)$ .

**Definition.** A *Banach space* is a complete normed vector space.  $\diamond$

## 2.2 Bounded linear operators

As usual in mathematics, once we introduce a new types of structure, we also want study maps between instances of those structures, with extra emphasis placed on the structure-preserving maps. We begin with linear maps from a normed space to a Banach space.

**Definition.** Let  $(V, \|\cdot\|_V)$  be a normed space and  $(W, \|\cdot\|_W)$  a Banach space. A linear map, also called a linear operator,  $A: V \rightarrow W$  is said to be *bounded* if

$$\sup_{f \in V} \frac{\|Af\|_W}{\|f\|_V} < \infty.$$

$\diamond$

Note that the quotient is not defined for  $f = 0$ . Hence, to be precise, we should write  $V \setminus \{0\}$  instead of just  $V$ . Let us agree that is what mean in the above definition. There are several equivalent characterisations of the boundedness property.

**Proposition 2.1.** *A linear operator  $A: V \rightarrow W$  is bounded if, and only if, any of the following conditions are satisfied.*

- (i)  $\sup_{\|f\|_V=1} \|Af\|_W < \infty$
- (ii)  $\exists k > 0 : \forall f \in V : \|f\|_V \leq 1 \Rightarrow \|Af\|_W \leq k$
- (iii)  $\exists k > 0 : \forall f \in V : \|Af\|_W \leq k\|f\|_V$
- (iv) *the map  $A: V \rightarrow W$  is continuous with respect to the topologies induced by the respective norms on  $V$  and  $W$*
- (v) *the map  $A$  is continuous at  $0 \in V$ .*

The first one of these follows immediately from the homogeneity of the norm. Indeed, suppose that  $\|f\|_V \neq 1$ . Then

$$\frac{\|Af\|_W}{\|f\|_V} = \|f\|_V^{-1} \|Af\|_W = \|A(\|f\|_V^{-1} f)\|_W = \|A\tilde{f}\|_W$$

where  $\tilde{f} := \|f\|_V^{-1} f$  is such that  $\|\tilde{f}\|_V = 1$ . Hence, the boundedness property is equivalent to condition (i) above.

**Definition.** Let  $A: V \rightarrow W$  be a bounded operator. The *operator norm* of  $A$  is defined as

$$\|A\| := \sup_{\|f\|_V=1} \|Af\|_W = \sup_{f \in V} \frac{\|Af\|_W}{\|f\|_V}.$$

◇

Example 2.2. Let  $\text{id}_W: W \rightarrow W$  be the identity operator on a Banach space  $W$ . Then

$$\sup_{f \in W} \frac{\|\text{id}_W f\|_W}{\|f\|_W} = \sup_{f \in W} 1 = 1 < \infty.$$

Hence,  $\text{id}_W$  is a bounded operator and has unit norm.

Example 2.3. Denote by  $C_{\mathbb{C}}^1[0, 1]$  the complex vector space of once continuously differentiable complex-valued functions on  $[0, 1]$ . Since differentiability implies continuity, this is a subset of  $C_{\mathbb{C}}^0[0, 1]$ . Moreover, since sums and scaling by a complex number of continuously differentiable functions are again continuously differentiable, this is, in fact, a vector subspace of  $C_{\mathbb{C}}^0[0, 1]$ , and hence also a normed space with the supremum norm  $\|\cdot\|_{\infty}$ .

Consider the first derivative operator

$$\begin{aligned} D: C_{\mathbb{C}}^1[0, 1] &\rightarrow C_{\mathbb{C}}^0[0, 1] \\ f &\mapsto f'. \end{aligned}$$

We know from undergraduate real analysis that  $D$  is a linear operator. We will now show that  $D$  is an unbounded<sup>4</sup> linear operator. That is,

$$\sup_{f \in C_{\mathbb{C}}^1[0,1]} \frac{\|Df\|_{\infty}}{\|f\|_{\infty}} = \infty.$$

Note that, since the norm is a function into the real numbers, both  $\|Df\|_{\infty}$  and  $\|f\|_{\infty}$  are always finite for any  $f \in C_{\mathbb{C}}^1[0,1]$ . Recall that the supremum of a set of real numbers is its least upper bound and, in particular, it need not be an element of the set itself. What we have to show is that the set

$$\left\{ \frac{\|Df\|_{\infty}}{\|f\|_{\infty}} \mid f \in C_{\mathbb{C}}^1[0,1] \right\} \subset \mathbb{R}$$

contains arbitrarily large elements. One way to do this is to exhibit a positively divergent (or unbounded from above) sequence within the set.

Consider the sequence  $\{f_n\}_{n \geq 1}$  where  $f_n(x) := \sin(2\pi nx)$ . We know that sine is continuously differentiable, hence  $f_n \in C_{\mathbb{C}}^1[0,1]$  for each  $n \geq 1$ , with

$$Df_n(x) = D(\sin(2\pi nx)) = 2\pi n \cos(2\pi nx).$$

We have

$$\|f_n\|_{\infty} = \sup_{x \in [0,1]} |f_n(x)| = \sup_{x \in [0,1]} |\sin(2\pi nx)| = \sup[-1,1] = 1$$

and

$$\|Df_n\|_{\infty} = \sup_{x \in [0,1]} |Df_n(x)| = \sup_{x \in [0,1]} |2\pi n \cos(2\pi nx)| = \sup[-2\pi n, 2\pi n] = 2\pi n.$$

Hence, we have

$$\sup_{f \in C_{\mathbb{C}}^1[0,1]} \frac{\|Df\|_{\infty}}{\|f\|_{\infty}} \geq \sup_{\{f_n\}_{n \geq 1}} \frac{\|Df_n\|_{\infty}}{\|f_n\|_{\infty}} = \sup_{n \geq 1} 2\pi n = \infty,$$

which is what we wanted. As an aside, we note that  $C_{\mathbb{C}}^1[0,1]$  is not complete with respect to the supremum norm, but it is complete with respect to the norm

$$\|f\|_{C^1} := \|f\|_{\infty} + \|f'\|_{\infty}.$$

While the derivative operator is still unbounded with respect to this new norm, in general, the boundedness of a linear operator does depend on the choice of norms on its domain and target, as does the numerical value of the operator norm.

*Remark 2.4.* Apart from the “minor” detail that in quantum mechanics we deal with Hilbert spaces, use a different norm than the supremum norm and that the (one-dimensional) momentum operator acts as  $P(\psi) := -i\hbar\psi'$ , the previous example is a harbinger of the fact that the momentum operator in quantum mechanics is unbounded. This will be the case for the position operator  $Q$  as well.

---

<sup>4</sup>Some people take the term unbounded to mean “not necessarily bounded”. We take it to mean “definitely not bounded” instead.

**Lemma 2.5.** *Let  $(V, \|\cdot\|)$  be a normed space. Then, addition, scalar multiplication, and the norm are all sequentially continuous. That is, for any sequences  $\{f_n\}_{n \in \mathbb{N}}$  and  $\{g_n\}_{n \in \mathbb{N}}$  in  $V$  converging to  $f \in V$  and  $g \in V$  respectively, and any sequence  $\{z_n\}_{n \in \mathbb{N}}$  in  $\mathbb{C}$  converging to  $z \in \mathbb{C}$ , we have*

$$(i) \quad \lim_{n \rightarrow \infty} (f_n + g_n) = f + g$$

$$(ii) \quad \lim_{n \rightarrow \infty} z_n f_n = z f.$$

$$(iii) \quad \lim_{n \rightarrow \infty} \|f_n\| = \|f\|$$

*Proof.* (i) Let  $\varepsilon > 0$ . Since  $\lim_{n \rightarrow \infty} f_n = f$  and  $\lim_{n \rightarrow \infty} g_n = g$  by assumption, there exist  $N_1, N_2 \in \mathbb{N}$  such that

$$\begin{aligned} \forall n \geq N_1 : \|f - f_n\| &< \frac{\varepsilon}{2} \\ \forall n \geq N_2 : \|g - g_n\| &< \frac{\varepsilon}{2}. \end{aligned}$$

Let  $N := \max\{N_1, N_2\}$ . Then, for all  $n \geq N$ , we have

$$\begin{aligned} \|(f_n + g_n) - (f + g)\| &= \|f_n - f + g_n - g\| \\ &\leq \|f_n - f\| + \|g_n - g\| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ &= \varepsilon. \end{aligned}$$

Hence  $\lim_{n \rightarrow \infty} (f_n + g_n) = f + g$ .

(ii) Since  $\{z_n\}_{n \in \mathbb{N}}$  is a convergent sequence in  $\mathbb{C}$ , it is bounded. That is,

$$\exists k > 0 : \forall n \in \mathbb{N} : |z_n| \leq k.$$

Let  $\varepsilon > 0$ . Since  $\lim_{n \rightarrow \infty} f_n = f$  and  $\lim_{n \rightarrow \infty} z_n = z$ , there exist  $N_1, N_2 \in \mathbb{N}$  such that

$$\begin{aligned} \forall n \geq N_1 : \|f - f_n\| &< \frac{\varepsilon}{2k} \\ \forall n \geq N_2 : \|z - z_n\| &< \frac{\varepsilon}{2\|f\|}. \end{aligned}$$

Let  $N := \max\{N_1, N_2\}$ . Then, for all  $n \geq N$ , we have

$$\begin{aligned} \|z_n f_n - z f\| &= \|z_n f_n - z_n f + z_n f - z f\| \\ &= \|z_n(f_n - f) + (z_n - z)f\| \\ &\leq \|z_n(f_n - f)\| + \|(z_n - z)f\| \\ &= |z_n| \|f_n - f\| + |z_n - z| \|f\| \\ &< k \frac{\varepsilon}{2k} + \frac{\varepsilon}{2\|f\|} \|f\| \\ &= \varepsilon. \end{aligned}$$

Hence  $\lim_{n \rightarrow \infty} z_n f_n = z f$ .

(iii) Let  $\varepsilon > 0$ . Since  $\lim_{n \rightarrow \infty} f_n = f$ , there exists  $N \in \mathbb{N}$  such that

$$\forall n \geq N : \|f_n - f\| < \varepsilon.$$

By the triangle inequality, we have

$$\|f_n\| = \|f_n - f + f\| \leq \|f_n - f\| + \|f\|$$

so that  $\|f_n\| - \|f\| \leq \|f_n - f\|$ . Similarly,  $\|f\| - \|f_n\| \leq \|f - f_n\|$ . Since

$$\|f - f_n\| = \|-(f_n - f)\| = |-1| \|f_n - f\| = \|f_n - f\|,$$

we have  $-\|f_n - f\| \leq \|f_n\| - \|f\| \leq \|f_n - f\|$  or, by using the modulus,

$$|\|f_n\| - \|f\|| \leq \|f_n - f\|.$$

Hence, for all  $n \geq N$ , we have  $|\|f_n\| - \|f\|| < \varepsilon$  and thus  $\lim_{n \rightarrow \infty} \|f_n\| = \|f\|$ .  $\square$

Note that by taking  $\{z_n\}_{n \in \mathbb{N}}$  to be the constant sequence whose terms are all equal to some fixed  $z \in \mathbb{C}$ , we have  $\lim_{n \rightarrow \infty} z f_n = z f$  as a special case of (ii).

This lemma will take care of some of the technicalities involved in proving the following crucially important result.

**Theorem 2.6.** *The set  $\mathcal{L}(V, W)$  of bounded linear operators from a normed space  $(V, \|\cdot\|_V)$  to a Banach space  $(W, \|\cdot\|_W)$ , equipped with pointwise addition and scalar multiplication and the operator norm, is a Banach space.*

*Proof.* (a) Define addition and scalar multiplication on  $\mathcal{L}(V, W)$  by

$$(A + B)f := Af + Bf \quad (zA)f := zAf.$$

It is clear that both  $A + B$  and  $zA$  are linear operators. Moreover, we have

$$\begin{aligned} \sup_{f \in V} \frac{\|(A + B)f\|_W}{\|f\|_V} &:= \sup_{f \in V} \frac{\|Af + Bf\|_W}{\|f\|_V} \\ &\leq \sup_{f \in V} \frac{\|Af\|_W + \|Bf\|_W}{\|f\|_V} \\ &= \sup_{f \in V} \frac{\|Af\|_W}{\|f\|_V} + \sup_{f \in V} \frac{\|Bf\|_W}{\|f\|_V} \\ &< \infty \end{aligned}$$

since  $A$  and  $B$  are bounded. Hence,  $A + B$  is also bounded and we have

$$\|A + B\| \leq \|A\| + \|B\|.$$

Similarly, for  $zA$  we have

$$\begin{aligned} \sup_{f \in V} \frac{\|(zA)f\|_W}{\|f\|_V} &:= \sup_{f \in V} \frac{\|zAf\|_W}{\|f\|_V} \\ &= \sup_{f \in V} \frac{|z| \|Af\|_W}{\|f\|_V} \\ &= |z| \sup_{f \in V} \frac{\|Af\|_W}{\|f\|_V} \\ &< \infty \end{aligned}$$

since  $A$  is bounded and  $|z|$  is finite. Hence,  $zA$  is bounded and we have

$$\|zA\| = |z|\|A\|.$$

Thus, we have two operations

$$\begin{aligned} +: \mathcal{L}(V, W) \times \mathcal{L}(V, W) &\rightarrow \mathcal{L}(V, W) & \cdot: \mathbb{C} \times \mathcal{L}(V, W) &\rightarrow \mathcal{L}(V, W) \\ (A, B) &\mapsto A + B & (z, A) &\mapsto zA \end{aligned}$$

and it is immediate to check that the vector space structure of  $W$  induces a vector space structure on  $\mathcal{L}(V, W)$  with these operations.

(b) We need to show that  $(\mathcal{L}(V, W), \|\cdot\|)$  is a normed space, i.e. that  $\|\cdot\|$  satisfies the properties of a norm. We have already shown two of these in part (a), namely

$$(b.iii) \quad \|zA\| = |z|\|A\|$$

$$(b.iv) \quad \|A + B\| \leq \|A\| + \|B\|.$$

The remaining two are easily checked.

$$(b.i) \quad \|A\| := \sup_{f \in V} \frac{\|Af\|_W}{\|f\|_V} \geq 0 \text{ since } \|\cdot\|_V \text{ and } \|\cdot\|_W \text{ are norms.}$$

(b.ii) Again, by using the fact that  $\|\cdot\|_W$  is a norm,

$$\begin{aligned} \|A\| = 0 &\Leftrightarrow \sup_{f \in V} \frac{\|Af\|_W}{\|f\|_V} = 0 \\ &\Leftrightarrow \forall f \in V : \|Af\|_W = 0 \\ &\Leftrightarrow \forall f \in V : Af = 0 \\ &\Leftrightarrow A = 0. \end{aligned}$$

Hence,  $(\mathcal{L}(V, W), \|\cdot\|)$  is a normed space.

(c) The heart of the proof is showing that  $(\mathcal{L}(V, W), \|\cdot\|)$  is complete. We will proceed in three steps, analogously to the case of  $C_{\mathbb{C}}^0[0, 1]$ .

(c.i) Let  $\{A_n\}_{n \in \mathbb{N}}$  be a Cauchy sequence in  $\mathcal{L}(V, W)$ . Fix  $f \in V$  and let  $\varepsilon > 0$ . Then, there exists  $N \in \mathbb{N}$  such that

$$\forall n, m \geq N : \|A_n - A_m\| < \frac{\varepsilon}{\|f\|_V}.$$

Then, for all  $n, m \geq N$ , we have

$$\begin{aligned} \|A_n f - A_m f\|_W &= \|(A_n - A_m)f\|_W \\ &= \|f\|_V \frac{\|(A_n - A_m)f\|_W}{\|f\|_V} \\ &\leq \|f\|_V \sup_{f \in V} \frac{\|(A_n - A_m)f\|_W}{\|f\|_V} \\ &=: \|f\|_V \|A_n - A_m\| \\ &< \|f\|_V \frac{\varepsilon}{\|f\|_V} \\ &= \varepsilon. \end{aligned}$$

(Note that if  $f = 0$ , we simply have  $\|A_n f - A_m f\|_W = 0 < \varepsilon$  and, in the future, we will not mention this case explicitly.) Hence, the sequence  $\{A_n f\}_{n \in \mathbb{N}}$  is a Cauchy sequence in  $W$ . Since  $W$  is a Banach space, the limit  $\lim_{n \rightarrow \infty} A_n f$  exists and is an element of  $W$ . Thus, we can define the operator

$$\begin{aligned} A: V &\rightarrow W \\ f &\mapsto \lim_{n \rightarrow \infty} A_n f, \end{aligned}$$

called the pointwise limit of  $\{A_n\}_{n \in \mathbb{N}}$ .

(c.ii) We now need to show that  $A \in \mathcal{L}(V, W)$ . This is where the previous lemma comes in handy. For linearity, let  $f, g \in V$  and  $z \in \mathbb{C}$ . Then

$$\begin{aligned} A(zf + g) &:= \lim_{n \rightarrow \infty} A_n(zf + g) \\ &= \lim_{n \rightarrow \infty} (zA_n f + A_n g) \\ &= z \lim_{n \rightarrow \infty} A_n f + \lim_{n \rightarrow \infty} A_n g \\ &=: zAf + Ag \end{aligned}$$

where we have used the linearity of each  $A_n$  and part (i) and (ii) of Lemma 2.5. For boundedness, part (ii) and (iii) of Lemma 2.5 yield

$$\begin{aligned} \|Af\|_W &= \lim_{n \rightarrow \infty} \|A_n f\|_W \\ &= \lim_{n \rightarrow \infty} \|f\|_V \frac{\|A_n f\|_W}{\|f\|_V} \\ &\leq \lim_{n \rightarrow \infty} \|f\|_V \sup_{f \in V} \frac{\|A_n f\|_W}{\|f\|_V} \\ &= \|f\|_V \lim_{n \rightarrow \infty} \|A_n\| \end{aligned}$$

for any  $f \in V$ . By rearranging, we have

$$\forall f \in V : \frac{\|Af\|_W}{\|f\|_V} \leq \lim_{n \rightarrow \infty} \|A_n\|.$$

Hence, to show that  $A$  is bounded, it suffices to show that the limit on the right hand side is finite. Let  $\varepsilon > 0$ . Since  $\{A_n\}_{n \in \mathbb{N}}$  is a Cauchy sequence, there exists  $N \in \mathbb{N}$  such that

$$\forall n, m \geq N : \|A_n - A_m\| < \varepsilon.$$

Then, by the proof of part (i) of Lemma 2.5, we have

$$|\|A_n\| - \|A_m\|| \leq \|A_n - A_m\| < \varepsilon$$

for all  $n, m \geq N$ . Hence, the sequence of real numbers  $\{\|A_n\|\}_{n \in \mathbb{N}}$  is a Cauchy sequence. Since  $\mathbb{R}$  is complete, this sequence converges to some real number  $r \in \mathbb{R}$ . Therefore

$$\sup_{f \in V} \frac{\|Af\|_W}{\|f\|_V} \leq \lim_{n \rightarrow \infty} \|A_n\| = r < \infty$$

and thus  $A \in \mathcal{L}(V, W)$ .

(c.iii) To conclude, we have to show that  $\lim_{n \rightarrow \infty} A_n = A$ . Let  $\varepsilon > 0$ . Then

$$\|A_n - A\| = \|A_n + A_m - A_m - A\| \leq \|A_n - A_m\| + \|A_m - A\|.$$

Since  $\{A_n\}_{n \in \mathbb{N}}$  is Cauchy, there exists  $N_1 \in \mathbb{N}$  such that

$$\forall n, m \geq N_1 : \|A_n - A_m\| < \frac{\varepsilon}{2}.$$

Moreover, since  $A$  is the pointwise limit of  $\{A_n\}_{n \in \mathbb{N}}$ , for any  $f \in V$  there exists  $N_2 \in \mathbb{N}$  such that

$$\forall m \geq N_2 : \|A_m f - A f\|_W < \frac{\varepsilon \|f\|_V}{2}$$

and hence, for all  $m \geq N_2$

$$\|A_m - A\| := \sup_{f \in V} \frac{\|A_m f - A f\|_W}{\|f\|_V} \leq \frac{\frac{\varepsilon \|f\|_V}{2}}{\|f\|_V} = \frac{\varepsilon}{2}$$

Let  $N := \max\{N_1, N_2\}$  and fix  $m \geq N$ . Then, for all  $n \geq N$ , we have

$$\|A_n - A\| \leq \|A_n - A_m\| + \|A_m - A\| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

Thus,  $\lim_{n \rightarrow \infty} A_n = A$  and we call  $A$  the uniform limit of  $\{A_n\}_{n \in \mathbb{N}}$ .

This concludes the proof that  $(\mathcal{L}(V, W), \|\cdot\|)$  is a Banach space.  $\square$

Remark 2.7. Note that if  $V$  and  $W$  are normed spaces, then  $\mathcal{L}(V, W)$  is again a normed space, while for  $\mathcal{L}(V, W)$  to be a Banach space it suffices that  $W$  be a Banach space.

Remark 2.8. In the proof that  $\mathcal{L}(V, W)$  is a Banach space, we have shown a useful inequality which we restate here for further emphasis. If  $A: V \rightarrow W$  is bounded, then

$$\forall f \in V : \|A f\|_W \leq \|A\| \|f\|_V$$

The following is an extremely important special case of  $\mathcal{L}(V, W)$ .

**Definition.** Let  $V$  be a normed space. Then  $V^* := \mathcal{L}(V, \mathbb{C})$  is called the *dual* of  $V$ .  $\diamond$

Note that, since  $\mathbb{C}$  is a Banach space, the dual of a normed space is a Banach space. The elements of  $V^*$  are variously called *covectors* or *functionals* on  $V$ .

Remark 2.9. You may recall from undergraduate linear algebra that the dual of a vector space was defined to be the vector space of *all* linear maps  $V \rightarrow \mathbb{C}$ , rather than just the bounded ones. This is because, in finite dimensions, all linear maps are bounded. So the two definitions agree as long as we are in finite dimensions. If we used the same definition for the infinite-dimensional case, then  $V^*$  would lack some very desirable properties, such as that of being a Banach space.

The dual space can be used to define a weaker notion of convergence called, rather unimaginatively, weak convergence.



**Definition.** A sequence  $\{f_n\}_{n \in \mathbb{N}}$  is said to *converge weakly* to  $f \in V$  if

$$\forall \varphi \in V^* : \lim_{n \rightarrow \infty} \varphi(f_n) = \varphi(f).$$

◇

Note that  $\{\varphi(f_n)\}_{n \in \mathbb{N}}$  is just a sequence of complex numbers. To indicate that the sequence  $\{f_n\}_{n \in \mathbb{N}}$  converges weakly to  $f \in V$  we write

$$\text{w-lim}_{n \rightarrow \infty} f_n = f.$$

In order to further emphasise the distinction with weak convergence, we may say that  $\{f_n\}_{n \in \mathbb{N}}$  converges *strongly* to  $f \in V$  if it converges according to the usual definition, and we will write accordingly

$$\text{s-lim}_{n \rightarrow \infty} f_n = f.$$

**Proposition 2.10.** *Let  $\{f_n\}_{n \in \mathbb{N}}$  be a sequence in a normed space  $(V, \|\cdot\|_V)$ . If  $\{f_n\}_{n \in \mathbb{N}}$  converges strongly to  $f \in V$ , then it also converges weakly to  $f \in V$ , i.e.*

$$\text{s-lim}_{n \rightarrow \infty} f_n = f \quad \Rightarrow \quad \text{w-lim}_{n \rightarrow \infty} f_n = f.$$

*Proof.* Let  $\varepsilon > 0$  and let  $\varphi \in V^*$ . Since  $\{f_n\}_{n \in \mathbb{N}}$  converges strongly to  $f \in V$ , there exists  $N \in \mathbb{N}$  such that

$$\forall n \geq N : \|f_n - f\|_V < \frac{\varepsilon}{\|\varphi\|}.$$

Then, since  $\varphi \in V^*$  is bounded, we have

$$\begin{aligned} |\varphi(f_n) - \varphi(f)| &= |\varphi(f_n - f)| \\ &\leq \|\varphi\| \|f_n - f\|_V \\ &< \|\varphi\| \frac{\varepsilon}{\|\varphi\|} \\ &= \varepsilon \end{aligned}$$

for any  $n \geq N$ . Hence,  $\lim_{n \rightarrow \infty} \varphi(f_n) = \varphi(f)$ . That is,  $\text{w-lim}_{n \rightarrow \infty} f_n = f$ . □

### 2.3 Extension of bounded linear operators

Note that, so far, we have only considered bounded linear maps  $A: \mathcal{D}_A \rightarrow W$  where  $\mathcal{D}_A$  is the whole of  $V$ , rather than a subspace thereof. The reason for this is that we will only consider densely defined linear maps in general, and any bounded linear map from a dense subspace of  $V$  can be extended to a bounded linear map from the whole of  $V$ . Moreover, the extension is unique. This is the content of the so-called BLT<sup>5</sup> theorem.

**Lemma 2.11.** *Let  $(V, \|\cdot\|)$  be a normed space and let  $\mathcal{D}_A$  be a dense subspace of  $V$ . Then, for any  $f \in V$ , there exists a sequence  $\{\alpha_n\}_{n \in \mathbb{N}}$  in  $\mathcal{D}_A$  which converges to  $f$ .*

<sup>5</sup>Bounded Linear Transformation, *not* Bacon, Lettuce, Tomato.



*Proof.* Let  $f \in V$ . Clearly, there exists a sequence  $\{f_n\}_{n \in \mathbb{N}}$  in  $V$  which converges to  $f$  (for instance, the constant sequence). Let  $\varepsilon > 0$ . Then, there exists  $N \in \mathbb{N}$  such that

$$\forall n \geq N : \|f_n - f\| < \frac{\varepsilon}{2}.$$

Since  $\mathcal{D}_A$  is dense in  $V$  and each  $f_n \in V$ , we have

$$\forall n \in \mathbb{N} : \exists \alpha_n \in \mathcal{D}_A : \|\alpha_n - f_n\| < \frac{\varepsilon}{2}.$$

The sequence  $\{\alpha_n\}_{n \in \mathbb{N}}$  is a sequence in  $\mathcal{D}_A$  and we have

$$\begin{aligned} \|\alpha_n - f\| &= \|\alpha_n - f_n + f_n - f\| \\ &\leq \|\alpha_n - f_n\| + \|f_n - f\| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ &= \varepsilon \end{aligned}$$

for all  $n \geq N$ . Hence  $\lim_{n \rightarrow \infty} \alpha_n = f$ . □

**Definition.** Let  $V, W$  be vector spaces and let  $A : \mathcal{D}_A \rightarrow W$  be a linear map, where  $\mathcal{D}_A \subseteq V$ . An *extension* of  $A$  is a linear map  $\hat{A} : V \rightarrow W$  such that

$$\forall \alpha \in \mathcal{D}_A : \hat{A}\alpha = A\alpha.$$

◇

**Theorem 2.12** (BLT theorem). *Let  $V$  be a normed space and  $W$  a Banach space. Any densely defined linear map  $A : \mathcal{D}_A \rightarrow W$  has a unique extension  $\hat{A} : V \rightarrow W$  such that  $\hat{A}$  is bounded. Moreover,  $\|\hat{A}\| = \|A\|$ .*

*Proof.* (a) Let  $A \in \mathcal{L}(\mathcal{D}_A, W)$ . Since  $\mathcal{D}_A$  is dense in  $V$ , for any  $f \in V$  there exists a sequence  $\{\alpha_n\}_{n \in \mathbb{N}}$  in  $\mathcal{D}_A$  which converges to  $f$ . Moreover, since  $A$  is bounded, we have

$$\forall n \in \mathbb{N} : \|A\alpha_n - A\alpha_m\|_W \leq \|A\| \|\alpha_n - \alpha_m\|_V,$$

from which it quickly follows that  $\{A\alpha_n\}_{n \in \mathbb{N}}$  is Cauchy in  $W$ . As  $W$  is a Banach space, this sequence converges to an element of  $W$  and thus we can define

$$\begin{aligned} \hat{A} : V &\rightarrow W \\ f &\mapsto \lim_{n \rightarrow \infty} A\alpha_n, \end{aligned}$$

where  $\{\alpha_n\}_{n \in \mathbb{N}}$  is any sequence in  $\mathcal{D}_A$  which converges to  $f$ .

(b) First, let us show that  $\hat{A}$  is well-defined. Let  $\{\alpha_n\}_{n \in \mathbb{N}}$  and  $\{\beta_n\}_{n \in \mathbb{N}}$  be two sequences in  $\mathcal{D}_A$  which converge to  $f \in V$  and let  $\varepsilon > 0$ . Then, there exist  $N_1, N_2 \in \mathbb{N}$  such that

$$\begin{aligned} \forall n \geq N_1 : \|\alpha_n - f\|_V &< \frac{\varepsilon}{2\|A\|} \\ \forall n \geq N_2 : \|\beta_n - f\|_V &< \frac{\varepsilon}{2\|A\|}. \end{aligned}$$

Let  $N := \max\{N_1, N_2\}$ . Then, for all  $n \geq N$ , we have

$$\begin{aligned}
\|A\alpha_n - A\beta_n\|_W &= \|A(\alpha_n - \beta_n)\|_W \\
&\leq \|A\| \|\alpha_n - \beta_n\|_V \\
&= \|A\| \|\alpha_n - f + f - \beta_n\|_V \\
&\leq \|A\| (\|\alpha_n - f\|_V + \|f - \beta_n\|_V) \\
&< \|A\| \left( \frac{\varepsilon}{2\|A\|} + \frac{\varepsilon}{2\|A\|} \right) \\
&= \varepsilon,
\end{aligned}$$

where we have used the fact that  $A$  is bounded. Thus, we have shown

$$\lim_{n \rightarrow \infty} (A\alpha_n - A\beta_n) = 0.$$

Then, by using Lemma 2.5 and rearranging, we find

$$\lim_{n \rightarrow \infty} A\alpha_n = \lim_{n \rightarrow \infty} A\beta_n,$$

that is,  $\hat{A}$  is indeed well-defined.

- (c) To see that  $\hat{A}$  is an extension of  $A$ , let  $\alpha \in \mathcal{D}_A$ . The constant sequence  $\{\alpha_n\}_{n \in \mathbb{N}}$  with  $\alpha_n = \alpha$  for all  $n \in \mathbb{N}$  is a sequence in  $\mathcal{D}_A$  converging to  $\alpha$ . Hence

$$\hat{A}\alpha := \lim_{n \rightarrow \infty} A\alpha_n = \lim_{n \rightarrow \infty} A\alpha = A\alpha.$$

- (d) We now check that  $A \in \mathcal{L}(V, W)$ . For linearity, let  $f, g \in V$  and  $z \in \mathbb{C}$ . As  $\mathcal{D}_A$  is dense in  $V$ , there exist sequences  $\{\alpha_n\}_{n \in \mathbb{N}}$  and  $\{\beta_n\}_{n \in \mathbb{N}}$  in  $\mathcal{D}_A$  converging to  $f$  and  $g$ , respectively. Moreover, as  $\mathcal{D}_A$  is a subspace of  $V$ , the sequence  $\{\gamma_n\}_{n \in \mathbb{N}}$  given by

$$\gamma_n := z\alpha_n + \beta_n$$

is again a sequence in  $\mathcal{D}_A$  and, by Lemma 2.5,

$$\lim_{n \rightarrow \infty} \gamma_n = zf + g.$$

Then, we have

$$\begin{aligned}
\hat{A}(zf + g) &:= \lim_{n \rightarrow \infty} A\gamma_n \\
&= \lim_{n \rightarrow \infty} A(z\alpha_n + \beta_n) \\
&= \lim_{n \rightarrow \infty} (zA\alpha_n + A\beta_n) \\
&= z \lim_{n \rightarrow \infty} A\alpha_n + \lim_{n \rightarrow \infty} A\beta_n \\
&=: z\hat{A}f + \hat{A}g.
\end{aligned}$$

For boundedness, let  $f \in V$  and  $\{\alpha_n\}_{n \in \mathbb{N}}$  a sequence in  $\mathcal{D}_A$  which converges to  $f$ . Then, since  $A$  is bounded,

$$\begin{aligned}\|\widehat{A}f\|_W &:= \left\| \lim_{n \rightarrow \infty} A\alpha_n \right\|_W \\ &= \lim_{n \rightarrow \infty} \|A\alpha_n\|_W \\ &\leq \lim_{n \rightarrow \infty} \|A\| \|\alpha_n\|_V \\ &= \|A\| \lim_{n \rightarrow \infty} \|\alpha_n\|_V \\ &= \|A\| \|f\|_V.\end{aligned}$$

Therefore

$$\sup_{f \in V} \frac{\|\widehat{A}f\|_W}{\|f\|_V} \leq \sup_{f \in V} \frac{\|A\| \|f\|_V}{\|f\|_V} = \sup_{f \in V} \|A\| = \|A\| < \infty$$

and hence  $\widehat{A}$  is bounded.

- (e) For uniqueness, suppose that  $\widetilde{A} \in \mathcal{L}(V, W)$  is another extension of  $A$ . Let  $f \in V$  and  $\{\alpha_n\}_{n \in \mathbb{N}}$  a sequence in  $\mathcal{D}_A$  which converges to  $f$ . Then, we have

$$\|\widetilde{A}f - A\alpha_n\|_W = \|\widetilde{A}f - \widetilde{A}\alpha_n\|_W \leq \|\widetilde{A}\| \|f - \alpha_n\|_V.$$

It follows that

$$\lim_{n \rightarrow \infty} (\widetilde{A}f - A\alpha_n) = 0$$

and hence, for all  $f \in V$ ,

$$\widetilde{A}f = \lim_{n \rightarrow \infty} A\alpha_n =: \widehat{A}f.$$

Therefore,  $\widetilde{A} = \widehat{A}$ .

- (f) Finally, we have already shown in part (d) that

$$\|\widehat{A}\| := \sup_{f \in V} \frac{\|\widehat{A}f\|_W}{\|f\|_V} \leq \|A\|.$$

On the other hand, since  $\mathcal{D}_A \subseteq V$ , we must also have

$$\|A\| := \sup_{f \in \mathcal{D}_A} \frac{\|Af\|_W}{\|f\|_V} = \sup_{f \in \mathcal{D}_A} \frac{\|\widehat{A}f\|_W}{\|f\|_V} \leq \sup_{f \in V} \frac{\|\widehat{A}f\|_W}{\|f\|_V} =: \|\widehat{A}\|.$$

Hence, we also have  $\|A\| \leq \|\widehat{A}\|$ . Thus,  $\|\widehat{A}\| = \|A\|$ .  $\square$

*Remark 2.13.* Note a slight abuse of notation in the equality  $\|\widehat{A}\| = \|A\|$ . The linear maps  $\widehat{A}$  and  $A$  belong to  $\mathcal{L}(V, W)$  and  $\mathcal{L}(\mathcal{D}_A, W)$ , respectively. These are different normed (in fact, Banach) spaces and, in particular, carry *different* norms. To be more precise, we should have written

$$\|\widehat{A}\|_{\mathcal{L}(V, W)} = \|A\|_{\mathcal{L}(\mathcal{D}_A, W)},$$

where

$$\|\widehat{A}\|_{\mathcal{L}(V, W)} := \sup_{f \in V} \frac{\|\widehat{A}f\|_W}{\|f\|_V} \quad \text{and} \quad \|A\|_{\mathcal{L}(\mathcal{D}_A, W)} := \sup_{f \in \mathcal{D}_A} \frac{\|Af\|_W}{\|f\|_V}.$$

### 3 Separable Hilbert Spaces

#### 3.1 Relationship between norms and inner products

A Hilbert space is a vector space  $(\mathcal{H}, +, \cdot)$  equipped with a sesqui-linear inner product  $\langle \cdot | \cdot \rangle$  which induces a norm  $\|\cdot\|_{\mathcal{H}}$  with respect to which  $\mathcal{H}$  is a Banach space. Note that by “being induced by  $\langle \cdot | \cdot \rangle$ ” we specifically mean that the norm is defined as

$$\begin{aligned} \|\cdot\|: V &\rightarrow \mathbb{R} \\ f &\mapsto \sqrt{\langle f | f \rangle}. \end{aligned}$$

Recall that a sesqui-linear inner product on  $\mathcal{H}$  is a map  $\langle \cdot | \cdot \rangle: \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{C}$  which is conjugate symmetric, linear in the second argument and positive-definite. Note that conjugate symmetry together with linearity in the second argument imply conjugate linearity in the first argument:

$$\begin{aligned} \langle z\psi_1 + \psi_2 | \varphi \rangle &= \overline{\langle \varphi | z\psi_1 + \psi_2 \rangle} \\ &= \overline{z\langle \varphi | \psi_1 \rangle + \langle \varphi | \psi_2 \rangle} \\ &= \bar{z}\overline{\langle \varphi | \psi_1 \rangle} + \overline{\langle \varphi | \psi_2 \rangle} \\ &= \bar{z}\langle \psi_1 | \varphi \rangle + \langle \psi_2 | \varphi \rangle. \end{aligned}$$

Of course, since Hilbert spaces are a special case of Banach spaces, everything that we have learned about Banach spaces also applies to Hilbert spaces. For instance,  $\mathcal{L}(\mathcal{H}, \mathcal{H})$ , the collection of all bounded linear maps  $\mathcal{H} \rightarrow \mathcal{H}$ , is a Banach space with respect to the operator norm. In particular, the dual of a Hilbert space  $\mathcal{H}$  is just  $\mathcal{H}^* := \mathcal{L}(\mathcal{H}, \mathbb{C})$ . We will see that the operator norm on  $\mathcal{H}^*$  is such that there exists an inner product on  $\mathcal{H}^*$  which induces it, so that the dual of a Hilbert space is again a Hilbert space.

First, in order to check that the norm induced by an inner product on  $V$  is indeed a norm on  $V$ , we need one of the most important inequalities in mathematics.

**Proposition 3.1** (Cauchy-Schwarz inequality<sup>6</sup>). *Let  $\langle \cdot | \cdot \rangle$  be a sesqui-linear inner product on  $V$ . Then, for any  $f, g \in V$ , we have*

$$|\langle f | g \rangle|^2 \leq \langle f | f \rangle \langle g | g \rangle.$$

*Proof.* If  $f = 0$  or  $g = 0$ , then equality holds. Hence suppose that  $f \neq 0$  and let

$$z := \frac{\langle f | g \rangle}{\langle f | f \rangle} \in \mathbb{C}.$$

---

<sup>6</sup>Also known as the Cauchy-Bunyakovsky-Schwarz inequality in the Russian literature.

Then, by positive-definiteness of  $\langle \cdot | \cdot \rangle$ , we have

$$\begin{aligned}
0 &\leq \langle zf - g | zf - g \rangle \\
&= |z|^2 \langle f | f \rangle - \bar{z} \langle f | g \rangle - z \langle g | f \rangle + \langle g | g \rangle \\
&= \frac{|\langle f | g \rangle|^2}{\langle f | f \rangle^2} \langle f | f \rangle - \frac{\overline{\langle f | g \rangle}}{\langle f | f \rangle} \langle f | g \rangle - \frac{\langle f | g \rangle}{\langle f | f \rangle} \overline{\langle f | g \rangle} + \langle g | g \rangle \\
&= \frac{|\langle f | g \rangle|^2}{\langle f | f \rangle} - \frac{|\langle f | g \rangle|^2}{\langle f | f \rangle} - \frac{|\langle f | g \rangle|^2}{\langle f | f \rangle} + \langle g | g \rangle \\
&= -\frac{|\langle f | g \rangle|^2}{\langle f | f \rangle} + \langle g | g \rangle.
\end{aligned}$$

By rearranging, since  $\langle f | f \rangle > 0$ , we obtain the desired inequality.  $\square$

Note that, by defining  $\|f\| := \sqrt{\langle f | f \rangle}$ , and using the fact that  $|\langle f | g \rangle| \geq 0$ , we can write the Cauchy-Schwarz inequality as

$$|\langle f | g \rangle| \leq \|f\| \|g\|.$$

**Proposition 3.2.** *The induced norm on  $V$  is a norm.*

*Proof.* Let  $f, g \in V$  and  $z \in \mathbb{C}$ . Then

- (i)  $\|f\| := \sqrt{\langle f | f \rangle} \geq 0$
- (ii)  $\|f\| = 0 \Leftrightarrow \|f\|^2 = 0 \Leftrightarrow \langle f | f \rangle = 0 \Leftrightarrow f = 0$  by positive-definiteness
- (iii)  $\|zf\| := \sqrt{\langle zf | zf \rangle} = \sqrt{z\bar{z}\langle f | f \rangle} = \sqrt{|z|^2 \langle f | f \rangle} = |z| \sqrt{\langle f | f \rangle} =: |z| \|f\|$
- (iv) Using the fact that  $z + \bar{z} = 2 \operatorname{Re} z$  and  $\operatorname{Re} z \leq |z|$  for any  $z \in \mathbb{C}$  and the Cauchy-Schwarz inequality, we have

$$\begin{aligned}
\|f + g\|^2 &:= \langle f + g | f + g \rangle \\
&= \langle f | f \rangle + \langle f | g \rangle + \langle g | f \rangle + \langle g | g \rangle \\
&= \langle f | f \rangle + \langle f | g \rangle + \overline{\langle f | g \rangle} + \langle g | g \rangle \\
&= \langle f | f \rangle + 2 \operatorname{Re} \langle f | g \rangle + \langle g | g \rangle \\
&\leq \langle f | f \rangle + 2 |\langle f | g \rangle| + \langle g | g \rangle \\
&\leq \langle f | f \rangle + 2 \|f\| \|g\| + \langle g | g \rangle \\
&= (\|f\| + \|g\|)^2.
\end{aligned}$$

By taking the square root of both sides, we have  $\|f + g\| \leq \|f\| + \|g\|$ .  $\square$

Hence, we see that any inner product space (i.e. a vector space equipped with a sesqui-linear inner product) is automatically a normed space under the induced norm. It is only natural to wonder whether the converse also holds, that is, whether every norm is induced by some sesqui-linear inner product. Unfortunately, the answer is negative in general. The following theorem gives a necessary and sufficient condition for a norm to be induced by a sesqui-linear inner product and, in fact, by a unique such.

**Theorem 3.3** (Jordan-von Neumann). *Let  $V$  be a vector space. A norm  $\|\cdot\|$  on  $V$  is induced by a sesqui-linear inner product  $\langle\cdot|\cdot\rangle$  on  $V$  if, and only if, the parallelogram identity*

$$\|f + g\|^2 + \|f - g\|^2 = 2\|f\|^2 + 2\|g\|^2$$

*holds for all  $f, g \in V$ , in which case,  $\langle\cdot|\cdot\rangle$  is determined by the polarisation identity*

$$\begin{aligned}\langle f|g\rangle &= \frac{1}{4} \sum_{k=0}^3 i^k \|f + i^{4-k}g\|^2 \\ &= \frac{1}{4} (\|f + g\|^2 - \|f - g\|^2 + i\|f - ig\|^2 - i\|f + ig\|^2).\end{aligned}$$

*Proof.* ( $\Rightarrow$ ) If  $\|\cdot\|$  is induced by  $\langle\cdot|\cdot\rangle$ , then by direct computation

$$\begin{aligned}\|f + g\|^2 + \|f - g\|^2 &:= \langle f + g|f + g\rangle + \langle f - g|f - g\rangle \\ &= \langle f|f\rangle + \langle f|g\rangle + \langle g|f\rangle + \langle g|g\rangle \\ &\quad + \langle f|f\rangle - \langle f|g\rangle - \langle g|f\rangle + \langle g|g\rangle \\ &= 2\langle f|f\rangle + 2\langle g|g\rangle \\ &=: 2\|f\|^2 + 2\|g\|^2,\end{aligned}$$

so the parallelogram identity is satisfied. We also have

$$\begin{aligned}\|f + g\|^2 - \|f - g\|^2 &:= \langle f + g|f + g\rangle - \langle f - g|f - g\rangle \\ &= \langle f|f\rangle + \langle f|g\rangle + \langle g|f\rangle + \langle g|g\rangle \\ &\quad - \langle f|f\rangle + \langle f|g\rangle + \langle g|f\rangle - \langle g|g\rangle \\ &= 2\langle f|g\rangle + 2\langle g|f\rangle\end{aligned}$$

and

$$\begin{aligned}i\|f - ig\|^2 - i\|f + ig\|^2 &:= i\langle f - ig|f - ig\rangle - i\langle f + ig|f + ig\rangle \\ &= i\langle f|f\rangle + \langle f|g\rangle - \langle g|f\rangle + i\langle g|g\rangle \\ &\quad - i\langle f|f\rangle + \langle f|g\rangle - \langle g|f\rangle - i\langle g|g\rangle \\ &= 2\langle f|g\rangle - 2\langle g|f\rangle.\end{aligned}$$

Therefore

$$\|f + g\|^2 - \|f - g\|^2 + i\|f - ig\|^2 - i\|f + ig\|^2 = 4\langle f|g\rangle.$$

that is, the inner product is determined by the polarisation identity.

( $\Leftarrow$ ) Suppose that  $\|\cdot\|$  satisfies the parallelogram identity. Define  $\langle\cdot|\cdot\rangle$  by

$$\langle f|g\rangle := \frac{1}{4} (\|f + g\|^2 - \|f - g\|^2 + i\|f - ig\|^2 - i\|f + ig\|^2).$$

We need to check that this satisfies the defining properties of a sesqui-linear inner product.

(i) For conjugate symmetry

$$\begin{aligned}
\overline{\langle f|g \rangle} &:= \frac{1}{4} (\|f+g\|^2 - \|f-g\|^2 + i\|f-ig\|^2 - i\|f+ig\|^2) \\
&= \frac{1}{4} (\|f+g\|^2 - \|f-g\|^2 - i\|f-ig\|^2 + i\|f+ig\|^2) \\
&= \frac{1}{4} (\|f+g\|^2 - \|f-g\|^2 - i\|(-i)(f+g)\|^2 + i\|i(-f+g)\|^2) \\
&= \frac{1}{4} (\|g+f\|^2 - \|g-f\|^2 - i\|(-i)^2\|g+if\|^2 + i\|i\|^2\|g-if\|^2) \\
&= \frac{1}{4} (\|g+f\|^2 - \|g-f\|^2 - i\|g+if\|^2 + i\|g-if\|^2) \\
&=: \langle g|f \rangle
\end{aligned}$$

(ii) We will now show linearity in the second argument. This is fairly non-trivial and quite lengthy. We will focus on additivity first. We have

$$\langle f|g+h \rangle := \frac{1}{4} (\|f+g+h\|^2 - \|f-g-h\|^2 + i\|f-ig-ih\|^2 - i\|f+ig+ih\|^2).$$

Consider the real part of  $\langle f|g+h \rangle$ . By successive applications of the parallelogram identity, we find

$$\begin{aligned}
\operatorname{Re}\langle f|g+h \rangle &= \frac{1}{4} (\|f+g+h\|^2 - \|f-g-h\|^2) \\
&= \frac{1}{4} (\|f+g+h\|^2 + \|f+g-h\|^2 - \|f+g-h\|^2 - \|f-g-h\|^2) \\
&= \frac{1}{4} (2\|f+g\|^2 + 2\|h\|^2 - 2\|f-h\|^2 - 2\|g\|^2) \\
&= \frac{1}{4} (2\|f+g\|^2 + 2\|f\|^2 + 2\|h\|^2 - 2\|f-h\|^2 - 2\|f\|^2 - 2\|g\|^2) \\
&= \frac{1}{4} (2\|f+g\|^2 + \|f+h\|^2 + \|f-h\|^2 - 2\|f-h\|^2 - \|f+g\|^2 - \|f-g\|^2) \\
&= \frac{1}{4} (\|f+g\|^2 + \|f+h\|^2 - \|f-h\|^2 - \|f-g\|^2) \\
&= \operatorname{Re}\langle f|g \rangle + \operatorname{Re}\langle f|h \rangle.
\end{aligned}$$

Replacing  $g$  and  $h$  with  $-ig$  and  $-ih$  respectively, we obtain

$$\operatorname{Im}\langle f|g+h \rangle = \operatorname{Im}\langle f|g \rangle + \operatorname{Im}\langle f|h \rangle.$$

Hence, we have

$$\begin{aligned}
\langle f|g+h \rangle &= \operatorname{Re}\langle f|g+h \rangle + i\operatorname{Im}\langle f|g+h \rangle \\
&= \operatorname{Re}\langle f|g \rangle + \operatorname{Re}\langle f|h \rangle + i(\operatorname{Im}\langle f|g \rangle + \operatorname{Im}\langle f|h \rangle) \\
&= \operatorname{Re}\langle f|g \rangle + i\operatorname{Im}\langle f|g \rangle + \operatorname{Re}\langle f|h \rangle + i\operatorname{Im}\langle f|h \rangle \\
&= \langle f|g \rangle + \langle f|h \rangle,
\end{aligned}$$

which proves additivity.

For scaling invariance, we will proceed in several steps.

(a) First, note that

$$\langle f|0 \rangle := \frac{1}{4} (\|f\|^2 - \|f\|^2 + i\|f\|^2 - i\|f\|^2) = 0$$

and hence  $\langle f|0g \rangle = 0\langle f|g \rangle$  holds.



(b) Suppose that  $\langle f|ng\rangle = n\langle f|g\rangle$  for some  $n \in \mathbb{N}$ . Then, by additivity

$$\begin{aligned}\langle f|(n+1)g\rangle &= \langle f|ng+g\rangle \\ &= \langle f|ng\rangle + \langle f|g\rangle \\ &= n\langle f|g\rangle + \langle f|g\rangle \\ &= (n+1)\langle f|g\rangle.\end{aligned}$$

Hence, by induction on  $n$  with base case (a), we have

$$\forall n \in \mathbb{N} : \langle f|ng\rangle = n\langle f|g\rangle.$$

(c) Note that by additivity

$$\langle f|g\rangle + \langle f|-g\rangle = \langle f|g-g\rangle = \langle f|0\rangle \stackrel{(a)}{=} 0.$$

Hence  $\langle f|-g\rangle = -\langle f|g\rangle$ .

(d) Then, for any  $n \in \mathbb{N}$

$$\langle f|-ng\rangle \stackrel{(c)}{=} -\langle f|ng\rangle \stackrel{(b)}{=} -n\langle f|g\rangle$$

and thus

$$\forall n \in \mathbb{Z} : \langle f|ng\rangle = n\langle f|g\rangle.$$

(e) Now note that for any  $m \in \mathbb{Z} \setminus \{0\}$

$$m\langle f|\frac{1}{m}g\rangle \stackrel{(d)}{=} \langle f|m\frac{1}{m}g\rangle = \langle f|g\rangle$$

and hence, by dividing by  $m$ , we have  $\langle f|\frac{1}{m}g\rangle = \frac{1}{m}\langle f|g\rangle$ .

(f) Therefore, for any  $r = \frac{n}{m} \in \mathbb{Q}$ , we have

$$\langle f|rg\rangle = \langle f|\frac{n}{m}g\rangle \stackrel{(d)}{=} n\langle f|\frac{1}{m}g\rangle \stackrel{(e)}{=} \frac{n}{m}\langle f|g\rangle = r\langle f|g\rangle$$

and hence

$$\forall r \in \mathbb{Q} : \langle f|rg\rangle = r\langle f|g\rangle.$$

(g) Before we turn to  $\mathbb{R}$ , we need to show that  $|\langle f|g\rangle| \leq \sqrt{2}\|f\|\|g\|$ . Note that here we *cannot* invoke the Cauchy-Schwarz inequality (which would also provide a better estimate) since we don't know that  $\langle \cdot | \cdot \rangle$  is an inner product yet. First, consider the real part of  $\langle f|g\rangle$ .

$$\begin{aligned}\operatorname{Re}\langle f|g\rangle &= \frac{1}{4}(\|f+g\|^2 - \|f-g\|^2) \\ &= \frac{1}{4}(2\|f+g\|^2 - \|f+g\|^2 - \|f-g\|^2) \\ &= \frac{1}{4}(2\|f+g\|^2 - 2\|f\|^2 - 2\|g\|^2) \\ &\leq \frac{1}{4}(2(\|f\| + \|g\|)^2 - 2\|f\|^2 - 2\|g\|^2) \\ &= \frac{1}{4}(2\|f\|^2 + 4\|f\|\|g\| + 2\|g\|^2 - 2\|f\|^2 - 2\|g\|^2) \\ &= \|f\|\|g\|.\end{aligned}$$

Replacing  $g$  with  $-ig$  and noting that  $\| -ig \| = | -i \| \|g\| = \|g\|$ , we also have

$$\operatorname{Im}\langle f|g\rangle \leq \|f\|\|g\|.$$

Hence, we find

$$\begin{aligned} |\langle f|g\rangle| &= |\operatorname{Re}\langle f|g\rangle + i\operatorname{Im}\langle f|g\rangle| \\ &= \sqrt{(\operatorname{Re}\langle f|g\rangle)^2 + (\operatorname{Im}\langle f|g\rangle)^2} \\ &\leq \sqrt{(\|f\|\|g\|)^2 + (\|f\|\|g\|)^2} \\ &= \sqrt{2}\|f\|\|g\|. \end{aligned}$$

- (h) Let  $r \in \mathbb{R}$ . Since  $\mathbb{R}$  is the completion of  $\mathbb{Q}$  (equivalently,  $\mathbb{Q}$  is dense in  $\mathbb{R}$ ), there exists a sequence  $\{r_n\}_{n \in \mathbb{N}}$  in  $\mathbb{Q}$  which converges to  $r$ . Let  $\varepsilon > 0$ . Then, there exist  $N_1, N_2 \in \mathbb{N}$  such that

$$\begin{aligned} \forall n \geq N_1 : |r_n - r| &< \frac{\varepsilon}{2\sqrt{2}\|f\|\|g\|} \\ \forall n, m \geq N_2 : |r_n - r_m| &< \frac{\varepsilon}{2\sqrt{2}\|f\|\|g\|}. \end{aligned}$$

Let  $N := \max\{N_1, N_2\}$  and fix  $m \geq N$ . Then, for all  $n \geq N$ , we have

$$\begin{aligned} |r_n\langle f|g\rangle - \langle f|rg\rangle| &= |r_n\langle f|g\rangle - r_m\langle f|g\rangle + r_m\langle f|g\rangle - \langle f|rg\rangle| \\ &\stackrel{(f)}{=} |r_n\langle f|g\rangle - r_m\langle f|g\rangle + \langle f|r_mg\rangle - \langle f|rg\rangle| \\ &= |(r_n - r_m)\langle f|g\rangle + \langle f|(r_m - r)g\rangle| \\ &\leq |(r_n - r_m)\langle f|g\rangle| + |\langle f|(r_m - r)g\rangle| \\ &\stackrel{(g)}{\leq} \sqrt{2}|r_n - r_m|\|f\|\|g\| + \sqrt{2}\|f\|\|(r_m - r)g\| \\ &= \sqrt{2}|r_n - r_m|\|f\|\|g\| + \sqrt{2}|r_m - r|\|f\|\|g\| \\ &< \sqrt{2}\frac{\varepsilon}{2\sqrt{2}\|f\|\|g\|}\|f\|\|g\| + \sqrt{2}\frac{\varepsilon}{2\sqrt{2}\|f\|\|g\|}\|f\|\|g\| \\ &= \varepsilon, \end{aligned}$$

that is,  $\lim_{n \rightarrow \infty} r_n\langle f|g\rangle = \langle f|rg\rangle$ .

- (i) Hence, for any  $r \in \mathbb{R}$ , we have

$$r\langle f|g\rangle = \left(\lim_{n \rightarrow \infty} r_n\right)\langle f|g\rangle = \lim_{n \rightarrow \infty} r_n\langle f|g\rangle \stackrel{(h)}{=} \langle f|rg\rangle$$

and thus

$$\forall r \in \mathbb{R} : r\langle f|g\rangle = \langle f|rg\rangle.$$

- (j) We now note that

$$\begin{aligned} \langle f|ig\rangle &:= \frac{1}{4}(\|f + ig\|^2 - \|f - ig\|^2 + i\|f - i^2g\|^2 - i\|f + i^2g\|^2) \\ &= \frac{1}{4}i(-i\|f + ig\|^2 + \|f - ig\|^2 + i\|f + g\|^2 - \|f - g\|^2) \\ &=: i\langle f|g\rangle \end{aligned}$$

and hence  $\langle f|ig\rangle = i\langle f|g\rangle$ .

(k) Let  $z \in \mathbb{C}$ . By additivity, we have

$$\begin{aligned}
\langle f|zg\rangle &= \langle f|(\operatorname{Re} z + i \operatorname{Im} z)g\rangle \\
&= \langle f|(\operatorname{Re} z)g\rangle + \langle f|i(\operatorname{Im} z)g\rangle \\
&\stackrel{(j)}{=} \langle f|(\operatorname{Re} z)g\rangle + i\langle f|(\operatorname{Im} z)g\rangle \\
&\stackrel{(i)}{=} \operatorname{Re} z\langle f|g\rangle + i\operatorname{Im} z\langle f|g\rangle \\
&= (\operatorname{Re} z + i \operatorname{Im} z)\langle f|g\rangle \\
&= z\langle f|g\rangle,
\end{aligned}$$

which shows scaling invariance in the second argument.

Combining additivity and scaling invariance in the second argument yields linearity in the second argument.

(iii) For positive-definiteness

$$\begin{aligned}
\langle f|f\rangle &:= \frac{1}{4}(\|f + f\|^2 - \|f - f\|^2 + i\|f - if\|^2 - i\|f + if\|^2) \\
&= \frac{1}{4}(4\|f\|^2 + i|1 - i|^2\|f\|^2 - i|1 + i|^2\|f\|^2) \\
&= \frac{1}{4}(4 + i|1 - i|^2 - i|1 + i|^2)\|f\|^2 \\
&= \frac{1}{4}(4 + 2i - 2i)\|f\|^2 \\
&= \|f\|^2.
\end{aligned}$$

Thus,  $\langle f|f\rangle \geq 0$  and  $\langle f|f\rangle = 0 \Leftrightarrow f = 0$ .

Hence,  $\langle \cdot | \cdot \rangle$  is indeed a sesqui-linear inner product. Note that, from part (iii) above, we have

$$\sqrt{\langle f|f\rangle} = \|f\|.$$

That is, the inner product  $\langle \cdot | \cdot \rangle$  does induce the norm from which we started, and this completes the proof.  $\square$

*Remark 3.4.* Our proof of linearity is based on the hints given in Section 6.1, Exercise 27, from *Linear Algebra* (4th Edition) by Friedberg, Insel, Spence. Other proofs of the Jordan-von Neumann theorem can be found in

- Kadison, Ringrose, *Fundamentals of the Theory of Operator Algebras: Volume I: Elementary Theory*, American Mathematical Society 1997
- Kutateladze, *Fundamentals of Functional Analysis*, Springer 1996.
- Day, *Normed Linear Spaces*, Springer 1973.

*Remark 3.5.* Note that, often in the more mathematical literature, a sesqui-linear inner product is defined to be linear in the *first* argument rather than the second. In that case, the polarisation identity takes the form

$$\begin{aligned}
\langle f|g\rangle &= \frac{1}{4} \sum_{k=0}^3 i^k \|f + i^k g\|^2 \\
&= \frac{1}{4}(\|f + g\|^2 - \|f - g\|^2 + i\|f + ig\|^2 - i\|f - ig\|^2).
\end{aligned}$$

Example 3.6. Consider  $C_{\mathbb{C}}^0[0, 1]$  and let  $f(x) = x$  and  $g(x) = 1$ . Then

$$\|f\|_{\infty} = 1, \quad \|g\|_{\infty} = 1, \quad \|f + g\|_{\infty} = 2, \quad \|f - g\|_{\infty} = 1$$

and hence

$$\|f + g\|_{\infty}^2 + \|f - g\|_{\infty}^2 = 5 \neq 4 = 2\|f\|_{\infty}^2 + 2\|g\|_{\infty}^2.$$

Thus, by the Jordan-von Neumann theorem, there is no inner product on  $C_{\mathbb{C}}^0[0, 1]$  which induces the supremum norm. Therefore,  $(C_{\mathbb{C}}^0[0, 1], \|\cdot\|_{\infty})$  cannot be a Hilbert space.

**Proposition 3.7.** *Let  $\mathcal{H}$  be a Hilbert space. Then,  $\mathcal{H}^*$  is a Hilbert space.*

*Proof.* We already know that  $\mathcal{H}^* := \mathcal{L}(\mathcal{H}, \mathbb{C})$  is a Banach space. The norm on  $\mathcal{H}^*$  is just the usual operator norm

$$\|f\|_{\mathcal{H}^*} := \sup_{\varphi \in \mathcal{H}} \frac{|f(\varphi)|}{\|\varphi\|_{\mathcal{H}}}$$

where, admittedly somewhat perversely, we have reversed our previous notation for the dual elements. Since the modulus is induced by the standard inner product on  $\mathbb{C}$ , i.e.  $|z| = \sqrt{z\bar{z}}$ , it satisfies the parallelogram identity. Hence, we have

$$\begin{aligned} \|f + g\|_{\mathcal{H}^*}^2 + \|f - g\|_{\mathcal{H}^*}^2 &:= \left( \sup_{\varphi \in \mathcal{H}} \frac{|(f + g)(\varphi)|}{\|\varphi\|_{\mathcal{H}}} \right)^2 + \left( \sup_{\varphi \in \mathcal{H}} \frac{|(f - g)(\varphi)|}{\|\varphi\|_{\mathcal{H}}} \right)^2 \\ &= \sup_{\varphi \in \mathcal{H}} \frac{|(f + g)(\varphi)|^2}{\|\varphi\|_{\mathcal{H}}^2} + \sup_{\varphi \in \mathcal{H}} \frac{|(f - g)(\varphi)|^2}{\|\varphi\|_{\mathcal{H}}^2} \\ &= \sup_{\varphi \in \mathcal{H}} \frac{|f(\varphi) + g(\varphi)|^2 + |f(\varphi) - g(\varphi)|^2}{\|\varphi\|_{\mathcal{H}}^2} \\ &= \sup_{\varphi \in \mathcal{H}} \frac{2|f(\varphi)|^2 + 2|g(\varphi)|^2}{\|\varphi\|_{\mathcal{H}}^2} \\ &= 2 \sup_{\varphi \in \mathcal{H}} \frac{|f(\varphi)|^2}{\|\varphi\|_{\mathcal{H}}^2} + 2 \sup_{\varphi \in \mathcal{H}} \frac{|g(\varphi)|^2}{\|\varphi\|_{\mathcal{H}}^2} \\ &=: 2\|f\|_{\mathcal{H}^*}^2 + 2\|g\|_{\mathcal{H}^*}^2, \end{aligned}$$

where several steps are justified by the fact that the quantities involved are non-negative. Hence, by the Jordan-von Neumann theorem, the inner product on  $\mathcal{H}^*$  defined by the polarisation identity induces  $\|\cdot\|_{\mathcal{H}^*}$ . Hence,  $\mathcal{H}^*$  is a Hilbert space.  $\square$

The following useful fact is an immediate application of the Cauchy-Schwarz inequality.

**Proposition 3.8.** *Inner products on a vector space are sequentially continuous.*

*Proof.* Let  $\langle \cdot | \cdot \rangle$  be an inner product on  $V$ . Fix  $\varphi \in V$  and let  $\lim_{n \rightarrow \infty} \psi_n = \psi$ . Then

$$\begin{aligned} |\langle \varphi | \psi_n \rangle - \langle \varphi | \psi \rangle| &= |\langle \varphi | \psi_n - \psi \rangle| \\ &\leq \|\varphi\| \|\psi_n - \psi\| \end{aligned}$$

and hence  $\lim_{n \rightarrow \infty} \langle \varphi | \psi_n \rangle = \langle \varphi | \psi \rangle$ .  $\square$

### 3.2 Hamel versus Schauder<sup>7</sup>

Choosing a basis on a vector space is normally regarded as mathematically inelegant. The reason for this is that most statements about vector spaces are much clearer and, we maintain, aesthetically pleasing when expressed without making reference to a basis. However, in addition to the fact that some statements are more easily and usefully written in terms of a basis, bases provide a convenient way to specify the elements of a vector space in terms of components. The notion of basis for a vector space that you most probably met in your linear algebra course is more properly known as Hamel basis.

**Definition.** A *Hamel basis* of a vector space  $V$  is a subset  $\mathcal{B} \subseteq V$  such that

- (i) any finite subset  $\{e_1, \dots, e_n\} \subseteq \mathcal{B}$  is *linearly independent*, i.e.

$$\sum_{i=1}^n \lambda^i e_i = 0 \quad \Rightarrow \quad \lambda^1 = \dots = \lambda^n = 0$$

- (ii) the set  $\mathcal{B}$  is a *generating* (or *spanning*) set for  $V$ . That is, for any element  $v \in V$ , there exist a finite subset  $\{e_1, \dots, e_n\} \subseteq \mathcal{B}$  and  $\lambda^1, \dots, \lambda^n \in \mathbb{C}$  such that

$$v = \sum_{i=1}^n \lambda^i e_i.$$

Equivalently, by defining the *linear span* of a subset  $U \subseteq V$  as

$$\text{span } U := \left\{ \sum_{i=1}^n \lambda^i u_i \mid \lambda^1, \dots, \lambda^n \in \mathbb{C}, u_1, \dots, u_n \in U \text{ and } n \geq 1 \right\},$$

i.e. the set of all finite linear combinations of elements of  $U$  with complex coefficients, we can restate this condition simply as  $V = \text{span } \mathcal{B}$ .

◇

Given a basis  $\mathcal{B}$ , one can show that for each  $v \in V$  the  $\lambda^1, \dots, \lambda^n$  appearing in (ii) above are uniquely determined. They are called the *components* of  $v$  with respect to  $\mathcal{B}$ .

One can also show that if a vector space admits a finite Hamel basis  $\mathcal{B}$ , then any other basis of  $V$  is also finite and, in fact, of the same cardinality as  $\mathcal{B}$ .

**Definition.** If a vector space  $V$  admits a finite Hamel basis, then it is said to be *finite-dimensional* and its *dimension* is  $\dim V := |\mathcal{B}|$ . Otherwise, it is said to be *infinite-dimensional* and we write  $\dim V = \infty$ . ◇

**Theorem 3.9.** *Every vector space admits a Hamel basis.*

---

<sup>7</sup>Not a boxing competition.



For a proof of (a slightly more general version of) this theorem, we refer the interested reader to Dr Schuller's *Lectures on the Geometric Anatomy of Theoretical Physics*.

Note that the proof that every vector space admits a Hamel basis relies on the axiom of choice and, hence, it is non-constructive. By a corollary to Baire's category theorem, a Hamel basis on a Banach space is either finite or *uncountably* infinite. Thus, while every Banach space admits a Hamel basis, such bases on infinite-dimensional Banach spaces are difficult to construct explicitly and, hence, not terribly useful to express vectors in terms of components and perform computations. Thankfully, we can use the extra structure of a Banach space to define a more useful type of basis.

**Definition.** Let  $(W, \|\cdot\|)$  be a Banach space. A *Schauder basis* of  $W$  is a sequence  $\{e_n\}_{n \in \mathbb{N}}$  in  $W$  such that, for any  $f \in W$ , there exists a unique sequence  $\{\lambda^n\}_{n \in \mathbb{N}}$  in  $\mathbb{C}$  such that

$$f = \lim_{n \rightarrow \infty} \sum_{i=0}^n \lambda^i e_i =: \sum_{i=0}^{\infty} \lambda^i e_i$$

or, by explicitly using the definition of limit in  $W$ ,

$$\lim_{n \rightarrow \infty} \left\| f - \sum_{i=0}^n \lambda^i e_i \right\| = 0.$$

◇

We note the following points.

- Since Schauder bases require a notion of convergence, they can only be defined on a vector space equipped with a (compatible) topological structure, of which Banach spaces are a special case.
- Unlike Hamel bases, Schauder bases need not exist.
- Since the convergence of a series may depend on the order of its terms, Schauder bases must be considered as ordered bases. Hence, two Schauder bases that merely differ in the ordering of their elements are different bases, and permuting the elements of a Schauder basis doesn't necessarily yield another Schauder basis.
- The uniqueness requirement in the definition immediately implies that the zero vector cannot be an element of a Schauder basis.
- Schauder bases satisfy a stronger linear independence property than Hamel bases, namely

$$\sum_{i=0}^{\infty} \lambda^i e_i = 0 \Rightarrow \forall i \in \mathbb{N} : \lambda^i = 0.$$

- At the same time, they satisfy a weaker spanning condition. Rather than the linear span of the basis being equal to  $W$ , we only have that it is dense in  $W$ . Equivalently,

$$W = \overline{\text{span}\{e_n \mid n \in \mathbb{N}\}},$$

where the *topological closure*  $\overline{U}$  of a subset  $U \subseteq W$  is defined as

$$\overline{U} := \left\{ \lim_{n \rightarrow \infty} u_n \mid \forall n \in \mathbb{N} : u_n \in U \right\}.$$

**Definition.** A Schauder basis  $\{e_n\}_{n \in \mathbb{N}}$  of  $(W, \|\cdot\|)$  is said to be *normalised* if

$$\forall n \in \mathbb{N} : \|e_n\| = 1.$$

◇

Multiplying an element of a Schauder basis by a complex number gives again a Schauder basis (not the same one, of course). Since Schauder bases do not contain the zero vector, any Schauder basis  $\{e_n\}_{n \in \mathbb{N}}$  gives rise to a normalised Schauder basis  $\{\tilde{e}_n\}_{n \in \mathbb{N}}$  by defining

$$\tilde{e}_n := \frac{e_n}{\|e_n\|}.$$

### 3.3 Separable Hilbert spaces

Separability is a topological property. A topological space is said to be separable if it contains a dense subset which is also countable. A Banach space is said to be separable if it is separable as a topological space with the topology induced by the norm. Similarly, a Hilbert space is said to be separable if it is separable as a topological space with the topology induced by the norm induced in turn by the inner product.

For infinite-dimensional Hilbert spaces, there is a much more useful characterisation of separability, which we will henceforth take as our definition.

**Proposition 3.10.** *An infinite-dimensional Hilbert space is separable if, and only if, it admits an orthonormal Schauder basis. That is, a Schauder basis  $\{e_n\}_{n \in \mathbb{N}}$  such that*

$$\forall i, j \in \mathbb{N} : \langle e_i | e_j \rangle = \delta_{ij} := \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}.$$

Whether this holds for Banach spaces or not was a famous open problem in functional analysis, problem 153 from the Scottish book. It was solved in 1972, more than three decades after it was first posed, when Swedish mathematician Enflo constructed an infinite-dimensional separable Banach space which lacks a Schauder basis. That same year, he was awarded a live goose<sup>8</sup> for his effort.



*Remark 3.11.* The *Kronecker symbol*  $\delta_{ij}$  appearing above does not represent the components of the identity map on  $\mathcal{H}$ . Instead,  $\delta_{ij}$  are the components of the sesqui-linear form  $\langle \cdot | \cdot \rangle$ , which is a map  $\mathcal{H} \times \mathcal{H} \rightarrow \mathbb{C}$ , unlike  $\text{id}_{\mathcal{H}}$  which is a map  $\mathcal{H} \rightarrow \mathcal{H}$ . If not immediately understood, this remark may be safely ignored.

*Remark 3.12.* In finite-dimensions, since every vector space admits (by definition) a finite Hamel basis, every inner product space admits an orthonormal basis by the Gram-Schmidt orthonormalisation process.

---

<sup>8</sup>[https://en.wikipedia.org/wiki/Per\\_Enflo#Basis\\_problem\\_of\\_Banach](https://en.wikipedia.org/wiki/Per_Enflo#Basis_problem_of_Banach)

From now on, we will only consider orthonormal Schauder bases, sometimes also called Hilbert bases, and just call them bases.

**Lemma 3.13.** *Let  $\mathcal{H}$  be a Hilbert space with basis  $\{e_n\}_{n \in \mathbb{N}}$ . The unique sequence in the expansion of  $\psi \in \mathcal{H}$  in terms of this basis is  $\{\langle e_n | \psi \rangle\}_{n \in \mathbb{N}}$ .*

*Proof.* By using the continuity of the inner product, we have

$$\begin{aligned} \langle e_i | \psi \rangle &= \left\langle e_i \left| \sum_{j=0}^{\infty} \lambda^j e_j \right. \right\rangle \\ &= \left\langle e_i \left| \lim_{n \rightarrow \infty} \sum_{j=0}^n \lambda^j e_j \right. \right\rangle \\ &= \lim_{n \rightarrow \infty} \left\langle e_i \left| \sum_{j=0}^n \lambda^j e_j \right. \right\rangle \\ &= \lim_{n \rightarrow \infty} \sum_{j=0}^n \lambda^j \langle e_i | e_j \rangle \\ &= \sum_{j=0}^{\infty} \lambda^j \delta_{ij} \\ &= \lambda^i, \end{aligned}$$

which is what we wanted.  $\square$

While we have already used the term orthonormal, let us note that this means both orthogonal and normalised. Two vectors  $\varphi, \psi \in \mathcal{H}$  are said to be *orthogonal* if  $\langle \varphi | \psi \rangle = 0$ , and a subset of  $\mathcal{H}$  is called orthogonal if its elements are pairwise orthogonal.

**Lemma 3.14** (Pythagoras' theorem). *Let  $\mathcal{H}$  be a Hilbert space and let  $\{\psi_0, \dots, \psi_n\} \subset \mathcal{H}$  be a finite orthogonal set. Then*

$$\left\| \sum_{i=0}^n \psi_i \right\|^2 = \sum_{i=0}^n \|\psi_i\|^2.$$

*Proof.* Using the pairwise orthogonality of  $\{\psi_0, \dots, \psi_n\}$ , we simply calculate

$$\left\| \sum_{i=0}^n \psi_i \right\|^2 := \left\langle \sum_{i=0}^n \psi_i \left| \sum_{j=0}^n \psi_j \right. \right\rangle = \sum_{i=0}^n \langle \psi_i | \psi_i \rangle + \sum_{i \neq j} \langle \psi_i | \psi_j \rangle =: \sum_{i=0}^n \|\psi_i\|^2. \quad \square$$

**Corollary 3.15.** *Let  $\psi \in \mathcal{H}$  and let  $\{e_n\}_{n \in \mathbb{N}}$  be a basis of  $\mathcal{H}$ . Then*

$$\|\psi\|^2 = \sum_{i=0}^{\infty} |\langle e_i | \psi \rangle|^2.$$

*Proof.* By continuity of the norm, we have

$$\|\psi\|^2 = \lim_{n \rightarrow \infty} \left\| \sum_{i=0}^n \langle e_i | \psi \rangle e_i \right\|^2 = \lim_{n \rightarrow \infty} \sum_{i=0}^n |\langle e_i | \psi \rangle|^2 \|e_i\|^2 = \sum_{i=0}^{\infty} |\langle e_i | \psi \rangle|^2. \quad \square$$



### 3.4 Unitary maps

An insightful way to study a structure in mathematics is to consider maps between different instances  $A, B, C, \dots$  of that structure, and especially the structure-preserving maps. If a certain structure-preserving map  $A \rightarrow B$  is invertible and its inverse is also structure-preserving, then both these maps are generically called *isomorphisms* and  $A$  and  $B$  are said to be *isomorphic* instances of that structure. Isomorphic instances of a structure are essentially the *same* instance of that structure, just dressed up in different ways. Typically, there are infinitely many concrete instances of any given structure. The highest form of understanding of a structure that we can hope to achieve is that of a classification of its instances up to isomorphism. That is, we would like to know how many different, non-isomorphic instances of a given structure there are.

In linear algebra, the structure of interest is that of vector space over some field  $\mathbb{F}$ . The structure-preserving maps are just the linear maps and the isomorphisms are the linear bijections (whose inverses are automatically linear). Finite-dimensional vector spaces over  $\mathbb{F}$  are completely classified by their dimension, i.e. there is essentially only one vector space over  $\mathbb{F}$  for each  $n \in \mathbb{N}$ , and  $\mathbb{F}^n$  is everyone's favourite. Assuming the axiom of choice, infinite-dimensional vector spaces over  $\mathbb{F}$  are classified in the same way, namely, there is, up to linear isomorphism, only one vector space over  $\mathbb{F}$  for each infinite cardinal.

Of course, one could do better and also classify the base fields themselves. The classification of finite fields (i.e. fields with a finite number of elements) was achieved in 1893 by Moore, who proved that the order (i.e. cardinality) of a finite field is necessarily a power of some prime number, and there is only one finite field of each order, up to the appropriate notion of isomorphism. The classification of infinite fields remains an open problem.

A classification with far-reaching implications in physics is that of finite-dimensional, semi-simple, complex Lie algebras, which is discussed in some detail in Dr Schuller's *Lectures on the Geometric Anatomy of Theoretical Physics*.

The structure-preserving maps between Hilbert spaces are those that preserve both the vector space structure and the inner product. The Hilbert space isomorphisms are called unitary maps.

For further clarification see Dr Schuller's table below:

"space"	structure	structure-preserving maps	terminology
set $M$	$ \cdot $ size/cardinality	bijection $M \xrightarrow{\phi} N$	(set-) isomorphisms
vector space ( $M, +, \cdot$ )	$+: M \times M \rightarrow M,$ $\cdot: \mathbb{C} \times M \rightarrow M$	<i>linear</i> bijections: $\phi: M \rightarrow N,$ $\phi(m + \lambda \cdot n) = \phi(m) + \lambda \cdot \phi(n)$	(vector space-) isomorphisms
Hilbert space ( $M, +, \cdot, \langle \cdot   \cdot \rangle$ )	as above and $\langle \cdot   \cdot \rangle: M \times M \rightarrow \mathbb{C}$	linear bijections which satisfy: $\langle \phi(m)   \phi(n) \rangle = \langle m   n \rangle$	unitary maps (bounded)

**Definition.** Let  $\mathcal{H}$  and  $\mathcal{G}$  be Hilbert spaces. A bounded bijection  $U \in \mathcal{L}(\mathcal{H}, \mathcal{G})$  is called a

unitary map (or unitary operator) if

$$\forall \psi, \varphi \in \mathcal{H} : \langle U\psi | U\varphi \rangle_{\mathcal{G}} = \langle \psi | \varphi \rangle_{\mathcal{H}}.$$

If there exists a unitary map  $\mathcal{H} \rightarrow \mathcal{G}$ , then  $\mathcal{H}$  and  $\mathcal{G}$  are said to be *unitarily equivalent* and we write  $\mathcal{H} \cong_{\text{Hil}} \mathcal{G}$ .  $\diamond$

There are a number of equivalent definitions of unitary maps (we will later see one involving adjoints) and, in fact, our definition is fairly redundant.

**Proposition 3.16.** *Let  $U : \mathcal{H} \rightarrow \mathcal{G}$  be a surjective map which preserves the inner product. Then,  $U$  is a unitary map.*

*Proof.* (i) First, let us check that  $U$  is linear. Let  $\psi, \varphi \in \mathcal{H}$  and  $z \in \mathbb{C}$ . Then

$$\begin{aligned} \|U(z\psi + \varphi) - zU\psi - U\varphi\|_{\mathcal{G}}^2 &= \langle U(z\psi + \varphi) - zU\psi - U\varphi | U(z\psi + \varphi) - zU\psi - U\varphi \rangle_{\mathcal{G}} \\ &= \langle U(z\psi + \varphi) | U(z\psi + \varphi) \rangle_{\mathcal{G}} + |z|^2 \langle U\psi | U\psi \rangle_{\mathcal{G}} + \langle U\varphi | U\varphi \rangle_{\mathcal{G}} \\ &\quad - z \langle U(z\psi + \varphi) | U\psi \rangle_{\mathcal{G}} - \langle U(z\psi + \varphi) | U\varphi \rangle_{\mathcal{G}} + \bar{z} \langle U\psi | U\varphi \rangle_{\mathcal{G}} \\ &\quad - \bar{z} \langle U\psi | U(z\psi + \varphi) \rangle_{\mathcal{G}} - \langle U\varphi | U(z\psi + \varphi) \rangle_{\mathcal{G}} + z \langle U\varphi | U\psi \rangle_{\mathcal{G}} \\ &= \langle z\psi + \varphi | z\psi + \varphi \rangle_{\mathcal{H}} + |z|^2 \langle \psi | \psi \rangle_{\mathcal{H}} + \langle \varphi | \varphi \rangle_{\mathcal{H}} \\ &\quad - z \langle z\psi + \varphi | \psi \rangle_{\mathcal{H}} - \langle z\psi + \varphi | \varphi \rangle_{\mathcal{H}} + \bar{z} \langle \psi | \varphi \rangle_{\mathcal{H}} \\ &\quad - \bar{z} \langle \psi | z\psi + \varphi \rangle_{\mathcal{H}} - \langle \varphi | z\psi + \varphi \rangle_{\mathcal{H}} + z \langle \varphi | \psi \rangle_{\mathcal{H}} \\ &= 2|z|^2 \langle \psi | \psi \rangle_{\mathcal{H}} + \bar{z} \langle \psi | \varphi \rangle_{\mathcal{H}} + z \langle \varphi | \psi \rangle_{\mathcal{H}} + 2\langle \varphi | \varphi \rangle_{\mathcal{H}} \\ &\quad - |z|^2 \langle \psi | \psi \rangle_{\mathcal{H}} - z \langle \varphi | \psi \rangle_{\mathcal{H}} - \bar{z} \langle \psi | \varphi \rangle_{\mathcal{H}} - \langle \varphi | \varphi \rangle_{\mathcal{H}} + \bar{z} \langle \psi | \varphi \rangle_{\mathcal{H}} \\ &\quad - |z|^2 \langle \psi | \psi \rangle_{\mathcal{H}} - \bar{z} \langle \psi | \varphi \rangle_{\mathcal{H}} - z \langle \varphi | \psi \rangle_{\mathcal{H}} - \langle \varphi | \varphi \rangle_{\mathcal{H}} + z \langle \varphi | \psi \rangle_{\mathcal{H}} \\ &= 0. \end{aligned}$$

Hence  $\|U(z\psi + \varphi) - zU\psi - U\varphi\|_{\mathcal{G}} = 0$ , and thus

$$U(z\psi + \varphi) = zU\psi + U\varphi.$$

(ii) For boundedness, simply note that since for any  $\psi \in \mathcal{H}$

$$\|U\psi\|_{\mathcal{G}} := \sqrt{\langle U\psi | U\psi \rangle_{\mathcal{G}}} = \sqrt{\langle \psi | \psi \rangle_{\mathcal{H}}} =: \|\psi\|_{\mathcal{H}},$$

we have

$$\sup_{\psi \in \mathcal{H}} \frac{\|U\psi\|_{\mathcal{G}}}{\|\psi\|_{\mathcal{H}}} = 1 < \infty.$$

Hence  $U$  is bounded and, in fact, has unit operator norm.

(iii) Finally, recall that a linear map is injective if, and only if, its kernel is trivial. Suppose that  $\psi \in \ker U$ . Then, we have

$$\langle \psi | \psi \rangle_{\mathcal{H}} = \langle U\psi | U\psi \rangle_{\mathcal{G}} = \langle 0 | 0 \rangle_{\mathcal{G}} = 0.$$

Hence, by positive-definiteness,  $\psi = 0$  and thus,  $U$  is injective. Since  $U$  is also surjective by assumption, it satisfies our definition of unitary map.  $\square$

Note that a map  $U: \mathcal{H} \rightarrow \mathcal{G}$  is called an *isometry* if

$$\forall \psi \in \mathcal{H} : \|U\psi\|_{\mathcal{G}} = \|\psi\|_{\mathcal{H}}.$$

Linear isometries are, of course, the structure-preserving maps between normed spaces. We have shown that every unitary map is an isometry has unit operator norm, hence the name *unitary operator*.

Example 3.17. Consider the set of all *square-summable* complex sequences

$$\ell^2(\mathbb{N}) := \left\{ a: \mathbb{N} \rightarrow \mathbb{C} \mid \sum_{i=0}^{\infty} |a_i|^2 < \infty \right\}.$$

We define addition and scalar multiplication of sequences termwise, that is, for all  $n \in \mathbb{N}$  and all complex numbers  $z \in \mathbb{C}$ ,

$$(a + b)_n := a_n + b_n$$

$$(z \cdot a)_n := za_n.$$

These are, of course, just the discrete analogues of pointwise addition and scalar multiplication of maps. The triangle inequality and homogeneity of the modulus, together with the vector space structure of  $\mathbb{C}$ , imply that  $(\ell^2(\mathbb{N}), +, \cdot)$  is a complex vector space.

The standard inner product on  $\ell^2(\mathbb{N})$  is

$$\langle a | b \rangle_{\ell^2} := \sum_{i=0}^{\infty} \bar{a}_i b_i.$$

This inner product induces the norm

$$\|a\|_{\ell^2} := \sqrt{\langle a | a \rangle_{\ell^2}} = \sqrt{\sum_{i=0}^{\infty} |a_i|^2},$$

with respect to which  $\ell^2(\mathbb{N})$  is complete. Hence,  $(\ell^2(\mathbb{N}), +, \cdot, \langle \cdot | \cdot \rangle_{\ell^2})$  is a Hilbert space.

Consider the sequence of sequences  $\{e_n\}_{n \in \mathbb{N}}$  where

$$e_0 = (1, 0, 0, 0, \dots)$$

$$e_1 = (0, 1, 0, 0, \dots)$$

$$e_2 = (0, 0, 1, 0, \dots)$$

$$\vdots$$

i.e. we have  $(e_n)_m = \delta_{nm}$ . Each  $a \in \ell^2(\mathbb{N})$  can be written uniquely as

$$a = \sum_{i=0}^{\infty} \lambda^i e_i,$$

where  $\lambda^i = \langle e_i | a \rangle_{\ell^2} = a_i$ . The sequences  $e_n$  are clearly square-summable and, in fact, they are orthonormal with respect to  $\langle \cdot | \cdot \rangle_{\ell^2}$

$$\langle e_n | e_m \rangle_{\ell^2} := \sum_{i=0}^{\infty} \overline{(e_n)_i} (e_m)_i = \sum_{i=0}^{\infty} \delta_{ni} \delta_{mi} = \delta_{nm}.$$

Hence, the sequence  $\{e_n\}_{n \in \mathbb{N}}$  is an orthonormal Schauder basis of  $\ell^2(\mathbb{N})$ , which is therefore an infinite-dimensional separable Hilbert space.

**Theorem 3.18** (Classification of separable Hilbert spaces). *Every infinite-dimensional separable Hilbert space is unitarily equivalent to  $\ell^2(\mathbb{N})$ .*

*Proof.* Let  $\mathcal{H}$  be a separable Hilbert space with basis  $\{e_n\}_{n \in \mathbb{N}}$ . Consider the map

$$U: \mathcal{H} \rightarrow \ell^2(\mathbb{N})$$

$$\psi \mapsto \{\langle e_n | \psi \rangle_{\mathcal{H}}\}_{n \in \mathbb{N}}.$$

Note that, for any  $\psi \in \mathcal{H}$ , the sequence  $\{\langle e_n | \psi \rangle_{\mathcal{H}}\}_{n \in \mathbb{N}}$  is indeed square-summable since we have

$$\sum_{i=0}^{\infty} |\langle e_i | \psi \rangle_{\mathcal{H}}|^2 = \|\psi\|_{\mathcal{H}}^2 < \infty.$$

By our previous proposition, in order to show that  $U$  is a unitary map, it suffices to show that it is surjective and preserves the inner product. For surjectivity, let  $\{a_n\}_{n \in \mathbb{N}}$  be a complex square-summable sequence. Then, by elementary analysis, we know that  $\lim_{n \rightarrow \infty} |a_n| = 0$ . This implies that, for any  $\varepsilon > 0$ , there exists  $N \in \mathbb{N}$  such that

$$\forall n \geq m \geq N : \sum_{i=m}^n |a_i|^2 < \varepsilon.$$

Then, for all  $n, m \geq N$  (without loss of generality, assume  $n > m$ ), we have

$$\left\| \sum_{i=0}^n a_i e_i - \sum_{j=0}^m a_j e_j \right\|_{\mathcal{H}}^2 = \left\| \sum_{i=m+1}^n a_i e_i \right\|_{\mathcal{H}}^2 = \sum_{i=m+1}^n |a_i|^2 \|e_i\|_{\mathcal{H}}^2 = \sum_{i=m+1}^n |a_i|^2 < \varepsilon.$$

That is,  $\{\sum_{i=0}^n a_i e_i\}_{n \in \mathbb{N}}$  is a Cauchy sequence in  $\mathcal{H}$ . Hence, by completeness, there exists  $\psi \in \mathcal{H}$  such that

$$\psi = \sum_{i=0}^{\infty} a_i e_i$$

and we have  $U\psi = \{a_n\}_{n \in \mathbb{N}}$ , so  $U$  is surjective. Moreover, we have

$$\begin{aligned} \langle \psi | \varphi \rangle_{\mathcal{H}} &= \left\langle \sum_{i=0}^{\infty} \langle e_i | \psi \rangle_{\mathcal{H}} e_i \left| \sum_{j=0}^{\infty} \langle e_j | \varphi \rangle_{\mathcal{H}} e_j \right\rangle_{\mathcal{H}} \\ &= \sum_{j=0}^{\infty} \sum_{i=0}^{\infty} \overline{\langle e_i | \psi \rangle_{\mathcal{H}}} \langle e_j | \varphi \rangle_{\mathcal{H}} \langle e_i | e_j \rangle_{\mathcal{H}} \\ &= \sum_{j=0}^{\infty} \sum_{i=0}^{\infty} \overline{\langle e_i | \psi \rangle_{\mathcal{H}}} \langle e_j | \varphi \rangle_{\mathcal{H}} \delta_{ij} \\ &= \sum_{j=0}^{\infty} \overline{\langle e_j | \psi \rangle_{\mathcal{H}}} \langle e_j | \varphi \rangle_{\mathcal{H}} \\ &=: \langle \{\langle e_n | \psi \rangle_{\mathcal{H}}\}_{n \in \mathbb{N}} | \{\langle e_n | \varphi \rangle_{\mathcal{H}}\}_{n \in \mathbb{N}} \rangle_{\ell^2} \\ &=: \langle U\psi | U\varphi \rangle_{\ell^2}. \end{aligned}$$

Hence,  $U$  preserves the inner product, and it is therefore a unitary map.  $\square$

## 4 Projectors, bras and kets

### 4.1 Projectors

Projectors play a key role in quantum theory, as you can see from Axioms 3 and 5.

**Definition.** Let  $\mathcal{H}$  be a separable Hilbert space. Fix a unit vector  $e \in \mathcal{H}$  (that is,  $\|e\| = 1$ ) and let  $\psi \in \mathcal{H}$ . The *projection* of  $\psi$  to  $e$  is

$$\psi_{\parallel} := \langle e | \psi \rangle e$$

while the *orthogonal complement* of  $\psi$  is

$$\psi_{\perp} := \psi - \psi_{\parallel}.$$

◇

We can extend these definitions to a countable orthonormal subset  $\{e_i\}_{i \in \mathbb{N}} \subset \mathcal{H}$ , i.e. a subset of  $\mathcal{H}$  whose elements are pairwise orthogonal and have unit norm. Note that  $\{e_i\}_{i \in \mathbb{N}}$  need not be a basis of  $\mathcal{H}$ .

**Proposition 4.1.** *Let  $\psi \in \mathcal{H}$  and let  $\{e_i\}_{i \in \mathbb{N}} \subset \mathcal{H}$  be an orthonormal subset. Then*

(a) *we can write  $\psi = \psi_{\parallel} + \psi_{\perp}$ , where*

$$\psi_{\parallel} := \sum_{i=0}^{\infty} \langle e_i | \psi \rangle e_i, \quad \psi_{\perp} := \psi - \psi_{\parallel}$$

*and we have*

$$\forall i \in \mathbb{N} : \langle \psi_{\perp} | e_i \rangle = 0.$$

(b) *Pythagoras' theorem holds:*

$$\|\psi\|^2 = \|\psi_{\parallel}\|^2 + \|\psi_{\perp}\|^2.$$

*Note that this is an extension to the finite-dimensional case.*

(c) *for any  $\gamma \in \text{span}\{e_i \mid i \in \mathbb{N}\}$ , we have the estimate*

$$\|\psi - \gamma\| \geq \|\psi_{\perp}\|$$

*with equality if, and only if,  $\gamma = \psi_{\parallel}$ .*

*Proof.* First consider the case of a finite orthonormal subset  $\{e_0, \dots, e_n\} \subset \mathcal{H}$ .

(a) Let  $\psi_{\parallel}$  and  $\psi_{\perp}$  be defined as in the proposition. Then  $\psi_{\parallel} + \psi_{\perp} = \psi$  and

$$\begin{aligned} \langle \psi_{\perp} | e_i \rangle &= \left\langle \psi - \sum_{j=0}^n \langle e_j | \psi \rangle e_j \middle| e_i \right\rangle \\ &= \langle \psi | e_i \rangle - \sum_{j=0}^n \overline{\langle e_j | \psi \rangle} \langle e_j | e_i \rangle \\ &= \langle \psi | e_i \rangle - \sum_{j=0}^n \langle \psi | e_j \rangle \delta_{ji} \\ &= \langle \psi | e_i \rangle - \langle \psi | e_i \rangle \\ &= 0 \end{aligned}$$

for all  $0 \leq i \leq n$ .

(b) From part (a), we have

$$\langle \psi_{\perp} | \psi_{\parallel} \rangle = \left\langle \psi_{\perp} \left| \sum_{i=0}^n \langle e_i | \psi \rangle e_i \right. \right\rangle = \sum_{i=0}^n \langle e_i | \psi \rangle \langle \psi_{\perp} | e_i \rangle = 0.$$

Hence, by (the finite-dimensional) Pythagoras' theorem

$$\|\psi\|^2 = \|\psi_{\parallel} + \psi_{\perp}\|^2 = \|\psi_{\parallel}\|^2 + \|\psi_{\perp}\|^2.$$

(c) Let  $\gamma \in \text{span}\{e_i \mid 0 \leq i \leq n\}$ . Then  $\gamma = \sum_{i=0}^n \gamma_i e_i$  for some  $\gamma_0, \dots, \gamma_n \in \mathbb{C}$ . Hence

$$\begin{aligned} \|\psi - \gamma\|^2 &= \|\psi_{\perp} + \psi_{\parallel} - \gamma\|^2 \\ &= \left\| \psi_{\perp} + \sum_{i=0}^n \langle e_i | \psi \rangle e_i - \sum_{i=0}^n \gamma_i e_i \right\|^2 \\ &= \left\| \psi_{\perp} + \sum_{i=0}^n (\langle e_i | \psi \rangle - \gamma_i) e_i \right\|^2 \\ &= \|\psi_{\perp}\|^2 + \sum_{i=0}^n |\langle e_i | \psi \rangle - \gamma_i|^2 \end{aligned}$$

and thus  $\|\psi - \gamma\| \geq \|\psi_{\perp}\|$  since  $|\langle e_i | \psi \rangle - \gamma_i|^2 \geq 0$  for all  $0 \leq i \leq n$ . Moreover, we have equality if, and only if,  $|\langle e_i | \psi \rangle - \gamma_i| = 0$  for all  $0 \leq i \leq n$ , that is  $\gamma = \psi_{\parallel}$ .

To extend this to a countably infinite orthonormal set  $\{e_i\}_{i \in \mathbb{N}}$ , note that by part (b) and Bessel's inequality, we have

$$\left\| \sum_{i=0}^n \langle e_i | \psi \rangle e_i \right\|^2 = \sum_{i=0}^n |\langle e_i | \psi \rangle|^2 \leq \|\psi\|^2.$$

Since  $|\langle e_i | \psi \rangle|^2 \geq 0$ , the sequence of partial sums  $\{\sum_{i=0}^n |\langle e_i | \psi \rangle|^2\}_{n \in \mathbb{N}}$  is monotonically increasing and bounded from above by  $\|\psi\|^2$ . Hence, it converges and this implies that

$$\psi_{\parallel} := \sum_{i=0}^{\infty} \langle e_i | \psi \rangle e_i$$

exists as an element of  $\mathcal{H}$ . The extension to the countably infinite case then follows by continuity of the inner product.  $\square$

## 4.2 Closed linear subspaces

We will often be interested in looking at linear subspaces of a Hilbert space  $\mathcal{H}$ , i.e. subsets  $\mathcal{M} \subseteq \mathcal{H}$  such that

$$\forall \psi, \varphi \in \mathcal{M} : \forall z \in \mathbb{C} : z\psi + \varphi \in \mathcal{M}.$$

Note that while every linear subspace  $\mathcal{M} \subset \mathcal{H}$  inherits the inner product on  $\mathcal{H}$  to become an inner product space, it may fail to be complete with respect to this inner product. In other words, not every linear subspace of a Hilbert space is necessarily a sub-Hilbert space.

The following definitions are with respect to the norm topology on a normed space and can, of course, be given more generally on an arbitrary topological space.

**Definition.** Let  $\mathcal{H}$  be a normed space. A subset  $\mathcal{M} \subset \mathcal{H}$  is said to be *open* if

$$\forall \psi \in \mathcal{M} : \exists r > 0 : \forall \varphi \in \mathcal{H} : \|\psi - \varphi\| < r \Rightarrow \varphi \in \mathcal{M}.$$

◇

Equivalently, by defining the *open ball* of radius  $r > 0$  and centre  $\psi \in \mathcal{H}$

$$B_r(\psi) := \{\varphi \in \mathcal{H} \mid \|\psi - \varphi\| < r\},$$

we can define  $\mathcal{M} \subset \mathcal{H}$  to be open if

$$\forall \psi \in \mathcal{M} : \exists r > 0 : B_r(\psi) \subseteq \mathcal{M}.$$

**Definition.** A subset  $\mathcal{M} \subset \mathcal{H}$  is said to be *closed* if its complement  $\mathcal{H} \setminus \mathcal{M}$  is open. ◇

**Proposition 4.2.** A closed subset  $\mathcal{M}$  of a complete normed space  $\mathcal{H}$  is complete.

*Proof.* Let  $\{\psi_n\}_{n \in \mathbb{N}}$  be a Cauchy sequence in the closed subset  $\mathcal{M}$ . Then,  $\{\psi_n\}_{n \in \mathbb{N}}$  is also a Cauchy sequence in  $\mathcal{H}$ , and hence it converges to some  $\psi \in \mathcal{H}$  since  $\mathcal{H}$  is complete. We want to show that, in fact,  $\psi \in \mathcal{M}$ . Suppose, for the sake of contradiction, that  $\psi \notin \mathcal{M}$ , i.e.  $\psi \in \mathcal{H} \setminus \mathcal{M}$ . Since  $\mathcal{M}$  is closed,  $\mathcal{H} \setminus \mathcal{M}$  is open. Hence, there exists  $r > 0$  such that

$$\forall \varphi \in \mathcal{H} : \|\varphi - \psi\| < r \Rightarrow \varphi \in \mathcal{H} \setminus \mathcal{M}.$$

However, since  $\psi$  is the limit of  $\{\psi_n\}_{n \in \mathbb{N}}$ , there exists  $N \in \mathbb{N}$  such that

$$\forall n \geq N : \|\psi_n - \psi\| < r.$$

Hence, for all  $n \geq N$ , we have  $\psi_n \in \mathcal{H} \setminus \mathcal{M}$ , i.e.  $\psi_n \notin \mathcal{M}$ , contradicting the fact that  $\{\psi_n\}_{n \in \mathbb{N}}$  is a sequence in  $\mathcal{M}$ . Thus, we must have  $\psi \in \mathcal{M}$ . □

**Corollary 4.3.** A closed linear subspace  $\mathcal{M}$  of a Hilbert space  $\mathcal{H}$  is a sub-Hilbert space with the inner product on  $\mathcal{H}$ . Moreover, if  $\mathcal{H}$  is separable, then so is  $\mathcal{M}$ .

Knowing that a linear subspace of a Hilbert space is, in fact, a sub-Hilbert space can be very useful. For instance, we know that there exists an orthonormal basis for the linear subspace. Note that the converse to the corollary does not hold: a sub-Hilbert space need not be a closed linear subspace.

### 4.3 Orthogonal projections

**Definition.** Let  $\mathcal{M} \subseteq \mathcal{H}$  be a (not necessarily closed) linear subspace of  $\mathcal{H}$ . The set

$$\mathcal{M}^\perp := \{\psi \in \mathcal{H} \mid \forall \varphi \in \mathcal{M} : \langle \varphi | \psi \rangle = 0\}$$

is called the *orthogonal complement* of  $\mathcal{M}$  in  $\mathcal{H}$ . ◇

**Proposition 4.4.** Let  $\mathcal{M} \subseteq \mathcal{H}$  be a linear subspace of  $\mathcal{H}$ . Then,  $\mathcal{M}^\perp$  is a closed linear subspace of  $\mathcal{H}$ .

*Proof.* Let  $\psi_1, \psi_2 \in \mathcal{M}^\perp$  and  $z \in \mathbb{C}$ . Then, for all  $\varphi \in \mathcal{M}$

$$\langle \varphi | z\psi_1 + \psi_2 \rangle = z\langle \varphi | \psi_1 \rangle + \langle \varphi | \psi_2 \rangle = 0$$

and hence  $z\psi_1 + \psi_2 \in \mathcal{M}^\perp$ . Thus,  $\mathcal{M}^\perp$  is a linear subspace of  $\mathcal{H}$ . It remains to be shown that it is also closed. Define the maps

$$\begin{aligned} f_\varphi: \mathcal{H} &\rightarrow \mathbb{C} \\ \psi &\mapsto \langle \varphi | \psi \rangle. \end{aligned}$$

Then, we can write

$$\mathcal{M}^\perp = \bigcap_{\varphi \in \mathcal{M}} \text{preim}_{f_\varphi}(\{0\}).$$

Since the inner product is continuous (in each slot), the maps  $f_\varphi$  are continuous. Hence, the pre-images of closed sets are closed. As the singleton  $\{0\}$  is closed in the standard topology on  $\mathbb{C}$ , the sets  $\text{preim}_{f_\varphi}(\{0\})$  are closed for all  $\varphi \in \mathcal{M}$ . Thus,  $\mathcal{M}^\perp$  is closed since arbitrary intersections of closed sets are closed.  $\square$

*Remark 4.5.* Note that  $\mathcal{M}^\perp$  is also not open (which is not necessarily the same as closed). To clarify, equally the preimage of  $\{0\}$  is an open set (as  $\{0\}$  is open), however it is only *finite* intersections of open sets that are open. So the inclusion of the intersection plays an important role.

Note that by Pythagoras' theorem, we have the decomposition

$$\mathcal{H} = \mathcal{M} \oplus \mathcal{M}^\perp := \{\psi + \varphi \mid \psi \in \mathcal{M}, \varphi \in \mathcal{M}^\perp\}$$

for any closed linear subspace  $\mathcal{M}$ .

**Proposition 4.6.** *For any closed linear subspace  $X \subset \mathcal{H}$  its true that  $X^{\perp\perp} = X$ .*

*Proof.* Let  $x \in X$ , then for all  $y \in X^\perp$  we have  $\langle x, y \rangle = 0$  and so  $x \in X^{\perp\perp}$ . This gives  $X \subseteq X^{\perp\perp}$ .

Now consider  $z \in X^{\perp\perp}$ . As  $X$  is closed from the above note we know it can be decomposed as  $z = x + y$  for  $x \in X$  and  $y \in X^\perp$ . We then have

$$\begin{aligned} 0 &= \langle y, z \rangle \\ &= \langle y, x + y \rangle \\ &= \langle y, x \rangle + \langle y, y \rangle \\ &= \|y\|^2 \\ \implies y &= 0 \end{aligned}$$

where the last step comes from the definiteness of  $\|\cdot\|$ . So we have  $z \in X$  and  $X^{\perp\perp} \subseteq X$ .  $\square$

**Proposition 4.7.** *For any linear subspace  $\mathcal{M} \subseteq \mathcal{H}$  it is true that  $\mathcal{M}^{\perp\perp} = \overline{\mathcal{M}}$ , where the latter is the topological closure of the set.*



*Proof.* We start with two observations:

- (i)  $\mathcal{M} \subseteq \mathcal{M}^{\perp\perp}$ , which was shown at the start of the last proof (as there was no use of the fact that  $X$  was closed there).
- (ii) If  $\mathcal{M}_1 \subseteq \mathcal{M}_2$  then  $\mathcal{M}_2^\perp \subseteq \mathcal{M}_1^\perp$ , which can be shown easily.

First let's show that  $\mathcal{M}^{\perp\perp} \subseteq \overline{\mathcal{M}}$ . Clearly  $\mathcal{M} \subseteq \overline{\mathcal{M}}$  (where the equality holds only if  $\mathcal{M}$  is closed), and so from (ii) we have  $\overline{\mathcal{M}}^\perp \subseteq \mathcal{M}^\perp$ , which in turn gives  $\mathcal{M}^{\perp\perp} \subseteq \overline{\mathcal{M}}^{\perp\perp}$ . But  $\overline{\mathcal{M}}$  is closed and so from the previous proposition we have  $\mathcal{M}^{\perp\perp} \subseteq \overline{\mathcal{M}}$ .

Now we need to show the reverse inclusion. From Proposition 4.4 we know that  $\mathcal{M}^{\perp\perp}$  is closed. Then (i) instantly tells us that  $\overline{\mathcal{M}} \subseteq \mathcal{M}^{\perp\perp}$ .  $\square$

**Definition.** Let  $\mathcal{M}$  be a closed linear subspace of a separable Hilbert space  $\mathcal{H}$  and fix some orthonormal basis of  $\mathcal{M}$ . The map

$$\begin{aligned} P_{\mathcal{M}}: \mathcal{H} &\rightarrow \mathcal{M} \\ \psi &\mapsto \psi_{\parallel} \end{aligned}$$

is called the *orthogonal projector* to  $\mathcal{M}$ .  $\diamond$

**Proposition 4.8.** Let  $P_{\mathcal{M}}: \mathcal{H} \rightarrow \mathcal{M}$  be an orthogonal projector to  $\mathcal{M} \subseteq \mathcal{H}$ . Then

- (i)  $P_{\mathcal{M}} \circ P_{\mathcal{M}} = P_{\mathcal{M}}$ , sometimes also written as  $P_{\mathcal{M}}^2 = P_{\mathcal{M}}$
- (ii)  $\forall \psi, \varphi \in \mathcal{H}: \langle P_{\mathcal{M}}\psi | \varphi \rangle = \langle \psi | P_{\mathcal{M}}\varphi \rangle$
- (iii)  $P_{\mathcal{M}^\perp}\psi = \psi_{\perp}$
- (iv)  $P_{\mathcal{M}} \in \mathcal{L}(\mathcal{H}, \mathcal{M})$ .

*Proof.* Let  $\{e_i\}_{i \in I}$  and  $\{e_i\}_{i \in J}$  be bases of  $\mathcal{M}$  and  $\mathcal{M}^\perp$  respectively, where  $I, J$  are disjoint and either finite or countably infinite, such that  $\{e_i\}_{i \in I \cup J}$  is a basis of  $\mathcal{H}$  (Note that we should think of  $I \cup J$  as having a definite ordering).

- (i) Let  $\psi \in \mathcal{H}$ . Then

$$\begin{aligned} P_{\mathcal{M}}(P_{\mathcal{M}}\psi) &:= P_{\mathcal{M}}\left(\sum_{i \in I} \langle e_i | \psi \rangle e_i\right) \\ &:= \sum_{j \in I} \left\langle e_j \left| \sum_{i \in I} \langle e_i | \psi \rangle e_i \right. \right\rangle e_j \\ &= \sum_{j \in I} \sum_{i \in I} \langle e_i | \psi \rangle \langle e_j | e_i \rangle e_j \\ &= \sum_{i \in I} \langle e_i | \psi \rangle e_i \\ &=: P_{\mathcal{M}}\psi. \end{aligned}$$

(ii) Let  $\psi, \varphi \in \mathcal{H}$ . Then

$$\begin{aligned}
\langle P_{\mathcal{M}}\psi | \varphi \rangle &:= \left\langle \sum_{i \in I} \langle e_i | \psi \rangle e_i \middle| \varphi \right\rangle \\
&= \sum_{i \in I} \overline{\langle e_i | \psi \rangle} \langle e_i | \varphi \rangle \\
&= \sum_{i \in I} \langle e_i | \varphi \rangle \langle \psi | e_i \rangle \\
&= \left\langle \psi \middle| \sum_{i \in I} \langle e_i | \varphi \rangle e_i \right\rangle \\
&=: \langle \psi | P_{\mathcal{M}}\varphi \rangle.
\end{aligned}$$

(iii) Let  $\psi \in \mathcal{H}$ . Then

$$P_{\mathcal{M}}\psi + P_{\mathcal{M}^\perp}\psi = \sum_{i \in I} \langle e_i | \psi \rangle e_i + \sum_{i \in J} \langle e_i | \psi \rangle e_i = \sum_{i \in I \cup J} \langle e_i | \psi \rangle e_i = \psi.$$

Hence

$$P_{\mathcal{M}^\perp}\psi = \psi - P_{\mathcal{M}}\psi = \psi - \psi_{\parallel} =: \psi_{\perp}.$$

(iv) Let  $\psi \in \mathcal{H}$ . Then, by Pythagoras' theorem,

$$\sup_{\psi \in \mathcal{H}} \frac{\|P_{\mathcal{M}}\psi\|}{\|\psi\|} = \sup_{\psi \in \mathcal{H}} \frac{\|\psi_{\parallel}\|}{\|\psi\|} = \sup_{\psi \in \mathcal{H}} \frac{\|\psi\| - \|\psi_{\perp}\|}{\|\psi\|} \leq 1 < \infty \quad \square$$

Quite interesting, and heavily used, is the converse.

**Theorem 4.9.** *Let  $P \in \mathcal{L}(\mathcal{H}, \mathcal{H})$  have the properties*

$$(i) \quad P \circ P = P$$

$$(ii) \quad \forall \psi, \varphi \in \mathcal{H} : \langle P\psi | \varphi \rangle = \langle \psi | P\varphi \rangle.$$

*Then, the range  $P(\mathcal{H})$  of  $P$  is closed and*

$$P = P_{P(\mathcal{H})}.$$

*In other words, every projector is the orthogonal projector to some closed linear subspace.*

#### 4.4 Riesz representation theorem, bras and kets

Let  $\mathcal{H}$  be a Hilbert space. Consider again the map

$$\begin{aligned}
f_{\varphi} : \mathcal{H} &\rightarrow \mathbb{C} \\
\psi &\mapsto \langle \varphi | \psi \rangle.
\end{aligned}$$

for  $\varphi \in \mathcal{H}$ . The linearity in the second argument of the inner product implies that this map is linear. Moreover, by the Cauchy-Schwarz inequality, we have

$$\sup_{\psi \in \mathcal{H}} \frac{|f_{\varphi}(\psi)|}{\|\psi\|} = \sup_{\psi \in \mathcal{H}} \frac{|\langle \varphi | \psi \rangle|}{\|\psi\|} \leq \sup_{\psi \in \mathcal{H}} \frac{\|\varphi\| \|\psi\|}{\|\psi\|} = \|\varphi\| < \infty.$$

Hence,  $f_{\varphi} \in \mathcal{L}(\mathcal{H}, \mathbb{C}) = \mathcal{H}^*$ . Therefore, to every element  $\varphi$  of  $\mathcal{H}$ , there is associated an element  $f_{\varphi}$  in the dual space  $\mathcal{H}^*$ . In fact, the converse is also true.

**Theorem 4.10** (Riesz representation). *Every  $f \in \mathcal{H}^*$  is of the form  $f_\varphi$  for a unique  $\varphi \in \mathcal{H}$ .*

*Proof.* First, suppose that  $f = 0$ , i.e.  $f$  is the zero functional on  $\mathcal{H}$ . Then, clearly,  $f = f_0$  with  $0 \in \mathcal{H}$ . Hence, suppose that  $f \neq 0$ . Since,  $\ker f := \text{preim}_f(\{0\})$  is a closed linear subspace, we can write

$$\mathcal{H} = \ker f \oplus (\ker f)^\perp.$$

As  $f \neq 0$ , there exists some  $\psi \in \mathcal{H}$  such that  $\psi \notin \ker f$ . Hence,  $\ker f \neq \mathcal{H}$ , and thus  $(\ker f)^\perp \neq \{0\}$ . Let  $\xi \in (\ker f)^\perp \setminus \{0\}$  and assume, w.l.o.g., that  $\|\xi\| = 1$ . Define

$$\varphi := \overline{f(\xi)}\xi \in (\ker f)^\perp.$$

Then, for any  $\psi \in \mathcal{H}$ , we have

$$\begin{aligned} f_\varphi(\psi) - f(\psi) &:= \langle \varphi | \psi \rangle - f(\psi) \\ &:= \langle \overline{f(\xi)}\xi | \psi \rangle - f(\psi)\langle \xi | \xi \rangle \\ &= \langle \xi | f(\xi)\psi \rangle - \langle \xi | f(\psi)\xi \rangle \\ &= \langle \xi | f(\xi)\psi - f(\psi)\xi \rangle. \end{aligned}$$

Note that

$$f(f(\xi)\psi - f(\psi)\xi) = f(\xi)f(\psi) - f(\psi)f(\xi) = 0,$$

that is,  $f(\xi)\psi - f(\psi)\xi \in \ker f$ . Since  $\xi \in (\ker f)^\perp$ , we have

$$\langle \xi | f(\xi)\psi - f(\psi)\xi \rangle = 0$$

and hence  $f_\varphi(\psi) = f(\psi)$  for all  $\psi \in \mathcal{H}$ , i.e.  $f = f_\varphi$ . For uniqueness, suppose that

$$f = f_{\varphi_1} = f_{\varphi_2}$$

for some  $\varphi_1, \varphi_2 \in \mathcal{H}$ . Then, for any  $\psi \in \mathcal{H}$ ,

$$\begin{aligned} 0 &= f_{\varphi_1}(\psi) - f_{\varphi_2}(\psi) \\ &= \langle \varphi_1 | \psi \rangle - \langle \varphi_2 | \psi \rangle \\ &= \langle \varphi_1 - \varphi_2 | \psi \rangle \end{aligned}$$

and hence,  $\varphi_1 = \varphi_2$  by positive-definiteness. □

Therefore, the so-called *Riesz map*

$$\begin{aligned} R: \mathcal{H} &\rightarrow \mathcal{H}^* \\ \varphi &\mapsto f_\varphi \equiv \langle \varphi | \cdot \rangle \end{aligned}$$

is a linear isomorphism, and  $\mathcal{H}$  and  $\mathcal{H}^*$  be identified with one another as vector spaces. This lead Dirac to suggest the following notation for the elements of the dual space

$$f_\varphi \equiv \langle \varphi | \cdot \rangle.$$

Correspondingly, he wrote  $|\psi\rangle$  for the element  $\psi \in \mathcal{H}$ . Since  $\langle \cdot | \cdot \rangle$  is “a bracket”, the first half  $\langle \cdot |$  is called a *bra*, while the second half  $|\cdot\rangle$  is called a *ket* (nobody knows where the missing *c* is). With this notation, we have

$$f_\varphi(\psi) \equiv \langle \varphi | (|\psi\rangle) \equiv \langle \varphi | \psi \rangle.$$

The notation makes evident the fact that, for any  $\varphi, \psi \in \mathcal{H}$ , we can always consider the inner product  $\langle \varphi | \psi \rangle \in \mathbb{C}$  as the result of applying  $f_\varphi \in \mathcal{H}^*$  to  $\psi \in \mathcal{H}$ .

The advantage of this notation is that some formulæ become more intuitive and hence are more easily memorised. For a concrete example, consider

$$\psi = \sum_{i=0}^{\infty} \langle e_i | \psi \rangle e_i$$

where  $\{e_n\}_{n \in \mathbb{N}}$  is a basis of  $\mathcal{H}$ . This becomes

$$|\psi\rangle = \sum_{i=0}^{\infty} \langle e_i | \psi \rangle |e_i\rangle.$$

By allowing the scalar multiplication of kets also from the right, defined to yield the same result as that on the left, we have

$$= \sum_{i=0}^{\infty} |e_i\rangle \langle e_i | \psi \rangle.$$

“Quite obviously”, we can bracket this as

$$= \left( \sum_{i=0}^{\infty} |e_i\rangle \langle e_i| \right) |\psi\rangle,$$

where by “quite obviously”, we mean that we have a suppressed tensor product (see section 8 of the *Lectures on the Geometric Anatomy of Theoretical Physics* for more details on tensors)

$$= \left( \sum_{i=0}^{\infty} |e_i\rangle \otimes \langle e_i| \right) |\psi\rangle.$$

Then, the sum in the round brackets is an element of  $\mathcal{H} \otimes \mathcal{H}^*$ . While  $\mathcal{H} \otimes \mathcal{H}^*$  is isomorphic to  $\text{End}(\mathcal{H})$ , its elements are maps  $\mathcal{H}^* \times \mathcal{H} \rightarrow \mathbb{C}$ . Hence, one needs to either make this isomorphism explicit or, equivalently,

$$= \left( \sum_{i=0}^{\infty} |e_i\rangle \otimes \langle e_i| \right) (\cdot, |\psi\rangle).$$

All of this to be able to write

$$\sum_{i=0}^{\infty} |e_i\rangle \langle e_i| = \text{id}_{\mathcal{H}}$$

and hence interpret the expansion of  $|\psi\rangle$  in terms of the basis as the “insertion” of an identity

$$|\psi\rangle = \text{id}_{\mathcal{H}} |\psi\rangle = \left( \sum_{i=0}^{\infty} |e_i\rangle\langle e_i| \right) |\psi\rangle = \sum_{i=0}^{\infty} \langle e_i|\psi\rangle |e_i\rangle.$$

But the original expression was already clear in the first place, without the need to add hidden tensor products and extra rules. Of course, part of the appeal of this notation is that one can intuitively think of something like  $|e_i\rangle\langle e_i|$  as a map  $\mathcal{H} \rightarrow \mathcal{H}$ , by imagining that the bra on the right acts on a ket in  $\mathcal{H}$ , thereby producing a complex number which becomes the coefficient of the remaining ket

$$(|e_i\rangle\langle e_i|)|\psi\rangle = |e_i\rangle\langle e_i|\psi\rangle = \langle e_i|\psi\rangle |e_i\rangle.$$

The major drawback of this notation, and the reason why we will *not* adopt it, is that in many places (for instance, if we consider self-adjoint operators, or Hermitian operators) this notation doesn’t produce inconsistencies only if certain conditions are satisfied. While these conditions will indeed be satisfied most of the times, it becomes extremely confusing to formulate conditions on our objects by using a notation that only makes sense if the objects already satisfy conditions.

Of course, as this notation is heavily used in physics and related applied sciences, it is necessary to be able to recognise it and become fluent in it. But note that it does *not* make things clearer. If anything, it makes things more complicated.

## 5 Measure theory

This and the next section will be a short recap of basic notions from measure theory and Lebesgue integration. These are inescapable subjects if one wants to understand quantum mechanics since

- (i) the spectral theorem requires the notion of (projection-valued) measures
- (ii) the most commonly emerging separable Hilbert space in quantum mechanics is the space  $L^2(\mathbb{R}^d)$ , whose definition needs the notion of Lebesgue integral.

### 5.1 General measure spaces and basic results

**Definition.** Let  $M$  be a non-empty set. A collection  $\Sigma \subseteq \mathcal{P}(M)$  of subsets of  $M$  is called a  $\sigma$ -algebra for  $M$  if the following conditions are satisfied

- (i)  $M \in \Sigma$
- (ii) if  $A \in \Sigma$ , then  $M \setminus A \in \Sigma$
- (iii) for any sequence  $\{A_n\}_{n \in \mathbb{N}}$  in  $\Sigma$  we have  $\bigcup_{n=0}^{\infty} A_n \in \Sigma$ .

◇

*Remark 5.1.* If we relax the third condition so that it applies only to finite (rather than countable) unions, we obtain the notion of an *algebra*, often called an *algebra of sets* in order to distinguish it from the notion of algebra as a vector space equipped with a bilinear product, with which it has nothing to do.

*Remark 5.2.* Note that by condition (ii) and De Morgan's laws, condition (iii) can be equivalently stated in terms of intersections rather than unions. Recall that De Morgan's laws "interchange" unions with intersections and vice-versa under the complement operation. That is, if  $M$  is a set and  $\{A_i\}_{i \in I}$  is a collection of sets, then

$$M \setminus \left( \bigcup_{i \in I} A_i \right) = \bigcap_{i \in I} (M \setminus A_i), \quad M \setminus \left( \bigcap_{i \in I} A_i \right) = \bigcup_{i \in I} (M \setminus A_i).$$

A  $\sigma$ -algebra is closed under countably infinite unions (by definition) but also under countably infinite intersections and finite unions and intersections.

**Proposition 5.3.** Let  $M$  be a set and let  $\Sigma$  be a  $\sigma$ -algebra on  $M$ . Let  $\{A_n\}_{n \in \mathbb{N}}$  be a sequence in  $\Sigma$ . Then, for all  $k \in \mathbb{N}$ , we have

- (i)  $\bigcup_{n=0}^k A_n \in \Sigma$
- (ii)  $\bigcap_{n=0}^{\infty} A_n \in \Sigma$  and  $\bigcap_{n=0}^k A_n \in \Sigma$ .

*Proof.* (i) Let the sequence  $\{B_n\}_{n \in \mathbb{N}}$  be defined as follows:

$$B_n = \begin{cases} A_n & \text{if } 0 \leq n \leq k \\ \emptyset & \text{if } n > k. \end{cases}$$

Then,  $\{B_n\}_{n \in \mathbb{N}}$  is a sequence in  $\Sigma$ , so  $\bigcup_{n=0}^{\infty} B_n \in \Sigma$ . Hence, we have:

$$\bigcup_{n=0}^{\infty} B_n = \left( \bigcup_{n=0}^k B_n \right) \cup \left( \bigcup_{n=k+1}^{\infty} B_n \right) = \bigcup_{n=0}^k A_n$$

and thus  $\bigcup_{n=0}^k A_n \in \Sigma$ .

(ii) As  $\{A_n\}_{n \in \mathbb{N}}$  is a sequence in  $\Sigma$ , so is  $\{M \setminus A_n\}_{n \in \mathbb{N}}$  and hence  $\bigcup_{n=0}^{\infty} (M \setminus A_n) \in \Sigma$ . Thus, we also have

$$M \setminus \left( \bigcup_{n=0}^{\infty} (M \setminus A_n) \right) \in \Sigma$$

and since  $M \setminus (M \setminus A_n) = A_n$ , by De Morgan's laws,  $\bigcap_{n=0}^{\infty} A_n \in \Sigma$ . That this holds for finite intersections is shown by defining  $\{B_n\}_{n \in \mathbb{N}}$  as above.  $\square$

**Definition.** A *measurable space* is a pair  $(M, \Sigma)$  where  $M$  is a set and  $\Sigma$  is a  $\sigma$ -algebra on  $M$ . The elements of  $\Sigma$  are called *measurable subsets* of  $M$ .  $\diamond$

Our goal is to assign volumes (i.e. measures) to subsets of a given set. Of course, we would also like this assignment to satisfy some sensible conditions. However, it turns out that one cannot sensibly assign volumes to any arbitrary collection of subsets of a given set<sup>9</sup>. It is necessary that the collection of subsets be a  $\sigma$ -algebra. In addition, just like in topology openness and closeness are not properties of subsets but properties of subsets with respect to a choice of topology, so does measurability of subsets only make sense with respect to a choice of  $\sigma$ -algebra. In particular, a given subset could be measurable with respect to some  $\sigma$ -algebra and not measurable with respect to some other  $\sigma$ -algebra.

Example 5.4. The pair  $(M, \mathcal{P}(M))$  is a measurable space for any set  $M$ . Of course, just like the discrete topology is not a very useful topology, the power set  $\mathcal{P}(M)$  is not a very useful  $\sigma$ -algebra on  $M$ , unless  $M$  is countable.

**Definition.** The *extended real line* is  $\overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, +\infty\}$ , where the symbols  $-\infty$  and  $+\infty$  (the latter often denoted simply by  $\infty$ ) satisfy

$$\forall r \in \overline{\mathbb{R}} : -\infty \leq r \leq \infty$$

with strict inequalities if  $r \in \mathbb{R}$ . The symbols  $\pm\infty$  satisfy the following arithmetic rules

$$(i) \quad \forall r \in \mathbb{R} : \pm\infty + r = r \pm \infty = \pm\infty$$

$$(ii) \quad \forall r > 0 : r(\pm\infty) = \pm\infty r = \pm\infty$$

---

<sup>9</sup>[https://en.wikipedia.org/wiki/Non-measurable\\_set](https://en.wikipedia.org/wiki/Non-measurable_set)

(iii)  $\forall r < 0 : r(\pm\infty) = \pm\infty r = \mp\infty$

(iv)  $0(\pm\infty) = \pm\infty 0 = 0$ .

Note that expressions such as  $\infty - \infty$  or  $-\infty + \infty$  are not defined.  $\diamond$

**Definition.** Let  $(M, \Sigma)$  be a measurable space. A *measure* on  $(M, \Sigma)$  is a function

$$\mu: \Sigma \rightarrow [0, \infty],$$

where  $[0, \infty] := \{r \in \overline{\mathbb{R}} \mid r \geq 0\}$ , such that

(i)  $\mu(\emptyset) = 0$

(ii) for any sequence  $\{A_n\}_{n \in \mathbb{N}}$  in  $\Sigma$  with  $A_i \cap A_j = \emptyset$  whenever  $i \neq j$ , we have

$$\mu\left(\bigcup_{n=0}^{\infty} A_n\right) = \sum_{n=0}^{\infty} \mu(A_n).$$

$\diamond$

A sequence  $\{A_n\}_{n \in \mathbb{N}}$  that satisfies the condition that  $A_i \cap A_j = \emptyset$  for all  $i \neq j$  is called a *pairwise disjoint* sequence.

Remark 5.5. Both sides of the equation in part (ii) of the definition of measure might take the value  $\infty$ . There are two possible reasons why  $\sum_{n=0}^{\infty} \mu(A_n)$  might be infinite. It could be that  $\mu(A_n) = \infty$  for some  $n \in \mathbb{N}$  or, alternatively, it could be that  $\mu(A_n) < \infty$  for all  $n \in \mathbb{N}$  but the sequence of partial sums  $\{\sum_{i=0}^n \mu(A_i)\}_{n \in \mathbb{N}}$ , which is an increasing sequence since  $\mu$  is non-negative by definition, is not bounded above.

**Definition.** A *measure space* is a triple  $(M, \Sigma, \mu)$  where  $(M, \Sigma)$  is a measurable space and  $\mu: \Sigma \rightarrow [0, \infty]$  is a measure on  $M$ .  $\diamond$

Example 5.6. Let  $M = \mathbb{N}$  and  $\Sigma = \mathcal{P}(\mathbb{N})$ . Define  $\mu: \Sigma \rightarrow [0, \infty]$  by:

$$\mu(A) = |A| := \begin{cases} n & \text{if } A \text{ is a finite set with } n \text{ elements} \\ \infty & \text{if } A \text{ is not a finite set} \end{cases}$$

Then, by definition,  $\mu(\emptyset) = 0$ . Moreover, if  $\{A_n\}_{n \in \mathbb{N}}$  is a pairwise disjoint sequence in  $\Sigma$  such that all but a finite number of  $A_n$ 's are empty and each  $A_n$  has a finite number of elements, we have:

$$\mu\left(\bigcup_{n=0}^{\infty} A_n\right) = \sum_{n=0}^{\infty} \mu(A_n)$$

by counting elements. Otherwise, that is, if an infinite number of  $A_n$ 's are non-empty or if at least one  $A_n$  is infinite, then

$$\mu\left(\bigcup_{n=0}^{\infty} A_n\right) = \infty = \sum_{n=0}^{\infty} \mu(A_n)$$

and thus, the triple  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), \mu)$  is a measure space. The measure  $\mu$  on  $(\mathbb{N}, \mathcal{P}(\mathbb{N}))$  is called *counting measure* and it is the usual measure on countable measurable spaces.



**Proposition 5.7.** *Let  $(M, \Sigma, \mu)$  be a measure space.*

(i) *If  $A_0, \dots, A_k \in \Sigma$  and  $A_i \cap A_j = \emptyset$  for all  $0 \leq i \neq j \leq k$ , then:*

$$\mu\left(\bigcup_{n=0}^k A_n\right) = \sum_{n=0}^k \mu(A_n)$$

(ii) *If  $A, B \in \Sigma$  and  $A \subseteq B$ , then  $\mu(A) \leq \mu(B)$*

(iii) *If  $A, B \in \Sigma$ ,  $A \subseteq B$  and  $\mu(A) < \infty$ , then  $\mu(B \setminus A) = \mu(B) - \mu(A)$ .*

*Proof.* (i) Let  $A_n = \emptyset$  for all  $n > k$ . Then,  $\{A_n\}_{n \in \mathbb{N}}$  is a pairwise disjoint sequence in  $\Sigma$  and hence, we have:

$$\mu\left(\bigcup_{n=0}^k A_n\right) = \mu\left(\bigcup_{n=0}^{\infty} A_n\right) = \sum_{n=0}^{\infty} \mu(A_n) = \sum_{n=0}^k \mu(A_n).$$

(ii) We have  $B = A \cup (B \setminus A)$  and  $A \cap (B \setminus A) = \emptyset$ . Hence, by part (i),

$$\mu(B) = \mu(A \cup (B \setminus A)) = \mu(A) + \mu(B \setminus A),$$

and since  $\mu(B \setminus A) \geq 0$ , we have  $\mu(A) \leq \mu(B)$ .

(iii) By decomposing  $B$  as above and rearranging, we immediately get

$$\mu(B \setminus A) = \mu(B) - \mu(A).$$

Note, however, that this only makes sense if  $\mu(A) < \infty$ , for otherwise we must also have  $\mu(B) = \infty$  by part (ii), and then  $\mu(B) - \mu(A)$  would not be defined.  $\square$

**Proposition 5.8.** *Let  $(M, \Sigma, \mu)$  be a measure space and let  $\{A_n\}_{n \in \mathbb{N}}$  be a sequence in  $\Sigma$ .*

(i) *If  $\{A_n\}_{n \in \mathbb{N}}$  is increasing, i.e.  $A_n \subseteq A_{n+1}$  for all  $n \in \mathbb{N}$ , then*

$$\mu\left(\bigcup_{n=0}^{\infty} A_n\right) = \lim_{n \rightarrow \infty} \mu(A_n).$$

(ii) *If  $\mu(A_0) < \infty$  and  $\{A_n\}_{n \in \mathbb{N}}$  is decreasing, i.e.  $A_{n+1} \subseteq A_n$  for all  $n \in \mathbb{N}$ , then*

$$\mu\left(\bigcap_{n=0}^{\infty} A_n\right) = \lim_{n \rightarrow \infty} \mu(A_n).$$

We say that  $\mu$  is (i) *continuous from below* and (ii) *continuous from above*.

*Proof.* (i) Define  $B_0 := A_0$  and  $B_n := A_n \setminus A_{n-1}$ . Then,  $\{B_n\}_{n \in \mathbb{N}}$  is a pairwise disjoint sequence in  $\Sigma$  such that

$$\bigcup_{i=0}^n B_i = A_n \quad \text{and} \quad \bigcup_{n=0}^{\infty} A_n = \bigcup_{n=0}^{\infty} B_n.$$

Hence, we have

$$\begin{aligned}
\mu\left(\bigcup_{n=0}^{\infty} A_n\right) &= \mu\left(\bigcup_{n=0}^{\infty} B_n\right) \\
&= \sum_{n=0}^{\infty} \mu(B_n) \\
&= \lim_{n \rightarrow \infty} \sum_{i=0}^n \mu(B_i) \\
&= \lim_{n \rightarrow \infty} \mu\left(\bigcup_{i=0}^n B_i\right) \\
&= \lim_{n \rightarrow \infty} \mu(A_n).
\end{aligned}$$

(ii) Define  $B_n := A_0 \setminus A_n$ . Then,  $B_n \subseteq B_{n+1}$  for all  $n \in \mathbb{N}$  and thus, by part (i), we have

$$\begin{aligned}
\mu\left(\bigcup_{n=0}^{\infty} B_n\right) &= \lim_{n \rightarrow \infty} \mu(B_n) \\
&= \lim_{n \rightarrow \infty} (\mu(A_0) - \mu(A_n)) \\
&= \mu(A_0) - \lim_{n \rightarrow \infty} \mu(A_n).
\end{aligned}$$

Note that, by definition of  $B_n$ , we have

$$\bigcup_{n=0}^{\infty} B_n = A_0 \setminus \bigcap_{n=0}^{\infty} A_n.$$

Since  $\mu(A_0) < \infty$ , it follows from the previous proposition that

$$\begin{aligned}
\mu(A_0) - \mu\left(\bigcap_{n=0}^{\infty} A_n\right) &= \mu\left(A_0 \setminus \bigcap_{n=0}^{\infty} A_n\right) \\
&= \mu\left(\bigcup_{n=0}^{\infty} B_n\right) \\
&= \mu(A_0) - \lim_{n \rightarrow \infty} \mu(A_n).
\end{aligned}$$

Therefore, we have


$$\mu\left(\bigcap_{n=1}^{\infty} A_n\right) = \lim_{n \rightarrow \infty} \mu(A_n). \quad \square$$

*Remark 5.9.* Note that the result in the second part of this proposition need not be true if  $\mu(A_0) = \infty$ . For example, consider  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), \mu)$ , where  $\mu$  is the counting measure on  $(\mathbb{N}, \mathcal{P}(\mathbb{N}))$ . If  $A_n = \{n, n+1, n+2, \dots\}$ , then  $\{A_n\}_{n \in \mathbb{N}}$  is a decreasing sequence. Since  $\mu(A_n) = \infty$  for all  $n \in \mathbb{N}$ , we have  $\lim_{n \rightarrow \infty} \mu(A_n) = \infty$ . On the other hand,  $\bigcap_{n=0}^{\infty} A_n = \emptyset$  and thus  $\mu(\bigcap_{n=0}^{\infty} A_n) = 0$ .

**Proposition 5.10.** *Let  $(M, \Sigma, \mu)$  be a measure space. Then,  $\mu$  is countably sub-additive. That is, for any sequence  $\{A_n\}_{n \in \mathbb{N}}$  in  $\Sigma$ , we have*

$$\mu\left(\bigcup_{n=0}^{\infty} A_n\right) \leq \sum_{n=0}^{\infty} \mu(A_n).$$

*Proof.* (a) First, we show that  $\mu(A \cup B) \leq \mu(A) + \mu(B)$  for any  $A, B \in \Sigma$ . Note that, for any pair of sets  $A$  and  $B$ , the sets  $A \setminus B$ ,  $B \setminus A$  and  $A \cap B$  are pairwise disjoint and their union is  $A \cup B$ .



By writing  $A = (A \setminus B) \cup (A \cap B)$  and  $B = (B \setminus A) \cup (A \cap B)$  and using the additivity and positivity of  $\mu$ , we have

$$\begin{aligned} \mu(A) + \mu(B) &= \mu((A \setminus B) \cup (A \cap B)) + \mu((B \setminus A) \cup (A \cap B)) \\ &= \mu(A \setminus B) + 2\mu(A \cap B) + \mu(B \setminus A) \\ &= \mu(A \cup B) + \mu(A \cap B) \\ &\geq \mu(A \cup B). \end{aligned}$$

(b) We now extend this to finite unions by induction. Let  $\{A_n\}_{n \in \mathbb{N}}$  be a sequence in  $\Sigma$  and suppose that

$$\mu\left(\bigcup_{i=0}^n A_i\right) \leq \sum_{i=0}^n \mu(A_i)$$

for some  $n \in \mathbb{N}$ . Then, by part (a), we have

$$\begin{aligned} \mu\left(\bigcup_{i=0}^{n+1} A_i\right) &= \mu\left(A_{n+1} \cup \bigcup_{i=0}^n A_i\right) \\ &\leq \mu(A_{n+1}) + \mu\left(\bigcup_{i=0}^n A_i\right) \\ &\leq \mu(A_{n+1}) + \sum_{i=0}^n \mu(A_i) \\ &= \sum_{i=0}^{n+1} \mu(A_i). \end{aligned}$$

Hence, by induction on  $n$  with base case  $n = 1$  and noting that the case  $n = 0$  is trivial (it reduces to  $\mu(A_0) = \mu(A_0)$ ), we have

$$\forall n \in \mathbb{N} : \mu\left(\bigcup_{i=0}^n A_i\right) \leq \sum_{i=0}^n \mu(A_i).$$

- (c) Let  $\{A_n\}_{n \in \mathbb{N}}$  be a sequence in  $\Sigma$ . Define  $B_n := \bigcup_{i=0}^n A_i$ . Then,  $\{B_n\}_{n \in \mathbb{N}}$  is an increasing sequence in  $\Sigma$ . Hence, by continuity from above of  $\mu$ , we have

$$\begin{aligned} \mu\left(\bigcup_{n=0}^{\infty} A_n\right) &= \mu\left(\bigcup_{n=0}^{\infty} B_n\right) \\ &= \lim_{n \rightarrow \infty} \mu(B_n) \\ &= \lim_{n \rightarrow \infty} \mu\left(\bigcup_{i=0}^n A_i\right) \\ &\leq \lim_{n \rightarrow \infty} \sum_{i=0}^n \mu(A_i) \\ &= \sum_{i=0}^{\infty} \mu(A_i) \end{aligned}$$

which is what we wanted.  $\square$

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space. The measure  $\mu$  is said to be *finite* if there exists a sequence  $\{A_n\}_{n \in \mathbb{N}}$  in  $\Sigma$  such that  $\bigcup_{n=0}^{\infty} A_n = M$  and

$$\forall n \in \mathbb{N} : \mu(A_n) < \infty.$$

$\diamond$

Example 5.11. The counting measure on  $(\mathbb{N}, \mathcal{P}(\mathbb{N}))$  is finite. To see this, define  $A_n := \{n\}$ . Then, clearly  $\bigcup_{n=0}^{\infty} A_n = \mathbb{N}$  and  $\mu(A_n) = |\{n\}| = 1 < \infty$  for all  $n \in \mathbb{N}$ .

## 5.2 Borel $\sigma$ -algebras

We have already remarked the parallel between topologies and  $\sigma$ -algebras. A further similarity stems from the fact that, just like for topologies, interesting  $\sigma$ -algebras are hardly ever given explicitly, except in some simple cases. In general, they are defined implicitly by some membership condition.

**Proposition 5.12.** Let  $M$  be a set and let  $\{\Sigma_i : i \in I\}$  be a collection of  $\sigma$ -algebras on  $M$ . Define the set

$$\Sigma := \bigcap_{i \in I} \Sigma_i = \{A \in \mathcal{P}(M) \mid A \in \Sigma_i, \forall i \in I\}.$$

Then,  $\Sigma$  is a  $\sigma$ -algebra on  $M$ .

*Proof.* We simply check that  $\Sigma$  satisfies the defining properties of a  $\sigma$ -algebra.

- (i) We have  $M \in \Sigma_i$  for all  $i \in I$  and hence  $M \in \Sigma$ .
- (ii) Let  $A \in \Sigma$ . Then,  $A \in \Sigma_i$  for all  $i \in I$  and, since each  $\Sigma_i$  is a  $\sigma$ -algebra, we also have  $M \setminus A \in \Sigma_i$  for all  $i \in I$ . Hence,  $M \setminus A \in \Sigma$ .

(iii) Let  $\{A_n\}_{n \in \mathbb{N}}$  be a sequence in  $\Sigma$ . Then,  $\{A_n\}_{n \in \mathbb{N}}$  is a sequence in each  $\Sigma_i$ . Thus,

$$\forall i \in I : \bigcup_{n=0}^{\infty} A_n \in \Sigma_i.$$

Hence, we also have  $\bigcup_{n=0}^{\infty} A_n \in \Sigma$ .  $\square$

**Definition.** Let  $M$  be a set and let  $\mathcal{E} \subseteq \mathcal{P}(M)$  be a collection of subsets of  $M$ . The  $\sigma$ -algebra *generated* by  $\mathcal{E}$ , denoted  $\sigma(\mathcal{E})$ , is the smallest  $\sigma$ -algebra on  $M$  containing all the sets in  $\mathcal{E}$ . That is,

$$A \in \sigma(\mathcal{E}) \iff \text{for all } \sigma\text{-algebras } \Sigma \text{ on } M : \mathcal{E} \subseteq \Sigma \Rightarrow A \in \Sigma$$

or, by letting  $\{\Sigma_i \mid i \in I\}$  be the collection of  $\sigma$ -algebras on  $M$  such that  $\mathcal{E} \subseteq \Sigma$ ,

$$\sigma(\mathcal{E}) := \bigcap_{i \in I} \Sigma_i.$$

$\diamond$

The set  $\mathcal{E}$  is called a *generating set* for  $\sigma(\mathcal{E})$ . Observe that the second characterisation makes it manifest that  $\sigma(\mathcal{E})$  is indeed a  $\sigma$ -algebra on  $M$  by the previous proposition.

**Theorem 5.13.** Let  $(M, \Sigma)$  be a measurable space. Then,  $\Sigma = \sigma(\mathcal{E})$  for some  $\mathcal{E} \subseteq \mathcal{P}(M)$ .

This generating construction immediately allows us to link the notions of topology and  $\sigma$ -algebra on a set  $M$  via the following definition.

**Definition.** Let  $(M, \mathcal{O})$  be a topological space. The *Borel  $\sigma$ -algebra* on  $(M, \mathcal{O})$  is  $\sigma(\mathcal{O})$ .  $\diamond$

Recall that a topology on  $M$  is a collection  $\mathcal{O} \subseteq \mathcal{P}(M)$  of subsets of  $M$  which contains  $\emptyset$  and  $M$  and is closed under finite intersections and arbitrary (even uncountable) unions. The elements of the topology are called *open sets*. Of course, while there many choices of  $\sigma$ -algebra on  $M$ , if we already have a topology  $\mathcal{O}$  on  $M$ , then the associated Borel  $\sigma$ -algebra is very convenient choice of  $\sigma$ -algebra since, as we will soon see, it induces a measurable structure which is “compatible” with the already given topological structure.

This is, in fact, the usual philosophy in mathematics: we always let the stronger structures induce the weaker ones, unless otherwise specified. For instance, once we have chosen an inner product on a space, we take the norm to be the induced norm, which induces a metric, which in turn induces a topology on that space, from which we now know how to obtain a canonical  $\sigma$ -algebra.

We remark that, while the Borel  $\sigma$ -algebra on a topological space is generated by the open sets, in general, it contains much more than just the open sets.

Example 5.14. Recall that the standard topology on  $\mathbb{R}$ , denoted  $\mathcal{O}_{\mathbb{R}}$ , is defined by

$$A \in \mathcal{O}_{\mathbb{R}} \iff \forall a \in A : \exists \varepsilon > 0 : \forall r \in \mathbb{R} : |r - a| < \varepsilon \Rightarrow r \in A.$$

In fact, the elements of  $\mathcal{O}_{\mathbb{R}}$  are at most countable unions of open intervals in  $\mathbb{R}$ . Consider now the Borel  $\sigma$ -algebra on  $(\mathbb{R}, \mathcal{O}_{\mathbb{R}})$ . Let  $a < b$ . Then, for any  $n \in \mathbb{N}$ , the interval  $(a - \frac{1}{n}, b)$

is open. Hence,  $\{(a - \frac{1}{n}, b)\}_{n \in \mathbb{N}}$  is a sequence in  $\sigma(\mathcal{O}_{\mathbb{R}})$ . Since  $\sigma$ -algebras are closed under countable intersections, we have

$$\bigcap_{n=0}^{\infty} (a - \frac{1}{n}, b) = [a, b) \in \sigma(\mathcal{O}_{\mathbb{R}}).$$

Hence,  $\sigma(\mathcal{O}_{\mathbb{R}})$  contains, in addition to all open intervals, also all half-open intervals. It is not difficult to show that it contains all closed intervals as well. In particular, since singletons are closed,  $\sigma(\mathcal{O}_{\mathbb{R}})$  also contains all countable subsets of  $\mathbb{R}$ . In fact, it is non-trivial<sup>10</sup> to produce a subset of  $\mathbb{R}$  which is not contained in  $\sigma(\mathcal{O}_{\mathbb{R}})$ .

### 5.3 Lebesgue measure on $\mathbb{R}^d$

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space. If  $A \in \Sigma$  is such that  $\mu(A) = 0$ , then  $A$  is called a *null set* or a *set of measure zero*.  $\diamond$

The following definition is not needed for the construction of the Lebesgue measure. However, since it is closely connected with that of null set and will be used a lot in the future, we chose to present it here.

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space and let  $P$  be some property or statement. We say that  $P$  holds *almost everywhere* on  $M$  if

$$\exists Z \in \Sigma : \mu(Z) = 0 \text{ and } \forall m \in M \setminus Z : P(m).$$

$\diamond$

In other words, the property  $P$  is said to hold almost everywhere on  $M$  if it holds everywhere on  $M$  except for a null subset of  $M$ .

Example 5.15. Let  $(M, \Sigma, \mu)$  be a measure space and let  $f, g : M \rightarrow N$  be maps. We say that  $f$  and  $g$  are *almost everywhere equal*, and we write  $f =_{\text{a.e.}} g$ , if there exists a null set  $Z \in \Sigma$  such that

$$\forall m \in M \setminus Z : f(m) = g(m).$$

The case  $f = g$  corresponds to  $Z = \emptyset$ .

**Definition.** A measure  $\mu : \Sigma \rightarrow [0, \infty]$  is said to be *complete* if every subset of every null set is measurable, i.e.

$$\forall A \in \Sigma : \forall B \in \mathcal{P}(A) : \mu(A) = 0 \Rightarrow B \in \Sigma.$$

$\diamond$

Note that since for any  $A, B \in \Sigma$ ,  $B \subseteq A$  implies  $\mu(B) \leq \mu(A)$ , it follows that every subset of a null set, if measurable, must also be a null set.

<sup>10</sup>[https://en.wikipedia.org/wiki/Borel\\_set#Non-Borel\\_sets](https://en.wikipedia.org/wiki/Borel_set#Non-Borel_sets)

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space and let  $(M, +, \cdot)$  be a vector space. The measure  $\mu$  is said to be *translation-invariant* if

$$\forall m \in M : \forall A \in \Sigma : A + m \in \Sigma \text{ and } \mu(A + m) = \mu(A),$$

where  $A + m := \{a + m \mid a \in A\}$ . ◇

**Theorem 5.16.** Let  $\mathcal{O}_{\mathbb{R}^d}$  be the standard topology on  $\mathbb{R}^d$ . There exists a unique complete, translation-invariant measure

$$\lambda^d : \sigma(\mathcal{O}_{\mathbb{R}^d}) \rightarrow [0, \infty]$$

such that for all  $a_i, b_i \in \mathbb{R}$  with  $1 \leq i \leq d$  and  $a_i < b_i$ , we have

$$\lambda^d([a_1, b_1) \times \cdots \times [a_d, b_d)) = \prod_{i=1}^d (b_i - a_i).$$

**Definition.** The measure  $\lambda^d$  is called the *Lebesgue measure* on  $\mathbb{R}^d$ . ◇


The superscript  $d$  in  $\lambda^d$  may be suppressed if there is no risk of confusion. Note that the Lebesgue measure on  $\mathbb{R}$ ,  $\mathbb{R}^2$  and  $\mathbb{R}^3$  coincides with the standard notions of length, area and volume, with the further insight that these are only defined for the elements of the respective Borel  $\sigma$ -algebras.

**Proposition 5.17.** The Lebesgue measure on  $\mathbb{R}$  is finite.

*Proof.* Consider the sequence  $\{[a_n, a_n + 1)\}_{n \in \mathbb{N}}$  where

$$a_n = \begin{cases} -\frac{1}{2}n & \text{if } n \text{ is even} \\ \frac{1}{2}(n+1) & \text{if } n \text{ is odd.} \end{cases}$$

That is,  $\{a_n\}_{n \in \mathbb{N}}$  is the sequence  $(0, 1, -1, 2, -2, 3, -3, \dots)$ .



Clearly, we have  $\bigcup_{n=0}^{\infty} [a_n, a_n + 1) = \mathbb{R}$ . Since, for all  $n \in \mathbb{N}$ ,  $[a_n, a_n + 1) \in \sigma(\mathcal{O}_{\mathbb{R}})$  and  $\lambda([a_n, a_n + 1)) = 1 < \infty$ , the Lebesgue measure  $\lambda$  is finite. □

This can easily be generalised to show that  $\lambda^d$  is finite for all  $d \geq 1$ .

## 5.4 Measurable maps

As we have remarked earlier, once we introduce a new structure, we should immediately think about the associated structure-preserving maps. In the case of measurable spaces, a measurable map is one that preserves the “measurability” structure.

**Definition.** Let  $(M, \Sigma_M)$  and  $(N, \Sigma_N)$  be measurable spaces. A map  $f: M \rightarrow N$  is said to be *measurable* if

$$\forall A \in \Sigma_N : \text{preim}_f(A) \in \Sigma_M.$$

◇

Note that this is exactly the definition of continuous map between topological spaces, with “continuous” replaced by “measurable” and topologies replaced by  $\sigma$ -algebras.

**Lemma 5.18.** *Let  $(M, \Sigma_M)$  and  $(N, \Sigma_N)$  be measurable spaces. A map  $f: M \rightarrow N$  is measurable if, and only if,*

$$\forall A \in \mathcal{E} : \text{preim}_f(A) \in \Sigma_M,$$

where  $\mathcal{E} \subseteq \mathcal{P}(N)$  is a generating set of  $\Sigma_N$ .

**Corollary 5.19.** *Let  $(M, \mathcal{O}_M)$  and  $(N, \mathcal{O}_N)$  be topological spaces. Any continuous map  $M \rightarrow N$  is measurable with respect to the Borel  $\sigma$ -algebras on  $M$  and  $N$ .*

Recall that a map  $\mathbb{R} \rightarrow \mathbb{R}$  is monotonic if it is either increasing or decreasing.

**Corollary 5.20.** *Any monotonic map  $\mathbb{R} \rightarrow \mathbb{R}$  is measurable with respect to the Borel  $\sigma$ -algebra (with respect to  $\mathcal{O}_{\mathbb{R}}$ ).*

**Proposition 5.21.** *Let  $(M, \Sigma_M)$ ,  $(N, \Sigma_N)$  and  $(P, \Sigma_P)$  be measurable spaces. If  $f: M \rightarrow N$  and  $g: N \rightarrow P$  are both measurable, then so is their composition  $g \circ f: M \rightarrow P$ .*

*Proof.* Let  $A \in \Sigma_P$ . As  $g$  is measurable, we have  $\text{preim}_g(A) \in \Sigma_N$ . Then, since  $f$  is measurable, it follows that

$$\text{preim}_f(\text{preim}_g(A)) = \text{preim}_{g \circ f}(A) \in \Sigma_M.$$

Hence,  $g \circ f$  is measurable. □

**Proposition 5.22.** *Let  $(M, \Sigma_M)$  and  $(N, \Sigma_N)$  be measurable spaces and let  $\{f_n\}_{n \in \mathbb{N}}$  be a sequence of measurable maps from  $M$  to  $N$  whose pointwise limit is  $f$ . Then,  $f$  is measurable.*

Recall that  $\{f_n\}_{n \in \mathbb{N}}$  converges pointwise to  $f: M \rightarrow N$  if

$$\forall m \in M : \lim_{n \rightarrow \infty} f_n(m) = f(m).$$

This is in contrast with continuity, as pointwise convergence of a sequence of continuous maps is not a sufficient condition for the continuity of the pointwise limit. In the case of real or complex-valued maps, a sufficient condition is convergence with respect to the supremum norm.



## 5.5 Push-forward of a measure

If we have a structure-preserving map  $f$  between two instances  $A$  and  $B$  of some structure, and an object on  $A$  (which depends in some way on the structure), we can often use  $f$  to induce a similar object on  $B$ . This is generically called the push-forward of that object along the map  $f$ .

**Proposition 5.23.** *Let  $(M, \Sigma_M, \mu)$  be a measure space, let  $(N, \Sigma_N)$  be a measurable space and let  $f: M \rightarrow N$  be a measurable map. Then, the map*

$$\begin{aligned} f_*\mu: \Sigma_N &\rightarrow [0, \infty] \\ A &\mapsto \mu(\text{preim}_f(A)) \end{aligned}$$

*is a measure on  $(N, \Sigma_N)$  called the push-forward of  $\mu$  along  $f$ .*

That  $f_*\mu$  is a measure follows easily from the fact that  $\mu$  is a measure and basic properties of pre-images of maps, namely

$$\begin{aligned} \text{preim}_f(A \setminus B) &= \text{preim}_f(A) \setminus \text{preim}_f(B) \\ \text{preim}_f\left(\bigcup_{i \in I} A_i\right) &= \bigcup_{i \in I} \text{preim}_f(A_i). \end{aligned}$$

## 6 Integration of measurable functions

We will now focus on measurable functions  $M \rightarrow \overline{\mathbb{R}}$  and define their integral on a subset of  $M$  with respect to some measure on  $M$ , which is called the Lebesgue integral. Note that, even if  $M \subseteq \mathbb{R}^d$ , the Lebesgue integral of a function need not be with respect to the Lebesgue measure.

The key application of this material is the definition of the Banach spaces of (classes of) Lebesgue integrable functions  $L^p$ . The case  $p = 2$  is especially important since  $L^2$  is, in fact, a Hilbert space. It appears a lot in quantum mechanics where it is loosely referred to as the space of square-integrable functions.


### 6.1 Characteristic and simple functions

**Definition.** Let  $M$  be a set and let  $A \in \mathcal{P}(M)$ . The *characteristic function* of  $A$ , denoted  $\chi_A: M \rightarrow \mathbb{R}$ , is defined by

$$\chi_A(m) = \begin{cases} 1 & \text{if } m \in A \\ 0 & \text{if } m \notin A. \end{cases}$$

◇

Example 6.1. Below is the graph of  $\chi_{(1,2]}: \mathbb{R} \rightarrow \mathbb{R}$ .



The following properties of characteristic functions are immediate.

**Proposition 6.2.** Let  $M$  be a set and let  $A, B \in \mathcal{P}(M)$ . Then

- (i)  $\chi_\emptyset = 0$
- (ii)  $\chi_{A \cup B} = \chi_A + \chi_B - \chi_{A \cap B}$
- (iii)  $\chi_{A \cap B} = \chi_A \chi_B$
- (iv)  $\chi_{M \setminus A} + \chi_A = 1$

where the addition and multiplication are pointwise and the 0 and 1 in parts (i) and (iv) are the constant functions  $M \rightarrow \mathbb{R}$  mapping every  $m \in M$  to  $0 \in \mathbb{R}$  and  $1 \in \mathbb{R}$ , respectively.

**Definition.** Let  $M$  be a set. A function  $s: M \rightarrow \mathbb{R}$  is *simple* if  $s(M) = \{r_1, \dots, r_n\}$  for some  $n \in \mathbb{N}$ .


◇

Equivalently,  $s: M \rightarrow \mathbb{R}$  is simple if there exist  $r_1, \dots, r_n \in \mathbb{R}$  and  $A_1, \dots, A_n \in \mathcal{P}(M)$ , for some  $n \in \mathbb{N}$ , such that

$$s = \sum_{i=1}^n r_i \chi_{A_i}.$$

So  $s$  is simple if it is a linear combination of characteristic functions.

Example 6.3. Consider the simple function  $s: \mathbb{R} \rightarrow \mathbb{R}$  given by  $s := \chi_{[1,3]} + 2\chi_{[2,5]}$ .



By observing the graph, we see that we can re-write  $s$  as

$$s = \chi_{[1,2)} + 3\chi_{[2,3)} + 2\chi_{[3,5]}.$$

**Definition.** A simple function is said to be in its *standard form* if

$$s = \sum_{i=1}^n r_i \chi_{A_i},$$

where  $A_i \cap A_j = \emptyset$  whenever  $i \neq j$ . ◇

Any simple function can be written in standard form. It is clear that if  $s$  is in its standard form, then  $A_i = \text{preim}_s(\{r_i\})$ .

**Proposition 6.4.** Let  $(M, \Sigma)$  be a measurable space and let  $A, A_1, \dots, A_n \in \mathcal{P}(M)$ . Then

- (i)  $\chi_A$  is measurable if, and only if,  $A \in \Sigma$
- (ii) if  $s = \sum_{i=1}^n r_i \chi_{A_i}$  is a simple function in its standard form, then  $s$  is measurable if, and only if, we have  $A_i \in \Sigma$  for all  $1 \leq i \leq n$ .

*Proof.* (i) We have  $\chi_A: M \rightarrow \mathbb{R}$  with possible values  $\{0, 1\}$ . Equipping  $\mathbb{R}$  with its Borel  $\sigma$ -algebra w.r.t. the standard topology,  $\sigma(\mathcal{O}_{\mathbb{R}})$ , we have:

(a) ( $\Rightarrow$ ) We have

$$A = M \setminus \text{preim}_{\chi_A}([0, 1)),$$

but  $[0, 1) \in \sigma(\mathcal{O}_{\mathbb{R}})$  and so  $\text{preim}_{\chi_A}([0, 1)) \in \Sigma$ . Then by property (ii) of a  $\sigma$ -algebra  $E \in \Sigma$ .

- (b) ( $\Leftarrow$ ) Let  $\alpha = [1, \infty)$ ,  $\beta = (-\infty, 0)$  and  $\gamma = [0, 1]$ . Clearly  $\alpha, \beta, \gamma \in \sigma(\mathcal{O}_{\mathbb{R}})$  and  $\alpha \cup \beta \cup \gamma = \mathbb{R}$ . Then

$$\begin{aligned}\text{preim}_{\chi_A}(\alpha) &= A, \\ \text{preim}_{\chi_A}(\beta) &= \emptyset, \\ \text{preim}_{\chi_A}(\gamma) &= M,\end{aligned}$$

all of which are measurable sets on  $M$  and so  $\chi_A$  is measurable.

- (ii) First define  $\tilde{\chi}_{A_i} := r_i \chi_{A_i}$ , which satisfies

$$\tilde{\chi}_{A_i}(m) = \begin{cases} r_i & \text{if } m \in A_i \\ 0 & \text{if } m \notin A_i. \end{cases}$$

As  $s$  is in its standard form we know  $\tilde{\chi}_{A_i} \cap \tilde{\chi}_{A_j} = \emptyset$  for all  $i \neq j$ . Combining these two things, and defining  $\alpha_i = [r_i, \infty)$ ,  $\gamma_i = [0, r_i]$ , the result follow from part (i).  $\square$

## 6.2 Integration of non-negative measurable simple functions

We begin with the definition of the integral of a non-negative, measurable, simple function.

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space and let  $s: M \rightarrow \mathbb{R}$  be a nowhere negative, measurable, simple function whose standard form is  $s = \sum_{i=1}^n r_i \chi_{A_i}$ . Then, we define

$$\int_M s \, d\mu := \sum_{i=1}^n r_i \mu(A_i).$$

$\diamond$

Note that the non-negativity condition is essential since  $\mu$  takes values in  $[0, \infty]$ , hence we could have  $\mu(A_i) = \infty$  for more than one  $A_i$ , and if the corresponding coefficients  $r_i$  have opposite signs, then we would have  $\int_M s \, d\mu = \infty - \infty$ , which is not defined. For the same reason, we are considering  $s: M \rightarrow [0, \infty)$  rather than  $s: M \rightarrow [0, \infty]$ .

*Example 6.5.* Consider the measure space  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), \mu)$ , where  $\mu$  is the counting measure, let  $f: \mathbb{N} \rightarrow \mathbb{R}$  be non-negative and suppose that there exists  $N \in \mathbb{N}$  such that  $f(n) = 0$  for all  $n > N$ . Then, we can write  $f$  as

$$f = \sum_{n=0}^N f(n) \chi_{\{n\}}.$$

That is,  $f$  is a non-negative, measurable, simple function and therefore

$$\int_{\mathbb{N}} f \, d\mu := \sum_{n=0}^N f(n) \mu(\{n\}) = \sum_{n=0}^N f(n).$$

Hence, the “integral” of  $f$  over  $\mathbb{N}$  with respect to the counting measure is just the sum.

The need for simple functions to be in their standard form, which was introduced to avoid any potential ambiguity in the definition of their integral, can be relaxed using the following lemma.

**Lemma 6.6.** *Let  $(M, \Sigma, \mu)$  be a measure space. Let  $s$  and  $t$  be non-negative, measurable, simple functions  $M \rightarrow \mathbb{R}$  and let  $c \in [0, \infty)$ . Then*

$$\int_M (cs + t) d\mu = c \int_M s d\mu + \int_M t d\mu.$$

**Proposition 6.7.** *Let  $(M, \Sigma, \mu)$  be a measure space and let  $s = \sum_{i=1}^n r_i \chi_{A_i}$  be a non-negative, measurable, simple function  $M \rightarrow \mathbb{R}$  not necessarily in its standard form. Then*

$$\int_M s d\mu = \sum_{i=1}^n r_i \mu(A_i).$$

**Corollary 6.8.** *Let  $(M, \Sigma, \mu)$  be a measure space. Let  $s$  and  $t$  be non-negative, measurable, simple functions  $M \rightarrow \mathbb{R}$  such that  $s \leq t$  (that is,  $s(m) \leq t(m)$  for all  $m \in M$ ). Then*

$$\int_M s d\mu \leq \int_M t d\mu.$$

**Lemma 6.9.** *Let  $(M, \Sigma, \mu)$  be a measure space and let  $s = \sum_{i=1}^n r_i \chi_{A_i}$  be a non-negative, measurable, simple function  $M \rightarrow \mathbb{R}$ . Define the map*

$$\begin{aligned} \nu_s: \Sigma &\rightarrow [0, \infty] \\ A &\mapsto \int_M s \chi_A d\mu, \end{aligned}$$

where  $s \chi_A$  is the pointwise product of  $s$  and  $\chi_A$ . Then,  $\nu_s$  is a measure on  $(M, \Sigma)$ .

*Proof.* First, note that we have

$$\int_M s \chi_A d\mu = \int_M \left( \sum_{i=1}^n r_i \chi_{A_i} \chi_A \right) d\mu = \int_M \left( \sum_{i=1}^n r_i \chi_{A_i \cap A} \right) d\mu = \sum_{i=1}^n r_i \mu(A_i \cap A).$$

We now check that  $\nu$  satisfies the defining properties of a measure.

$$(i) \quad \nu_s(\emptyset) := \int_M s \chi_{\emptyset} d\mu = \sum_{i=1}^n r_i \mu(A_i \cap \emptyset) = \sum_{i=1}^n r_i \mu(\emptyset) = 0$$

(ii) Let  $\{B_j\}_{j \in \mathbb{N}}$  be a pairwise disjoint sequence in  $\Sigma$ . Then

$$\begin{aligned}
\nu_s\left(\bigcup_{j=0}^{\infty} B_j\right) &= \int_M s \chi_{(\bigcup_{j=0}^{\infty} B_j)} d\mu \\
&= \sum_{i=1}^n r_i \mu\left(\bigcup_{j=0}^{\infty} (A_i \cap B_j)\right) \\
&= \sum_{j=0}^{\infty} \sum_{i=1}^n r_i \mu(A_i \cap B_j) \\
&= \sum_{j=0}^{\infty} \int_M s \chi_{B_j} d\mu \\
&= \sum_{j=0}^{\infty} \nu_s(B_j).
\end{aligned}$$

Thus,  $\nu_s$  is a measure on  $(M, \Sigma)$ . □

### 6.3 Integration of non-negative measurable functions

As we are interested in measurable functions  $M \rightarrow \overline{\mathbb{R}}$ , we need to define a  $\sigma$ -algebra on  $\overline{\mathbb{R}}$ . We cannot use the Borel  $\sigma$ -algebra since we haven't even defined a topology on  $\overline{\mathbb{R}}$ . In fact, we can easily get a  $\sigma$ -algebra on  $\overline{\mathbb{R}}$  as follows.

**Proposition 6.10.** *The set  $\overline{\Sigma} := \{A \in \mathcal{P}(\overline{\mathbb{R}}) \mid A \cap \mathbb{R} \in \sigma(\mathcal{O}_{\mathbb{R}})\}$  is a  $\sigma$ -algebra on  $\overline{\mathbb{R}}$ .*

In other words, we can simply ignore the infinities in a subset of  $\overline{\mathbb{R}}$  and consider it to be measurable if  $A \setminus \{-\infty, +\infty\}$  is in the Borel  $\sigma$ -algebra of  $\mathbb{R}$ . We will always consider  $\overline{\mathbb{R}}$  to be equipped with this  $\sigma$ -algebra.

**Lemma 6.11.** *Let  $(M, \Sigma)$  be a measurable space and let  $f, g: M \rightarrow \overline{\mathbb{R}}$  be measurable. Then, the following functions are measurable.*

- (i)  $cf + g$ , for any  $c \in \mathbb{R}$
- (ii)  $|f|$  and  $f^2$
- (iii)  $fg$  (pointwise product) if  $f$  and  $g$  are nowhere infinite
- (iv)  $\max(f, g)$  (defined pointwise).

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space and let  $f: M \rightarrow \overline{\mathbb{R}}$  be a non-negative, measurable function. Denote by  $S$  the set of all non-negative, measurable, simple functions  $s: M \rightarrow \mathbb{R}$  such that  $s \leq f$ . Then, we define

$$\int_M f d\mu := \sup_{s \in S} \int_M s d\mu.$$

◇

*Remark 6.12.* It is often very convenient to introduce the notation

$$\int_M f(x) \mu(dx) \equiv \int_M f \, d\mu,$$

where  $x$  is a dummy variable and could be replaced by any other symbol. The reason why this is a convenient notation is that, while some functions have standard symbols but cannot be easily represented by an algebraic expression (e.g. characteristic functions), others are easily expressed in terms of an algebraic formula but do not have a standard name. For instance, it is much easier to just write

$$\int_{\mathbb{R}} x^2 \mu(dx)$$

than having to first denote the function  $\mathbb{R} \rightarrow \mathbb{R}$ ,  $x \mapsto x^2$  by a generic  $f$  or, say, the more specific  $\text{sq}_{\mathbb{R}}$ , and then write

$$\int_{\mathbb{R}} \text{sq}_{\mathbb{R}} \, d\mu.$$

In computer programming, this is akin to defining *anonymous functions*.

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space and let  $f: M \rightarrow \overline{\mathbb{R}}$  be a non-negative, measurable function. For any  $A \in \Sigma$  (that is, any measurable subset of  $M$ ), we define

$$\int_A f \, d\mu := \int_M f \chi_A \, d\mu.$$

◇

Note that the product  $f \chi_A$  is measurable by part (iii) of Lemma 6.11.

**Lemma 6.13.** Let  $(M, \Sigma, \mu)$  be a measure space, let  $f, g: M \rightarrow \overline{\mathbb{R}}$  be non-negative, measurable functions such that  $f \leq g$ , and let  $A, B \in \Sigma$  be such that  $A \subseteq B$ . Then

$$(i) \quad \int_M f \, d\mu \leq \int_M g \, d\mu$$

$$(ii) \quad \int_A f \, d\mu \leq \int_B f \, d\mu.$$

*Proof.* (i) Denote by  $S_f$  and  $S_g$  the sets of non-negative, measurable, simple functions that are less than or equal to  $f$  and  $g$ , respectively. As  $f \leq g$ , we have  $S_f \subseteq S_g$  and hence

$$\int_M f \, d\mu := \sup_{s \in S_f} \int_M s \, d\mu \leq \sup_{s \in S_g} \int_M s \, d\mu =: \int_M g \, d\mu.$$

(ii) Since  $A \subseteq B$ , for any  $m \in M$  we have

$$f(m) \chi_A(m) \leq f(m) \chi_B(m).$$


In fact, we have equality whenever  $m \in A$  or  $m \in M \setminus B$ , while for  $m \in B \setminus A$  the left hand side is zero and the right-hand side is non-negative. Hence,  $f \chi_A \leq f \chi_B$  and thus, by part (i), we have

$$\int_A f \, d\mu := \int_M f \chi_A \, d\mu \leq \int_M f \chi_B \, d\mu =: \int_B f \, d\mu \quad \square$$

**Proposition 6.14** (Markov inequality). *Let  $(M, \Sigma, \mu)$  be a measure space and let  $f: M \rightarrow \overline{\mathbb{R}}$  be a non-negative, measurable function. For any  $z \in [0, \infty]$ , we have*

$$\int_M f \, d\mu \geq z \mu(\text{preim}_f([z, \infty])).$$

Equality is achieved whenever  $z$  is an upper bound for  $f$ .



The following is the pivotal theorem of Lebesgue integration.

**Theorem 6.15** (Monotone convergence theorem). *Let  $(M, \Sigma, \mu)$  be a measure space and let  $\{f_n\}_{n \in \mathbb{N}}$  be a sequence of non-negative, measurable functions  $M \rightarrow \overline{\mathbb{R}}$  such that  $f_{n+1} \geq f_n$  for all  $n \in \mathbb{N}$ . If there exists a function  $f: M \rightarrow \overline{\mathbb{R}}$  such that*

$$\forall m \in M : \lim_{n \rightarrow \infty} f_n(m) = f(m)$$

(i.e.  $f$  is the pointwise limit of  $\{f_n\}_{n \in \mathbb{N}}$ ), then  $f$  is measurable and

$$\lim_{n \rightarrow \infty} \int_M f_n \, d\mu = \int_M f \, d\mu.$$

*Remark 6.16.* Observe that this result is in stark contrast with what one may be used from Riemann integration, where pointwise converge of a sequence of integrable functions  $\{f_n\}_{n \in \mathbb{N}}$  is *not* a sufficient condition for the integral of the limit  $f$  to be equal to the limit of the integrals of  $f_n$  or, in fact, even for  $f$  to be integrable. For these, we need stronger conditions on the sequence  $\{f_n\}_{n \in \mathbb{N}}$ , such as uniform converge.

The definition of the integral as a supremum is clear and geometrically reasonable. However, it is in general very difficult to evaluate the integral of any particular function using it. The monotone convergence theorem provides a much simpler way to evaluate the integral. One can show that, for any non-negative, measurable function  $f$ , there exists an increasing sequence  $\{s_n\}_{n \in \mathbb{N}}$  of non-negative, measurable, simple functions (which can be explicitly constructed from  $f$ ) whose pointwise limit is  $f$ , and hence we have

$$\int_M f \, d\mu = \lim_{n \rightarrow \infty} \int_M s_n \, d\mu,$$

where the right-hand side can usually be evaluated fairly easily.



*Example 6.17.* Consider the measure space  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), \mu)$ , where  $\mu$  is the counting measure, and let  $f: \mathbb{N} \rightarrow \overline{\mathbb{R}}$  be non-negative. Note that the choice of  $\sigma$ -algebra  $\mathcal{P}(\mathbb{N})$  on  $\mathbb{N}$  makes every function on  $\mathbb{N}$  (to any measurable space) measurable. Define, for every  $n \in \mathbb{N}$ ,

$$s_n = \sum_{i=0}^n f(i) \chi_{\{i\}}.$$

Then,  $\{s_n\}_{n \in \mathbb{N}}$  is an increasing sequence of non-negative, measurable, simple functions whose pointwise limit is  $f$  and therefore, by the monotone convergence theorem,

$$\int_{\mathbb{N}} f \, d\mu = \lim_{n \rightarrow \infty} \int_{\mathbb{N}} s_n \, d\mu = \lim_{n \rightarrow \infty} \sum_{i=0}^n f(i) \mu(\{i\}) = \sum_{i=0}^{\infty} f(i).$$

If you ever wondered why series seem to share so many properties with integrals, the reason is that series are just integrals with respect to a discrete measure.

The monotone convergence theorem can be used to extend some of the properties of integrals of non-negative, measurable simple functions to non-negative, measurable functions which are not-necessarily simple.

**Lemma 6.18.** *Let  $(M, \Sigma, \mu)$  be a measure space, let  $f, g: M \rightarrow \overline{\mathbb{R}}$  be non-negative, measurable functions and let  $c \in [0, \infty)$ . Then*

$$(i) \quad \int_M (cf + g) \, d\mu = c \int_M f \, d\mu + \int_M g \, d\mu$$

$$(ii) \quad \text{the map } \nu_f: \Sigma \rightarrow [0, \infty] \text{ defined by } \nu_f(A) := \int_A f \, d\mu \text{ is a measure on } (M, \Sigma)$$

$$(iii) \quad \text{for any } A \in \Sigma, \text{ we have } \int_M f \, d\mu = \int_A f \, d\mu + \int_{M \setminus A} f \, d\mu.$$

*Proof.* (i) Let  $\{s_n\}_{n \in \mathbb{N}}$  and  $\{t_n\}_{n \in \mathbb{N}}$  be increasing sequences of non-negative, measurable, simple functions whose pointwise limits are  $f$  and  $g$ , respectively. Then, it is easy to see that  $\{cs_n + t_n\}_{n \in \mathbb{N}}$  is an increasing sequence of non-negative, measurable, simple functions whose pointwise limit is  $cf + g$ . Hence, by Lemma 6.6 and the monotone convergence theorem

$$\begin{aligned} \int_M (cf + g) \, d\mu &= \lim_{n \rightarrow \infty} \int_M (cs_n + t_n) \, d\mu \\ &= \lim_{n \rightarrow \infty} \left( c \int_M s_n \, d\mu + \int_M t_n \, d\mu \right) \\ &= c \lim_{n \rightarrow \infty} \int_M s_n \, d\mu + \lim_{n \rightarrow \infty} \int_M t_n \, d\mu \\ &= c \int_M f \, d\mu + \int_M g \, d\mu. \end{aligned}$$

(ii) To check that  $\nu_f$  is a measure on  $(M, \Sigma)$ , first note that we have

$$\nu_f(\emptyset) = \int_{\emptyset} f \, d\mu := \int_M f \chi_{\emptyset} \, d\mu = 0.$$

Let  $\{A_i\}_{i \in \mathbb{N}}$  be a pairwise disjoint sequence in  $\Sigma$ . Define, for any  $n \in \mathbb{N}$ ,

$$f_n := f \chi_{(\bigcup_{i=0}^n A_i)}.$$

Since, for all  $n \in \mathbb{N}$ , we have  $\bigcup_{i=0}^n A_i \subseteq \bigcup_{i=0}^{n+1} A_i$  and  $f$  is non-negative,  $\{f_n\}_{n \in \mathbb{N}}$  is an increasing sequence of non-negative, measurable, simple functions whose pointwise limits is  $f \chi_{(\bigcup_{i=0}^{\infty} A_i)}$ . Hence, by recalling Proposition 6.2, we have

$$\begin{aligned} \nu_f\left(\bigcup_{i=0}^{\infty} A_i\right) &:= \int_{\bigcup_{i=0}^{\infty} A_i} f \, d\mu \\ &= \int_M f \chi_{(\bigcup_{i=0}^{\infty} A_i)} \, d\mu \\ &= \lim_{n \rightarrow \infty} \int_M f \chi_{(\bigcup_{i=0}^n A_i)} \, d\mu \\ &= \lim_{n \rightarrow \infty} \int_M \left(f \sum_{i=0}^n \chi_{A_i}\right) \, d\mu \\ &= \lim_{n \rightarrow \infty} \left(\sum_{i=0}^n \int_M f \chi_{A_i} \, d\mu\right) \\ &= \sum_{i=0}^{\infty} \nu_f(A_i). \end{aligned}$$

(iii) Note that  $A \cap (M \setminus A) = \emptyset$ . Hence, by using the fact that  $\nu_f$  from part (ii) is a measure on  $(M, \Sigma)$ , we have

$$\int_M f \, d\mu = \nu_f(M) = \nu_f(A) + \nu_f(M \setminus A) = \int_A f \, d\mu + \int_{M \setminus A} f \, d\mu. \quad \square$$

Part (i) of the previous lemma and the monotone convergence theorem also imply that, for any sequence  $\{f_n\}_{n \in \mathbb{N}}$  of non-negative, measurable functions, we have

$$\int_M \left(\sum_{n=0}^{\infty} f_n\right) d\mu = \sum_{n=0}^{\infty} \int_M f_n \, d\mu.$$

Again, note that this result does *not* hold for the Riemann integral unless stronger conditions are places on the sequence  $\{f_n\}_{n \in \mathbb{N}}$ .

Finally, we have a simple but crucial result for Lebesgue integration.

**Theorem 6.19.** *Let  $(M, \Sigma, \mu)$  be a measure space and let  $f: M \rightarrow \overline{\mathbb{R}}$  be a non-negative, measurable function. Then*

$$\int_M f \, d\mu = 0 \quad \Leftrightarrow \quad f =_{\text{a.e.}} 0.$$

*Proof.*  $(\Rightarrow)$  Suppose that  $\int_M f \, d\mu = 0$ . Define  $A_n := \{m \in M \mid f(m) > \frac{1}{n+1}\}$  and let

$$s_n := \frac{1}{n+1} \chi_{A_n}.$$

By definition of  $A_n$ , we clearly have  $s_n \leq f$  for all  $n \in \mathbb{N}$ . Hence,

$$0 \leq \int_M s_n \, d\mu \leq \int_M f \, d\mu = 0$$

and thus,  $\int_M s_n \, d\mu = 0$  for all  $n \in \mathbb{N}$ . Since by definition

$$\int_M s_n \, d\mu = \frac{1}{n+1} \mu(A_n),$$

we must also have  $\mu(A_n) = 0$  for all  $n \in \mathbb{N}$ . Let  $A := \{m \in M \mid f(m) \neq 0\}$ . Then, as  $f$  is non-negative, we have

$$A = \bigcup_{n=0}^{\infty} A_n = \bigcup_{n=0}^{\infty} \{m \in M \mid f(m) > \frac{1}{n+1}\}$$

and, since  $A_n \subseteq A_{n+1}$  for all  $n \in \mathbb{N}$ , we have

$$\mu(A) = \mu\left(\bigcup_{n=0}^{\infty} A_n\right) = \lim_{n \rightarrow \infty} \mu(A_n) = 0.$$

Thus,  $f$  is zero except on the null set  $A$ . That is,  $f =_{\text{a.e.}} 0$ .

( $\Leftarrow$ ) Suppose that  $f =_{\text{a.e.}} 0$ . Let  $S$  be the set of non-negative, measurable, simple functions  $s$  such that  $s \leq f$ . As  $f =_{\text{a.e.}} 0$ , we have  $s =_{\text{a.e.}} 0$  for all  $s \in S$ . Thus, if

$$s = \sum_{i=1}^n r_i \chi_{A_i},$$

we must have either  $r_i = 0$  or  $\mu(A_i) = 0$  for all  $1 \leq i \leq n$ . Hence, for all  $s \in S$ ,

$$\int_M s \, d\mu := \sum_{i=1}^n r_i \mu(A_i) = 0.$$

Therefore, we have

$$\int_M f \, d\mu := \sup_{s \in S} \int_M s \, d\mu = 0. \quad \square$$

This means that, for the purposes of Lebesgue integration, null sets can be neglected as they do not change the value of an integral. The following are some examples of this.

**Corollary 6.20.** *Let  $(M, \Sigma, \mu)$  be a measure space, let  $A \in \Sigma$  and let  $f, g: M \rightarrow \overline{\mathbb{R}}$  be non-negative, measurable functions.*

- (i) *If  $\mu(A) = 0$ , then  $\int_A f \, d\mu = 0$*
- (ii) *If  $f =_{\text{a.e.}} g$ , then  $\int_M f \, d\mu = \int_M g \, d\mu$*
- (iii) *If  $f \leq_{\text{a.e.}} g$ , then  $\int_M f \, d\mu \leq \int_M g \, d\mu$ .*

*Proof.* (i) Clearly,  $f\chi_A =_{\text{a.e.}} 0$  and hence

$$\int_A f \, d\mu = \int_M f\chi_A \, d\mu = 0.$$

(ii) As  $f =_{\text{a.e.}} g$ , we have  $f - g =_{\text{a.e.}} 0$  and thus

$$0 = \int_M (f - g) \, d\mu = \int_M f \, d\mu - \int_M g \, d\mu.$$

(iii) Let  $B := \{m \in M \mid f(m) > g(m)\}$ . As  $f \leq_{\text{a.e.}} g$ , we have  $\mu(B) = 0$  and  $f \leq g$  on  $M \setminus B$ . Thus

$$\int_M f \, d\mu = \int_B f \, d\mu + \int_{M \setminus B} f \, d\mu \leq \int_{M \setminus B} g \, d\mu \leq \int_M g \, d\mu$$

where we used part (i) and Lemma 6.13.  $\square$

Example 6.21. Consider  $(\mathbb{R}, \sigma(\mathcal{O}_{\mathbb{R}}), \lambda)$  and let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be the *Dirichlet function*

$$f(r) := \begin{cases} 1 & \text{if } r \in \mathbb{Q} \\ 0 & \text{if } r \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

The Dirichlet function is the usual example of a function which is *not* Riemann integrable (on any real interval). We will now show that we can easily assign a numerical value to its integral on any measurable subset of  $\mathbb{R}$ . First, note that a set  $A \in \sigma(\mathcal{O}_{\mathbb{R}})$  is null with respect to the Lebesgue measure if, and only if,

$$\forall \varepsilon > 0 : \exists \{I_n\}_{n \in \mathbb{N}} : A \subset \bigcup_{n=0}^{\infty} I_n \text{ and } \sum_{n=0}^{\infty} \lambda(I_n) < \varepsilon$$

where  $\{I_n\}_{n \in \mathbb{N}}$  is a sequence of real intervals. From this, it immediately follows that any countable subset of  $\mathbb{R}$  has zero Lebesgue measure. Thus,  $\lambda(\mathbb{Q}) = 0$  and hence,  $f =_{\text{a.e.}} 0$ . Therefore, by the previous lemmas, we have

$$\int_A f \, d\lambda = 0$$

for any measurable subset  $A$  of  $\mathbb{R}$ .

## 6.4 Lebesgue integrable functions


Since the difference  $\infty - \infty$  is not defined, we cannot integrate all measurable functions. There is, however, a very small extra condition (beyond measurability) that determines the class of functions to which we can extend our previous definition.

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space and let  $f: M \rightarrow \overline{\mathbb{R}}$ . The function  $f$  is said to be (*Lebesgue*) *integrable* if it is measurable and

$$\int_M |f| \, d\mu < \infty.$$

We denote the set of all integrable functions  $M \rightarrow \overline{\mathbb{R}}$  by  $\mathcal{L}_{\mathbb{R}}^1(M, \Sigma, \mu)$ , or simply  $\mathcal{L}^1(M)$  if there is no risk of confusion.  $\diamond$

For any  $f: M \rightarrow \overline{\mathbb{R}}$ , we define  $f^+ := \max(f, 0)$  and  $f^- := \max(-f, 0)$ , which are measurable whenever  $f$  is measurable by part (iv) of Lemma 6.11.



Observe that  $f = f^+ - f^-$  and  $|f| = f^+ + f^-$ . Clearly, we have  $f^+ \leq |f|$  and  $f^- \leq |f|$ , and hence

$$\int_M |f| d\mu < \infty \quad \Leftrightarrow \quad \int_M f^+ d\mu < \infty \quad \text{and} \quad \int_M f^- d\mu < \infty.$$

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space and let  $f: M \rightarrow \overline{\mathbb{R}}$  be integrable. Then, the (Lebesgue) integral of  $f$  over  $M$  with respect to  $\mu$  is

$$\int_M f d\mu := \int_M f^+ d\mu - \int_M f^- d\mu.$$

◇

It should be clear that the role of the integrability condition  $\int_M |f| d\mu < \infty$  is to prevent the integral of  $f$  from being  $\infty - \infty$ , which is not defined.

In quantum mechanics, we usually deal with complex functions. The extension to the complex case of the integration theory presented thus far is straightforward.

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space. A complex function  $f: M \rightarrow \mathbb{C}$  is said to be *integrable* if the real functions  $\operatorname{Re}(f)$  and  $\operatorname{Im}(f)$  are measurable and

$$\int_M |f| d\mu < \infty,$$

where  $|f|$  denotes the complex modulus, i.e.  $|f|^2 = \operatorname{Re}(f)^2 + \operatorname{Im}(f)^2$ . We denote the set of all integrable complex functions by  $\mathcal{L}_{\mathbb{C}}^1(M, \Sigma, \mu)$ , or simply  $\mathcal{L}^1(M)$  if there is no risk of confusion. ◇

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space and let  $f: M \rightarrow \mathbb{C}$  be integrable. We define

$$\int_M f d\mu := \int_M \operatorname{Re}(f) d\mu + i \int_M \operatorname{Im}(f) d\mu.$$

◇

The following lemma gives the properties expected of sums and scalar multiples of integrals. Note, however, that before we show that, say, the integral of a sum is the sum of the integrals, it is necessary to first show that the sum of two functions in  $\mathcal{L}^1(M)$  is again in  $\mathcal{L}^1(M)$ .

**Lemma 6.22.** *Let  $(M, \Sigma, \mu)$  be a measure space, let  $f, g \in \mathcal{L}^1(M)$  and let  $c \in \mathbb{R}$ . Then*

- (i)  $|f| \in \mathcal{L}^1(M)$  and  $\left| \int_M f \, d\mu \right| \leq \int_M |f| \, d\mu$
- (ii)  $cf \in \mathcal{L}^1(M)$  and  $\int_M cf \, d\mu = c \int_M f \, d\mu$
- (iii)  $f + g \in \mathcal{L}^1(M)$  and  $\int_M (f + g) \, d\mu = \int_M f \, d\mu + \int_M g \, d\mu$
- (iv)  $\mathcal{L}^1(M)$  is a vector space.

*Proof.* (i) As  $||f|| = |f|$ , we have  $|f| \in \mathcal{L}^1(M)$ . Then, by the triangle inequality,

$$\begin{aligned}
 \left| \int_M f \, d\mu \right| &= \left| \int_M f^+ \, d\mu - \int_M f^- \, d\mu \right| \\
 &\leq \left| \int_M f^+ \, d\mu \right| + \left| \int_M f^- \, d\mu \right| \\
 &= \int_M f^+ \, d\mu + \int_M f^- \, d\mu \\
 &= \int_M (f^+ + f^-) \, d\mu \\
 &= \int_M |f| \, d\mu.
 \end{aligned}$$

(ii) We have

$$\int_M |cf| \, d\mu = \int_M |c||f| \, d\mu = |c| \int_M |f| \, d\mu < \infty$$

and hence, we have  $cf \in \mathcal{L}^1(M)$ .

Suppose  $c \geq 0$ . Then,  $(cf)^+ = cf^+$  and  $(cf)^- = cf^-$  and thus

$$\begin{aligned}
 \int_M cf \, d\mu &= \int_M (cf)^+ \, d\mu - \int_M (cf)^- \, d\mu \\
 &= \int_M cf^+ \, d\mu - \int_M cf^- \, d\mu \\
 &= c \int_M f^+ \, d\mu - c \int_M f^- \, d\mu \\
 &= c \int_M (f^+ - f^-) \, d\mu \\
 &= c \int_M f \, d\mu
 \end{aligned}$$

Now suppose  $c = -1$ . Then,  $(-f)^+ = f^-$  and  $(-f)^- = f^+$ . Thus

$$\begin{aligned}
\int_M (-f) \, d\mu &= \int_M (-f)^+ \, d\mu - \int_M (-f)^- \, d\mu \\
&= \int_M f^- \, d\mu - \int_M f^+ \, d\mu \\
&= -\left( \int_M f^+ \, d\mu - \int_M f^- \, d\mu \right) \\
&= -\int_M (f^+ - f^-) \, d\mu \\
&= -\int_M f \, d\mu.
\end{aligned}$$

The case  $c < 0$  follows by writing  $c = (-1)(-c)$  and applying the above results.

(iii) By the triangle inequality, we have  $|f + g| \leq |f| + |g|$  and thus

$$\int_M |f + g| \, d\mu \leq \int_M |f| \, d\mu + \int_M |g| \, d\mu < \infty.$$

Hence,  $f + g \in \mathcal{L}^1(M)$ . Moreover, we have

$$f + g = f^+ - f^- + g^+ - g^- = (f^+ + g^+) - (f^- + g^-),$$

where  $f^+ + g^+$  and  $f^- + g^-$  are non-negative and measurable. Note that, while  $(f + g)^+ \neq f^+ + g^+$  and  $(f + g)^- \neq f^- + g^-$ , one can show that  $f^+ + g^+$  and  $f^- + g^-$  give an equivalent splitting of  $f + g$  to define its integral. Therefore

$$\begin{aligned}
\int_M (f + g) \, d\mu &= \int_M (f^+ + g^+) \, d\mu - \int_M (f^- + g^-) \, d\mu \\
&= \int_M f^+ \, d\mu + \int_M g^+ \, d\mu - \int_M f^- \, d\mu - \int_M g^- \, d\mu \\
&= \int_M (f^+ - f^-) \, d\mu + \int_M (g^+ - g^-) \, d\mu \\
&= \int_M f \, d\mu + \int_M g \, d\mu.
\end{aligned}$$

(iv) The set of all functions from  $M$  to  $\overline{\mathbb{R}}$  is a vector space. By parts (ii) and (iii), we have that  $\mathcal{L}^1(M)$  is a vector subspace of this vector space and hence, a vector space in its own right.  $\square$

Some properties of the integrals of non-negative, measurable functions easily carry over to general integrable functions.

**Lemma 6.23.** *Let  $(M, \Sigma, \mu)$  be a measure space and let  $f, g \in \mathcal{L}^1(M)$ .*

(i) *If  $f =_{\text{a.e.}} g$ , then  $\int_M f \, d\mu = \int_M g \, d\mu$*

(ii) *If  $f \leq_{\text{a.e.}} g$ , then  $\int_M f \, d\mu \leq \int_M g \, d\mu$ .*

Just as the monotone convergence theorem was very important for integrals of non-negative, measurable functions, there is a similar theorem that is important for integrals of functions in  $\mathcal{L}^1(X)$ .

**Theorem 6.24** (Dominated convergence theorem). *Let  $(M, \Sigma, \mu)$  be a measure space and let  $\{f_n\}_{n \in \mathbb{N}}$  be a sequence of measurable functions which converges almost everywhere to a measurable function  $f$ . If there exists  $g \in \mathcal{L}^1(M)$  such that  $|f_n| \leq_{\text{a.e.}} g$  for all  $n \in \mathbb{N}$ , then*

(i)  $f \in \mathcal{L}^1(M)$  and  $f_n \in \mathcal{L}^1(M)$  for all  $n \in \mathbb{N}$

(ii)  $\lim_{n \rightarrow \infty} \int_M |f_n - f| d\mu = 0$

(iii)  $\lim_{n \rightarrow \infty} \int_M f_n d\mu = \int_M f d\mu$ .

Remark 6.25. By “ $\{f_n\}_{n \in \mathbb{N}}$  converges almost everywhere to  $f$ ” we mean, of course, that there exists a null set  $A \in \Sigma$  such that

$$\forall m \in M \setminus A : \lim_{n \rightarrow \infty} f_n(m) = f(m).$$

### 6.5 The function spaces $L^p(M, \Sigma, \mu)$

While in quantum mechanics we need the theory of  $L^p$  spaces only for  $p = 2$ , it is worthwhile to develop the theory in more generality, by considering  $L^p$  for all  $1 \leq p \leq \infty$ .

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space and let  $p \in [1, \infty)$ . We define

$$\mathcal{L}_{\mathbb{R}}^p(M, \Sigma, \mu) := \left\{ f : M \rightarrow \overline{\mathbb{R}} \mid f \text{ is measurable and } \int_M |f|^p d\mu < \infty \right\}$$

and, similarly,

$$\mathcal{L}_{\mathbb{C}}^p(M, \Sigma, \mu) := \left\{ f : M \rightarrow \mathbb{C} \mid \operatorname{Re}(f) \text{ and } \operatorname{Im}(f) \text{ are measurable and } \int_M |f|^p d\mu < \infty \right\}.$$

Whenever there is no risk of confusion, we lighten the notation to just  $\mathcal{L}^p$ .  $\diamond$

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space and let  $f : M \rightarrow \overline{\mathbb{R}}$ . The *essential supremum* of  $f$  is defined as

$$\operatorname{ess\,sup} f := \inf \{ c \in \overline{\mathbb{R}} \mid f \leq_{\text{a.e.}} c \}.$$

Then,  $f$  is said to be *almost everywhere bounded (from above)* if  $\operatorname{ess\,sup} f < \infty$ .  $\diamond$

Alternatively,  $f : M \rightarrow \overline{\mathbb{R}}$  is almost everywhere bounded if there exists a null set  $A \in \Sigma$  such that  $f$  restricted to  $M \setminus A$  is bounded.

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space. We define

$$\mathcal{L}_{\mathbb{R}}^{\infty}(M, \Sigma, \mu) := \left\{ f : M \rightarrow \overline{\mathbb{R}} \mid f \text{ is measurable and } \operatorname{ess\,sup} |f| < \infty \right\}$$

and, similarly,

$$\mathcal{L}_{\mathbb{C}}^{\infty}(M, \Sigma, \mu) := \left\{ f : M \rightarrow \mathbb{C} \mid \operatorname{Re}(f) \text{ and } \operatorname{Im}(f) \text{ are measurable and } \operatorname{ess\,sup} |f| < \infty \right\}.$$

Whenever there is no risk of confusion, we lighten the notation to just  $\mathcal{L}^p$ , for  $p \in [1, \infty]$ .  $\diamond$



All the  $\mathcal{L}^p$  spaces become vector spaces once equipped with pointwise addition and multiplication. Let us show this in detail for  $\mathcal{L}_{\mathbb{C}}^2$ .

**Proposition 6.26.** *Let  $(M, \Sigma, \mu)$  be a measure space. Then,  $\mathcal{L}_{\mathbb{C}}^2$  is a complex vector space.*

*Proof.* The set of all functions  $M \rightarrow \mathbb{C}$ , often denoted  $M^{\mathbb{C}}$ , is a vector space under pointwise addition and multiplication. Hence, it suffices to show that  $\mathcal{L}_{\mathbb{C}}^2$  is a subspace of  $M^{\mathbb{C}}$ .

(i) Let  $f \in \mathcal{L}_{\mathbb{C}}^2$  and  $z \in \mathbb{C}$ . As  $|z| \in \mathbb{R}$ , we have:

$$\int_M |zf|^2 d\mu = |z|^2 \int_M |f|^2 d\mu < \infty$$

and hence,  $zf \in \mathcal{L}_{\mathbb{C}}^2$ .

(ii) Let  $f, g \in \mathcal{L}_{\mathbb{C}}^2$ . Note that

$$\begin{aligned} |f+g|^2 &= (f+g)\overline{(f+g)} \\ &= (f+g)(\bar{f}+\bar{g}) \\ &= f\bar{f} + f\bar{g} + g\bar{f} + g\bar{g} \\ &= |f|^2 + f\bar{g} + g\bar{f} + |g|^2. \end{aligned}$$

Moreover, as

$$0 \leq |f-g|^2 = |f|^2 - f\bar{g} - g\bar{f} + |g|^2,$$

we have  $f\bar{g} + g\bar{f} \leq |f|^2 + |g|^2$ , and thus

$$|f+g|^2 \leq 2|f|^2 + 2|g|^2.$$

Therefore

$$\int_M |f+g|^2 d\mu \leq 2 \int_M |f|^2 d\mu + 2 \int_M |g|^2 d\mu < \infty,$$

and hence  $f+g \in \mathcal{L}_{\mathbb{C}}^2$ . □

Ideally, we would like to turn all these  $\mathcal{L}^p$  spaces into Banach spaces. Let us begin by equipping them with a weaker piece of extra structure.

**Proposition 6.27.** *Let  $(M, \Sigma, \mu)$  be a measure space and let  $p \in [0, \infty]$ . Then, the maps  $\|\cdot\|_p: \mathcal{L}^p \rightarrow \mathbb{R}$  defined by*

$$\|f\|_p := \begin{cases} \left( \int_M |f|^p d\mu \right)^{\frac{1}{p}} & \text{for } 1 \leq p < \infty \\ \text{ess sup } |f| & \text{for } p = \infty \end{cases}$$

*are semi-norms on  $\mathcal{L}^p$ . That is, for all  $z \in \mathbb{C}$  and  $f, g \in \mathcal{L}^p$ ,*

$$(i) \quad \|f\|_p \geq 0$$

$$(ii) \quad \|zf\|_p = |z| \|f\|_p$$

$$(iii) \|f + g\|_p \leq \|f\|_p + \|g\|_p.$$

In other words, the notion of semi-norm is a generalisation of that of norm obtained by relaxing the definiteness condition. If the measure space  $(M, \Sigma, \mu)$  is such that the empty set is the only null set, then  $\|\cdot\|_p$  is automatically definite and hence, a norm.

*Example 6.28.* Consider  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), \mu)$ , where  $\mu$  is the counting measure. Then, as  $\mu(A)$  is the cardinality of  $A$ , the only null set is the empty set. Thus, recalling that functions on  $\mathbb{N}$  are just sequences, the maps

$$\|\{a_n\}_{n \in \mathbb{N}}\|_p = \begin{cases} \left( \sum_{n=0}^{\infty} |a_n|^p \right)^{\frac{1}{p}} & \text{for } 1 \leq p < \infty \\ \sup\{|a_n| \mid n \in \mathbb{N}\} & \text{for } p = \infty \end{cases}$$

are norms on  $\mathcal{L}^p(\mathbb{N})$ . In particular, note that we have  $\mathcal{L}^2(\mathbb{N}) = \ell^2(\mathbb{N})$ .

However, in general measure spaces, we only have

$$\|f\|_p = 0 \quad \Leftrightarrow \quad f =_{\text{a.e.}} 0,$$

as we have shown in Theorem 6.19 for  $\mathcal{L}^1$ , and it is often very easy to produce an  $f \neq 0$  such that  $\|f\|_p = 0$ . The solution to this problem is to construct new spaces from the  $\mathcal{L}^p$  in which functions that are almost everywhere equal are, in fact, the same function. In other words, we need to consider the quotient space of  $\mathcal{L}^p$  by the equivalence relation “being almost everywhere equal”.

**Definition.** Let  $M$  be a set. An *equivalence relation* on  $M$  is a set  $\sim \subseteq M \times M$  such that, writing  $a \sim b$  for  $(a, b) \in \sim$ , we have

$$(i) \quad a \sim a \quad \text{(reflexivity)}$$

$$(ii) \quad a \sim b \Leftrightarrow b \sim a \quad \text{(symmetry)}$$

$$(iii) \quad (a \sim b \text{ and } b \sim c) \Rightarrow a \sim c \quad \text{(transitivity)}$$

for all  $a, b, c \in M$ . If  $\sim$  is an equivalence relation on  $M$ , we define the *equivalence class* of  $m \in M$  by  $[m] := \{a \in M \mid m \sim a\}$  and the *quotient set* of  $M$  by  $\sim$  by

$$M/\sim := \{[m] \mid m \in M\}.$$

◇

It is easy to show that  $M/\sim$  is a *partition* of  $M$ , i.e.

$$M = \bigcup_{m \in M} [m] \quad \text{and} \quad [a] \cap [b] = \emptyset \text{ whenever } a \neq b.$$

In fact, the notions of equivalence relation on  $M$  and partition of  $M$  are one and the same.

**Lemma 6.29.** Let  $(M, \Sigma, \mu)$  be a measure space and let  $\sim$  be defined by

$$f \sim g \quad :\Leftrightarrow \quad f =_{\text{a.e.}} g.$$

Then,  $\sim$  is an equivalence relation on  $\mathcal{L}^p$ .

*Proof.* Let  $f, g, h \in \mathcal{L}^p$ . Clearly,  $f \sim f$  and  $f \sim g \Leftrightarrow g \sim f$ . Now suppose that  $f \sim g$  and  $g \sim h$ . Then, there exist null sets  $A, B \in \Sigma$  such that  $f = g$  on  $M \setminus A$  and  $g = h$  on  $M \setminus B$ . Recall that  $\sigma$ -algebras are closed under intersections and hence,  $A \cap B \in \Sigma$ . Obviously, we have  $f = h$  on  $M \setminus (A \cap B)$  and, since

$$\mu(A \cap B) \leq \mu(A \cup B) \leq \mu(A) + \mu(B) = 0,$$

the set  $A \cap B$  is null. Thus,  $f \sim h$ . □

**Definition.** Let  $(M, \Sigma, \mu)$  be a measure space and let  $f \sim g \Leftrightarrow f =_{\text{a.e.}} g$ . We define

$$L^p := \mathcal{L}^p / \sim = \{[f] \mid f \in \mathcal{L}^p\}.$$

◇

**Lemma 6.30.** Let  $(M, \Sigma, \mu)$  be a measure space. Then, the maps

$$(i) \quad +: L^p \times L^p \rightarrow L^p \text{ defined by } [f] + [g] := [f + g]$$

$$(ii) \quad \cdot: \mathbb{C} \times L^p \rightarrow L^p \text{ defined by } z[f] := [zf]$$

$$(iii) \quad \|\cdot\|_p: L^p \rightarrow \mathbb{R} \text{ defined by } \|[f]\|_p := \|f\|_p$$

are well-defined. Moreover,  $\|\cdot\|_p: L^p \rightarrow \mathbb{R}$  is a norm on  $L^p$ .

**Lemma 6.31** (Hölder's inequality). Let  $(M, \Sigma, \mu)$  be a measure space and let  $p, q \in [1, \infty]$  be such that  $\frac{1}{p} + \frac{1}{q} = 1$  (where  $\frac{1}{\infty} := 0$ ). Then, for all measurable functions  $f, g: M \rightarrow \mathbb{C}$ , we have

$$\left| \int_M fg \, d\mu \right| \leq \left( \int_M |f|^p \, d\mu \right)^{\frac{1}{p}} \left( \int_M |g|^q \, d\mu \right)^{\frac{1}{q}}.$$

Hence, if  $[f] \in L^p$  and  $[g] \in L^q$ , with  $\frac{1}{p} + \frac{1}{q} = 1$ , then  $[fg] \in L^1$  and

$$\|[fg]\|_1 \leq \|[f]\|_p \|[g]\|_q.$$

The equality holds if and only if  $|f|^p$  and  $|g|^q$  are linearly dependent on  $L^1$ . That is, there exists non-negative real numbers  $\alpha, \beta \in \mathbb{R}$  such that

$$\alpha |f|^p =_{\text{a.e.}} \beta |g|^q$$

Note it is clear that  $\alpha$  and  $\beta$  cannot both vanish.

**Theorem 6.32.** The spaces  $L^p$  are Banach spaces for all  $p \in [0, \infty]$ .

We have already remarked that the case  $p = 2$  is special in that  $L^2$  is the only  $L^p$  space which can be made into a Hilbert space.

**Proposition 6.33.** *Let  $(M, \Sigma, \mu)$  be a measure space and define*

$$\begin{aligned} \langle \cdot | \cdot \rangle_{L^2} : L^2 \times L^2 &\rightarrow \mathbb{C} \\ ([f], [g]) &\mapsto \int_M \bar{f}g \, d\mu. \end{aligned}$$

*Then,  $\langle \cdot | \cdot \rangle_{L^2}$  is well-defined and it is a sesqui-linear inner product on  $L^2$ .*

*Proof.* First note that if  $[f] \in L^2$ , then  $[\bar{f}] \in L^2$  and hence, by Hölder inequality,  $[\bar{f}g] \in L^1$ . This ensures that  $\langle [f] | [g] \rangle_{L^2} \in \mathbb{C}$  for all  $[f], [g] \in L^2$ .

To show well-definedness, let  $f' =_{\text{a.e.}} f$  and  $g' =_{\text{a.e.}} g$ . Then,  $\bar{f}'g' =_{\text{a.e.}} \bar{f}g$  and thus

$$\langle [f'] | [g'] \rangle_{L^2} := \int_M \bar{f}'g' \, d\mu = \int_M \bar{f}g \, d\mu =: \langle [f] | [g] \rangle_{L^2}.$$

Now, let  $[f], [g], [h] \in L^2$  and  $z \in \mathbb{C}$ . Then

$$(i) \quad \overline{\langle [f] | [g] \rangle_{L^2}} = \overline{\int_M \bar{f}g \, d\mu} = \int_M f\bar{g} \, d\mu = \langle [g] | [f] \rangle_{L^2}$$

(ii) We have

$$\begin{aligned} \langle [f] | z[g] + [h] \rangle_{L^2} &= \int_M \bar{f}(zg + h) \, d\mu \\ &= z \int_M \bar{f}g \, d\mu + \int_M \bar{f}h \, d\mu \\ &= z \langle [f] | [g] \rangle_{L^2} + \langle [f] | [h] \rangle_{L^2} \end{aligned}$$

$$(iii) \quad \text{We have } \langle [f] | [f] \rangle_{L^2} = \int_M \bar{f}f \, d\mu = \int_M |f|^2 \, d\mu \geq 0 \text{ and}$$

$$\langle [f] | [f] \rangle_{L^2} = 0 \quad \Leftrightarrow \quad \int_M |f|^2 \, d\mu = 0 \quad \Leftrightarrow \quad \| [f] \|_2 = 0.$$

Thus,  $[f] = 0 := [0]$ . □

The last part of the proof also shows that  $\langle \cdot | \cdot \rangle_{L^2}$  induces the norm  $\| \cdot \|_2$ , with respect to which  $L^2$  is a Banach space. Hence,  $(L^2, \langle \cdot | \cdot \rangle_{L^2})$  is a Hilbert space.

Remark 6.34. The inner product  $\langle \cdot | \cdot \rangle_{L^2}$  on  $L^2_{\mathbb{C}}(\mathbb{N}, \mathcal{P}(\mathbb{N}), \mu)$  coincides with the inner product  $\langle \cdot | \cdot \rangle_{\ell^2}$  on  $\ell^2(\mathbb{N})$  defined in the section on separable Hilbert spaces.

## 7 Self-adjoint and essentially self-adjoint operators

While we have already given some of the following definitions in the introductory section on the axioms of quantum mechanics, we reproduce them here for completeness.

### 7.1 Adjoint operators

**Definition.** A linear map or operator  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  is said to be *densely defined* if  $\mathcal{D}_A$  is dense in  $\mathcal{H}$ , i.e.

$$\forall \varepsilon > 0 : \forall \psi \in \mathcal{H} : \exists \alpha \in \mathcal{D}_A : \|\alpha - \psi\| < \varepsilon.$$

◇

Equivalently,  $\overline{\mathcal{D}_A} = \mathcal{H}$ , i.e. for every  $\psi \in \mathcal{H}$  there exists a sequence  $\{\alpha_n\}_{n \in \mathbb{N}}$  in  $\mathcal{D}_A$  whose limit is  $\psi$ .

**Definition.** Let  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  be a densely defined operator on  $\mathcal{H}$ . The *adjoint* of  $A$  is the operator  $A^*: \mathcal{D}_{A^*} \rightarrow \mathcal{H}$  defined by

$$(i) \quad \mathcal{D}_{A^*} := \{\psi \in \mathcal{H} \mid \exists \eta \in \mathcal{H} : \forall \alpha \in \mathcal{D}_A : \langle \psi | A\alpha \rangle = \langle \eta | \alpha \rangle\}$$

$$(ii) \quad A^*\psi := \eta.$$

◇

**Proposition 7.1.** *The adjoint operator  $A^*: \mathcal{D}_{A^*} \rightarrow \mathcal{H}$  is well-defined.*

*Proof.* Let  $\psi \in \mathcal{H}$  and let  $\eta, \tilde{\eta} \in \mathcal{H}$  be such that

$$\forall \alpha \in \mathcal{D}_A : \quad \langle \psi | A\alpha \rangle = \langle \eta | \alpha \rangle \quad \text{and} \quad \langle \psi | A\alpha \rangle = \langle \tilde{\eta} | \alpha \rangle.$$

Then, for all  $\alpha$  in  $\mathcal{D}_A$ , we have

$$\langle \eta - \tilde{\eta} | \alpha \rangle = \langle \eta | \alpha \rangle - \langle \tilde{\eta} | \alpha \rangle = \langle \psi | A\alpha \rangle - \langle \psi | A\alpha \rangle = 0$$

and hence, by positive-definiteness,  $\eta = \tilde{\eta}$ . □

If  $A$  and  $B$  are densely defined and  $\mathcal{D}_A = \mathcal{D}_B$ , then the pointwise sum  $A + B$  is clearly densely defined and hence,  $(A + B)^*$  exists. However, we do *not* have  $(A + B)^* = A^* + B^*$  in general, unless one of  $A$  and  $B$  is bounded, but we do have the following result.

**Proposition 7.2.** *If  $A$  is densely defined, then*

$$(A + z \text{id}_{\mathcal{D}_A})^* = A^* + \bar{z} \text{id}_{\mathcal{D}_A}$$

for any  $z \in \mathbb{C}$ .

The identity operator  $\text{id}_{\mathcal{D}_A}$  is usually suppressed in the notation, so that the above equation reads  $(A + z)^* = A^* + \bar{z}$ .

**Definition.** Let  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  be a linear operator. The *kernel* and *range* of  $A$  are

$$\ker(A) := \{\alpha \in \mathcal{D}_A \mid A\alpha = 0\}, \quad \text{ran}(A) := \{A\alpha \mid \alpha \in \mathcal{D}_A\}.$$

The range is also called *image* and  $\text{im}(A)$  and  $A(\mathcal{D}_A)$  are alternative notations.  $\diamond$

**Proposition 7.3.** An operator  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  is

- (i) *injective if, and only if,  $\ker(A) = \{0\}$*
- (ii) *surjective if, and only if,  $\text{ran}(A) = \mathcal{H}$ .*

**Definition.** An operator  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  is *invertible* if there exists an operator  $B: \mathcal{H} \rightarrow \mathcal{D}_A$  such that  $A \circ B = \text{id}_{\mathcal{H}}$  and  $B \circ A = \text{id}_{\mathcal{D}_A}$ .  $\diamond$

**Proposition 7.4.** An operator  $A$  is invertible if, and only if,

$$\ker(A) = \{0\} \quad \text{and} \quad \overline{\text{ran}(A)} = \mathcal{H}.$$

**Proposition 7.5.** Let  $A$  be densely defined. Then,  $\ker(A^*) = \text{ran}(A)^\perp$ .

*Proof.* We have

$$\psi \in \ker(A^*) \Leftrightarrow A^*\psi = 0 \Leftrightarrow \forall \alpha \in \mathcal{D}_A : \langle \psi | A\alpha \rangle = 0 \Leftrightarrow \psi \in \text{ran}(A)^\perp. \quad \square$$

**Definition.** Let  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  and  $B: \mathcal{D}_B \rightarrow \mathcal{H}$  be operators. We say that  $B$  is an *extension* of  $A$ , and we write  $A \subseteq B$ , if

- (i)  $\mathcal{D}_A \subseteq \mathcal{D}_B$
- (ii)  $\forall \alpha \in \mathcal{D}_A : A\alpha = B\alpha$ .

$\diamond$

**Proposition 7.6.** Let  $A, B$  be densely defined. If  $A \subseteq B$ , then  $B^* \subseteq A^*$ .

*Proof.* (i) Let  $\psi \in \mathcal{D}_{B^*}$ . Then, there exists  $\eta \in \mathcal{H}$  such that

$$\forall \beta \in \mathcal{D}_B : \langle \psi | B\beta \rangle = \langle \eta | \beta \rangle.$$

In particular, as  $A \subseteq B$ , we have  $\mathcal{D}_A \subseteq \mathcal{D}_B$  and thus

$$\forall \alpha \in \mathcal{D}_A \subseteq \mathcal{D}_B : \langle \psi | B\alpha \rangle = \langle \psi | A\alpha \rangle = \langle \eta | \alpha \rangle.$$

Therefore,  $\psi \in \mathcal{D}_{A^*}$  and hence,  $\mathcal{D}_{B^*} \subseteq \mathcal{D}_{A^*}$ .

- (ii) From the above, we also have  $B^*\psi := \eta =: A^*\psi$  for all  $\psi \in \mathcal{D}_{B^*}$ .  $\square$

## 7.2 The adjoint of a symmetric operator

**Definition.** A densely defined operator  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  is called *symmetric* if

$$\forall \alpha, \beta \in \mathcal{D}_A : \langle \alpha | A\beta \rangle = \langle A\alpha | \beta \rangle.$$

◇

*Remark 7.7.* In the physics literature, symmetric operators are usually referred to as *Hermitian* operators. However, this notion is then confused with the that of self-adjointness when physicists say that observables in quantum mechanics correspond to Hermitian operators, which is not the case. On the other hand, if one decides to use Hermitian as a synonym of self-adjoint, it is then not true that all symmetric operators are Hermitian. In order to prevent confusion, we will avoid the term Hermitian altogether.

**Proposition 7.8.** *If  $A$  is symmetric, then  $A \subseteq A^*$ .*

*Proof.* Let  $\psi \in \mathcal{D}_A$  and let  $\eta := A\psi$ . Then, by symmetry, we have

$$\forall \alpha \in \mathcal{D}_A : \langle \psi | A\alpha \rangle = \langle \eta | \alpha \rangle$$

and hence  $\psi \in \mathcal{D}_{A^*}$ . Therefore,  $\mathcal{D}_A \subseteq \mathcal{D}_{A^*}$  and  $A^*\psi := \eta = A\psi$ . □

**Definition.** A densely defined operator  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  is *self-adjoint* if  $A = A^*$ . That is,

$$(i) \quad \mathcal{D}_A = \mathcal{D}_{A^*}$$

$$(ii) \quad \forall \alpha \in \mathcal{D}_A : A\alpha = A^*\alpha.$$

◇

*Remark 7.9.* Observe that any self-adjoint operator is also symmetric, but a symmetric operator need not be self-adjoint.

**Corollary 7.10.** *A self-adjoint operator is maximal with respect to self-adjoint extension.*

*Proof.* Let  $A, B$  be self-adjoint and suppose that  $A \subseteq B$ . Then

$$A \subseteq B = B^* \subseteq A^* = A$$

and hence,  $B = A$ . □

In fact, self-adjoint operators are maximal even with respect to symmetric extension, for we would have  $B \subseteq B^*$  instead of  $B = B^*$  in the above equation.

### 7.3 Closability, closure, closedness

**Definition.** (i) A densely defined operator  $A$  is called *closable* if its adjoint  $A^*$  is also densely defined.

(ii) The *closure* of a closable operator is  $\overline{A} := A^{**} = (A^*)^*$ .

(iii) An operator is called *closed* if  $A = \overline{A}$ .

◇

*Remark 7.11.* Note that we have used the overline notation in several contexts with different meanings. When applied to complex numbers, it denotes complex conjugation. When applied to subsets of a topological space, it denotes their topological closure. Finally, when applied to (closable) operators, it denotes their closure as defined above.

**Proposition 7.12.** *A symmetric operator is necessarily closable.*

*Proof.* Let  $A$  be symmetric. Then,  $A \subseteq A^*$  and hence,  $\mathcal{D}_A \subseteq \mathcal{D}_{A^*}$ . Since a symmetric operators are densely defined, we have

$$\mathcal{H} = \overline{\mathcal{D}_A} \subseteq \overline{\mathcal{D}_{A^*}} \subseteq \mathcal{H}.$$

Hence,  $\overline{\mathcal{D}_{A^*}} = \mathcal{H}$ .

□

Note carefully that the adjoint of a symmetric operator need *not* be symmetric. In particular, we cannot conclude that  $A^* \subseteq A^{**}$ . In fact, the reversed inclusion holds.

**Proposition 7.13.** *If  $A$  is symmetric, then  $A^{**} \subseteq A^*$ .*

*Proof.* Since  $A$  is symmetric, we have  $A \subseteq A^*$ . Hence,  $A^{**} \subseteq A^*$  by Proposition 7.6.

□

**Lemma 7.14.** *For any closable operator  $A$ , we have  $A \subseteq A^{**}$ .*

*Proof.* Recall that  $\mathcal{D}_{A^*} := \{\psi \in \mathcal{H} \mid \forall \alpha \in \mathcal{D}_A : \langle \psi | A\alpha \rangle = \langle \eta | \alpha \rangle\}$ . Then

$$\forall \psi \in \mathcal{D}_{A^*} : \forall \alpha \in \mathcal{D}_A : \langle \psi | A\alpha \rangle = \langle A^*\psi | \alpha \rangle.$$

Since  $\psi$  and  $\alpha$  above are “dummy variables”, and the order of quantifiers of the same type is immaterial, we have

$$\forall \psi \in \mathcal{D}_A : \forall \alpha \in \mathcal{D}_{A^*} : \langle \alpha | A\psi \rangle = \langle A^*\alpha | \psi \rangle.$$

Then, taking the conjugate on both inner product terms and moving them across the equal sign, we have

$$\forall \psi \in \mathcal{D}_A : \forall \alpha \in \mathcal{D}_{A^*} : \langle \psi | A^*\alpha \rangle = \langle A\psi | \alpha \rangle.$$

Now, letting  $\eta := A\psi \in \mathcal{H}$  we see that  $\psi \in \mathcal{D}_{A^{**}}$ , and so  $\mathcal{D}_A \subseteq \mathcal{D}_{A^{**}}$ . Moreover, by definition,  $A^{**}\psi := \eta := A\psi$  for all  $\psi \in \mathcal{D}_A$ , and thus  $A \subseteq A^{**}$ .

□

**Corollary 7.15.** *If  $A$  is symmetric, then  $A \subseteq \overline{A} \subseteq A^*$ .*



**Corollary 7.16.** *If  $A$  is symmetric, then  $\overline{A}$  is symmetric.*

**Theorem 7.17.** *Let  $A$  be a densely defined operator.*

- (i) *The operator  $A^*$  is closed*
- (ii) *If  $A$  is invertible, we have  $(A^{-1})^* = (A^*)^{-1}$*
- (iii) *If  $A$  is invertible and closable and  $\overline{A}$  is injective, then  $\overline{A^{-1}} = \overline{A}^{-1}$ .*

These theorems are proved using a graphical formulation which has not been presented here, so in order to save space all the proofs are not given. However as a example of the technique, we shall provide the proof for (i).<sup>11</sup>

*Proof.* Let  $\mathcal{H}$  and  $\mathcal{K}$  be two Hilbert spaces and let  $A : \mathcal{H} \rightarrow \mathcal{K}$  be densely defined. We define the *graph* of  $A$  as

$$\Gamma(A) := \{(h, Ah) \mid h \in \mathcal{D}_A\}.$$

In this context, we say  $A$  is closed if and only if  $\Gamma(A)$  is closed w.r.t. the product topology on  $\mathcal{H} \oplus \mathcal{K}$ . Next define the operator

$$\begin{aligned} \mathcal{J} : \mathcal{H} \oplus \mathcal{K} &\rightarrow \mathcal{H} \oplus \mathcal{K} \\ h \oplus k &\mapsto (-k) \oplus h. \end{aligned}$$

We now wish to show that  $\Gamma(A^*) \cong [\mathcal{J}(\Gamma(A))]^\perp$ , as the right hand side is closed due to the orthogonal projection, which gives us that  $A^*$  is closed.

( $\Rightarrow$ ) Let  $x \in \mathcal{D}_A$  and  $y \in \mathcal{D}_{A^*}$ . We have  $y \oplus A^*(y) \in \Gamma(A^*)$ , and

$$\begin{aligned} \langle y \oplus A^*(y), \mathcal{J}(x \oplus A(x)) \rangle &= \langle y \oplus A^*(y), -A(x) \oplus x \rangle \\ &= -\langle y, A(x) \rangle + \langle A^*(y), x \rangle \\ &= -\langle y, A(x) \rangle + \langle y, A(x) \rangle \\ &= 0 \end{aligned}$$

and so  $\Gamma(A^*) \subseteq [\mathcal{J}(\Gamma(A))]^\perp$ .

( $\Leftarrow$ ) Let  $x \in \mathcal{D}_A$  and  $y \oplus z \in [\mathcal{J}(\Gamma(A))]^\perp$ . Then,

$$\begin{aligned} \langle y \oplus z, -A(x) \oplus x \rangle &= 0 \\ \therefore \langle y, A(x) \rangle &= \langle z, x \rangle, \end{aligned}$$

which, from the definition of the adjoint, tells us that  $y \in \mathcal{D}_{A^*}$  and  $z = A^*(y)$  and so  $y \oplus z \in \Gamma(A^*)$  and  $[\mathcal{J}(\Gamma(A))]^\perp \subseteq \Gamma(A^*)$ .

□

---

<sup>11</sup>Many thanks to Alfredo Sepulveda-Ximenez for providing [a brilliant answer to this on Quora](#).

## 7.4 Essentially self-adjoint operators

Usually, checking that an operator is symmetric is easy. By contrast, checking that an operator is self-adjoint (directly from the definition) requires the construction of the adjoint, which is not always easy. However, since every self-adjoint operator is symmetric, we can first check the symmetry property, and then determine criteria for when a symmetric operator is self-adjoint, or for when a self-adjoint extension exists.

Two complications with the extension approach are that, given a symmetric operator, there could be *no* self-adjoint extension at all, or there may be *several* different self-adjoint extensions. There is, however, a class of operators for which the situation is much nicer.

**Definition.** A symmetric operator  $A$  is called *essentially self-adjoint* if  $\overline{A}$  is self-adjoint.  $\diamond$

This is weaker than the self-adjointness condition.

**Proposition 7.18.** *If  $A$  is self-adjoint, then it is essentially self-adjoint.*

*Proof.* If  $A = A^*$ , then  $A \subseteq A^*$  and  $A^* \subseteq A$ . Hence,  $A^{**} \subseteq A^*$  and  $A^* \subseteq A^{**}$ , so  $A^* = A^{**}$ . Similarly, we have  $A^{**} = A^{***}$ , which is just  $\overline{A} = \overline{A}^*$ .  $\square$

**Theorem 7.19.** *If  $A$  is essentially self-adjoint, then there exists a unique self-adjoint extension of  $A$ , namely  $\overline{A}$ .*

*Proof.* (i) Since  $A$  is symmetric, it is closable and hence  $\overline{A}$  exists.

(ii) By Lemma 7.14, we have  $A \subseteq \overline{A}$ , so  $\overline{A}$  is an extension of  $A$ .

(iii) Suppose that  $B$  is another self-adjoint extension of  $A$ . Then,  $A \subseteq B = B^*$ , hence  $B^{**} \subseteq A^*$ , and thus  $A^{**} \subseteq B^{***} = B$ . This means that  $\overline{A} \subseteq B$ , i.e.  $B$  is a self-adjoint extension of the self-adjoint operator  $\overline{A}$ . Hence,  $B = \overline{A}$  by Corollary 7.10.  $\square$

**Remark 7.20.** One may get the feeling at this point that checking for essential self-adjointness of an operator  $A$ , i.e. checking that  $A^{**} = A^{***}$ , is hardly easier than checking whether  $A$  is self-adjoint, that is, whether  $A = A^*$ . However, this is not so. While we will show below that there is a sufficient criterion for self-adjointness which does not require to calculate the adjoint, we will see that there is, in fact, a necessary and sufficient criterion to check for essential self-adjointness of an operator without calculating a single adjoint.

**Remark 7.21.** If a symmetric operator  $A$  fails to even be essentially self-adjoint, then there is either no self-adjoint extension of  $A$  or there are several.

**Definition.** Let  $A$  be a densely defined operator. The *defect indices* of  $A$  are

$$d_+ := \dim(\ker(A^* - i)), \quad d_- := \dim(\ker(A^* + i)),$$

where by  $A^* \pm i$  we mean, of course,  $A^* \pm i \cdot \text{id}_{\mathcal{D}_{A^*}}$ .  $\diamond$

**Theorem 7.22.** *A symmetric operator has a self-adjoint extension if its defect indices coincide. Otherwise, there exist no self-adjoint extension.*

**Remark 7.23.** We will later see that if  $d_+ = d_- = 0$ , then  $A$  is essentially self-adjoint.

## 7.5 Criteria for self-adjointness and essential self-adjointness

**Theorem 7.24.** *A symmetric operator  $A$  is self-adjoint if (but not only if)*

$$\exists z \in \mathbb{C} : \operatorname{ran}(A + z) = \mathcal{H} = \operatorname{ran}(A + \bar{z}).$$

*Proof.* Since  $A$  is symmetric, by Proposition 7.8, we have  $A \subseteq A^*$ . Hence, it remains to be shown that  $A^* \subseteq A$ . To that end, let  $\psi \in \mathcal{D}_{A^*}$  and let  $z \in \mathbb{C}$ . Clearly,

$$A^*\psi + \bar{z}\psi \in \mathcal{H}.$$

Now suppose that  $z$  satisfies the hypothesis of the theorem. Then, as  $\operatorname{ran}(A + \bar{z}) = \mathcal{H}$ ,

$$\exists \alpha \in \mathcal{D}_A : A^*\psi + \bar{z}\psi = (A + \bar{z})\alpha.$$

By using the symmetry of  $A$ , we have that, for any  $\beta \in \mathcal{D}_A$ ,

$$\begin{aligned} \langle \psi | (A + z)\beta \rangle &= \langle (A + z)^*\psi | \beta \rangle \\ &= \langle A^*\psi + \bar{z}\psi | \beta \rangle \\ &= \langle (A + \bar{z})\alpha | \beta \rangle \\ &= \langle A\alpha | \beta \rangle + z\langle \alpha | \beta \rangle \\ &= \langle \alpha | A\beta \rangle + \langle \alpha | z\beta \rangle \\ &= \langle \alpha | (A + z)\beta \rangle. \end{aligned}$$

Since  $\beta \in \mathcal{D}_A$  was arbitrary and  $\operatorname{ran}(A + z) = \mathcal{H}$ , we have

$$\forall \varphi \in \mathcal{H} : \langle \psi | \varphi \rangle = \langle \alpha | \varphi \rangle.$$

Hence, by positive-definiteness of the inner product, we have  $\psi = \alpha$ , thus  $\psi \in \mathcal{D}_A$  and therefore,  $A^* \subseteq A$ .  $\square$

**Theorem 7.25.** *A symmetric operator  $A$  is essentially self-adjoint if, and only if,*

$$\exists z \in \mathbb{C} \setminus \mathbb{R} : \overline{\operatorname{ran}(A + z)} = \mathcal{H} = \overline{\operatorname{ran}(A + \bar{z})}.$$

The following criterion for essential self-adjointness, which does require the calculation of  $A^*$ , is equivalent to the previous result and, in some situations, it can be easier to check.

**Theorem 7.26.** *A symmetric operator  $A$  is essentially self-adjoint if, and only if,*

$$\exists z \in \mathbb{C} \setminus \mathbb{R} : \ker(A^* + z) = \{0\} = \ker(A^* + \bar{z}).$$

*Proof.* We show that this is equivalent to the previous condition. Recall that if  $\mathcal{M}$  is a linear subspace of  $\mathcal{H}$ , then  $\mathcal{M}^\perp$  is closed and hence,  $\mathcal{M}^{\perp\perp} = \overline{\mathcal{M}}$  (Proposition 4.7). Thus, by Proposition 7.5, we have

$$\overline{\operatorname{ran}(A + z)} = \operatorname{ran}(A + z)^{\perp\perp} = \ker(A^* + \bar{z})^\perp$$

and, similarly,

$$\overline{\operatorname{ran}(A + \bar{z})} = \ker(A^* + z)^\perp.$$

Since  $\mathcal{H}^\perp = \{0\}$ , the above condition is equivalent to

$$\overline{\operatorname{ran}(A + z)} = \mathcal{H} = \overline{\operatorname{ran}(A + \bar{z})}. \quad \square$$

## 8 Spectra and perturbation theory

We will now focus on the spectra of operators and on the decomposition of the spectra of self-adjoint operators. The significance of spectra is that the axioms of quantum mechanics prescribe that the possible measurement values of an observable (which is, in particular, a self-adjoint operator) are those in the so-called spectrum of the operator.

A common task in almost any quantum mechanical problem that you might wish to solve is to determine the spectrum of some observable. This is usually the Hamiltonian, or energy operator, since the time evolution of a quantum system is governed by the exponential of the Hamiltonian, which is more practically determined by first determining its spectrum.

More often than not, it is not possible to determine the spectrum of an operator exactly (i.e. analytically). One then resorts to perturbation theory which consists in expressing the operator whose spectrum we want to determine as the sum of an operator whose spectrum can be determined analytically and another whose contribution is “small” in some sense to be made precise.

### 8.1 Resolvent map and spectrum

**Definition.** The *resolvent map* of an operator  $A$  is the map

$$R_A: \rho(A) \rightarrow \mathcal{L}(\mathcal{H})$$

$$z \mapsto (A - z)^{-1},$$

where  $\mathcal{L}(\mathcal{H}) \equiv \mathcal{L}(\mathcal{H}, \mathcal{H})$  and  $\rho(A)$  is the *resolvent set* of  $A$ , defined as

$$\rho(A) := \{z \in \mathbb{C} \mid (A - z)^{-1} \in \mathcal{L}(\mathcal{H})\}.$$

◇

*Remark 8.1.* Checking whether a complex number  $z$  belongs to  $\rho(A)$  may seem like a daunting task and, in general, it is. However, we will almost exclusively be interested in closed operators, and the closed graph theorem states that if  $A$  is closed, then  $(A - z)^{-1} \in \mathcal{L}(\mathcal{H})$  if, and only if,  $A - z$  is bijective.

**Definition.** The *spectrum* of an operator  $A$  is  $\sigma(A) := \mathbb{C} \setminus \rho(A)$ .

◇

**Definition.** A complex number  $\lambda \in \mathbb{C}$  is said to be an *eigenvalue* of  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  if

$$\exists \psi \in \mathcal{D}_A \setminus \{0\} : A\psi = \lambda\psi.$$

Such an element  $\psi$  is called an *eigenvector* of  $A$  associated to the eigenvalue  $\lambda$ .

◇

**Corollary 8.2.** Let  $\lambda \in \mathbb{C}$  be an eigenvalue of  $A$ . Then,  $\lambda \in \sigma(A)$ .

*Proof.* If  $\lambda$  is an eigenvalue of  $A$ , then there exists  $\psi \in \mathcal{D}_A \setminus \{0\}$  such that  $A\psi = \lambda\psi$ , i.e.

$$(A - \lambda)\psi = 0.$$

Thus,  $\psi \in \ker(A - \lambda)$  and hence, since  $\psi \neq 0$ , we have

$$\ker(A - \lambda) \neq \{0\}.$$

This means that  $A - \lambda$  is not injective, hence not invertible and thus,  $\lambda \notin \rho(A)$ . Then, by definition,  $\lambda \in \sigma(A)$ .  $\square$

*Remark 8.3.* If  $\mathcal{H}$  is finite-dimensional, then the converse of the above corollary holds and hence, the spectrum coincides with the set of eigenvalues. However, in infinite-dimensional spaces, the spectrum of an operator contains more than just the eigenvalues of the operator.

## 8.2 The spectrum of a self-adjoint operator

Recall that a self-adjoint operator is necessarily closed since  $A = A^*$  implies  $A = A^{**}$ . While the following refinement of the notion of spectrum can be made in greater generality, we will primarily be interested in the case of self-adjoint operators.

**Definition.** Let  $A$  be a self-adjoint operator. Then, we define

- (i) the *pure point spectrum* of  $A$

$$\sigma_{\text{pp}}(A) := \{z \in \mathbb{C} \mid \text{ran}(A - z) = \overline{\text{ran}(A - z)} \neq \mathcal{H}\}$$

- (ii) the *point embedded in continuum spectrum* of  $A$

$$\sigma_{\text{pec}}(A) := \{z \in \mathbb{C} \mid \text{ran}(A - z) \neq \overline{\text{ran}(A - z)} \neq \mathcal{H}\}$$

- (iii) the *purely continuous spectrum* of  $A$

$$\sigma_{\text{pc}}(A) := \{z \in \mathbb{C} \mid \text{ran}(A - z) \neq \overline{\text{ran}(A - z)} = \mathcal{H}\}.$$

$\diamond$

These form a partition of  $\sigma(A)$ , i.e. they are pairwise disjoint and their union is  $\sigma(A)$ .

**Definition.** Let  $A$  be a self-adjoint operator. Then, we further define

- (i) the *point spectrum* of  $A$

$$\sigma_{\text{p}}(A) := \sigma_{\text{pp}}(A) \cup \sigma_{\text{pec}}(A) = \{z \in \mathbb{C} \mid \overline{\text{ran}(A - z)} \neq \mathcal{H}\}$$

- (ii) the *continuous spectrum* of  $A$

$$\sigma_{\text{c}}(A) := \sigma_{\text{pec}}(A) \cup \sigma_{\text{pc}}(A) = \{z \in \mathbb{C} \mid \text{ran}(A - z) \neq \overline{\text{ran}(A - z)}\}.$$

$\diamond$

Clearly,  $\sigma_{\text{p}}(A) \cup \sigma_{\text{c}}(A) = \sigma(A)$  but, since  $\sigma_{\text{p}}(A) \cap \sigma_{\text{c}}(A) = \sigma_{\text{pec}}(A)$  is not necessarily empty, the point and continuous spectra do not form a partition of the spectrum in general.

**Lemma 8.4.** *Let  $A$  be self-adjoint and let  $\lambda$  be an eigenvalue of  $A$ . Then,  $\lambda \in \mathbb{R}$ .*

*Proof.* Let  $\psi \in \mathcal{D}_A \setminus \{0\}$  be an eigenvector of  $A$  associated to  $\lambda$ . By self-adjointness of  $A$ ,

$$\lambda \langle \psi | \psi \rangle = \langle \psi | \lambda \psi \rangle = \langle \psi | A \psi \rangle = \langle A \psi | \psi \rangle = \langle \lambda \psi | \psi \rangle = \bar{\lambda} \langle \psi | \psi \rangle.$$

Thus, we have

$$(\lambda - \bar{\lambda}) \langle \psi | \psi \rangle = 0$$

and since  $\psi \neq 0$ , it follows that  $\lambda = \bar{\lambda}$ . That is,  $\lambda \in \mathbb{R}$ .  $\square$

**Theorem 8.5.** *If  $A$  is a self-adjoint operator, then the elements of  $\sigma_p(A)$  are precisely the eigenvalues of  $A$ .*

*Proof.* ( $\Leftarrow$ ) Suppose that  $\lambda$  is an eigenvalue of  $A$ . Then, by self-adjointness of  $A$ ,

$$\{0\} \neq \ker(A - \lambda) = \ker(A^* - \lambda) = \ker((A - \bar{\lambda})^*) = \text{ran}(A - \bar{\lambda})^\perp = \text{ran}(A - \lambda)^\perp,$$

where we made use of our previous lemma. Hence, we have

$$\overline{\text{ran}(A - \lambda)} = \text{ran}(A - \lambda)^{\perp\perp} \neq \{0\}^\perp = \mathcal{H}$$

and thus,  $\lambda \in \sigma_p(A)$ .

( $\Rightarrow$ ) We now need to show that if  $\lambda \in \sigma_p(A)$ , then  $\lambda$  is an eigenvalue of  $A$ . By contraposition, suppose that  $\lambda \in \mathbb{C}$  is not an eigenvalue of  $A$ . Note that if  $\lambda$  is real, then  $\lambda = \bar{\lambda}$  while if  $\lambda$  is not real, then  $\bar{\lambda}$  is not real. Hence, if  $\lambda$  is not an eigenvalue of  $A$ , then neither is  $\bar{\lambda}$ . Therefore, there exists no non-zero  $\psi$  in  $\mathcal{D}_A$  such that  $A\psi = \bar{\lambda}\psi$ . Thus, we have

$$\{0\} = \ker(A - \bar{\lambda}) = \ker(A^* - \bar{\lambda}) = \ker((A - \lambda)^*) = \text{ran}(A - \lambda)^\perp$$

and hence

$$\overline{\text{ran}(A - \lambda)} = \text{ran}(A - \lambda)^{\perp\perp} = \{0\}^\perp = \mathcal{H}.$$

Therefore,  $\lambda \notin \sigma_p(A)$ .  $\square$

**Remark 8.6.** The *contrapositive* of the statement  $P \Rightarrow Q$  is the statement  $\neg Q \Rightarrow \neg P$ , where the symbol  $\neg$  denotes logical negation. A statement and its contrapositive are logically equivalent and “proof by contraposition” simply means “proof of the contrapositive”.

### 8.3 Perturbation theory for point spectra of self-adjoint operators

Before we move on to perturbation theory, we will need some preliminary definitions. First, note that if  $\psi$  and  $\varphi$  are both eigenvectors of an operator  $A$  associated to some eigenvalue  $\lambda$ , then, for any  $z \in \mathbb{C}$ , the vector  $z\psi + \varphi$  is either zero or it is again an eigenvector of  $A$  associated to  $\lambda$ .

**Definition.** Let  $A$  be an operator and let  $\lambda$  be an eigenvalue of  $A$ .

(i) The *eigenspace* of  $A$  associated to  $\lambda$  is

$$\text{Eig}_A(\lambda) := \{\psi \in \mathcal{D}_A \mid A\psi = \lambda\psi\}.$$

(ii) The eigenvalue  $\lambda$  is said to be *non-degenerate* if  $\dim \text{Eig}_A(\lambda) = 1$ , and *degenerate* if  $\dim \text{Eig}_A(\lambda) > 1$ .

(iii) The *degeneracy* of  $\lambda$  is  $\dim \text{Eig}_A(\lambda)$ .

◇

*Remark 8.7.* Of course, it is possible that  $\dim \text{Eig}_A(\lambda) = \infty$  in general. However, in this section, we will only consider operators whose eigenspaces are finite-dimensional.

**Lemma 8.8.** *Eigenvectors associated to distinct eigenvalues of a self-adjoint operator are orthogonal.*

*Proof.* Let  $\lambda, \lambda'$  be distinct eigenvalues of a self-adjoint operator  $A$  and let  $\psi, \varphi \in \mathcal{D}_A \setminus \{0\}$  be eigenvectors associated to  $\lambda$  and  $\lambda'$ , respectively. As  $A$  is self-adjoint, we already know that  $\lambda, \lambda' \in \mathbb{R}$ . Then, note that

$$\begin{aligned} (\lambda - \lambda')\langle \psi | \varphi \rangle &= \lambda \langle \psi | \varphi \rangle - \lambda' \langle \psi | \varphi \rangle \\ &= \langle \lambda \psi | \varphi \rangle - \langle \psi | \lambda' \varphi \rangle \\ &= \langle A\psi | \varphi \rangle - \langle \psi | A\varphi \rangle \\ &= \langle \psi | A\varphi \rangle - \langle \psi | A\varphi \rangle \\ &= 0. \end{aligned}$$

Since  $\lambda - \lambda' \neq 0$ , we must have  $\langle \psi | \varphi \rangle = 0$ . □

## A. Unperturbed spectrum

Let  $H_0$  be a self-adjoint operator whose eigenvalues and eigenvectors are known and satisfy

$$H_0 e_{n\delta} = h_n e_{n\delta},$$

where

- the index  $n$  varies either over  $\mathbb{N}$  or some finite range  $1, 2, \dots, N$
- the real numbers  $h_n$  are the eigenvalues of  $H_0$
- the index  $\delta$  varies over the range  $1, 2, \dots, d(n)$ , with  $d(n) := \dim \text{Eig}_{H_0}(h_n)$
- for each fixed  $n$ , the set

$$\{e_{n\delta} \in \mathcal{D}_{H_0} \mid 1 \leq \delta \leq d(n)\}$$

is a linearly independent subset (in fact, a Hamel basis) of  $\text{Eig}_{H_0}(h_n)$ .

Note that, since we are assuming that all eigenspaces of  $H_0$  are finite-dimensional,  $\text{Eig}_{H_0}(h_n)$  is a sub-Hilbert space of  $\mathcal{H}$  and hence, for each fixed  $n$ , we can choose the  $e_{n\delta}$  so that

$$\langle e_{n\alpha} | e_{n\beta} \rangle = \delta_{\alpha\beta}.$$

In fact, thanks to our previous lemma, we can choose the eigenvectors of  $H_0$  so that

$$\langle e_{n\alpha} | e_{m\beta} \rangle = \delta_{nm} \delta_{\alpha\beta}.$$

Let  $W: \mathcal{D}_{H_0} \rightarrow \mathcal{H}$  be a not necessarily self-adjoint operator. Let  $\lambda \in (-\varepsilon, \varepsilon) \subseteq \mathbb{R}$  and consider the real one-parameter family of operators  $\{H_\lambda \mid \lambda \in (-\varepsilon, \varepsilon)\}$ , where

$$H_\lambda := H_0 + \lambda W.$$

Further assume that  $H_\lambda$  is self-adjoint for all  $\lambda \in (-\varepsilon, \varepsilon)$ . Recall, however, that this assumption does *not* force  $W$  to be self-adjoint.

We seek to understand the eigenvalue equation for  $H_\lambda$ ,

$$H_\lambda e_{n\delta}(\lambda) = h_{n\delta}(\lambda) e_{n\delta}(\lambda),$$

by exploiting the fact that it coincides with the eigenvalue equation for  $H_0$  when  $\lambda = 0$ . In particular, we will be interested in the lifting of the degeneracy of  $h_n$  (for some fixed  $n$ ) once the perturbation  $W$  is “switched on”, i.e. when  $\lambda \neq 0$ . Indeed, it is possible, for instance, that while the two eigenvectors  $e_{n1}$  and  $e_{n2}$  are associated to the same (degenerate) eigenvalue  $h_n$  of  $H_0$ , the “perturbed” eigenvectors  $e_{n1}(\lambda)$  and  $e_{n2}(\lambda)$  may be associated to *different* eigenvalues of  $H_\lambda$ . Hence the reason why we added a  $\delta$ -index to the eigenvalue in the above equation. Of course, when  $\lambda = 0$ , we have  $h_{n\delta}(\lambda) = h_n$  for all  $\delta$ .

## B. Formal power series ansatz<sup>12</sup>

In order to determine  $h_{n\delta}(\lambda)$  and  $e_{n\delta}(\lambda)$ , we make, for both, the following ansatz

$$\begin{aligned} h_{n\delta}(\lambda) &=: h_n + \lambda \theta_{n\delta}^{(1)} + \lambda^2 \theta_{n\delta}^{(2)} + \mathcal{O}(\lambda^3) \\ e_{n\delta}(\lambda) &=: e_{n\delta} + \lambda \epsilon_{n\delta}^{(1)} + \lambda^2 \epsilon_{n\delta}^{(2)} + \mathcal{O}(\lambda^3), \end{aligned}$$

where  $\theta_{n\delta}^{(1)}, \theta_{n\delta}^{(2)} \in \mathbb{R}$  and  $\epsilon_{n\delta}^{(1)}, \epsilon_{n\delta}^{(2)} \in \mathcal{D}_{H_0}$ .

*Remark 8.9.* Recall that the *Big O notation* is defined as follows. If  $f$  and  $g$  are functions  $I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  and  $a \in I$ , then we write

$$f(x) = \mathcal{O}(g(x)) \quad \text{as } x \rightarrow a$$

to mean

$$\exists k, M > 0 : \forall x \in I : 0 < |x - a| < k \Rightarrow |f(x)| < M|g(x)|.$$

The qualifier “as  $x \rightarrow a$ ” can be omitted when the value of  $a$  is clear from the context. In our expressions above, we obviously have “as  $\lambda \rightarrow 0$ ”.

---

<sup>12</sup>German for “educated guess”.



### C. Fixing phase and normalisation of perturbed eigenvectors

Eigenvectors in a complex vector space are only defined up to a complex scalar or, alternatively, up to phase and magnitude. Hence, we impose the following conditions relating the perturbed eigenvalues and eigenvectors to the unperturbed ones.

We require, for all  $\lambda \in (-\varepsilon, \varepsilon)$ , all  $n$  and all  $\delta$ ,

$$(i) \quad \text{Im}\langle e_{n\delta} | e_{n\delta}(\lambda) \rangle = 0$$

$$(ii) \quad \|e_{n\delta}(\lambda)\|^2 = 1.$$

Inserting the formal power series ansatz into these conditions yields

$$(i) \quad \text{Im}\langle e_{n\delta} | \epsilon_{n\delta}^{(k)} \rangle = 0 \text{ for } k = 1, 2, \dots$$

$$(ii) \quad 0 = 2\lambda \text{Re}\langle e_{n\delta} | \epsilon_{n\delta}^{(1)} \rangle + \lambda^2 (2 \text{Re}\langle e_{n\delta} | \epsilon_{n\delta}^{(2)} \rangle + \|\epsilon_{n\delta}^{(1)}\|^2) + \mathcal{O}(\lambda^3).$$

Since (ii) holds for all  $\lambda \in (-\varepsilon, \varepsilon)$ , we must have

$$\text{Re}\langle e_{n\delta} | \epsilon_{n\delta}^{(1)} \rangle = 0, \quad 2 \text{Re}\langle e_{n\delta} | \epsilon_{n\delta}^{(2)} \rangle + \|\epsilon_{n\delta}^{(1)}\|^2 = 0.$$

Since we know from (i) that  $\text{Im}\langle e_{n\delta} | \epsilon_{n\delta}^{(1)} \rangle = 0$  and  $\text{Im}\langle e_{n\delta} | \epsilon_{n\delta}^{(2)} \rangle = 0$ , we can conclude

$$\langle e_{n\delta} | \epsilon_{n\delta}^{(1)} \rangle = 0, \quad \langle e_{n\delta} | \epsilon_{n\delta}^{(2)} \rangle = -\frac{1}{2} \|\epsilon_{n\delta}^{(1)}\|^2.$$

That is,  $\epsilon_{n\delta}^{(1)}$  is orthogonal to  $e_{n\delta}$  and

$$\epsilon_{n\delta}^{(2)} = -\frac{1}{2} \|\epsilon_{n\delta}^{(1)}\|^2 e_{n\delta} + e$$

for some  $e \in \text{span}(\{e_{n\delta}\})^\perp$ .

### D. Order-by-order decomposition of the perturbed eigenvalue problem

Let us insert our formal power series ansatz into the perturbed eigenvalue equation. On the left-hand side, we find

$$\begin{aligned} H_\lambda e_{n\delta}(\lambda) &= (H_0 + \lambda W)(e_{n\delta} + \lambda \epsilon_{n\delta}^{(1)} + \lambda^2 \epsilon_{n\delta}^{(2)} + \mathcal{O}(\lambda^3)) \\ &= H_0 e_{n\delta} + \lambda(W e_{n\delta} + H_0 \epsilon_{n\delta}^{(1)}) + \lambda^2(W \epsilon_{n\delta}^{(1)} + H_0 \epsilon_{n\delta}^{(2)}) + \mathcal{O}(\lambda^3), \end{aligned}$$

while, on the right-hand side, we have

$$\begin{aligned} h_{n\delta}(\lambda) e_{n\delta}(\lambda) &= (h_n + \lambda \theta_{n\delta}^{(1)} + \lambda^2 \theta_{n\delta}^{(2)} + \mathcal{O}(\lambda^3))(e_{n\delta} + \lambda \epsilon_{n\delta}^{(1)} + \lambda^2 \epsilon_{n\delta}^{(2)} + \mathcal{O}(\lambda^3)) \\ &= h_n e_{n\delta} + \lambda(h_n \epsilon_{n\delta}^{(1)} + \theta_{n\delta}^{(1)} e_{n\delta}) + \lambda^2(h_n \epsilon_{n\delta}^{(2)} + \theta_{n\delta}^{(1)} \epsilon_{n\delta}^{(1)} + \theta_{n\delta}^{(2)} e_{n\delta}) + \mathcal{O}(\lambda^3). \end{aligned}$$

Comparing terms order-by-order yields

$$\begin{aligned} (H_0 - h_n) e_{n\delta} &= 0 \\ (H_0 - h_n) \epsilon_{n\delta}^{(1)} &= -(W - \theta_{n\delta}^{(1)}) e_{n\delta} \\ (H_0 - h_n) \epsilon_{n\delta}^{(2)} &= -(W - \theta_{n\delta}^{(1)}) \epsilon_{n\delta}^{(1)} + \theta_{n\delta}^{(2)} e_{n\delta}. \end{aligned}$$

Of course, one may continue this expansion up to the desired order. Note that the zeroth order equation is just our unperturbed eigenvalue equation.

### E. First-order correction

To extract information from the first-order equation, let us project both sides onto the unperturbed eigenvectors  $e_{n\alpha}$  (i.e. apply  $\langle e_{n\alpha} | \cdot \rangle$  to both sides). This yields

$$\langle e_{n\alpha} | (H_0 - h_n) \epsilon_{n\delta}^{(1)} \rangle = -\langle e_{n\alpha} | (W - \theta_{n\delta}^{(1)}) e_{n\delta} \rangle.$$

By self-adjointness of  $H_0$ , we have

$$\langle e_{n\alpha} | (H_0 - h_n) \epsilon_{n\delta}^{(1)} \rangle = \langle (H_0 - h_n)^* e_{n\alpha} | \epsilon_{n\delta}^{(1)} \rangle = \langle (H_0 - h_n) e_{n\alpha} | \epsilon_{n\delta}^{(1)} \rangle = 0.$$

Therefore,

$$0 = -\langle e_{n\alpha} | W e_{n\delta} \rangle + \langle e_{n\alpha} | \theta_{n\delta}^{(1)} e_{n\delta} \rangle = -\langle e_{n\alpha} | W e_{n\delta} \rangle + \theta_{n\delta}^{(1)} \delta_{\alpha\delta}$$

and thus, the first-order eigenvalue correction is

$$\theta_{n\delta}^{(1)} = \langle e_{n\delta} | W e_{n\delta} \rangle.$$

Note that the right-hand side of the first-order equation is now completely known and hence, if  $H_0 - h_n$  were invertible, we could determine  $\epsilon_{n\delta}^{(1)}$  immediately. However, this is only possible if the unperturbed eigenvalue  $h_n$  is non-degenerate. More generally, we proceed as follows. Let  $E := \text{Eig}_{H_0}(h_n)$ . Then, we can rewrite the right-hand side of the first-order equation as

$$\begin{aligned} -(W - \theta_{n\delta}^{(1)}) e_{n\delta} &= -\text{id}_{\mathcal{H}}(W - \theta_{n\delta}^{(1)}) e_{n\delta} \\ &= -(\text{P}_E + \text{P}_{E^\perp})(W - \theta_{n\delta}^{(1)}) e_{n\delta} \\ &= -\sum_{\beta=1}^{d(n)} \langle e_{n\beta} | (W - \theta_{n\delta}^{(1)}) e_{n\delta} \rangle e_{n\beta} - \text{P}_{E^\perp} W e_{n\delta} + \theta_{n\delta}^{(1)} \text{P}_{E^\perp} e_{n\delta} \\ &= -\text{P}_{E^\perp} W e_{n\delta} \end{aligned}$$

so that we have  $(H_0 - h_n) \epsilon_{n\delta}^{(1)} \in E^\perp$ . Note that the operator

$$\text{P}_{E^\perp} \circ (H_0 - h_n): E^\perp \rightarrow E^\perp$$

is invertible. Hence, the equation

$$\text{P}_{E^\perp} (H_0 - h_n) \text{P}_{E^\perp} \epsilon_{n\delta}^{(1)} = -\text{P}_{E^\perp} W e_{n\delta}$$

is solved by

$$\text{P}_{E^\perp} \epsilon_{n\delta}^{(1)} = -\text{P}_{E^\perp} (H_0 - h_n)^{-1} \text{P}_{E^\perp} W e_{n\delta}.$$

The “full” eigenvector correction  $\epsilon_{n\delta}^{(1)}$  is given by

$$\text{id}_{\mathcal{H}} \epsilon_{n\delta}^{(1)} = (\text{P}_E + \text{P}_{E^\perp}) \epsilon_{n\delta}^{(1)} = \sum_{\beta=1}^{d(n)} c_{\delta\beta} e_{n\beta} - \text{P}_{E^\perp} (H_0 - h_n)^{-1} \text{P}_{E^\perp} W e_{n\delta},$$

where the coefficients  $c_{\delta\beta}$  cannot be fully determined at this order in the perturbation. What we do know is that our previous fixing of the phase and normalisation of the perturbed eigenvectors implies that  $\epsilon_{n\delta}^{(1)}$  is orthogonal to  $e_{n\delta}$ , and hence we must have  $c_{\delta\delta} = 0$ .

## F. Second-order eigenvalue correction

Here we will content ourselves with calculating the second-order correction to the eigenvalues only, since that is the physically interesting formula. As before, we proceed by projecting both sides of the second-order equation onto an unperturbed eigenvector, this time specifically  $e_{n\delta}$ . We find

$$\langle e_{n\delta} | (H_0 - h_n) \epsilon_{n\delta}^{(2)} \rangle = -\langle e_{n\delta} | W \epsilon_{n\delta}^{(1)} \rangle - \theta_{n\delta}^{(1)} \langle e_{n\delta} | \epsilon_{n\delta}^{(1)} \rangle + \theta_{n\delta}^{(2)} \langle e_{n\delta} | e_{n\delta} \rangle.$$

Noting, as before, that

$$\langle e_{n\delta} | (H_0 - h_n) \epsilon_{n\delta}^{(2)} \rangle = 0$$

and recalling that  $\langle e_{n\delta} | \epsilon_{n\delta}^{(1)} \rangle = 0$  and  $\langle e_{n\delta} | e_{n\delta} \rangle = 1$ , we have

$$\theta_{n\delta}^{(2)} = \langle e_{n\delta} | W \epsilon_{n\delta}^{(1)} \rangle.$$

Plugging in our previous expression for  $\epsilon_{n\delta}^{(1)}$  yields

$$\begin{aligned} \theta_{n\delta}^{(2)} &= \left\langle e_{n\delta} \left| W \sum_{\beta=1}^{d(n)} c_{\delta\beta} e_{n\beta} - W P_{E^\perp} (H_0 - h_n)^{-1} P_{E^\perp} W e_{n\delta} \right. \right\rangle \\ &= \sum_{\beta=1}^{d(n)} c_{\delta\beta} \langle e_{n\delta} | W e_{n\beta} \rangle - \langle e_{n\delta} | W P_{E^\perp} (H_0 - h_n)^{-1} P_{E^\perp} W e_{n\delta} \rangle \\ &= \sum_{\beta=1}^{d(n)} c_{\delta\beta} \theta_{n\beta}^{(1)} \delta_{\delta\beta} - \langle e_{n\delta} | W P_{E^\perp} (H_0 - h_n)^{-1} P_{E^\perp} W e_{n\delta} \rangle \\ &= -\langle e_{n\delta} | W P_{E^\perp} (H_0 - h_n)^{-1} P_{E^\perp} W e_{n\delta} \rangle \end{aligned}$$

since  $c_{\delta\delta} = 0$ . One can show that the eigenvectors of  $H_0$  (or any other self-adjoint operator) form an orthonormal basis of  $\mathcal{H}$ . In particular, this implies that we can decompose the identity operator on  $\mathcal{H}$  as

$$\text{id}_{\mathcal{H}} = \sum_{n=1}^{\infty} \sum_{\beta=1}^{d(n)} \langle e_{n\beta} | \cdot \rangle e_{n\beta}.$$

By inserting this appropriately into our previous expression for  $\theta_{n\delta}^{(2)}$ , we obtain

$$\theta_{n\delta}^{(2)} = - \sum_{\substack{m=1 \\ m \neq n}}^{\infty} \sum_{\beta=1}^{d(m)} \frac{|\langle e_{m\beta} | W e_{n\delta} \rangle|}{h_m - h_n}.$$

Putting everything together, we have the following second-order expansion of the perturbed eigenvalues

$$\begin{aligned} h_{n\delta}(\lambda) &= h_n + \lambda \theta_{n\delta}^{(1)} + \lambda^2 \theta_{n\delta}^{(2)} + \mathcal{O}(\lambda^3) \\ &= h_n + \lambda \langle e_{n\delta} | W e_{n\delta} \rangle - \lambda^2 \sum_{\substack{m=1 \\ m \neq n}}^{\infty} \sum_{\beta=1}^{d(m)} \frac{|\langle e_{m\beta} | W e_{n\delta} \rangle|}{h_m - h_n} + \mathcal{O}(\lambda^3). \end{aligned}$$

Remark 8.10. Note that, while the first-order correction to the perturbed  $n\delta$  eigenvalue only depends on the unperturbed  $n\delta$  eigenvalue and eigenvector, the second-order correction draws information from *all* the unperturbed eigenvalues and eigenvectors. Hence, if we try to approximate a relativistic system as a perturbation of a non-relativistic system, then the second-order corrections may be unreliable.

## 9 Case study: momentum operator

We will now put the machinery developed so far to work by considering the so-called momentum operator for two cases: a compact interval  $[a, b] \subset \mathbb{R}$ , and a circle. As the name suggests, this operator is meant to be the QM observable whose eigenvalues are the momenta of the system. It is clear, therefore, that we require (recall  $\mathcal{H} = L^2(\mathbb{R}^d)$  up to unitary equivalence)

$$\mathcal{P}: \mathcal{D}_{\mathcal{P}} \rightarrow L^2(\mathbb{R}^d)$$

to be self adjoint.

We will specialise to the case of  $d = 1$  in order to simplify things, while also demonstrating the main ideas. The concepts can be extended to higher values of  $d$ . We will also set  $\hbar = 1$  throughout this section.

### 9.1 The Momentum Operator

**Definition.** The *momentum operator* is an operator  $\mathcal{P}$  given by

$$\begin{aligned} \mathcal{P}: \mathcal{D}_{\mathcal{P}} &\rightarrow L^2(\mathbb{R}) \\ \psi &\mapsto (-i)\psi', \end{aligned}$$

where the prime indicates a derivative. ◇

The first obvious question is why does this deserve its name? (i.e. how is it related to what we know classically as the momentum?) The answer, unfortunately, can not yet be provided in full detail as it requires us to know the spectral theorem and Stone-von Neumann theorem. However these details are provided here as they will help later when discussing these theorems. For now we must just take it in faith.<sup>13</sup>

There is yet another important question we must ask: how do we choose  $\mathcal{D}_{\mathcal{P}}$ ? The immediate response might be ‘such that the derivative is square integrable.’ However, this is not good enough. We also require that  $\mathcal{P}$  be self adjoint and, as we have seen previously, the concept of self adjointness depends heavily on the domains considered.

Luckily, not all hope is lost. The method will be as follows: guess a reasonable  $\mathcal{D}_{\mathcal{P}}$  and then search for a self adjoint extension, should one exist. Before doing so, though, we will first introduce some new definitions that will prove invaluable.

### 9.2 Absolutely Continuous Functions and Sobolev Spaces

During the calculations that follow we will naturally encounter three spaces:

- (i) The space of once-continuously differential functions over some interval,  $I$ ;  $C^1(I)$ ,
- (ii) The Sobolev space  $H^1(I)$ , and
- (iii) The space of absolutely continuous functions;  $AC(I)$ .

---

<sup>13</sup>Very unlike Dr. Schuller.

As we shall see they are related via

$$C^1(I) \subseteq H^1(I) \subseteq AC(I),$$

and so they will provide a convenient way to compare the domains  $\mathcal{D}_{\mathcal{P}}$ ,  $\mathcal{D}_{\mathcal{P}^*}$ , etc, to test for self adjointness.

**Definition.** Let  $I \subset \mathbb{R}$ . A function  $\psi: I \rightarrow \mathbb{C}$  is called *absolutely continuous* if there exists a Lebesgue integrable function  $\rho: I \rightarrow \mathbb{C}$  such that

$$\psi(x) = \psi(a) + \int_a^x \rho(y) dy,$$

for all compact subsets  $[a, x] \subseteq I$ . ◇

**Corollary 9.1.** *Given a absolutely continuous function, it is clear that  $\rho =_{a.e.} \psi'$ , where the almost everywhere condition comes from the fact that a Lebesgue integral does not distinguish two elements that differ by a measure zero.*

**Definition.** The set of *absolutely continuous functions* is simply

$$AC(I) := \{\psi \in L^2(I) \mid \psi \text{ is absolutely continuous}\}.$$
◇

**Definition.** Let  $\Omega \subset \mathbb{R}$  be open, and  $\psi: \Omega \rightarrow \mathbb{C}$  be Lebesgue measurable.  $\psi$  is called *p-locally integrable* if, for  $1 \leq p \leq \infty$ ,

$$\int_K |\psi(x)|^p dx < \infty,$$

for all compact subsets  $K \subset \Omega$ . The set of all functions is

$$L^p_{loc}(\Omega) := \{\psi: \Omega \rightarrow \mathbb{C} \mid \psi \text{ measurable}, \psi|_K \in L^p(K), \forall K \subset \Omega, K \text{ compact}\}.$$
◇

Remark 9.2. For case  $p = 1$ , we just call  $\psi$  locally integrable.

**Theorem 9.3.** *Every  $\psi \in L^p(\Omega)$  for  $1 \leq p \leq \infty$  is locally integrable. In other words  $L^p(\Omega) \subset L^1_{loc}(\Omega)$ .*

**Definition.** A function  $\psi \in L^1_{loc}(\Omega)$  is called *weakly differentiable* if there exists a  $\rho \in L^1_{loc}(\Omega)$  such that

$$\int_{\Omega} \psi(x) \varphi'(x) dx = - \int_{\Omega} \rho(x) \varphi(x) dx,$$

for all  $\varphi \in C_c^\infty(\Omega)$ <sup>14</sup>. This function is known as the *weak derivative* of  $\psi$  and is denoted by  $\rho := \psi'$ . ◇

---

<sup>14</sup>The subscript indicates that  $\varphi$  vanishes at the limits of integration.

**Corollary 9.4.** Note that for any weakly differentiable function the integration by parts result,

$$\int_{\Omega} \psi(x) \varphi'(x) dx = - \int_{\Omega} \psi'(x) \varphi(x) dx,$$

holds for all  $\varphi \in C_c^\infty(\Omega)$ .

**Corollary 9.5.** Given that  $\psi \in C^k(\Omega)$ ,  $\varphi \in C_c^\infty(\Omega)$ , we can show by induction that

$$\int_{\Omega} \psi(x) \varphi^{(\alpha)}(x) = (-1)^\alpha \int_{\Omega} \psi^{(\alpha)}(x) \varphi(x),$$

where  $\psi^{(\alpha)}(x)$  means the  $\alpha$ -order derivative of  $\psi(x)$ .

*Remark 9.6.* In the above Corollary we have used the fact that we are only considering one dimensional problems here. The expression is much the same for higher dimensional problems, however one has to take into account the different derivative directions.

**Definition.** Let  $\Omega \subset \mathbb{R}$  be open,  $k \in \mathbb{N}$  and  $1 \leq p \leq \infty$ . The *Sobolev Space* is the space with set

$$W^{k,p} := \{\psi \in L^p(\Omega) \cap W^k(\Omega) \mid \psi^{(\alpha)} \in L^p(\Omega), \forall |\alpha| \leq k\},$$

where  $W^k$  is the set of all locally integrable functions that also have weak derivatives of order  $\alpha$  for all  $|\alpha| \leq k$ . We introduce the notation  $H^k(\Omega) := W^{k,2}(\Omega)$ .  $\diamond$

*Remark 9.7.* Sobolev spaces can be made into Banach spaces by equipping them with a norm and  $H^k(\Omega)$  can be made into a Hilbert space.

**Proposition 9.8.** We can rewrite the space  $H^1(\Omega)$  as

$$H^1(\Omega) := \{\psi \in AC(\Omega) \mid \psi' \in L^2(\Omega)\},$$

where  $\psi'$  denotes the normal notion of derivative.

*Proof.* See Theorem 7.13 in ‘A first Course in Sobolev Spaces, Giovanni Leoni’.  $\square$

### 9.3 Momentum Operator on a Compact Interval

Let's now consider the case where the physical space is some compact interval in  $\mathbb{R}$  (i.e. we have a particle moving along the bottom of a well). W.l.o.g. take  $[0, 2\pi] =: I$ , the justification for which will be clear from the fact that next we will consider a circle.

We now need to come up with a reasonable guess for the domain  $\mathcal{D}_{\mathcal{P}}$ . First recall

$$\begin{aligned} \mathcal{P}: \mathcal{D}_{\mathcal{P}} &\rightarrow L^2(I) \\ \psi &\mapsto (-i)\psi'. \end{aligned}$$

It is therefore reasonable to restrict ourselves to  $\psi \in C^1(I)$ . Equally, physically we expect the function to vanish at the boundaries (the walls of the well). This gives us our first guess

$$\mathcal{D}_{\mathcal{P}} := \{\psi \in C^1(I) \mid \psi(0) = 0 = \psi(2\pi)\} =: C_c^1(I).$$

The question still remains, though, as to whether  $\mathcal{P}$  is self adjoint. Recalling the results of Lecture 7, it is first instructive to see if  $\mathcal{P}$  is symmetric.

### A. Symmetric?

Let  $\psi, \varphi \in \mathcal{D}_{\mathcal{P}}$ , then

$$\begin{aligned}
\langle \psi | \mathcal{P} \varphi \rangle &= \int_0^{2\pi} dx \bar{\psi}(x) (-i) \varphi'(x) \\
&= - \int_0^{2\pi} dx (-i) \bar{\psi}'(x) \varphi(x) - i [\bar{\psi}(x) \varphi(x)]_0^{2\pi} \\
&= \int_0^{2\pi} dx \overline{(-i) \psi'(x)} \varphi(x) \\
&= \langle \mathcal{P} \psi | \varphi \rangle,
\end{aligned}$$

where integration by parts was used. So, yes  $\mathcal{P}$  is symmetric.

### B. Self Adjoint?

From above we know that  $\mathcal{P} \subseteq \mathcal{P}^*$ , and so  $\mathcal{D}_{\mathcal{P}} \subseteq \mathcal{D}_{\mathcal{P}^*}$ , so we need to ask the question of how  $\mathcal{P}^*$  behaves outside the domain  $\mathcal{D}_{\mathcal{P}}$ . The obvious answer is to just extend the definition to be

$$\begin{aligned}
\mathcal{P}^*: \mathcal{D}_{\mathcal{P}^*} &\rightarrow L^2(I) \\
\psi &\mapsto (-i) \psi'.
\end{aligned}$$

Note that the  $\psi$  here is not necessarily the same as the  $\psi$  in above. The same symbol is just used and the context tells us where it lives.

All that is left to check is the domain  $\mathcal{D}_{\mathcal{P}^*}$ . From the definition of the adjoint we have

$$\psi \in \mathcal{D}_{\mathcal{P}^*} \implies \exists \eta \in L^2(I): \forall \varphi \in \mathcal{D}_{\mathcal{P}}: \langle \psi | \mathcal{P} \varphi \rangle = \langle \eta | \varphi \rangle,$$

with  $\eta := \mathcal{P}^* \psi$ . Before proceeding further with the calculation first introduce a function  $N: I \rightarrow \mathbb{C}$  such that  $\eta =_{a.e.} N'$ . Note  $N$  is Lebesgue integrable and that the almost everywhere condition is sufficient as  $\eta$  appears in a Lebesgue integral. Therefore we have,

$$\begin{aligned}
\int_0^{2\pi} dx \bar{\psi}(x) (-i) \varphi'(x) &= \int_0^{2\pi} dx \overline{N'}(x) \varphi(x) \\
\int_0^{2\pi} dx [\bar{\psi}(x) (-i) \varphi'(x) + \overline{N}(x) \varphi'(x)] &= [\overline{N}(x) \varphi(x)]_0^{2\pi} \\
-i \int_0^{2\pi} dx \overline{[\psi(x) - iN(x)]} \varphi'(x) &= 0 \\
\langle \psi - iN | \varphi' \rangle &= 0,
\end{aligned}$$

which tells us that

$$\psi - iN \in \{\varphi' \mid \varphi \in \mathcal{D}_{\mathcal{P}}\}^{\perp}.$$

This does not appear to have got us any closer to determining the domain  $\mathcal{D}_{\mathcal{P}^*}$ . However consider the following two Lemmas.

**Lemma 9.9.**  $\{\varphi' \mid \varphi \in \mathcal{D}_{\mathcal{P}}\} = \{\xi \in C^0(I) \mid \int_0^{2\pi} \xi(x) dx = 0\}.$



*Proof.* Let  $A := \{\varphi' \mid \varphi \in \mathcal{D}_{\mathcal{P}}\}$  and  $B := \{\xi \in C^0(I) \mid \int_0^{2\pi} \xi(x)dx = 0\}$ .

Now consider a  $\varphi' \in A$ , then

$$\int_0^{2\pi} \varphi'(x)dx = [\varphi(x)]_0^{2\pi} = 0,$$

so clearly  $\xi := \varphi$  and  $A \subseteq B$ .

Now consider a  $\xi \in B$  and define

$$\varphi(x) := \int_0^x \xi(y)dy.$$

Then, since  $\xi \in C^0(I)$  it follows that  $\varphi \in C^1(I)$ . It also follows that  $\varphi(0) = 0 = \varphi(2\pi)$  and so  $\varphi' \in A$  and  $B \subseteq A$ .  $\square$

**Lemma 9.10.** *Let  $\{1\}$  denote the set consisting of the element  $1 \in L^2(I)$  with  $1(x) = 1_{\mathbb{C}}$  for all  $x \in I$ . Then*

$$\overline{\{\varphi' \mid \varphi \in \mathcal{D}_{\mathcal{P}}\}} = \{1\}^{\perp}.$$

*Proof.* From the previous Lemma we have

$$\begin{aligned} \overline{\{\varphi' \mid \varphi \in \mathcal{D}_{\mathcal{P}}\}} &= \overline{\{\xi \in C^0(I) \mid \langle 1 \mid \xi \rangle = 0\}} \\ &= \overline{\{\xi \in \overline{C^0(I)} \mid \langle 1 \mid \xi \rangle = 0\}} \\ &= \{\xi \in L^2(I) \mid \langle 1 \mid \xi \rangle = 0\} \\ &= \{1\}^{\perp}, \end{aligned}$$

where the fact that  $C^0(I)$  is dense in  $L^2(I)$  to go from the second to third line.  $\square$

Putting this all together we have

$$\begin{aligned} \psi - iN &\in A^{\perp} = \overline{A^{\perp}} \\ &= (A^{\perp})^{\perp\perp} \\ &= (A^{\perp\perp})^{\perp} \\ &= \overline{A}^{\perp} \\ &= \{1\}^{\perp\perp} \\ &= \overline{\{1\}} \\ &= \{C : I \rightarrow \mathbb{C} \mid x \mapsto C_{\mathbb{C}}\}, \end{aligned}$$

where  $C_{\mathbb{C}}$  is a constant in  $\mathbb{C}$ . Recalling that  $N$  is Lebesgue integrable we see that

$$\psi(x) = C_{\mathbb{C}} + iN(x) \in AC(I),$$

and so

$$\mathcal{D}_{\mathcal{P}^*} \subseteq AC(I).$$

Now recalling

$$\begin{aligned} \mathcal{P}^* : \mathcal{D}_{\mathcal{P}^*} &\rightarrow L^2(I) \\ \psi &\mapsto (-i)\psi', \end{aligned}$$

and using Proposition 9.8, we have

$$\mathcal{D}_{\mathcal{P}^*} \subseteq H^1(I).$$

Finally we see that because all of the integration by parts results above were of the form

$$\int_I dx \psi'(x) \varphi(x) = \int_I \psi(x) \varphi'(x),$$

for arbitrary  $\varphi \in C_c^1(I)$ . Now since  $C_c^\infty(I) \subset C_c^1(I)$ , the integrals also hold for any  $\varphi \in C_c^\infty(I)$ , but this is just the condition for weak derivative and so we see that

$$\mathcal{D}_{\mathcal{P}^*} = H^1(I),$$

and

$$\mathcal{D}_{\mathcal{P}} \subsetneq \mathcal{D}_{\mathcal{P}^*},$$

so  $\mathcal{P}$  is not self adjoint.

### C. Essentially Self Adjoint?

We have managed to show that our initial guess for  $\mathcal{P}$  is not self adjoint. The next step is to ask if there is a self adjoint extension and if this extension is unique. Recall that a symmetric operator has a unique self adjoint extension if it is essentially self adjoint (i.e.  $\overline{\mathcal{P}} := \mathcal{P}^{**}$  is self adjoint). We follow the same method as above,

$$\psi \in \mathcal{D}_{\overline{\mathcal{P}}} \implies \forall \varphi \in \mathcal{D}_{\mathcal{P}^*}: \langle \psi | \mathcal{P}^* \varphi \rangle = \langle \overline{\mathcal{P}} \psi | \varphi \rangle.$$

Now, recall that for a symmetric operator  $\mathcal{P} \subseteq \overline{\mathcal{P}} \subseteq \mathcal{P}^*$  so it's clear that

$$\langle \overline{\mathcal{P}} \psi | \varphi \rangle = \langle \mathcal{P}^* \psi | \varphi \rangle$$

in the above. Writing as integrals we have

$$\begin{aligned} \int_0^{2\pi} dx \overline{\psi}(x) \varphi'(x) &= \int_0^{2\pi} dx \overline{(-i)\psi'}(x) \varphi(x) \\ -i \int_0^{2\pi} dx [\overline{\psi}(x) \varphi'(x) - \overline{\psi}'(x) \varphi(x)] &= i [\overline{\psi}(x) \varphi(x)]_0^{2\pi} \\ 0 &= \overline{\psi}(2\pi) \varphi(2\pi) - \overline{\psi}(0) \varphi(0), \end{aligned}$$

where again integration by parts has been used. We need to be careful in what conclusions we draw from this final statement, though.  $\varphi \in \mathcal{D}_{\mathcal{P}^*} = H^1(I)$ , which places no restrictions on the values of  $\varphi$  on the boundary, nor does it make any conditions between the two values  $\varphi(0)$  and  $\varphi(2\pi)$  — they are independently arbitrary. We must, therefore, conclude that

$$\overline{\psi}(2\pi) = \psi(2\pi) = 0 = \overline{\psi}(0) = \psi(0),$$

and so, at best,

$$\mathcal{D}_{\overline{\mathcal{P}}} = \{\psi \in H^1(I) \mid \psi(2\pi) = 0 = \psi(0)\} \subsetneq \mathcal{D}_{\mathcal{P}^*},$$

and so  $\overline{\mathcal{P}} \neq \mathcal{P}^*$ , which, after taking the adjoint of both sides, tells us that  $\overline{\mathcal{P}} \neq \overline{\mathcal{P}}^*$ , and so  $\mathcal{P}$  is not even essentially self adjoint.

## D. Defect Indices

We only have one tool left to check for a self adjoint extension of  $\mathcal{P}$ , check the defect indices to see if a self adjoint extension even exists. Recall

$$d_+ := \dim(\ker(\mathcal{P}^* - i)), \quad d_- := \dim(\ker(\mathcal{P}^* + i)),$$

and a symmetric operator has a (not necessarily unique) self adjoint extension if  $d_+ = d_-$ . We therefore need to determine how many  $\psi \in \mathcal{D}_{\mathcal{P}^*}$  lie in  $\ker(\mathcal{P}^* \mp i)$ :

$$\begin{aligned} (\mathcal{P}^* \mp i)\psi &= 0 \\ -i\psi' \mp i\psi &= 0 \\ \psi(x) &= a_{\pm} e^{\mp x} \end{aligned}$$

for  $a_+, a_- \in \mathbb{C}$ . There is only one solution for each and so  $d_+ = 1 = d_-$ . We therefore know that there does exist at least one self adjoint extension of  $\mathcal{P}$ <sup>15</sup>, however we don't know the form of any of them.<sup>16</sup>

*Remark 9.11.* If instead of a compact interval we take a half line  $I = [a, \infty)$ , then  $d_+ \neq d_-$  and so there is no self adjoint extension of  $\mathcal{P}$ , meaning there is no notion of a QM momentum in this case. Note however that people often talk about free particles along an infinite line in QM, however they always require the wave function ( $\psi$ ) to vanish at  $\pm\infty$ . This is clearly just the same as taking a large, yet finite, compact interval  $I = [a, b]$ .

## 9.4 Momentum Operator on a Circle

We now want to repeat all of the above but for a circle instead of a finite line segment. Fortunately almost all the work is done, the only slight difference is in the definition of  $\mathcal{D}_{\mathcal{P}}$ :

$$\mathcal{D}_{\mathcal{P}} := \{\psi \in C^1(I) \mid \psi(2\pi) = \psi(0)\},$$

which is exactly the same as before apart from now we do not require  $\psi$  to vanish at the boundary. We still have

$$\begin{aligned} \mathcal{P}: \mathcal{D}_{\mathcal{P}} &\rightarrow L^2(\mathbb{R}) \\ \psi &\mapsto (-i)\psi', \end{aligned}$$

so it follows that  $\mathcal{P}_I \subsetneq \mathcal{P}_c$ , where the  $I$  and  $c$  denote interval and circle respectively. In other words  $\mathcal{P}_c$  is an extension of  $\mathcal{P}_I$ .

## A. Symmetric?

Repeating the steps from above it is clear that  $\mathcal{P}$  is still symmetric. Note however, it is symmetric for a different reason: before we had  $[\psi(x)\varphi(x)] = 0$  as both  $\psi$  and  $\varphi$  vanished at the limits, whereas now it holds simply because  $\psi(2\pi) = \psi(0)$  and likewise for  $\varphi$ .

---

<sup>15</sup>Whew!

<sup>16</sup>Not whew!

## B. Self Adjoint?

As before we have

$$\psi \in \mathcal{D}_{\mathcal{P}_c^*} \implies \forall \varphi \in \mathcal{D}_{\mathcal{P}_c} : \langle \psi | \mathcal{P}_c \varphi \rangle = \langle \mathcal{P}_c^* \psi | \varphi \rangle.$$

However, recalling that the adjoint flips the inequality sign we have  $\mathcal{P}_c^* \subseteq \mathcal{P}_I^*$  and therefore  $\mathcal{D}_{\mathcal{P}_c^*} \subseteq H^1(I)$ . We can, therefore, replace the unknown  $\mathcal{P}_c^*$  with the known  $\mathcal{P}_I^*$  in the final part of the above, i.e.

$$\langle \mathcal{P}_c^* \psi | \varphi \rangle = \langle \mathcal{P}_I^* \psi | \varphi \rangle.$$

Then, following the exactly as before, we arrive at

$$\begin{aligned} 0 &= i [\overline{\psi} \varphi(x)]_0^{2\pi} \\ &= [\overline{\psi}(2\pi) - \overline{\psi}(0)] \varphi(0) \\ \implies \psi(2\pi) &= \psi(0), \end{aligned}$$

giving us<sup>17</sup>

$$\mathcal{D}_{\mathcal{P}_c^*} := \{\psi \in H^1(I) \mid \psi(2\pi) = \psi(0)\} =: H_{cyc}^1(I),$$

so  $\mathcal{D}_{\mathcal{P}_c} \subsetneq \mathcal{D}_{\mathcal{P}_c^*}$ , and therefore  $\mathcal{P}_c$  is not self adjoint.

## C. Essentially Self Adjoint?

Again, as before, we have  $\mathcal{P}_c \subseteq \overline{\mathcal{P}_c} \subseteq \mathcal{P}_c^*$  and

$$\psi \in \mathcal{D}_{\overline{\mathcal{P}_c}} \implies \forall \varphi \in \mathcal{D}_{\mathcal{P}_c^*} : \langle \psi | \mathcal{P}_c^* \varphi \rangle = \langle \overline{\mathcal{P}_c} \psi | \varphi \rangle = \langle \mathcal{P}_c^* \psi | \varphi \rangle,$$

which results in

$$\begin{aligned} 0 &= i [\overline{\psi} \varphi(x)]_0^{2\pi} \\ &= [\overline{\psi}(2\pi) - \overline{\psi}(0)] \varphi(0) \\ \implies \psi(2\pi) &= \psi(0), \end{aligned}$$

and so

$$\mathcal{D}_{\overline{\mathcal{P}_c}} := H_{cyc}^1(I) = \mathcal{D}_{\mathcal{P}_c^*},$$

so we conclude that  $\mathcal{P}_c$  is essentially self adjoint and  $\overline{\mathcal{P}_c}$  is the *unique* self adjoint extension.

To summarise, we have found *the* momentum operator on a circle:

$$\begin{aligned} \mathcal{P}_{S^1} : H_{cyc}^1(I) &\rightarrow L^2(\mathbb{R}) \\ \psi &\mapsto (-i)\psi'. \end{aligned}$$

---

<sup>17</sup>Note we are OK extend the domain to all of  $H^1(I)$  provided we impose the conditions above.

## 10 Inverse spectral theorem

This section is devoted to the development of all the notions and results necessary to understand and prove the spectral theorem, stated below.

**Theorem 10.1** (Spectral theorem). *For every self-adjoint operator  $A: \mathcal{D}_A \rightarrow \mathcal{H}$  there is a unique projection-valued measure  $P_A: \sigma(\mathcal{O}_{\mathbb{R}}) \rightarrow \mathcal{L}(\mathcal{H})$  such that*

$$A = \int_{\mathbb{R}} i_{\mathbb{R}} dP_A \equiv \int_{\mathbb{R}} \lambda P_A(d\lambda),$$

where  $i_{\mathbb{R}}: \mathbb{R} \hookrightarrow \mathbb{C}$  is the inclusion of  $\mathbb{R}$  into  $\mathbb{C}$ .

While useful in theory, existence results are often of limited use in practice since they usually only tell us that something exists, and not how to construct it. However, we should note here that the proof of the spectral theorem is, in fact, constructive in nature. Hence, given any self-adjoint operator  $A$ , we will be able to explicitly determine its associated projection-valued measure  $P_A$  along the following steps.

- (i) For each  $\psi \in \mathcal{H}$ , construct the real-valued Borel measure  $\mu_{\psi}^A: \sigma(\mathcal{O}_{\mathbb{R}}) \rightarrow \mathbb{R}$  given by

$$\mu_{\psi}^A((-\infty, \lambda]) := \lim_{\delta \rightarrow 0^+} \lim_{\varepsilon \rightarrow 0^+} \int_{-\infty}^{\lambda+\delta} dt \operatorname{Im} \langle \psi | R_A(t + i\varepsilon) \psi \rangle,$$

where  $R_A: \rho(A) \rightarrow \mathcal{L}(\mathcal{H})$  is the resolvent map of  $A$ . This is known as the *Stieltjes inversion formula*. Note that while not every element in  $\sigma(\mathcal{O}_{\mathbb{R}})$  is of the form  $(-\infty, \lambda]$ , such Borel measurable sets do generate the entire  $\sigma(\mathcal{O}_{\mathbb{R}})$  via unions, intersections and set differences. Hence, the value of  $\mu_{\psi}^A(\Omega)$  for  $\Omega \in \sigma(\mathcal{O}_{\mathbb{R}})$  can be determined by applying the corresponding formulae for measures, namely  $\sigma$ -additivity, continuity from above and measure of set differences.

- (ii) For all  $\psi, \varphi \in \mathcal{H}$ , define the complex-valued Borel measure  $\mu_{\psi, \varphi}^A: \sigma(\mathcal{O}_{\mathbb{R}}) \rightarrow \mathbb{C}$  by

$$\mu_{\psi, \varphi}^A(\Omega) := \frac{1}{4}(\mu_{\psi+\varphi}^A(\Omega) - \mu_{\psi-\varphi}^A(\Omega) + i\mu_{\psi-i\varphi}^A(\Omega) - i\mu_{\psi+i\varphi}^A(\Omega)).$$

- (iii) Define the projection-valued measure  $P_A: \sigma(\mathcal{O}_{\mathbb{R}}) \rightarrow \mathcal{L}(\mathcal{H})$  by requiring  $P_A(\Omega)$ , for each  $\Omega \in \sigma(\mathcal{O}_{\mathbb{R}})$ , to be the unique map in  $\mathcal{L}(\mathcal{H})$  satisfying

$$\forall \psi, \varphi \in \mathcal{H}: \langle \psi | P_A(\Omega) \varphi \rangle = \int_{\mathbb{R}} \chi_{\Omega} d\mu_{\psi, \varphi}^A.$$

We will now make all the notions and constructions used herein precise. In fact, we will present the relevant definitions and results by taking the inverse route, starting with projection-valued measures and arriving at their associated self-adjoint operators, obtaining (and proving) what we will call the inverse spectral theorem.

## 10.1 Projection-valued measures

Projection-valued measures are, unsurprisingly, objects sharing characteristics of both measures and projection operators.

**Definition.** A map  $P: \sigma(\mathcal{O}_{\mathbb{R}}) \rightarrow \mathcal{L}(\mathcal{H})$  is called a *projection-valued measure* if it satisfies the following properties.

- (i)  $\forall \Omega \in \sigma(\mathcal{O}_{\mathbb{R}}) : P(\Omega)^* = P(\Omega)$
- (ii)  $\forall \Omega \in \sigma(\mathcal{O}_{\mathbb{R}}) : P(\Omega) \circ P(\Omega) = P(\Omega)$
- (iii)  $P(\mathbb{R}) = \text{id}_{\mathcal{H}}$
- (iv) For any pairwise disjoint sequence  $\{\Omega_n\}_{n \in \mathbb{N}}$  in  $\sigma(\mathcal{O}_{\mathbb{R}})$  and any  $\psi \in \mathcal{H}$ ,

$$\sum_{n=0}^{\infty} P(\Omega_n)\psi = P\left(\bigcup_{n=0}^{\infty} \Omega_n\right)\psi.$$

◇

*Remark 10.2.* Note in the final condition we included  $\psi \in \mathcal{H}$  as, for the case countably infinite  $n \in \mathbb{N}$  we need to check convergence, which involves using the norm. Without the  $\psi$  we would need to use the norm on  $\mathcal{L}(\mathcal{H})$  which may prove difficult. However, by including the  $\psi$  we can work with the norm on  $\mathcal{H}$  itself.

**Lemma 10.3.** Let  $P: \sigma(\mathcal{O}_{\mathbb{R}}) \rightarrow \mathcal{L}(\mathcal{H})$  be a projection-valued measure. Then, for any  $\Omega, \Omega_1, \Omega_2 \in \sigma(\mathcal{O}_{\mathbb{R}})$ ,

- (i)  $P(\emptyset) = 0$ , where by 0 we mean  $0 \in \mathcal{L}(\mathcal{H})$
- (ii)  $P(\mathbb{R} \setminus \Omega) = \text{id}_{\mathcal{H}} - P(\Omega)$
- (iii)  $P(\Omega_1 \cup \Omega_2) = P(\Omega_1) + P(\Omega_2) - P(\Omega_1 \cap \Omega_2)$
- (iv)  $P(\Omega_1 \cap \Omega_2) = P(\Omega_1) \circ P(\Omega_2)$
- (v) if  $\Omega_1 \subseteq \Omega_2$ , then  $\text{ran}(P(\Omega_1)) \subseteq \text{ran}(P(\Omega_2))$ .

*Proof.* Let  $\Omega, \Omega_1, \Omega_2 \in \sigma(\mathcal{O}_{\mathbb{R}})$ ,

(i)

$$\begin{aligned} P(\emptyset)\psi &= P(\emptyset \cup \emptyset)\psi = (P(\emptyset) + P(\emptyset))\psi = 2P(\emptyset)\psi \\ \therefore P(\emptyset)\psi &= 0_{\mathcal{H}} \implies P(\emptyset) = 0. \end{aligned}$$

(ii)

$$\begin{aligned} P(\mathbb{R}) &= P((\mathbb{R} \setminus \Omega) \cup \Omega) = P(\mathbb{R} \setminus \Omega) + P(\Omega) \\ \therefore P(\mathbb{R} \setminus \Omega) &= \text{id}_{\mathcal{H}} - P(\Omega). \end{aligned}$$

where we used the fact that  $(\mathbb{R} \setminus \Omega) \cap \Omega = \emptyset$ .

(iii)

$$P(\Omega_1) = P((\Omega_1 \cap \Omega_2) \cup (\Omega_1 \setminus \Omega_2)) = P((\Omega_1 \cap \Omega_2)) + P((\Omega_1 \setminus \Omega_2)),$$

and similarly for  $P(\Omega_2)$ . Also

$$\begin{aligned} P(\Omega_1 \setminus \Omega_2) + P(\Omega_2 \setminus \Omega_1) + P(\Omega_1 \cap \Omega_2) &= P((\Omega_1 \setminus \Omega_2) \cup (\Omega_2 \setminus \Omega_1) \cup (\Omega_1 \cap \Omega_2)) \\ &= P(\Omega_1 \cup \Omega_2). \end{aligned}$$

Putting this all together gives the result.

(iv) First consider  $\Omega_1 \cap \Omega_2 = \emptyset$ . Then, using (ii) from the definition we have

$$\begin{aligned} [P(\Omega_1 \cup \Omega_2)]^2 &= [P(\Omega_1) + P(\Omega_2)]^2 \\ P(\Omega_1 \cup \Omega_2) &= P(\Omega_1) + P(\Omega_2) + P(\Omega_1) \circ P(\Omega_2) + P(\Omega_2) \circ P(\Omega_1) \\ \therefore P(\Omega_1) \circ P(\Omega_2) &= -P(\Omega_2) \circ P(\Omega_1) \\ \implies P(\Omega_1) \circ P(\Omega_2) \circ P(\Omega_2) &= -P(\Omega_2) \circ P(\Omega_1) \circ P(\Omega_2) \\ P(\Omega_1) \circ P(\Omega_2) &= P(\Omega_2) \circ P(\Omega_2) \circ P(\Omega_1) \\ P(\Omega_1) \circ P(\Omega_2) &= P(\Omega_2) \circ P(\Omega_1) \\ \therefore P(\Omega_1) \circ P(\Omega_2) &= 0 \quad \forall \Omega_1 \cap \Omega_2 = \emptyset. \end{aligned}$$

Now from  $P(\Omega_1) = P((\Omega_1 \setminus \Omega_2) \cup (\Omega_1 \cap \Omega_2)) = P(\Omega_1 \setminus \Omega_2) + P(\Omega_1 \cap \Omega_2)$ , we have

$$\begin{aligned} P(\Omega_1) \circ P(\Omega_2) &= [P(\Omega_1 \setminus \Omega_2) + P(\Omega_1 \cap \Omega_2)] \circ [P(\Omega_2 \setminus \Omega_1) + P(\Omega_1 \cap \Omega_2)] \\ &= P(\Omega_1 \cap \Omega_2) \circ P(\Omega_1 \cap \Omega_2) \\ &= P(\Omega_1 \cap \Omega_2) \end{aligned}$$

where we have made use of the fact that  $(\Omega_1 \setminus \Omega_2) \cap (\Omega_2 \setminus \Omega_1) = \emptyset$  etc.


(v) If  $\Omega_1 \subset \Omega_2$  we have

$$P(\Omega_2) = P((\Omega_2 \setminus \Omega_1) \cup \Omega_1) = P(\Omega_2 \setminus \Omega_1) + P(\Omega_1),$$

which along with the fact that  $P(\Omega) \geq 0$  gives the result.

□

Note most of these properties make sense simply by thinking of  $P(\Omega)$  as the area of the set  $\Omega \in \sigma(\mathcal{O}_{\mathbb{R}})$ . For example  $P(\Omega_1) = P(\Omega_1 \setminus \Omega_2) + P(\Omega_1 \cap \Omega_2)$  is



Remark 10.4. As noted before, it suffices to know  $P((-\infty, \lambda])$  for all  $\lambda \in \mathbb{R}$ , and for this reason a new notation is introduced;

$$P(\lambda) := P((-\infty, \lambda]).$$

However, as they are written they appear to have different domains. We therefore give a new name to  $P(\lambda)$ ; the *resolution of the identity*.

## 10.2 Real and Complex Valued Borel Measures Induced by a PVM

**Definition.** For all  $\psi, \varphi \in \mathcal{H}$  we define the  $\mathbb{C}$ -valued measure

$$\begin{aligned} \mu_{\psi, \varphi} : \sigma(\mathcal{O}_{\mathbb{R}}) &\rightarrow \mathbb{C} \\ \Omega &\mapsto \mu_{\psi, \varphi}(\Omega) := \langle \psi | P(\Omega) \varphi \rangle. \end{aligned}$$

◇

**Definition.** For all  $\psi \in \mathcal{H}$  we define the  $\mathbb{R}$ -valued measure

$$\mu_{\psi} := \mu_{\psi, \psi}$$

◇

*Proof.* that  $\mu_{\psi}(\Omega) \in \mathbb{R}$ .

$$\begin{aligned} \mu_{\psi}(\Omega) &= \langle \psi | P(\Omega) \psi \rangle \\ &= \langle P(\Omega) \psi | \psi \rangle \\ &= \overline{\langle \psi | P(\Omega) \psi \rangle} \\ &= \overline{\mu_{\psi}(\Omega)}, \end{aligned}$$

where we have used the fact that  $P$  is self adjoint to go from the first to the second line. □

## 10.3 Integration With Respect to a PVM

We now wish to make sense of the operator  $\int_{\mathbb{R}} f dP$  for measurable  $f : \mathbb{R} \rightarrow \mathbb{C}$ . We will build this up in three steps:

- (i) For simple  $f$ ,
- (ii) For bounded  $f$ , and
- (iii) For not necessarily bounded<sup>18</sup>  $f$ .

Remark 10.5. As we shall see, if  $f$  is bounded (in the sense that there exists a  $a \in \mathbb{R}$  such that  $|f(x)| < a$  for all  $x \in \mathbb{R}$ ) that the integral part of the operator will have nice properties. For example it will be linear in  $f$ , i.e.

$$\int_{\mathbb{R}} (\alpha f + g) dP = \alpha \int_{\mathbb{R}} f dP + \int_{\mathbb{R}} g dP,$$

---

<sup>18</sup>Recall footnote 5 from lecture 2: to us the term ‘unbounded’ means definitely *not* bounded.



for  $\alpha \in \mathbb{C}$ . However, if  $f$  is unbounded, domain issues will destroy the equality above. It is important to note, though, that it is exactly the latter case we need as

$$\begin{aligned} f &= i_{\mathbb{R}}: \mathbb{R} \hookrightarrow \mathbb{C} \\ x &\mapsto x \end{aligned}$$

in the Spectral theorem, which is clearly unbounded.

### A. Simple $f$

Recall a simple function is a measurable function that takes a finite number of results in the target, i.e. if  $f: \mathbb{R} \rightarrow \mathbb{C}$  then

$$f(\mathbb{R}) = \{f_1, \dots, f_N\} \subset \mathbb{C}$$

for some  $N \in \mathbb{N}$ . This allows us to rewrite  $f$  as

$$f = \sum_{n=1}^N f_n \chi_{\Omega_n},$$

where  $\Omega_n := \text{preim}_f(\{f_n\})$  and  $\chi$  is the characteristic function.

**Definition.** For simple  $f: \mathbb{R} \rightarrow \mathbb{C}$  and PVM  $P$  we define

$$\int_{\mathbb{R}} f dP := \sum_{n=1}^N f_n P(\Omega_n).$$

◇

**Proposition 10.6.** For simple  $f$ ,  $\int_{\mathbb{R}} f dP$  is linear in  $f$ .

*Proof.* Let  $S(\mathbb{R}, \mathbb{C})$  denote the set of all simple functions  $f: \mathbb{R} \rightarrow \mathbb{C}$ . We can make this set into a  $\mathbb{C}$ -vector space by inheriting the addition and  $s$ -multiplication from  $\mathbb{C}$ , namely define

$$(f + g)(x) := f(x) + g(x), \quad (\alpha \cdot f)(x) := \alpha \cdot f(x),$$

for all  $f, g \in S(\mathbb{R}, \mathbb{C})$ ,  $\alpha \in \mathbb{C}$ .

Now consider the preimage part: As  $f$  and  $g$  are both simple it follows that

$$(f + g)(x) = f_n + g_n$$

for some  $f_n, g_n \in \mathbb{C}$ , so the preimage term becomes

$$\begin{aligned} \text{preim}_{f+g}\{f_n + g_n\} &= \{x\} = \text{preim}_f\{f_n\} \\ &= \text{preim}_g\{g_n\}. \end{aligned}$$

It follows trivially, then, that

$$\int_{\mathbb{R}} (f + g) dP = \int_{\mathbb{R}} f dP + \int_{\mathbb{R}} g dP.$$

A similar method gives the  $\alpha \in \mathbb{C}$  condition of linearity. □

Remark 10.7. Observe that  $\chi_\Omega$  for any  $\Omega \in \sigma(\mathcal{O}_\mathbb{R})$  is simple (it only takes the values 0 or 1), and hence

$$\int_{\mathbb{R}} \chi_\Omega dP = 1 \cdot P(\Omega) + 0 \cdot P(\sigma \setminus \Omega) = P(\Omega).$$

Remark 10.8. Observe also that for any  $\psi, \varphi \in \mathcal{H}$ ,

$$\begin{aligned} \left\langle \psi \left| \left( \int_{\mathbb{R}} f dP \right) \varphi \right\rangle &= \left\langle \psi \left| \sum_{n=1}^N f_n P(\Omega_n) \varphi \right\rangle \\ &= \sum_{n=1}^N f_n \langle \psi | P(\Omega_n) \varphi \rangle \\ &= \sum_{n=1}^N f_n \mu_{\psi, \varphi}(\Omega_n) \\ &=: \int_{\mathbb{R}} f d\mu_{\psi, \varphi}, \end{aligned}$$

where Proposition 6.7 was used.

**Definition.** For simple  $f$  we can define the map

$$\begin{aligned} \left( \int_{\mathbb{R}} dP \right) : S(\mathbb{R}, \mathbb{C}) &\rightarrow \mathcal{L}(\mathcal{H}) \\ f &\mapsto \int_{\mathbb{R}} f dP, \end{aligned}$$

which, if we equip  $S(\mathbb{R}, \mathbb{C})$  with the supremum norm and  $\mathcal{L}(\mathcal{H})$  with its operator norm, has operator norm  $\| \int_{\mathbb{R}} dP \| = 1$ .  $\diamond$

*Proof.* We have already shown that  $\int_{\mathbb{R}} f dP \in \mathcal{L}(\mathcal{H})$  (i.e. it is linear), so we just need to

show the norm condition. First, let  $f \in S(\mathbb{R}, \mathbb{C})$  and  $\psi \in \mathcal{H}$ , then

$$\begin{aligned}
\left\| \left( \int_{\mathbb{R}} f dP \right) \psi \right\|_{\mathcal{H}}^2 &= \left\langle \left( \int_{\mathbb{R}} f dP \right) \psi \left| \left( \int_{\mathbb{R}} f dP \right) \psi \right\rangle \\
&= \left\langle \sum_{n=1}^N f_n P(\Omega_n) \psi \left| \sum_{m=1}^N f_m P(\Omega_m) \psi \right\rangle \\
&= \left\langle \psi \left| \sum_{n,m=1}^N \overline{f_n} f_m P(\Omega_n) P(\Omega_m) \psi \right\rangle \\
&= \left\langle \psi \left| \sum_{n,m=1}^N \overline{f_n} f_m \delta_{nm} P(\Omega_m) \psi \right\rangle \\
&= \sum_{n=1}^N |f|^2 \langle \psi | P(\Omega_n) \psi \rangle \\
&= \sum_{n=1}^N |f|^2 \mu_{\psi}(\Omega_n) \\
&=: \int_{\mathbb{R}} |f|^2 d\mu_{\psi} \\
\Rightarrow \left\| \left( \int_{\mathbb{R}} f dP \right) \psi \right\|_{\mathcal{H}} &\leq \|f\|_{\infty} \|\psi\|_{\mathcal{H}},
\end{aligned}$$

where we have used the definition of the norm in terms of the inner product, the fact that  $P$  is self adjoint, the fact that  $P(\Omega_n)P(\Omega_m) = \delta_{nm}$  for pairwise disjoint  $\Omega_n/\Omega_m$ , and Proposition 6.27 along with the fact that  $\|f\|_{\infty} := \sup_{x \in \mathbb{R}} f(x)$ . The equality in the last line can be assumed provided  $f$  and  $\psi$  are sufficiently chosen.

Thus we have

$$\begin{aligned}
\left\| \int_{\mathbb{R}} dP \right\| &:= \sup_{f \in S(\mathbb{R}, \mathbb{C})} \frac{\left\| \int_{\mathbb{R}} f dP \right\|_{\mathcal{L}(\mathcal{H})}}{\|f\|_{\infty}} \\
&:= \sup_{f \in S(\mathbb{R}, \mathbb{C})} \sup_{\psi \in \mathcal{H}} \frac{\left\| \left( \int_{\mathbb{R}} f dP \right) \psi \right\|_{\mathcal{H}}}{\|f\|_{\infty} \|\psi\|_{\mathcal{H}}} \\
&= 1.
\end{aligned}$$

□

## B. Bounded Borel Functions

**Definition.** The set of all bounded, measurable functions is denoted

$$\mathcal{B}(\mathbb{R}, \mathbb{C}) := \{f: \mathbb{R} \rightarrow \mathbb{C} \mid \text{measurable, } \|f\|_{\infty} < \infty\}.$$

◇

**Proposition 10.9.** *The set  $\mathcal{B}$  can be made into a Banach space by defining the norm*

$$\|f\|_{\mathcal{B}} := \sup_{x \in \mathbb{R}} |f(x)|.$$

*Proof.* We turn the set into a linear vector space in the usual manner; we inherit the addition and  $s$ -multiplication from  $\mathbb{C}$ .

Now prove  $\|f\|_{\mathcal{B}}$  is a norm. Comparing to the definition given at the bottom of Page 9, for  $f, g \in \mathcal{B}(\mathbb{R}, \mathbb{C})$  and  $z \in \mathbb{C}$  we have

(i) Clearly  $\|f\|_{\mathcal{B}} \geq 0$ .

(ii)

$$\begin{aligned}\|f\|_{\mathcal{B}} &= 0 \\ \Leftrightarrow \sup_{x \in \mathbb{R}} |f(x)| &= 0 \\ \Leftrightarrow f(x) &= 0 \quad \forall x \in \mathbb{R}\end{aligned}$$

so  $f = 0$ .

(iii)

$$\begin{aligned}\|z \cdot f\|_{\mathcal{B}} &:= \sup_{x \in \mathbb{R}} |z \cdot f(x)| \\ &= |z| \sup_{x \in \mathbb{R}} |f(x)| \\ &=: |Z| \cdot \|f\|_{\mathcal{B}}.\end{aligned}$$

(iv)

$$\begin{aligned}\|f + g\|_{\mathcal{B}} &:= \sup_{x \in \mathbb{R}} |(f + g)(x)| \\ &= \sup_{x \in \mathbb{R}} |f(x) + g(x)| \\ &\leq \sup_{x \in \mathbb{R}} |f(x)| + \sup_{x \in \mathbb{R}} |g(x)| \\ &=: \|f\|_{\mathcal{B}} + \|g\|_{\mathcal{B}}.\end{aligned}$$

Now let  $\{f_n\}_{n \in \mathbb{N}}$  be a Cauchy sequence in  $\mathcal{B}(\mathbb{R}, \mathbb{C})$ , that is:  $\forall \epsilon > 0, \exists N \in \mathbb{N}: \forall m, n \geq N$  we have

$$d(f_n, f_m) := \|f_n - f_m\|_{\mathcal{B}} := \sup_{x \in \mathbb{R}} |f_n(x) - f_m(x)| < \epsilon.$$

Now from the definition of the supremum we have

$$|f_n(x) - f_m(x)| \leq \|f - g\|_{\mathcal{B}},$$

so it follows that the sequence  $\{f_n(x)\}_{n \in \mathbb{N}}$  is a Cauchy sequence in  $\mathbb{C}$ . But  $\mathbb{C}$  is a complete metric space so we know that this Cauchy sequence converges in  $\mathbb{C}$ , i.e.

$$\lim_{n \rightarrow \infty} f_n(x) = z_x \in \mathbb{C}.$$

We can thus define a point-wise limit  $f$  of the sequence  $\{f_n\}_{n \in \mathbb{N}} \subset \mathcal{B}(\mathbb{R}, \mathbb{C})$  as

$$f(x) := \lim_{n \rightarrow \infty} f_n(x) = z_x$$

for all  $x \in \mathbb{R}$ . Then, by equipping  $\mathbb{C}$  and  $\mathbb{R}$  with their respective Borel  $\sigma$ -algebras, Proposition 5.22 tells us that  $f$  is measurable.

Finally from the fact that  $\mathcal{B}(\mathbb{R}, \mathbb{C}) \subset \mathcal{L}(\mathbb{R}, \mathbb{C})$ , Theorem 2.6 tells us that  $f$  is bounded and so  $\mathcal{B}(\mathbb{R}, \mathbb{C})$  is a Banach space.  $\square$

**Corollary 10.10.** *Observe that  $S(\mathbb{R}, \mathbb{C})$  is in-fact a dense, linear subspace of  $\mathcal{B}(\mathbb{R}, \mathbb{C})$ . Thus, the BLT theorem tells us that we have a unique extension of the operator*

$$\left( \int_{\mathbb{R}} dP \right) : S(\mathbb{R}, \mathbb{C}) \rightarrow \mathcal{L}(\mathcal{H})$$

*to the domain  $\mathcal{B}(\mathbb{R}, \mathbb{C})$  with equal operator norm. That is, we have an operator*

$$\left( \widehat{\int_{\mathbb{R}} dP} \right) : \mathcal{B}(\mathbb{R}, \mathbb{C}) \rightarrow \mathcal{L}(\mathcal{H})$$

*with  $\|\widehat{\int_{\mathbb{R}} dP}\| = 1$ .*

**Corollary 10.11.** *By suitable definition we can turn the space  $\mathcal{B}(\mathbb{R}, \mathbb{C})$  into an  $C^*$ -algebra and our operator then has the following properties*

(i)

$$\int_{\mathbb{R}} 1 dP = \text{id}_{\mathcal{H}}$$

(ii)

$$\int_{\mathbb{R}} (f \cdot g) dP = \left( \int_{\mathbb{R}} f dP \right) \circ \left( \int_{\mathbb{R}} g dP \right)$$

(iii)

$$\int_{\mathbb{R}} \bar{f} dP = \left( \int_{\mathbb{R}} f dP \right)^*$$

*These properties collectively make the operator a  $C^*$ -algebra homomorphism.*

## C. General Borel Function

We now want to allow for the case that  $f$  is unbounded. We will write the following such that it reduces to the above when  $f$  is bounded.

**Definition.** Let  $f : \mathbb{R} \rightarrow \mathbb{C}$  be measurable, then we define the linear map

$$\left( \int_{\mathbb{R}} f dP \right) : \mathcal{D}_{\int_{\mathbb{R}} f dP} \rightarrow \mathcal{H}$$

where

$$\mathcal{D}_{\int_{\mathbb{R}} f dP} := \left\{ \psi \in \mathcal{H} \mid \int_{\mathbb{R}} |f|^2 d\mu_{\psi} < \infty \right\} \subseteq \mathcal{H},$$

is a dense, linear subspace. The linear map is defined via

$$\left( \int_{\mathbb{R}} f dP \right) \psi := \lim_{n \rightarrow \infty} \left[ \left( \int_{\mathbb{R}} f_n dP \right) \psi \right],$$

where the sequence  $\{f_n\}_{n \in \mathbb{N}} \subseteq \mathcal{B}(\mathbb{R}, \mathbb{C})$  defined by

$$f_n := \chi_{\{x \in \mathbb{R} | f(x) < n\}} f.$$

◇


*Remark 10.12.* Note that the map above includes the  $f$ , it is not just the integral defined in the previous section. Note also for the case when  $f \in \mathcal{B}(\mathbb{R}, \mathbb{C})$  we just recover the case above and we have  $\mathcal{D}_{\int_{\mathbb{R}} f dP} = \mathcal{H}$  (i.e. we have  $\mathcal{L}(\mathcal{H})$ ). Otherwise it is a proper subset.

*Remark 10.13.* The literature often introduces the notation

$$\mathcal{D}_f := \mathcal{D}_{\int_{\mathbb{R}} f dP},$$

however this could lead one to think of the domain of  $f$  itself, which here is  $\mathbb{R}$ . We will avoid this notation.

*Remark 10.14.* The sequence  $\{f_n\}_{n \in \mathbb{N}}$  can be thought of as ‘chopping’  $f$  into bounded parts.



**Lemma 10.15.** *The sequence  $\{f_n\}_{n \in \mathbb{N}}$  is Cauchy in  $L^2(\mathbb{R})$ . This in turn implies that the sequence  $\{(\int_{\mathbb{R}} f_n dP)\psi\}_{n \in \mathbb{N}}$  is Cauchy in  $\mathcal{H}$ , which is required for the limit in the definition to make sense, i.e. the result lies in  $\mathcal{H}$ .*

For this general case of a measurable  $f$  we have:

(i) As before

$$\int_{\mathbb{R}} \bar{f} dP = \left( \int_{\mathbb{R}} f dP \right)^*.$$

(ii) For  $\alpha \in \mathbb{C}$  and  $f, g$  measurable,

$$\left( \alpha \int_{\mathbb{R}} f dP + \int_{\mathbb{R}} g dP \right) \subseteq \int_{\mathbb{R}} (\alpha f + g) dP,$$

where the equality holds only for bounded  $f$ . As explained earlier, the inequality arises due to domain issues. We can now see this more explicitly from the definition of  $\mathcal{D}_{\int_{\mathbb{R}} f dP}$ ; just because  $f$  and  $g$  are both measurable, it does not mean that their respective map domains will coincide. However, the domain for the LHS is  $\mathcal{D}_{\int_{\mathbb{R}} (|\alpha f| + |g|) dP}$ .

(iii)

$$\left( \int_{\mathbb{R}} f dP \right) \circ \left( \int_{\mathbb{R}} g dP \right) \subseteq \int_{\mathbb{R}} (f \cdot g) dP,$$

again where the equality holds only when  $f$  and  $g$  are bounded.

## 10.4 The Inverse Spectral Theorem

We are now in a place where we can understand the inverse spectral theorem.

**Definition.** Given a PVM,  $P$ , we can construct a self adjoint operator  $A_P$  as

$$A_P := \int_{\mathbb{R}} \text{id}_{\mathbb{R}} dP,$$

where  $\text{id}_{\mathbb{R}}: \mathbb{R} \hookrightarrow \mathbb{C}$  is the inclusion map. ◇

*Proof.*

$$\begin{aligned} (A_P)^* &= \int_{\mathbb{R}} \overline{\text{id}_{\mathbb{R}}} dP \\ &= \int_{\mathbb{R}} \text{id}_{\mathbb{R}} dP \\ &= A_P, \end{aligned}$$

so  $A_P$  is self adjoint. □

## 11 Spectral theorem

The inverse spectral theorem tells us how to construct a self adjoint operator  $A_P$  given a projection valued measure  $P$ . The aim of this lecture is to do the opposite; given a self adjoint operator  $A$  we want to find a PVM  $P_A$ . We want the two methods to be in unison — that is we want  $A_{P_A} = A$  and  $P_{A_P} = P$ . We shall start by assuming that  $A$  can be written in integral form, and then shall remove this restriction.

### 11.1 Measurable Function Applied To A Spectrally Decomposable Self Adjoint Operator

The term *spectrally decomposable* means that it has integral form.

**Definition.** Let the self adjoint operator  $A$  be *spectrally decomposable*; i.e. there exists a PVM  $P$  such that

$$A = \int_{\mathbb{R}} \text{id}_{\mathbb{R}} dP,$$

then for any measurable function  $f : \mathbb{R} \rightarrow \mathbb{C}$ , we define the operator

$$f(A) : \mathcal{D}_{\int f dP} \rightarrow \mathcal{H},$$

given by

$$f(A) := \int_{\mathbb{R}} (f \circ \text{id}_{\mathbb{R}}) dP \equiv \int_{\mathbb{R}} f(\lambda) P(d\lambda).$$

◇

*Remark 11.1.* The spectral theorem will show that every self adjoint operator  $A$  is spectrally decomposable by virtue of a uniquely determined  $P$ .

**Corollary 11.2.** *If  $f : \mathbb{R} \rightarrow \mathbb{R}$ , then  $f(A)$  is again self adjoint.*

*Proof.*

$$\begin{aligned} [f(A)]^* &:= \left[ \int_{\mathbb{R}} (f \circ \text{id}_{\mathbb{R}}) dP \right]^* \\ &= \int_{\mathbb{R}} (\bar{f} \circ \text{id}_{\mathbb{R}}) dP \\ &= \int_{\mathbb{R}} \bar{f} dP \\ &=: f(A). \end{aligned}$$

□

Let us now consider two important examples.

Example 11.3. Let  $A = \int_{\mathbb{R}} \lambda P(d\lambda)$  be self adjoint. Then

$$\exp(A) := \int_{\mathbb{R}} e^{\lambda} P(d\lambda)$$



is self adjoint due to the previous Corollary. However

$$\exp(iA) := \int_{\mathbb{R}} e^{i\lambda} P(d\lambda)$$

is *not* self adjoint. This latter case is of high importance in QM, as can be seen by revisiting Axiom 4 at the start.

Example 11.4. Let  $A = \int_{\mathbb{R}} \lambda P(d\lambda)$  be self adjoint and  $\Omega \in \sigma(\mathcal{O}_{\mathbb{R}})$ . Then

$$P(\Omega) = \int_{\mathbb{R}} \chi_{\Omega} dP$$

implies

$$\begin{aligned} \text{sq}(P(\Omega)) &:= \int_{\mathbb{R}} (\text{sq} \circ \chi_{\Omega}) dP \\ &\equiv \int_{\mathbb{R}} [\chi_{\Omega}(\lambda)]^2 P(d\lambda) \\ &= \int_{\mathbb{R}} \chi_{\Omega}(\lambda) P(d\lambda) \\ &= P(\Omega), \end{aligned}$$

which is one of the projection conditions.

## 11.2 Reconstruct PVM From a Spectrally Decomposable, Self Adjoint Operator

The key to reconstructing the associated PVM  $P$  is to consider the resolvents.

**Definition.** Given a spectrally decomposable operator  $A$  and an the resolvent set  $\rho(A)$ , we define

$$r_z(A) = R_Z(A) := (A - z \text{id}_{\mathcal{H}})^{-1},$$

which is rewritten as

$$\begin{aligned} r_z &: \mathbb{R} \rightarrow \mathbb{C} \\ \lambda &\mapsto \frac{1}{\lambda - z}, \end{aligned}$$

and, due to the fact that  $A$  is spectrally decomposable, satisfies

$$r_z(A) = \int_{\mathbb{R}} (r_z \circ \text{id}_{\mathbb{R}}) dP \equiv \int_{\mathbb{R}} \frac{1}{\lambda - z} P(d\lambda).$$

◇

Note, using the results in the previous lecture, we have that for any  $\psi \in \mathcal{H}$ ,

$$\begin{aligned} \langle \psi | R_A(z) \psi \rangle &= \left\langle \psi \left| \left( \int_{\mathbb{R}} (r_z \circ \text{id}_{\mathbb{R}}) dP \right) \psi \right. \right\rangle \\ &= \int_{\mathbb{R}} (r_z \circ \text{id}_{\mathbb{R}}) d\mu_{\psi} \\ &\equiv \int_{\mathbb{R}} \frac{1}{\lambda - z} \mu_{\psi}(d\lambda). \end{aligned}$$

**Definition.** A *Herglotz function* is an analytic complex function that maps the upper half plane into itself, but need not be surjective or injective. They are also known as *Nevanlinna/Pick/R* functions.  $\diamond$

**Theorem 11.5.** *The function*

$$\begin{aligned} \langle \psi | R_A(\cdot) \psi \rangle : \mathbb{C} &\rightarrow \mathbb{C} \\ z &\mapsto \int_{\mathbb{R}} \frac{1}{\lambda - z} \mu_{\psi}(d\lambda) \end{aligned}$$

is *Herglotz*.

*Proof.* Recall

$$\mu_{\psi} : \sigma(\mathcal{O}_{\mathbb{R}}) \rightarrow \mathbb{R}_0^+$$

is real-valued. Then using

$$\operatorname{Im}(r) = \frac{1}{2}(r - \bar{r}),$$

we have

$$\begin{aligned} \operatorname{Im} \langle \psi | R_A(z) \psi \rangle &= \int_{\mathbb{R}} \operatorname{Im} \left( \frac{1}{\lambda - z} \right) \mu_{\psi}(d\lambda) \\ &= \frac{1}{2} \int_{\mathbb{R}} \left[ \frac{1}{\lambda - z} - \frac{1}{\lambda - \bar{z}} \right] \mu_{\psi}(d\lambda) \\ &= \int_{\mathbb{R}} \frac{z - \bar{z}}{2|\lambda - z|^2} \mu_{\psi}(d\lambda) \\ &= \operatorname{Im}(z) \int_{\mathbb{R}} \frac{1}{|\lambda - z|^2} \mu_{\psi}(d\lambda). \end{aligned}$$

Then, since the fact that the integral is Lebesgue and the integrand is non-negative and so

$$\operatorname{Im} \langle \psi | R_A(z) \psi \rangle > 0 \quad \Leftrightarrow \quad \operatorname{Im}(z) > 0.$$

$\square$

Recalling the start of last lecture, if we can find a way to construct  $\mu_{\psi}$  from our  $A$  then we can use that to reconstruct  $P$ . The result of this is the previously mentioned Stieltjes Inversion Formula, and it is obtained as follows.

Let  $t, \epsilon \in \mathbb{R}$ . Then, since  $A$  is self adjoint and so its spectrum is purely real,  $t + i\epsilon \in \rho(A)$ . This allows us to act on it with  $R_A$ . Thus, consider

$$\begin{aligned} \lim_{\epsilon \rightarrow 0^+} \frac{1}{\pi} \int_{t_1}^{t_2} dt \operatorname{Im} \langle \psi | R_A(t + i\epsilon) \psi \rangle &= \lim_{\epsilon \rightarrow 0^+} \frac{1}{\pi} \int_{t_1}^{t_2} dt \int_{\mathbb{R}} \frac{\epsilon}{|\lambda - t - i\epsilon|^2} \mu_{\psi}(d\lambda) \\ &= \lim_{\epsilon \rightarrow 0^+} \int_{\mathbb{R}} \left( \frac{1}{\pi} \int_{t_1}^{t_2} dt \frac{\epsilon}{(\lambda - t)^2 + \epsilon^2} \right) \mu_{\psi}(d\lambda), \end{aligned}$$

where Fubini's Theorem<sup>19</sup> has been used. The inner integral is a standard integral, with result


$$\frac{1}{\pi} \int_{t_1}^{t_2} dt \frac{\epsilon}{(\lambda - t)^2 + \epsilon^2} = \frac{1}{\pi} \left[ \arctan \left( \frac{t - \lambda}{\epsilon} \right) \right]_{t_1}^{t_2}.$$

---


<sup>19</sup>See [Wiki](#)

Now strictly, at this stage, we cannot simply pull the  $\varepsilon$  limit into this expression; we would need to check that the above result is bounded first and then, by dominated convergence, we can pull it in. This will turn out to be true, and so in order to simplify the following we consider  $\varepsilon$  to be small here.

In order to work out the above expression, we can use the  $\lambda$ -graphs. Let's plot both terms (including the overall minus sign that comes with  $t_1$ ) on the same graph:



Taking the limit and adding gives



So we have

$$\lim_{\varepsilon \rightarrow 0^+} \frac{1}{\pi} \int_{t_1}^{t_2} dt \frac{\varepsilon}{(\lambda - t)^2 + \varepsilon^2} = \frac{1}{2} (\chi_{(t_1, t_2)} + \chi_{[t_1, t_2]}),$$

and

$$\lim_{\varepsilon \rightarrow 0^+} \frac{1}{\pi} \int_{t_1}^{t_2} dt \operatorname{Im} \langle \psi | R_A(t + i\varepsilon) \psi \rangle = \frac{1}{2} \int_{\mathbb{R}} (\chi_{(t_1, t_2)} + \chi_{[t_1, t_2]}) \mu_{\psi}(d\lambda).$$

Finally, we have the Stieltjes Inversion Formula.

**Theorem 11.6** (Stieltjes Inversion Formula). *Given a spectrally decomposable, self adjoint operator  $A$  and its associated resolvent map  $R_A$ , we can construct a real-valued measure*

$$\mu_{\psi}^A((-\infty, \lambda]) = \lim_{\delta \rightarrow 0^+} \lim_{\varepsilon \rightarrow 0^+} \frac{1}{\pi} \int_{-\infty}^{\lambda + \delta} dt \operatorname{Im} \langle \psi | R_A(t + i\varepsilon) \psi \rangle.$$

*Proof.*

$$\begin{aligned}
\lim_{\delta \rightarrow 0^+} \lim_{\varepsilon \rightarrow 0^+} \frac{1}{\pi} \int_{-\infty}^{\lambda+\delta} dt \operatorname{Im} \langle \psi | R_A(t + i\varepsilon) \psi \rangle &= \lim_{\delta \rightarrow 0^+} \frac{1}{2} \int_{\mathbb{R}} (\chi_{(-\infty, \lambda+\delta)} + \chi_{(-\infty, \delta]}) \mu_\psi(d\lambda) \\
&= \int_{\mathbb{R}} \chi_{(-\infty, \lambda]} \mu_\psi(d\lambda) \\
&= \mu_\psi((-\infty, \lambda]),
\end{aligned}$$

where we used the fact that the  $\chi(\Omega)$  is bounded to move the limit inside the integral along with the fact that

$$\lim_{\delta \rightarrow 0^+} (-\infty, \lambda + \delta) = (-\infty, \lambda].$$

□

*Remark 11.7.* Note the fact that  $\langle \psi | R_A(t + i\varepsilon) \psi \rangle$  is Herglotz with the fact that  $\varepsilon > 0$  gives us that  $\mu_\psi^A \geq 0$ , which is required for it to be a real-valued measure.

*Remark 11.8.* If we already know that  $A$  is spectrally decomposable w.r.t. some PVM  $P$ , then we can recover  $P$  from  $A$  by virtue of: for any  $\Omega \in \sigma(\mathcal{O}_{\mathbb{R}})$  and for all  $\psi, \varphi \in \mathcal{H}$

$$\langle \psi | P(\Omega) \varphi \rangle = \int_{\mathbb{R}} \chi_\Omega d\mu_{\psi, \varphi},$$

where  $\mu_{\psi, \varphi}$  is obtained from  $\mu_\psi$  using the method given at the start of the previous lecture.

### 11.3 Construction Of PVM From A Self Adjoint Operator

We need to free ourselves from the fact that  $A$  is known to be spectrally decomposable from the start. We could do this by trying to recreate the above method, i.e. arrive at the Stieltjes Inversion Formula for an operator  $A$ , by showing that

- (i)  $\langle \psi | R_A(\cdot) \psi \rangle : \mathbb{C} \rightarrow \mathbb{C}$  is Herglotz for any self adjoint  $A$
- (ii)  $\langle \psi | P(\Omega) \varphi \rangle := \int_{\mathbb{R}} \chi_\Omega d\mu_{\psi, \varphi}$  is indeed a PVM.

In order to prove these we first need a new theorem.

**Theorem 11.9** (First-Resolvent Formula). *For any operator  $A : \mathcal{D}_A \rightarrow \mathcal{H}$  and  $a, b \in \rho(A)$  we have*

$$R_A(a) - R_A(b) = (a - b)R_A(a)R_A(b) = (a - b)R_A(b)R_A(a).$$

*Proof.* Consider

$$\begin{aligned}
R_A(a) - (a - b)R_A(a)R_A(b) &:= (A - a)^{-1} - (a - b)(A - a)^{-1}(A - b)^{-1} \\
&= (A - a)^{-1} [\operatorname{id}_{\mathcal{H}} - (a - b)(A - b)^{-1}] \\
&= (A - a)^{-1} [\operatorname{id}_{\mathcal{H}} - (a - A + A - b)(A - b)^{-1}] \\
&= (A - a)^{-1} [\operatorname{id}_{\mathcal{H}} + (A - a)(A - b)^{-1} - (A - b)(A - b)^{-1}] \\
&= (A - b)^{-1} \\
&= R_A(b),
\end{aligned}$$

and similarly for the the other result. □

Conclusion, the Spectral Theorem, Theorem 10.1, together with the recipe for the construction of the PVM  $P$  from a self adjoint operator  $A$  holds.

### 11.4 Commuting Operators

The study of QM is teaming with so called *commutators*. However, they are not as simply defined as is often erroneously assumed. In particular, the commutator between the position and momentum given by

$$[Q^i, P_j] = i\hbar\delta^i_j$$

is not even defined, unless further provisions are given. This formula appears in the opening sections of almost all QM textbooks though, and so it's important we understand what is meant.

One can happily write the commutator provided the operators involved are bounded (which from the lecture 9 we see that at least  $P_j$  is not).

**Definition.** Let  $B_1, B_2 \in \mathcal{L}(\mathcal{H})$ , i.e. they are bounded linear operators from  $\mathcal{H}$  to  $\mathcal{H}$ . Then one may define

$$[B_1, B_2] := B_1 \circ B_2 - B_2 \circ B_1,$$

where

$$[B_1, B_2] \in \mathcal{L}(\mathcal{H}).$$

◇

*Remark 11.10.* The tuple  $(\mathcal{L}(\mathcal{H}), +, \cdot, [\cdot, \cdot])$  is a Lie algebra<sup>20</sup>.

**Corollary 11.11.** Let  $A, B, C \in \mathcal{L}(\mathcal{H})$  then the following holds

$$[A \circ B, C] = [A, C] \circ B + A \circ [B, C].$$

*Proof.* Consider the action on some arbitrary  $\psi \in \mathcal{H}$ .

$$\begin{aligned} [A \circ B, C]\psi &:= (A \circ B) \circ (C\psi) - C \circ (A \circ B\psi) \\ &= A \circ (B \circ C\psi) - C \circ (A \circ B\psi) \\ &= A \circ (C \circ B\psi + [B, C]\psi) - C \circ (A \circ B\psi) \\ &= (A \circ C) \circ B\psi + A \circ [B, C]\psi - C \circ (A \circ B\psi) \\ &= (C \circ A + [A, C]) \circ B\psi + A \circ [B, C]\psi - C \circ (A \circ B\psi) \\ &= [A, C] \circ B\psi + A \circ [B, C]\psi \\ &= ([A, C] \circ B + A \circ [B, C])\psi, \end{aligned}$$

where we used the associativity of the composition of maps. Then, as  $\psi$  was arbitrary, we have our result. □

**Corollary 11.12.** Let  $A$  and  $B$  be two operators. Then if one of the them is unbounded the domain  $\mathcal{D}_{[A, B]}$  may only have a trivial definition, i.e.  $\mathcal{D}_{[A, B]} = \{0_{\mathcal{H}}\}$ .

<sup>20</sup>See Dr. Schuller's Lectures on the Geometric Anatomy of Theoretical Physics

*Proof.* Let  $A : \mathcal{D}_A \rightarrow \mathcal{H}$  be unbounded with  $\mathcal{D}_A \subset \mathcal{H}$ , and define the bounded operator

$$\begin{aligned} B_\varphi : \mathcal{H} &\rightarrow \mathcal{H} \\ \alpha &\mapsto \langle \varphi | \alpha \rangle \psi =: \ell_\varphi(\alpha) \psi \end{aligned}$$

for some fixed  $\varphi, \psi \in \mathcal{H}$  where  $\psi \notin \mathcal{D}_A$ . So we have  $\text{ran}(B_\varphi) = \mathcal{H} \setminus \mathcal{D}_A$ . Then from the first term in

$$[A, B_\varphi] := A \circ B_\varphi - B_\varphi \circ A,$$

it follows that

$$\mathcal{D}_{[A, B_\varphi]} = \text{ran}(B_\varphi) \cap \mathcal{D}_A = \{0_{\mathcal{H}}\}.$$

□

**Definition.** Two bounded linear operators  $A, B \in \mathcal{L}(\mathcal{H})$  are said to *commute* if

$$[A, B] = 0.$$

◇

**Corollary 11.13.** *Let  $A, B \in \mathcal{L}(\mathcal{H})$  be commuting operators. Then if  $A$  is also non-degenerate then any  $\psi$  that is an eigenvector of  $A$  is also an eigenvector of  $B$ . In other words, the set of  $A$ 's eigenvectors is contained within the set of  $B$ 's.*

*Proof.* Let  $\psi \in \mathcal{D} \setminus \{0\}$  be an eigenvector of  $A$  with eigenvalue  $\lambda \in \mathbb{C}$ . Then we have

$$\begin{aligned} [A, B]\psi &:= (A \circ B - B \circ A)\psi \\ &= A(B\psi) - B(A\psi) \\ &= A(B\psi) - \lambda B(\psi), \end{aligned}$$

where we have used the fact that  $B$  is linear. Then from the fact that  $[A, B] = 0$  it follows that  $B\psi$  is also an eigenvector of  $A$  with eigenvalue  $\lambda$ . Finally from the fact that  $A$  is non-degenerate it follows that  $B\psi = \mu\psi$  must hold for some  $\mu \in \mathbb{C}$ . □

However, as highlighted at the start of this section, we also want to look at situations when one of the operators may not be bounded. In other words we want to know how to extend the idea of commuting to

- (i)  $A$  self adjoint and not necessarily bounded,  $B$  bounded.
- (ii) Both  $A$  and  $B$  self adjoint and not necessarily bounded.

As is often the case in maths/physics problems, the strategy is to reduce the problem to the known case. We then have three possible bounded, linear operators constructed from  $A$ :

- (i) From the Spectral Theorem we know that if  $A$  is self adjoint then there exists a unique PVM  $P$  such that  $A$  is spectrally decomposable. Recall that, from Remark 10.7,  $P_A(\Omega) \in \mathcal{L}(\mathcal{H})$  for any  $\Omega \in \sigma(\mathcal{O}_{\mathbb{R}})$ .

- (ii) From the definition of the resolvent set, we have  $R_A(z) \in \mathcal{L}(\mathcal{H})$  for any  $z \in \rho(A)$ .
- (iii) Again, as  $A$  is self adjoint it is spectrally decomposable and so we can consider  $\exp(itA)$  for some  $t \in \mathbb{R}$ , which was defined in Example 11.3. Again this is not a self adjoint operator, but it is unitary, which means  $\|\exp(itA)\| = 1$ .

**Definition.** Let  $A$  be self adjoint and  $B$  be bounded.  $A$  and  $B$  are said to commute if either of the following holds

- (i)  $[R_A(z), B] = 0$  for some  $z \in \rho(A)$ .
- (ii)  $[\exp(itA), B] = 0$  for some  $t \in \mathbb{R} \setminus \{0\}$ .

◇

**Definition.** Let  $A$  and  $B$  be self adjoint. They are said to commute is one of the following holds

- (i)  $[R_A(z_A), R_B(z_B)] = 0$  for some  $z_A \in \rho(A)$  and  $z_B \in \rho(B)$ . This is known as the *Resolvent way*.
- (ii)  $[\exp(itA), \exp(isB)] = 0$  for some  $t, s \in \mathbb{R} \setminus \{0\}$ . This is known as the *Weil way*.
- (iii)  $[P_A(\Omega), P_B(\Omega)] = 0$  for all  $\Omega \in \sigma(\mathcal{O}_{\mathbb{R}})$ . This is known as the *Projector way*.

◇

Remark 11.14. The literature normally uses a practical, yet misleading, notation at this point. For any of the above we simply write  $[A, B] = 0$  for commuting  $A$  and  $B$ . However this commutator is *not* the same as the one defined at the start — i.e. it does not correspond to  $A \circ B - B \circ A$ . Really we should write it slightly differently to highlight this, i.e. make it red;  $[A, B]$ .

**Theorem 11.15.** Let  $A$  and  $B$  be self adjoint and bounded. Then

$$[A, B] = 0 \quad \Leftrightarrow \quad [A, B] = 0.$$

## 12 Stone's Theorem and Construction of Observables

In this lecture we aim to answer two questions by deriving and using Stone's Theorem. They are

- (i) How arbitrary is the stipulation of Axiom 4; that the dynamics in the absence of a measurement be controlled by

$$U(t) := \exp(itH)$$

for some self adjoint operator  $H$ ?

- (ii) How does one *practically* construct observables, including the question of how to find the correct domain such that the operator is at least essentially self adjoint?

Remark 12.1. Clearly for (i) we want  $U(t) \circ U(s) = U(t + s)$  and  $U(0) = \text{id}_{\mathcal{H}}$ .

### 12.1 One Parameter Groups and Their Generators

**Definition.** A *group* is the double  $(G, \diamond)$ , where  $G$  is a set and  $\diamond: G \rightarrow G$  satisfying:

- (i) For all  $g, h, k \in G$ , (Associativity)

$$(g \diamond h) \diamond k = g \diamond (h \diamond k).$$

- (ii) There exists  $e \in G$  such that for all  $g \in G$  (Neutral Element)

$$g \diamond e = e \diamond g = g.$$

- (iii) For all  $g \in G$  there exists  $g^{-1} \in G$  such that (Inverse)

$$g \diamond g^{-1} = g^{-1} \diamond g = e.$$

◇

**Definition.** A *Abelian group* is a group is one whose group operation is symmetric. That is for all  $g, h \in G$

$$g \diamond h = h \diamond g.$$

◇

Remark 12.2. Abelian groups are also known as commutative groups and the condition is referred to as the commutativity of the elements with respect to the group operation.

Example 12.3. The real numbers equipped with addition form an Abelian group, with  $e = 0 \in \mathbb{R}$  and  $g^{-1} = -g$ .

Note it is important that we consider all of  $\mathbb{R}$ , and not just the positive numbers, as in the latter case the inverse would not lie in the group.



Example 12.4. The set  $\mathbb{R} \setminus \{0\}$  form an Abelian group with respect to multiplication, with  $e = 1$  and  $g^{-1} = 1/g$ .

Note here we have to exclude 0 as  $1/0$  is not an element of  $\mathbb{R}$ .

Example 12.5. The set  $\mathbb{R} \setminus \{0\}$  is *not* a group with respect to division as it fails to satisfy associativity.

**Definition.** A *one-parameter group* is a group whose underlying set is

$$G = \{U(t) \mid t \in \mathbb{R}\},$$

and whose group operation satisfies

$$U(t) \diamond U(s) = U(\delta(t, s)),$$

for some  $\delta: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ . ◇

Remark 12.6. Unless the group is Abelian then  $\delta(s, t) \neq \delta(t, s)$ .

We will only deal with Abelian one-parameter groups, in which case one can always reparameterise so that

$$U(t) \diamond U(s) = U(t + s)$$

where the commutativity with respect to  $\diamond$  is inherited from the commutativity with respect to  $+$ . We also choose the parameterisation such that  $U(0) = e$ . In particular, we will look at unitary one-parameter groups, i.e. those with

$$G = \{U(t) \in \mathcal{L}(\mathcal{H}) \mid t \in \mathbb{R}, U^*(t)U(t) = \text{id}_{\mathcal{H}}, \|U(t)\| = 1\},$$

that are *strongly continuous* in the parameter,

$$\forall \psi \in \mathcal{H} : \lim_{t \rightarrow t_0} (U(t)\psi) = U(t_0)\psi.$$

**Definition.** Let  $U(\cdot): \mathbb{R} \rightarrow \mathcal{L}(\mathcal{H})$  be a unitary, Abelian, one-parameter group (UAOPG). Then its (generator) is the linear map

$$A: \mathcal{D}_A^{\text{Stone}} \rightarrow \mathcal{H}$$

$$\psi \mapsto A\psi := \lim_{\varepsilon \rightarrow 0} \frac{i}{\varepsilon} (U(\varepsilon)\psi - \psi),$$

where

$$\mathcal{D}_A^{\text{Stone}} := \left\{ \psi \in \mathcal{H} \mid \lim_{\varepsilon \rightarrow 0} \frac{i}{\varepsilon} (U(\varepsilon)\psi - \psi) \text{ exists} \right\}.$$

◇

Remark 12.7. Note

$$\lim_{\varepsilon \rightarrow 0} \frac{i}{\varepsilon} (U(\varepsilon)\psi - \psi) = i \lim_{\varepsilon \rightarrow 0} \frac{U(0 + \varepsilon)\psi - U(0)\psi}{\varepsilon} := i[U(\cdot)\psi]'(0),$$

and so we can rewrite

$$\mathcal{D}_A^S \equiv \mathcal{D}_A^{\text{Stone}} := \left\{ \psi \in \mathcal{H} \mid i[U(\cdot)\psi]'(0) \text{ exists} \right\}.$$

Note also that  $[U(\cdot)\psi]': \mathbb{R} \rightarrow \mathcal{H}$ .

## 12.2 Stone's Theorem

**Theorem 12.8** (Stone's Theorem). *Let  $U(\cdot)$  be a UAOPG that is strongly continuous and whose group operation is the composition of maps, i.e.*

$$\begin{aligned} U(t) \circ U(s) &= U(t+s) \\ U(0) &= \text{id}_{\mathcal{H}}. \end{aligned}$$

*Then its generator  $A: \mathcal{D}_A^S \rightarrow \mathcal{H}$  is self adjoint on  $\mathcal{D}_A^S$ , and*

$$U(t) = \exp(-itA).$$

Before proving this, consider the following.

**Corollary 12.9.** *Given  $U(t) = \exp(-itA)$  for some self adjoint  $A$  then the Spectral Theorem tells us that  $U(t)$  is a UAOPG.*

*Proof.* (i) First show  $U(t) \circ U(s) = U(t+s)$ :

$$\begin{aligned} U(t) \circ U(s) &:= \exp(-itA) \circ \exp(-isA) \\ &:= \int_{\mathbb{R}} e^{-it\lambda} e^{-is\lambda} P(d\lambda) \\ &= \int_{\mathbb{R}} e^{-i(t+s)\lambda} P(d\lambda) \\ &=: U(t+s), \end{aligned}$$

where Example 11.3 has been used.

We also have

$$U(0) := \exp(0) = \text{id}_{\mathcal{H}}.$$

Now show Abelian property:

$$\begin{aligned} U^*(t) &:= \left( \int_{\mathbb{R}} e^{-it\lambda} P(d\lambda) \right)^* \\ &= \int_{\mathbb{R}} e^{+it\lambda} P(d\lambda) \\ &=: U(-t). \end{aligned}$$

Then from the above we have  $U^*(t)U(t) = U(-t+t) = U(0) = \text{id}_{\mathcal{H}}$ . Then noticing that  $\|U(t)\| = \|U^*(t)\|$  it follows that

$$\|U(t)\| = \sqrt{\|\text{id}_{\mathcal{H}}\|} = \sqrt{1} = 1,$$

where we have used the fact that the norm is strictly positive to remove the negative root, and so it is unitary.

Finally show that it is a group. This is easily done, and we have  $e = \text{id}_{\mathcal{H}} = U(0)$  and  $[U(t)]^{-1} = U(-t)$ .  $\square$

**Corollary 12.10.** *Let  $\psi \in \mathcal{D}_A^S$  for some generator  $A$  and  $t \in \mathbb{R}$ . Then*

$$[U(\cdot)\psi]'(t) = -iAU(t)\psi,$$

and

$$U(t)\mathcal{D}_A^S = \mathcal{D}_A^S.$$

Now we can proceed with the proof of Stone's Theorem. To do so, we will need to show:

- (i) The generator  $A$  is densely defined (otherwise we  $A^*$  wouldn't be defined properly).
- (ii)  $A$  is symmetric on  $\mathcal{D}_A^S$  and that it is essentially self adjoint.
- (iii)  $U(t) = \exp(-itA^{**})$ , from which it follows that  $A = A^{**}$  and so it is self adjoint (as  $A^{**}$  is).

*Proof.* (Stone's Theorem)

- (i) Let  $\psi \in \mathcal{H}$ . If we can show that an arbitrarily small neighbourhood  $\mathcal{N}$  around  $\psi$  contains a  $\varphi \in \mathcal{D}_A^S$ , then we know  $\mathcal{D}_A^S$  is dense in  $\mathcal{H}$ . Consider the real family, for all  $\tau \in \mathbb{R}$

$$\psi_\tau := \int_0^\tau dr U(r)\psi,$$

which satisfies

$$\lim_{\tau \rightarrow 0} \left( \frac{\psi_\tau}{\tau} \right) = \psi.$$

This is just the idea of the points in a neighbourhood, and so we know, therefore, that there exists a  $\tau_0$  such that  $\psi_{\tau_0} \in \mathcal{N}$ . Now consider

$$\begin{aligned} (U(\varepsilon)\psi_\tau - \psi_\tau) &:= U(\varepsilon) \int_0^\tau dr U(r)\psi - \int_0^\tau U(r)\psi \\ &= \int_0^\tau dr U(\varepsilon + r)\psi - \int_0^\tau dr U(r)\psi \\ &= \int_0^{\varepsilon+\tau} dr U(r)\psi - \int_0^\tau dr U(r)\psi \\ &= \int_\tau^{\tau+\varepsilon} dr U(r)\psi + \int_\varepsilon^\tau dr U(r)\psi - \int_0^\tau dr U(r)\psi \\ &= \int_\tau^{\tau+\varepsilon} dr U(r)\psi - \int_\tau^\varepsilon dr U(r)\psi - \int_0^\tau dr U(r)\psi \\ &= \int_\tau^{\tau+\varepsilon} dr U(r)\psi - \int_0^\varepsilon dr U(r)\psi \\ &= U(\tau) \int_0^\varepsilon dr U(r)\psi - \int_0^\varepsilon dr U(r)\psi \\ &= [U(\tau) - \text{id}_{\mathcal{H}}] \psi_\varepsilon. \end{aligned}$$

Now using the fact that  $\|U(\tau)\| = \|\text{id}_{\mathcal{H}}\| = 1$  and so they're bounded, we can take a limit and push it through the operators. Thus we have

$$\begin{aligned} i[U(\cdot)\psi_\tau]'(0) &\equiv \lim_{\varepsilon \rightarrow 0} \frac{i}{\varepsilon} (U(\varepsilon)\psi_\tau - \psi_\tau) \\ &= i[U(\tau) - \text{id}_{\mathcal{H}}] \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \psi_\varepsilon \\ &= i[U(\tau) - \text{id}_{\mathcal{H}}]\psi, \end{aligned}$$

which is an element of  $\mathcal{H}$ , and so we know that  $\psi_\tau \in \mathcal{D}_A^S$  and therefore there exists a  $\psi_{\tau_0} \in \mathcal{N} \cap \mathcal{D}_A^S$ .

(ii) Let  $\varphi, \psi \in \mathcal{D}_A^S$ , then

$$\begin{aligned} \langle \varphi | A\psi \rangle &:= \left\langle \varphi \left| \lim_{\varepsilon \rightarrow 0} \frac{i}{\varepsilon} (U(\varepsilon)\psi - \psi) \right. \right\rangle \\ &= \lim_{\varepsilon \rightarrow 0} \left\langle \frac{-i}{\varepsilon} (U^*(\varepsilon) - \text{id}_{\mathcal{H}}) \varphi \left| \psi \right. \right\rangle \\ &= \left\langle \lim_{\varepsilon \rightarrow 0} \frac{i}{-\varepsilon} (U(-\varepsilon) - \text{id}_{\mathcal{H}}) \varphi \left| \psi \right. \right\rangle \\ &= \left\langle \lim_{\varepsilon \rightarrow 0} \frac{i}{\varepsilon} (U(\varepsilon) - \text{id}_{\mathcal{H}}) \varphi \left| \psi \right. \right\rangle \\ &=: \langle A\varphi | \psi \rangle, \end{aligned}$$

where we have used the continuity of the inner product to move the limit in and out, the result  $U^*(t) = U(-t)$ , the fact that the identity is self adjoint and the fact that we're taking the limit to 'ignore' the minus signs on second to last line.

We now want to show that it is essentially self adjoint. Recalling Theorem 7.26, we need to check if: for  $z \in \mathbb{C} \setminus \mathbb{R}$  that

$$\ker(A^* - \bar{z}) = \{0_{\mathcal{H}}\} = \ker(A^* - z).$$

Let  $\varphi \in \ker(A^* - \bar{z}) \cap \mathcal{D}_{A^*}^S$ . Then for all  $\psi \in \mathcal{D}_A^S$

$$\begin{aligned} [\langle \varphi | U(\cdot)\psi \rangle]'(t) &= \langle \varphi | [U(\cdot)\psi]'(t) \rangle \\ &= \langle \varphi | -iAU(t)\psi \rangle \\ &= -i \langle A^*\varphi | U(t)\psi \rangle \\ &= -i \langle \bar{z}\varphi | U(t)\psi \rangle \\ &= -iz \langle \varphi | U(\cdot)\psi \rangle(t) \\ \implies \langle \varphi | U(\cdot)\psi \rangle(t) &= \langle \varphi | \psi \rangle e^{-izt}, \end{aligned}$$

where we have used Corollary 12.10 and the fact that  $U(0) = \text{id}_{\mathcal{H}}$ . But, since  $z$  is purely imaginary, the exponential is unbounded and so the RHS is unbounded. However, the LHS is bounded (as  $U(\cdot)$  is bounded) and so the only way the equality holds is if  $\langle \varphi | \psi \rangle = 0$ . Finally since we took all  $\psi \in \mathcal{H}$  it follows that  $\varphi = \{0_{\mathcal{H}}\}$  and so  $\ker(A^* - \bar{z}) = \{0_{\mathcal{H}}\}$ . The proof for  $\ker(A^* - z)$  follows trivially from this result — i.e. the RHS just becomes unbounded in the opposite direction.

- (iii) We know that  $A$  is essentially self adjoint, which means that  $A^{**}$  is self adjoint. Now construct

$$\tilde{U}(t) := \exp(-itA^{**}) = \int_{\mathbb{R}} e^{it\lambda} P_{A^{**}}(d\lambda).$$

Now let  $\psi \in \mathcal{D}_A^S \subseteq \mathcal{D}_{A^{**}}$  and consider the real family

$$\psi(t) := [\exp(-itA^{**}) - U(t)]\psi.$$

Then

$$\psi'(t) = [-tA^{**} \exp(-itA^{**}) + iAU(t)]\psi = -iA^{**}\psi(t),$$

where we have used the fact that  $A = A^{**}$  on  $\mathcal{D}_A^S$ . Then we have

$$\begin{aligned} (\|\psi(t)\|^2)' &:= \langle \psi(t) | \psi(t) \rangle' \\ &= 2 \operatorname{Re} \langle \psi(t) | \psi'(t) \rangle \\ &= 2 \operatorname{Re} (-i \langle \psi(t) | A^{**} \psi(t) \rangle) \\ &= 0, \end{aligned}$$

where we have used the fact that  $\langle \psi | A^{**} \psi(t) \rangle \in \mathbb{R}$  as  $A^{**}$  is self adjoint. So we have that  $\|\psi(t)\|$  is a constant w.r.t.  $t$ . From the definition, we have  $\psi(0) = 0$  and so  $\|\psi(t)\| = \|\psi(0)\| = 0$ , which from the definition of the norm tells us  $\psi(t) = 0$  for all  $t$ . Finally it follows that

$$\exp(-itA) =: U(t) = \exp(-itA^{**}) \implies A = A^{**}.$$

□

### 12.3 Domains of Essential Self Adjointness ("Cores")

Stone's Theorem showed us that the generator  $A: \mathcal{D}_A^S \rightarrow \mathcal{H}$  is self adjoint. Sometimes a compromise in choosing the domain is in order, as we shall see in the two section's time.

**Corollary 12.11.** *Inspection of the part (ii) of the proof shows that if one considers  $A: \mathcal{D} \rightarrow \mathcal{H}$  for some dense  $\mathcal{D} \subseteq \mathcal{D}_A^S$  that also satisfies  $U(t)\mathcal{D} = \mathcal{D}_A$ , then we  $A$  is essentially self adjoint on  $\mathcal{D}$ .*

### 12.4 Position, Momentum and Angular Momentum

Employ Stone's Theorem to properly and easily define these three operators in quantum mechanical systems. For the rest of this lecture we shall take  $\mathcal{H} = L^2(\mathbb{R}^3, \lambda) =: L^2$  where  $\lambda$  is the Lebesgue measure.

**Definition.** The *position operators*, denoted  $Q^j$  for  $j = 1, 2, 3$ , are defined as the generators of

$$U^j(\cdot): L^2 \rightarrow L^2,$$

with

$$(U^j(t)\psi)(x) := \psi(x)e^{-itx^j},$$

for  $(x) := (x_1, x_2, x_3)$ . That is, they are the self adjoint

$$Q^j: \mathcal{D}_{Q^j}^S \rightarrow L^2$$

with

$$(Q^j \psi)(x) = x^j \psi(x).$$

◇

Remark 12.12. Note clearly  $U(t) \circ U(s) = U(t+s)$ ,  $U(0) = \text{id}_{L^2}$  and  $\|U(t)\psi\| = \|\psi\|$  which tells us  $\|U(t)\| = 1$ , all of which are required for  $U(t)$  to be a UAOPG.

**Definition.** The *momentum operators*, denoted  $P_j$  for  $j = 1, 2, 3$ , are the generators of

$$U_j(\cdot): L^2 \rightarrow L^2,$$

with

$$(U_j(a)\psi)(x) := \psi(\dots, x^j - a, \dots),$$

i.e. they shift the  $j^{\text{th}}$  slot to the right by  $a$ . That is they are the self adjoint operators on their Stone domain that satisfy

$$P_j \psi = -i \partial_j \psi.$$

◇

Remark 12.13. Note this is exactly the definition we used for the action of the operator in Lecture 9.

**Definition.** The *orbital angular momentum operators*, denoted  $L_j$  for  $j = 1, 2, 3$ , are the generators of

$$U_j(\cdot): L^2 \rightarrow L^2,$$

with

$$(U_j(\alpha)\psi)(x) := \psi(D_j(\alpha)x),$$

where  $D_j(\alpha): \mathbb{R}^3 \rightarrow \mathbb{R}^3$  is the operator that describes the rotation about the  $j^{\text{th}}$  axis by angle  $\alpha$ . They satisfy

$$(L_1 \psi)(x) = -i(x^2 \partial_3 \psi - x^3 \partial_2 \psi)$$

$$(L_2 \psi)(x) = -i(x^3 \partial_1 \psi - x^1 \partial_3 \psi)$$

$$(L_3 \psi)(x) = -i(x^1 \partial_2 \psi - x^2 \partial_1 \psi)$$

◇

**Corollary 12.14.** *The spectrum for the orbital angular momentum is contained within the integers;  $\sigma(L_j) \subseteq \mathbb{Z}$  for  $j = 1, 2, 3$ .*

*Proof.* From Stone's theorem we have

$$U_j(\alpha) = \exp(-i\alpha L_j),$$

which together with  $D_j(\alpha + 2\pi) = D_j(\alpha)$  gives

$$\exp(-i2\pi L_j) = \text{id}_{\mathcal{H}}.$$

Then, from the fact that  $L_j$  is self adjoint, we can use the Spectral theorem to decompose both sides

$$\int_{\mathbb{R}} e^{-i2\pi\lambda} P_{L_j}(d\lambda) = \int_{\mathbb{R}} P_{L_j}(d\lambda),$$

and so  $\lambda \in \mathbb{Z}$ . □

## 12.5 Schwartz Space $S(\mathbb{R}^d)$

As we have just seen, Stone's theorem gives us a nice way to define the position, momentum and orbital angular momentum operators. However there are two problems with the definitions we have, both of which relate to their Stone domains. They are

- (i)  $\mathcal{D}_Q^S \neq \mathcal{D}_P^S \neq \mathcal{D}_L^S \neq \mathcal{D}_Q^S$
- (ii)  $\mathcal{D}_Q^S \neq \mathcal{D}_{Q \circ Q}^S \neq \mathcal{D}_{Q \circ Q \circ Q}^S \neq \dots$  and similarly for  $P$  and  $L$ .

This, at first, might not seem like such a big deal but on a second look we see that it means havoc when it comes to trying to define the QM version of kinetic energy as  $(P \circ P)/2m$ . The problem is especially bad when it comes to considering commutators, as highlighted before.

We get around this problem using the compromise given in Corollary 12.11.

**Definition.** The *Schwartz Space* on  $\mathbb{R}^d$ , denoted  $S(\mathbb{R}^d)$ , is the vector space with set

$$S(\mathbb{R}^d) := \{\psi \in C^\infty(\mathbb{R}^d) \mid \sup_{x \in \mathbb{R}^d} |x^\alpha (\partial_\beta \psi)(x)| < \infty, \forall \alpha, \beta \in \mathbb{N}_0^{\times d}\},$$

where

$$\mathbb{N}_0^{\times d} := \underbrace{N_0 \times \dots \times N_0}_{d\text{-fold}},$$

and

$$\begin{aligned} x^\alpha &\equiv x^{(\alpha_1, \dots, \alpha_d)} := (x^1)^{\alpha_1} \dots (x^d)^{\alpha_d}, \\ \partial_\beta &\equiv \partial_{(\beta_1, \dots, \beta_d)} := (\partial_1)^{\beta_1} \dots (\partial_d)^{\beta_d}. \end{aligned}$$

◇

Remark 12.15. The Schwartz Space is also known as the space of rapidly decaying test functions.

Remark 12.16. Clearly the space  $C_c^\infty(\mathbb{R}^d)$ , as defined in footnote 15 in Lecture 9, is a contained within  $S(\mathbb{R}^d)$ .

**Lemma 12.17.** *The Schwartz space is closed under pointwise multiplication; if  $\psi, \varphi \in S(\mathbb{R}^d)$  then  $\psi \bullet \varphi \in S(\mathbb{R}^d)$ . In fact we have the Schwartz algebra  $(S(\mathbb{R}^d), +, \cdot, \bullet)$ .*

*Proof.* This result follows simply from the so called *Leibniz Rule*, which is an extension of the product rule.<sup>21</sup>

$$\begin{aligned} \sup_{x \in \mathbb{R}^d} |x^\alpha (\partial_\beta (\psi \bullet \varphi))(x)| &= \sup_{x \in \mathbb{R}^d} |x^\alpha (\partial_\beta (\psi) \bullet \varphi + \psi \bullet \partial_\beta (\varphi))(x)| \\ &\leq \sup_{x \in \mathbb{R}^d} |x^\alpha (\partial_\beta (\psi) \bullet \varphi)(x)| + \sup_{x \in \mathbb{R}^d} |x^\alpha (\psi \bullet \partial_\beta (\varphi))(x)| \\ &< \infty. \end{aligned}$$

Then, using the fact that the pointwise multiplication of two smooth functions is smooth, we have  $\psi \bullet \varphi \in S(\mathbb{R}^d)$ . Finally using the linearity of everything involved we get the algebra.  $\square$

**Lemma 12.18.** *For  $1 \leq p \leq \infty$  we have  $S(\mathbb{R}^d) \subset L^p(\mathbb{R}^d)$ .*

*Proof.* Let  $\psi \in S(\mathbb{R}^d)$ . Then  $|\psi(x)| < \infty$  for all  $x \in \mathbb{R}^d$ , and so it is integrable. Then Corollary 5.19 tells us that it is measurable with respect to the Borel  $\sigma$ -algebras, so  $\psi \in L^1(\mathbb{R}^d)$ . Then finally from  $L^1(\mathbb{R}^d) \subset L^p(\mathbb{R}^d)$  for all  $p > 1$ , the result follows.  $\square$

**Lemma 12.19.** *One can show that the Fourier Transform is a linear isomorphism from  $S(\mathbb{R}^d)$  onto itself.<sup>22</sup>*

**Theorem 12.20.** *The Schwartz space as defined above satisfies*

- (i)  $S(\mathbb{R}^d) \subseteq L^2(\mathbb{R}^d, \lambda)$  is dense,
- (ii)  $S(\mathbb{R}^3) \subseteq \mathcal{D}_Q^S, \mathcal{D}_P^S, \mathcal{D}_L^S$  is dense,
- (iii)  $Q^j: S(\mathbb{R}^3) \rightarrow S(\mathbb{R}^3)$  is essentially self adjoint. Same for  $P_j$  and  $L_j$ .

Remark 12.21. From the last condition we see that we can repeatedly apply the operators, in any order, to a system. This fixes our problem above.

---

<sup>21</sup>See Dr. Schuller's Lecture's on the Geometric Anatomy of Theoretical Physics for a definition in context.

<sup>22</sup>See lecture 18.



### 13 Spin

In the previous lecture we defined *orbital* angular momentum. The emphasis on ‘orbital’ was not a mistake; this lecture aims to discuss what is referred to as *general* angular momentum (or just angular momentum) in QM. This latter case gets its name from the fact that any concrete set of three operators,  $\{J_1, J_2, J_3\}$  say, obey analogous commutation relations to  $\{L_1, L_2, L_3\}$ .

Recall, for  $\psi \in S(\mathbb{R}^3)$  we have

$$L_i: S(\mathbb{R}^3) \rightarrow S(\mathbb{R}^3),$$

essentially self adjoint with

$$\begin{aligned}(L_1\psi)(x) &= -i(x^2\partial_3\psi - x^3\partial_2\psi), \\ (L_2\psi)(x) &= -i(x^3\partial_1\psi - x^1\partial_3\psi), \\ (L_3\psi)(x) &= -i(x^1\partial_2\psi - x^2\partial_1\psi).\end{aligned}$$

We can, therefore, calculate their commutation relations.

**Lemma 13.1.** *The orbital angular momentum operators obey the following commutation relations:*

$$\begin{aligned}[L_1, L_2] &= iL_3, \\ [L_2, L_3] &= iL_1, \\ [L_3, L_1] &= iL_2.\end{aligned}$$

*Proof.* The proof follows from direct computation. Consider

$$\begin{aligned}[L_1, L_2]\psi &= (L_1 \circ L_2 - L_2 \circ L_1)\psi \\ &= (-i)^2(x^2\partial_3 - x^3\partial_2)(x^3\partial_1\psi - x^1\partial_3\psi) - (-i)^2(x^3\partial_1 - x^1\partial_3)(x^2\partial_3\psi - x^3\partial_2\psi) \\ &= -(x^2\partial_1\psi + x^2x^3\partial_3\partial_1\psi - x^2x^1\partial_3^2\psi - (x^3)^2\partial_2\partial_1\psi + x^3x^1\partial_2\partial_3\psi) \\ &\quad + (x^2x^3\partial_1\partial_3\psi - (x^3)^2\partial_1\partial_2\psi - x^1x^2\partial_3^2\psi + x^1x^3\partial_3\partial_2\psi + x^1\partial_2\psi) \\ &= x^1\partial_2\psi - x^2\partial_1\psi \\ &= iL_3\psi,\end{aligned}$$

where we have used the fact that  $S(\mathbb{R}^3) \subset C^\infty(\mathbb{R}^3) \subset C^2(\mathbb{R}^3)$ , and so we can swap derivative order, i.e.

$$\partial_1\partial_2\psi = \partial_2\partial_1\psi.$$

The same method is used for the other two commutation relations.  $\square$

**Remark 13.2.** The vector space with set  $V := \text{span}_{\mathbb{C}}\{L_1, L_2, L_3\}$  can be defined, and we have that  $iL_j \in V$ , and so the above tells us that  $(V, +, \cdot, [\cdot, \cdot])$  is the orbital angular momentum *Lie algebra*.

Remark 13.3. We can re-write the commutation relations in the compact and convenient form

$$[L_i, L_j] = i\epsilon_{ijk}L_k,$$

where  $\epsilon_{ijk}$  is the totally antisymmetric *Levi-Civita symbol*, defined as

$$\epsilon_{ijk} = \begin{cases} +1 & \text{if } (i, j, k) \text{ is an even permutation of } (1, 2, 3) \\ -1 & \text{if } (i, j, k) \text{ is an odd permutation of } (1, 2, 3) \\ 0 & \text{otherwise.} \end{cases}$$

**Proposition 13.4.** *It is not possible have a common eigenvector between the operators  $(L_1, L_2, L_3)$ .*

*Proof.* Let  $V := \text{span}_{\mathbb{C}}\{L_1, L_2, L_3\}$ . Now assume that  $\psi \in \mathcal{D} \setminus \{0_{\mathcal{H}}\}$  is an eigenvalue of both  $L_1$  and  $L_2$  with eigenvalues  $\lambda, \mu \in \mathbb{C}$ , respectively. Then we have

$$\begin{aligned} [L_1, L_2]\psi &:= L_1(L_2\psi) - L_2(L_1\psi) \\ &= \mu L_1(\psi) - \lambda L_2\psi \\ &= \mu\lambda\psi - \lambda\mu\psi \\ &= 0, \end{aligned}$$

where we have used the linearity of the operators. It follows from the commutation relations that  $L_3\psi = 0$ . However, from the other commutation relations we then have

$$\begin{aligned} [L_2, L_3]\psi &= iL_1\psi \\ L_2(L_3\psi) - L_3(L_2\psi) &= i\lambda\psi \\ L_2(0) - \mu L_3\psi &= i\lambda\psi \\ 0 - \mu 0 &= i\lambda\psi \\ \implies \lambda &= 0, \end{aligned}$$

and similarly you can show  $\mu = 0$ . It follows then that for *any*  $D \in V$  we have  $D\psi = 0$ , which can only be true is  $\psi = 0_{\mathcal{H}}$ . But this contradicts the opening assumption and so it can't be true.  $\square$

Remark 13.5. People often say that "two non-commuting operators have no common eigenvectors", however this statement is not strictly true. What is meant is "two operators, whose commutator does *not* contain the zero vector in its range, do not have common eigenvectors." This is subtly different, however the distinction is important. For example, in the previous proposition if we instead had  $[L_1, L_2] = iL_3$  and  $[L_2, L_3] = 0 = [L_3, L_1]$ , we would not need to require  $\lambda = 0 = \mu$ , and so, unless further constraints were placed on the system of operators, it is possible that  $\psi$  is a common eigenvector to  $L_1$  and  $L_2$ .

### 13.1 General Spin

At this point we might ask why we are bothering to work out these commutation relations? After all they appear to be of no use when it comes to calculating things such as the spectra of the operators (as is evident by Corollary 12.14). The answer to this is that we want to see what information we can obtain about the system (specifically its spectrum) using solely the commutation relations, as then any other set of observables that shares these commutation relations immediately obey the same results.

To emphasise, given a Lie algebra that contains three operators,  $S_1, S_2, S_3$  say, with

$$S_j : \mathcal{D} \rightarrow \mathcal{D},$$

for *some*  $\mathcal{D} \subseteq \mathcal{H}$ , that obey commutation relations analogous to those of Lemma 13.1, will instantly satisfy any results we derive, using only the commutation relations, for the orbital angular momentum.

Example 13.6. An example of such a set of operators are the so-called *Pauli spin algebra*, which has

$$S_i := \frac{1}{2}\sigma_i,$$

with  $\mathcal{D} = \mathcal{H} = \mathbb{C}^2$ , where

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

known as the *Pauli spin matrices*. It is easily checked, through the rules of matrix multiplication, that this algebra obeys the correct commutation relations. This is an example of a so-called spin- $\frac{1}{2}$  system.

Remark 13.7. We can not expect the commutation relations to necessarily tell us everything about the spectrum (or any other quantity we try to calculate) as they can be derived by several different operator sets with potentially differing spectra. But, as said above, whatever we can infer from the commutation relations alone must hold for all the operator sets.

### 13.2 Derivation of Pure Point Spectrum

We start from a general Lie algebra with our required conditions. We shall denote the operators by  $J_1, J_2, J_3$ , however they need not be the orbital angular momentum operators. Equally the domain,  $\mathcal{D}$ , is left arbitrary, up the condition that the operators are at least essentially self adjoint on them.

**Definition.** A Casimir operator for the algebra  $(V, +, \cdot, [\cdot, \cdot])$ , is a symmetric operator

$$\Omega : \mathcal{D} \rightarrow \mathcal{D}$$

that commutes with every element in  $V$ . ◇

*Remark 13.8.* Note, due to the bilinearity of the commutator, we only need to check that  $\Omega$  commutes with the basis elements of  $V$ .

**Proposition 13.9.** *The operator*

$$\Omega := J_1 \circ J_1 + J_2 \circ J_2 + J_3 \circ J_3$$

*is a Casimir operator for the algebras we're considering.*

*Proof.* The symmetric part follows trivially from the fact that Corollary 12.11 tells us  $J_1, J_2, J_3$  are all symmetric. Next let  $\psi \in \mathcal{D}$  and consider

$$\begin{aligned} [\Omega, J_1]\psi &:= [J_1 \circ J_1 + J_2 \circ J_2 + J_3 \circ J_3, J_1]\psi \\ &= [J_1 \circ J_1, J_1]\psi + [J_2 \circ J_2, J_1]\psi + [J_3 \circ J_3, J_1]\psi \\ &= (J_1 \circ J_1 \circ J_1)\psi - (J_1 \circ J_1 \circ J_1)\psi \\ &\quad + (J_2 \circ J_2 \circ J_1)\psi - (J_1 \circ J_2 \circ J_2)\psi \\ &\quad + (J_3 \circ J_3 \circ J_1)\psi - (J_1 \circ J_3 \circ J_3)\psi \\ &= (J_2 \circ J_1 \circ J_2)\psi + (J_2 \circ [J_2, J_1])\psi - (J_1 \circ J_2 \circ J_2)\psi \\ &\quad + (J_3 \circ J_1 \circ J_3)\psi + (J_3 \circ [J_3, J_1])\psi - (J_1 \circ J_3 \circ J_3)\psi \\ &= (J_1 \circ J_2 \circ J_2)\psi + ([J_2, J_1] \circ J_2)\psi + (J_2 \circ [J_2, J_1])\psi - (J_1 \circ J_2 \circ J_2)\psi \\ &\quad + (J_1 \circ J_3 \circ J_3)\psi + ([J_3, J_1] \circ J_3)\psi + (J_3 \circ [J_3, J_1])\psi - (J_1 \circ J_3 \circ J_3)\psi \\ &= -i(J_3 \circ J_2)\psi - i(J_2 \circ J_3)\psi + i(J_2 \circ J_3)\psi + i(J_3 \circ J_2)\psi \\ &= 0, \end{aligned}$$

which because  $\psi \in \mathcal{D}$  was arbitrary tells us  $[\Omega, J_1] = 0$ . The same method gives  $[\Omega, J_2] = 0 = [\Omega, J_3]$ .  $\square$

**Definition.** Let  $J_1, J_2, J_3$  be three operators that satisfy our conditions. Then define

$$\begin{aligned} J_+ &:= J_1 + iJ_2, \\ J_- &:= J_1 - iJ_2, \end{aligned}$$

known as the *ladder operators*, for a reason that will soon become clear.  $\diamond$

*Remark 13.10.* We can choose to consider the set  $\{J_+, J_-, J_3\}$  in place of the set  $\{J_1, J_2, J_3\}$  while still keeping all the information — as we can simply reconstruct  $J_1$  and  $J_2$  from  $J_+$  and  $J_-$ . Note, however, in doing this we have broken the symmetry of the algebra (in the sense that none of the  $J_j$ s are special, they all obey the same commutation relations) by singling out  $J_3$ , while taking linear combinations of  $J_1$  and  $J_2$ . Indeed we did not need to make this choice of symmetry breaking, but in fact we could have chosen to keep  $J_1$  while defining  $J_+$  and  $J_-$  as linear combinations of  $J_2$  and  $J_3$ . Importantly, the results that follow will hold equally for whichever  $J$  we choose, and so in order to stick with convention we pick  $J_3$ . Note also that we no longer have a set of observables as  $(J_+)^* = J_-$  and  $(J_-)^* = J_+$ .

**Lemma 13.11.**  $J_+$  and  $J_-$  satisfy the following commutation relations

$$\begin{aligned}[J_+, J_-] &= 2J_3, \\ [J_3, J_\pm] &= \pm J_\pm, \\ [\Omega, J_\pm] &= 0.\end{aligned}$$

*Proof.* From direct computation. Not done here to save space.  $\square$

**Lemma 13.12.** We can rewrite our Casimir operator as

$$\Omega = J_+ \circ J_- + J_3 \circ (J_3 - \text{id}_{\mathcal{D}}),$$

or equally as

$$\Omega = J_- \circ J_+ + J_3 \circ (J_3 + \text{id}_{\mathcal{D}}).$$

*Proof.* We shall just show the first one:

$$\begin{aligned}J_+ \circ J_- + J_3 \circ (J_3 - \text{id}_{\mathcal{D}}) &:= (J_1 + iJ_2) \circ (J_1 - iJ_2) + J_3 \circ (J_3 - \text{id}_{\mathcal{D}}) \\ &= J_1 \circ J_1 + J_2 \circ J_2 - i(J_1 \circ J_2 - J_2 \circ J_1) + J_3 \circ J_3 - J_3 \\ &= J_1 \circ J_1 + J_2 \circ J_2 + J_3 \circ J_3 - i[J_1, J_2] - J_3 \\ &= J_1 \circ J_1 + J_2 \circ J_2 + J_3 \circ J_3 - (i)^2 J_3 - J^3 \\ &= \Omega.\end{aligned}$$

$\square$

*Remark 13.13.* Both of the expressions in the above definition are *always* true. It is not that one is true under certain circumstances and then the other is true. This is an important observation that we shall use.

At this point we might wonder why we are going through so much effort introducing the Casimir, when what we're looking for is the spectra of the operators  $J_1, J_2, J_3$ . The answer is to make the problem *seemingly* more complicated by now considering only eigenvectors that are *common* to both  $J_3$  and  $\Omega$ . Note it is necessary that they commute if they are to have common eigenvectors — as is easily verified from the definition of the commutator. That is we want to find a  $\psi_{\lambda, \mu} \in \mathcal{D} \setminus \{0\}$ <sup>23</sup> such that

$$\begin{aligned}J_3 \psi_{\lambda, \mu} &= \mu \psi_{\lambda, \mu} \\ \Omega \psi_{\lambda, \mu} &= \lambda \psi_{\lambda, \mu},\end{aligned}$$

where the subscript is included in order to label the eigenvector by its eigenvalues.

Again this appears to be a more complicated problem — we now not only need to check our  $\psi$  is an eigenvector of  $J_3$  but we also need to check that it's an eigenvalue of  $\Omega$ . However, we can show that every eigenvector of  $J_3$  (and equally for  $J_1$  and  $J_2$ ) is also an eigenvector of  $\Omega$ . For a proof of this see [Peter Ferguson's well detailed answer on Quora](#).

---

<sup>23</sup>Recall that an eigenvector can not be the zero-vector by definition. We shall use this later.

We might also ask whether this will give us the spectrum of  $J_3$  anyways, as  $J_3$  is only essentially self adjoint on our Schwartz domain,  $\mathcal{D}$ . We are OK, however as  $(J_3)^{**}$  is self adjoint on the  $\mathcal{D}$  and so we could just use this instead, where the operation is defined to be the same as  $J_3$  — just as we did for the momentum operator in Lecture 9.

**Lemma 13.14.** *The eigenvalues for these common eigenvectors satisfy*

$$\lambda \geq |\mu|(|\mu| + 1).$$

*Proof.* We shall consider both cases for the rewriting of  $\Omega$ , however, as explained above, the cases bracket does not mean one is true under certain conditions and the other otherwise, they are both true. We shall also drop the  $\circ$  symbols to lighten notation. Thus we have

$$\begin{aligned} \lambda \langle \psi_{\lambda,\mu} | \psi_{\lambda,\mu} \rangle &= \langle \psi_{\lambda,\mu} | \Omega \psi_{\lambda,\mu} \rangle \\ &= \begin{cases} \langle \psi_{\lambda,\mu} | (J_+ J_- + J_3(J_3 - \text{id}_{\mathcal{D}})) \psi_{\lambda,\mu} \rangle \\ \langle \psi_{\lambda,\mu} | (J_- J_+ + J_3(J_3 + \text{id}_{\mathcal{D}})) \psi_{\lambda,\mu} \rangle \end{cases} \\ &= \begin{cases} \langle J_- \psi_{\lambda,\mu} | J_- \psi_{\lambda,\mu} \rangle + \mu(\mu - 1) \langle \psi_{\lambda,\mu} | \psi_{\lambda,\mu} \rangle \\ \langle J_+ \psi_{\lambda,\mu} | J_+ \psi_{\lambda,\mu} \rangle + \mu(\mu + 1) \langle \psi_{\lambda,\mu} | \psi_{\lambda,\mu} \rangle, \end{cases} \end{aligned}$$

where we have used the fact that  $(J_-)^* = J_+$  and vice versa.<sup>24</sup>

Now recalling that  $\psi_{\lambda,\mu}$  is an eigenvector, and so, by definition, not the zero vector we know the inner product is positive definite, and thus we can divide by it, giving

$$\lambda = \begin{cases} \frac{\|J_- \psi_{\lambda,\mu}\|^2}{\|\psi\|^2} + \mu(\mu - 1) \\ \frac{\|J_+ \psi_{\lambda,\mu}\|^2}{\|\psi\|^2} + \mu(\mu + 1). \end{cases}$$

Finally, from the fact that the norm is non-negative definite (i.e. the first term in each case is either positive or vanishes) we have

$$\begin{aligned} \lambda &\geq \begin{cases} \mu(\mu - 1) \\ \mu(\mu + 1) \end{cases} \\ &= \begin{cases} -\mu(-\mu + 1) \\ \mu(\mu + 1), \end{cases} \end{aligned}$$

and so it follows that  $\lambda \geq |\mu|(|\mu| + 1)$ . □

**Lemma 13.15.** *The elements  $J_{\pm} \psi_{\lambda,\mu}$  are common ‘eigenvectors’ of  $\Omega$  and  $J_3$  with eigenvalues  $\lambda$  and  $(\mu \pm 1)$ , respectively.*

*Proof.* First consider  $\Omega$ . We have

$$\begin{aligned} \Omega J_{\pm} \psi_{\lambda,\mu} &= J_{\pm} \Omega \psi_{\lambda,\mu} + [\Omega, J_{\pm}] \psi_{\lambda,\mu} \\ &= J_{\pm} (\lambda \psi_{\lambda,\mu}) \\ &= \lambda (J_{\pm} \psi_{\lambda,\mu}), \end{aligned}$$

---

<sup>24</sup>Really what you need to do is expand out  $J_-$  and  $J_+$  in terms of  $J_1$  and  $J_2$  and then take the adjoint. The result holds.

where we used the fact that  $[\Omega, J_\pm] = 0$ , and the linearity of  $J_\pm$ .

Now consider  $J_3$ ,


$$\begin{aligned} J_3 J_\pm \psi_{\lambda, \mu} &= J_\pm J_3 \psi_{\lambda, \mu} + [J_3, J_\pm] \psi_{\lambda, \mu} \\ &= \mu(J_\pm \psi_{\lambda, \mu}) \pm J_\pm \psi_{\lambda, \mu} \\ &= (\mu \pm 1)(J_\pm \psi_{\lambda, \mu}), \end{aligned}$$

where again we used the commutator,  $[J_3, J_\pm] = \pm J_\pm$ . □

Remark 13.16. The previous allows us to conclude that  $J_\pm \psi_{\lambda, \mu} \propto \psi_{\lambda, \mu \pm 1}$ .

Remark 13.17. This is why  $J_\pm$  are known as *ladder operators*, with  $J_+$  known as the *raising* operator and  $J_-$  the *lowering* operator. As we see these names derive from the eigenvalues they produce as eigenvectors of  $J_z$ .

If the  $J_z$  eigenvalue of  $\psi_{\lambda, \mu}$  corresponds to the  $\mu$ -th rung of a ladder then the eigenvalue of  $J_+ \psi_{\lambda, \mu}$  corresponds to the  $(\mu + 1)$ -th rung, and  $J_- \psi_{\lambda, \mu}$  the  $(\mu - 1)$ -th. Note each one of the rungs is separated by exactly the same distance, and that we get the  $(\mu + n)$ -th rung from  $(J_+)^n \psi_{\lambda, \mu}$  and similarly for the  $(\mu - n)$ -th rung.



The next question would be is this a ‘proper’ ladder; that is does it have a top and bottom rung or does it continue forever? The answer comes in the form of the next lemma.

**Lemma 13.18.** *There exists a  $\psi_{\lambda, \bar{\mu}}$  such that  $J_+ \psi_{\lambda, \bar{\mu}} = 0$ . Equally there exists a  $\psi_{\lambda, \underline{\mu}}$  such that  $J_- \psi_{\lambda, \underline{\mu}} = 0$ .*

*Proof.* We know from Lemma 13.14 that  $|\mu|(|\mu| + 1) \leq \lambda$  holds for *any* common eigenvector of  $\Omega$  and  $J_3$ . We see from Lemma 13.15 that  $(J_\pm)^n \psi_{\lambda, \mu}$  is such a common eigenvector, and so must obey  $|\mu \pm n|(|\mu \pm n| + 1) \leq \lambda$ . However,  $\lambda$  is unchanged by this repeated application of the ladder operators, and so, unless remedied, this inequality will eventually be broken – that is we need to somehow cap the available  $n$  values.

Consider first the raising operator. In this case  $\mu + n$  gets bigger and bigger, and so we need to cap  $n$  from above. In other words, we require there to be an  $m \in \mathbb{N}$  such that for all  $n > m$ ,  $(J_+)^n \psi_{\lambda, \mu} = 0$ . This fixes our problem as this corresponds to the zero vector

and so, by definition, it cannot be an eigenvector, and the  $\lambda$  inequality no longer need hold. We define

$$\psi_{\lambda, \bar{\mu}} := (J_+)^m \psi_{\lambda, \mu}.$$

The idea is exactly the same is true for the lowering operator, however now  $\mu - n$  is getting smaller and smaller, and so its modulus (after  $n > \mu$  is reached) gets bigger and bigger. So again we need to cap  $n$  from above. We require there to be a  $\ell \in \mathbb{N}$  such that for all  $n > \ell$ ,  $(J_-)^n \psi_{\lambda, \mu} = 0$ . We define

$$\psi_{\lambda, \underline{\mu}} := (J_-)^\ell \psi_{\lambda, \mu}.$$

Note we do not have any a priori relation between the values of  $m$  and  $\ell$ . To use the ladder analogy,  $m$  is the number of rungs above  $\mu$ -th rung and  $\ell$  is the number of rungs below the  $\mu$ -th rung. For a given  $\lambda$ , the highest value of  $\mu$  is denoted  $\bar{\mu}(\lambda)$  and the lowest value  $\underline{\mu}(\lambda)$ .  $\square$

*Remark 13.19.* Note the above tells us that  $J_\pm \psi$  are not strictly eigenvalues, as it could be the zero vector. This is why we wrote ‘eigenvectors’ in inverted commas in Lemma 13.15.

**Proposition 13.20.** *The maximum and minimum values of  $\mu$  satisfy*

- (i)  $\lambda = \bar{\mu}(\lambda)(\bar{\mu}(\lambda) + 1)$ ,
- (ii)  $\underline{\mu}(\lambda) = -\bar{\mu}(\lambda)$ ,
- (iii)  $\bar{\mu}(\lambda) \in \frac{\mathbb{N}_0}{2}$ .

*Proof.* (i) From the proof of Lemma 13.14, and the fact that  $J_+ \psi_{\lambda, \bar{\mu}(\lambda)} = 0$ , and so  $\|J_+ \psi_{\lambda, \bar{\mu}(\lambda)}\| = 0$ , we have

$$\lambda = \bar{\mu}(\lambda)(\bar{\mu}(\lambda) + 1)$$

(ii) Repeating the above argument but with the fact that  $\|J_- \psi_{\lambda, \underline{\mu}(\lambda)}\| = 0$ , we have

$$\begin{aligned} \lambda &= \underline{\mu}(\lambda)(\underline{\mu}(\lambda) - 1) \\ &= -\underline{\mu}(\lambda)(-\underline{\mu}(\lambda) + 1), \end{aligned}$$

and so  $\underline{\mu}(\lambda) = -\bar{\mu}$ .

(iii) From the previous, along with the fact that  $\bar{\mu}(\lambda) - \underline{\mu}(\lambda) \in \mathbb{N}_0$  this result follows trivially.  $\square$

*Remark 13.21.* In order to be consistent with the literature we shall introduce the following relabelling

$$j := \bar{\mu}(\lambda), \quad m := \mu.$$

Note we have  $j \in \frac{\mathbb{N}_0}{2}$ .



**Theorem 13.22.** *The common eigenvectors of  $\Omega$  and  $J_3$  come as families  $\psi_{j(j+1),m}$ , where  $m = -j, -j+1, \dots, j-1, j$ . The eigenvalue  $j(j+1)$  is associated to  $\Omega$  and  $m$  is associated to  $J_3$ .*

One normally normalises these eigenvectors and defines

$$\Phi_{j,m} := \frac{\psi_{j(j+1),m}}{\|\psi_{j(j+1),m}\|}.$$

Then, from Lemma 8.8 and the fact that the eigenvectors have distinct eigenvalues, we have

$$\langle \Phi_{j,m} | \Phi_{k,n} \rangle = \delta_{jk} \delta_{mn}.$$

**Corollary 13.23.** *We have  $m \in \frac{\mathbb{Z}}{2}$ .*

*Proof.* This comes from just allowing  $j \in \frac{\mathbb{N}_0}{2}$  to be any element and then using  $m = -j, \dots, j$ .  $\square$

**Proposition 13.24.** *The common eigenvectors  $\Phi_{j,m}$  satisfy*

$$J_{\pm} \Phi_{j,m} = \sqrt{j(j+1) - m(m \pm 1)} \Phi_{j,m \pm 1}.$$

*Proof.* We shall show this for  $J_+$ , the method of  $J_-$  follows analogously. Recall from Lemma 13.12 that

$$J_- J_+ = \Omega - J_3(J_3 + \text{id}_{\mathcal{D}}).$$

Now consider

$$\begin{aligned} \langle J_+ \Phi_{j,m} | J_+ \Phi_{j,m} \rangle &= \langle \Phi_{j,m} | J_- J_+ \Phi_{j,m} \rangle \\ &= \langle \Phi_{j,m} | \Omega \Phi_{j,m} \rangle - \langle \Phi_{j,m} | J_3(J_3 + \text{id}_{\mathcal{D}}) \Phi_{j,m} \rangle \\ &= j(j+1) \langle \Phi_{j,m} | \Phi_{j,m} \rangle - m(m+1) \langle \Phi_{j,m} | \Phi_{j,m} \rangle \\ &= j(j+1) - m(m+1). \end{aligned}$$

Combining this with Remark 13.16 allows us to conclude the result.  $\square$

### 13.3 Pure Spin- $j$ Systems

**Definition.** A quantum mechanical system is called a *pure spin- $j$*  system if its Hilbert space is  $(2j+1)$ -dimensional that possesses an orthonormal eigenbasis  $\{\Phi_{j,m}\}$  for the three operators  $J_1, J_2, J_3$  defined on  $\mathcal{H}$ .  $\diamond$

**Corollary 13.25.** *The Hilbert space is isomorphic to  $\mathbb{C}^{2j+1}$ .*

**Corollary 13.26.** *For a pure spin- $j$  system the spectrum of the operators is*

$$\sigma(J_i) = \{-j, -j+1, \dots, j-1, j\},$$

for  $i = 1, 2, 3$ .

Example 13.27.

$j$	$\sigma(J_i), i = 1, 2, 3$
0	0
1/2	$\{-1/2, 1/2\}$
1	$\{-1, 0, 1\}$

Remark 13.28. When you introduce spin to a particle, its Hilbert space becomes a product space. For example for an electron (spin-1/2) in  $\mathbb{R}^3$  its Hilbert space is

$$\mathcal{H}_e = L^2(\mathbb{R}^3) \otimes \mathbb{C}^2.$$

We shall return to this and expand on it in the next lecture.

Remark 13.29. You can also have non-pure spin systems. For the orbital angular momentum, you take a direct sum of the Hilbert spaces. That is if  $\mathcal{H}_j$  is the Hilbert space associated to the pure spin- $j$  system then the composite system's Hilbert space is

$$\mathcal{H}_{comp} = \bigoplus_j \mathcal{H}_j.$$

We shall return to this in two lectures time.

## 14 Composite Systems

Recall Axiom 1, which says that to every quantum system there is an underlying Hilbert space. The question we now want to ask is: Let  $\mathcal{H}_1$  be the Hilbert space associated to one system and  $\mathcal{H}_2$  be the Hilbert space associated to another. What is the underlying Hilbert space associated to the composite system?

To clarify what we mean, imagine having a proton and an electron. We first look at the proton by itself and call this system one. We then look at the electron separately and call that system two. We now want to look at both of them together, but we wish to use the fact that we have already studied them separately to simplify the problem. It may seem ‘natural’ to model the composite  $\mathcal{H}$  as the so called *direct sum*, which as a set is<sup>25</sup>

$$\mathcal{H}_1 \oplus \mathcal{H}_2 := \{(\psi, \varphi) \mid \psi \in \mathcal{H}_1, \varphi \in \mathcal{H}_2\},$$

and where the linearity is inherited from  $\mathcal{H}_1$  and  $\mathcal{H}_2$ , namely

$$(a\psi_1 + \psi_2, b\varphi_1 + \varphi_2) = ab(\psi_1, \varphi_1) + a(\psi_1, \varphi_2) + b(\psi_2, \varphi_1) + (\psi_2, \varphi_2).$$

This is what we do in classical systems and it tells us that if we know everything about the states<sup>26</sup> of our two systems, then we also know everything about the states of the composite system.

However, as with all things quantum, things are more complicated, and the above is not the case. The main problem comes from the fact that not all linear combinations of elements of the form  $(\psi, \varphi)$  can also be written in that form.

*Example 14.1.* Let  $\psi_1, \psi_2 \in \mathcal{H}_1$ ,  $\varphi_1, \varphi_2 \in \mathcal{H}_2$  and  $a, b \in \mathbb{C}$ . Then, assuming the linearity as above, we have

$$\begin{aligned} a(\psi_1, \varphi_1) + b(\psi_2, \varphi_2) &= (a\psi_1, \varphi_1) + (\psi_2, b\varphi_2) \\ &= (a\psi_1 + \psi_2, \varphi_1 + b\varphi_2) \\ &= a(\psi_1, \varphi_1) + b(\psi_2, \varphi_2) + ab(\psi_1, \varphi_2) + (\psi_2, \varphi_1), \end{aligned}$$

a clear problem.

Note this example actually tells us that we  $\mathcal{H}_1 \oplus \mathcal{H}_2$  is not closed under the linearity, and so would not be a vector space. We could just restrict ourselves to elements that do obey these rules, however, as we shall see when considering entanglement, we require elements of this form in our underlying Hilbert space.

This calls for a slight refinement of axiom one. We add the addendum<sup>27</sup>:

If a quantum system is composed of two (and hence, by induction, any finite number) of ‘sub’systems, then its underlying Hilbert space is the *tensor product space*  $\mathcal{H}_1 \otimes \mathcal{H}_2$ , equipped with a inner product.

<sup>25</sup>This definition holds as we are only taking the direct product of two spaces, and so the index set is finite. See [wiki](#) for details on this.

<sup>26</sup>Recall that the elements of the Hilbert space are *not* the states, but are associated to them. We shall return to this at the end of the lecture.

<sup>27</sup>We shall define what these new terms are in the next section.

Example 14.2. Remark 13.28 is an example of such a composite system.

### 14.1 Tensor Product of Hilbert Spaces

In order to give a nice definition for the tensor product of two vector spaces, we first need to introduce the so called *free vector space*.

**Definition.** Let  $V$  be a  $\mathbb{F}$ -vector space and let  $B \subseteq V$  be a generating subset of  $V$  (i.e. any element of  $V$  can be obtained via finite linear combinations of elements of  $B$ ). The *free vector space* is

$$F(B) := \text{span}_{\mathbb{F}}(B),$$

i.e. the set of all linear combinations of elements of  $B$ . ◇

**Lemma 14.3.** *Every vector space is a free vector space with  $B$  being a Hamel basis.*

Remark 14.4. Note it need not be true that  $F(B) = V$ , as it might be the case that the same element in  $V$  is reached via two different linear combinations of elements of  $B$ . In fact if  $F(B) = V$ , then  $B$  is just a Hamel basis.

The free vector space for vector spaces might seem almost redundant, given that every vector space has a basis. However if your vector space is countably infinite then such a basis might be incredibly difficult to construct. However you can simply take the entire set for  $B$  and construct the free vector space  $F(V)$ , which will be a huge set, mind. Note, then, that any linear combination of elements in this set is automatically still in the set, and so it is indeed a vector space.

**Definition.** Let  $V$  and  $W$  be two  $\mathbb{F}$ -vector spaces, and let  $A \subseteq V$  and  $B \subseteq W$  be generating subsets. Then we define their tensor product as the vector space with set

$$V \otimes W := F(A \times B) / \sim,$$

where  $\times$  is the Cartesian product and  $\sim$  is an equivalence relation such that: if  $a, a_1, a_2 \in A$ ,  $b, b_1, b_2 \in B$  and  $f \in \mathbb{F}$  then

- (i)  $(a, b) \sim (a, b)$ ,
- (ii)  $(a_1, b_1) + (a_1, b_2) \sim (a_1, b_1 + b_2)$  and  $(a_1, b_1) + (a_2, b_1) \sim (a_1 + a_2, b_1)$ , and continued by induction,
- (iii)  $f(a, b) \sim (fa, b)$  and  $f(a, b) \sim (a, fb)$ .
- (iv) Combinations of (ii) and (iii), e.g.  $(a_1, b_1) + f(a_1, b_2) \sim (a_1, b_1 + fb_2)$ .

◇

Remark 14.5. Note the equivalence relation looks a lot like a linearity condition on  $V \otimes W$ , however on closer inspection it is not quite. The linearity condition that make  $V \otimes W$  into a vector space is simply

$$f(a_1, b_1) + (a_2, b_2) \in V \otimes W.$$

This, in itself, does not need to satisfy the equivalence relation. However, if we did not include it we could end up with a huge redundancy in elements, as a repercussion of Remark 14.4. This equivalence relation makes the corresponding set of equivalence classes a vector space in the way we normally think of them (there is no repeated elements).

This is exactly the type of structure we need to overcome the problem highlighted before (that not all linear combinations can be expressed as a single term), as now we only require that linear combinations of linear combinations are linear combinations, which they obviously are.

**Proposition 14.6.** *Let  $\mathcal{H}_1$  and  $\mathcal{H}_2$  be our two vector spaces. We can define the map*

$$+_{\mathcal{H}_1 \otimes \mathcal{H}_2} : (\mathcal{H}_1 \otimes \mathcal{H}_2) \times (\mathcal{H}_1 \otimes \mathcal{H}_2) \rightarrow (\mathcal{H}_1 \otimes \mathcal{H}_2) \\ [(\psi, \varphi_1), [(\psi, \varphi_2)]] \mapsto [(\psi, \varphi_1)] +_{\mathcal{H}_1 \otimes \mathcal{H}_2} [(\psi, \varphi_2)] := [(\psi, \varphi_1) + (\psi, \varphi_2)],$$

where the additions inside the brackets are w.r.t.  $\mathcal{H}_1$  and  $\mathcal{H}_2$ .

*Proof.* We need to show this is well defined. We shall write  $+_{12}$  now in order to lighten notation. Consider it case by case.

(i) Assume  $(\widetilde{\psi}, \widetilde{\varphi}_1) = (\psi, \varphi_1)$ .

(a) If  $(\widetilde{\psi}, \widetilde{\varphi}_2) = (\psi, \varphi_2)$ , then it follows trivially that

$$[(\widetilde{\psi}, \widetilde{\varphi}_1) + (\widetilde{\psi}, \widetilde{\varphi}_2)] = [(\psi, \varphi_1) + (\psi, \varphi_2)],$$

and so

$$[(\widetilde{\psi}, \widetilde{\varphi}_1)] +_{12} [(\widetilde{\psi}, \widetilde{\varphi}_2)] = [(\psi, \varphi_1)] +_{12} [(\psi, \varphi_2)].$$

(b) If  $(\widetilde{\psi}, \widetilde{\varphi}_2) = (\psi, \varphi_2^1) + (\psi, \varphi_2^2)$ , where  $\varphi_2 = \varphi_2^1 + \varphi_2^2$ , we have

$$\begin{aligned} [(\widetilde{\psi}, \widetilde{\varphi}_1) + (\widetilde{\psi}, \widetilde{\varphi}_2)] &= [(\psi, \varphi_1) + (\psi, \varphi_2^1) + (\psi, \varphi_2^2)] \\ &= [(\psi, \varphi_1 + \varphi_2^1 + \varphi_2^2)] \\ &= [(\psi, \varphi_1 + \varphi_2)] \\ &= [(\psi, \varphi_1) + (\psi, \varphi_2)], \end{aligned}$$

and so

$$[(\widetilde{\psi}, \widetilde{\varphi}_1)] +_{12} [(\widetilde{\psi}, \widetilde{\varphi}_2)] = [(\psi, \varphi_1)] +_{12} [(\psi, \varphi_2)].$$

(c) If  $(\widetilde{\psi}, \widetilde{\varphi}_2) = f(\psi, \varphi_2^3)$ , where  $\varphi_2 = f\varphi_2^3$ , then

$$\begin{aligned} [(\widetilde{\psi}, \widetilde{\varphi}_1) + (\widetilde{\psi}, \widetilde{\varphi}_2)] &= [(\psi, \varphi_1) + f(\psi, \varphi_2^3)] \\ &= [(\psi, \varphi_1) + (\psi, f\varphi_2^3)] \\ &= [(\psi, \varphi_1) + (\psi, \varphi_2)], \end{aligned}$$

and so

$$[(\widetilde{\psi}, \widetilde{\varphi}_1)] +_{12} [(\widetilde{\psi}, \widetilde{\varphi}_2)] = [(\psi, \varphi_1)] +_{12} [(\psi, \varphi_2)].$$

(ii) Assume  $(\tilde{\psi}, \tilde{\varphi}_1) = (\psi, \varphi_1^1) + (\psi, \varphi_1^2)$  where  $\varphi_1 = \varphi_1^1 + \varphi_1^2$ .

(a) If  $(\tilde{\psi}, \tilde{\varphi}_2) = (\psi, \varphi_2)$  then we have essentially the same as (i)(b), so we wont re-write it here.

(b) If  $(\tilde{\psi}, \tilde{\varphi}_2) = (\psi, \varphi_2^1) + (\psi, \varphi_2^2)$ , where  $\varphi_2 = \varphi_2^1 + \varphi_2^2$ , we have

$$\begin{aligned} [(\tilde{\psi}, \tilde{\varphi}_1) + (\tilde{\psi}, \tilde{\varphi}_2)] &= [(\psi, \varphi_1^1) + (\psi, \varphi_1^2) + (\psi, \varphi_2^1) + (\psi, \varphi_2^2)] \\ &= [(\psi, \varphi_1^1 + \varphi_1^2 + \varphi_2^1 + \varphi_2^2)] \\ &= [(\psi, \varphi_1 + \varphi_2)] \\ &= [(\psi, \varphi_1) + (\psi, \varphi_2)], \end{aligned}$$

and so

$$[(\tilde{\psi}, \tilde{\varphi}_1)] +_{12} [(\tilde{\psi}, \tilde{\varphi}_2)] = [(\psi, \varphi_1)] +_{12} [(\psi, \varphi_2)].$$

(c) If  $(\tilde{\psi}, \tilde{\varphi}_2) = f(\psi, \varphi_2^3)$  where  $\varphi_2 = f\varphi_2^3$ , then we have

$$\begin{aligned} [(\tilde{\psi}, \tilde{\varphi}_1) + (\tilde{\psi}, \tilde{\varphi}_2)] &= [(\psi, \varphi_1^1) + (\psi, \varphi_1^2) + f(\psi, \varphi_2^3)] \\ &= [(\psi, \varphi_1^1 + \varphi_1^2 + f\varphi_2^3)] \\ &= [(\psi, \varphi_1 + \varphi_2)] \\ &= [(\psi, \varphi_1) + (\psi, \varphi_2)] \end{aligned}$$

and so

$$[(\tilde{\psi}, \tilde{\varphi}_1)] +_{12} [(\tilde{\psi}, \tilde{\varphi}_2)] = [(\psi, \varphi_1)] +_{12} [(\psi, \varphi_2)].$$

(iii) Assume  $(\tilde{\psi}, \tilde{\varphi}_1) \sim g(\psi, \varphi_1^3)$ , where  $\varphi_1 = g\varphi_1^3$ .

(a) If  $(\tilde{\psi}, \tilde{\varphi}_2) = (\psi, \varphi_2)$ , then we have basically same as (i)(c) and so we wont write it again.

(b) If  $(\tilde{\psi}, \tilde{\varphi}_2) = (\psi, \varphi_2^1) + (\psi, \varphi_2^2)$ , where  $\varphi_2 = \varphi_2^1 + \varphi_2^2$ , then we are basically the same as (ii)(c) and so we wont write it again.

(c) If  $(\tilde{\psi}, \tilde{\varphi}_2) = f(\psi, \varphi_2^3)$ , where  $\varphi_2 = f\varphi_2^3$  then

$$\begin{aligned} [(\tilde{\psi}, \tilde{\varphi}_1) + (\tilde{\psi}, \tilde{\varphi}_2)] &= [g(\psi, \varphi_1^3) + f(\psi, \varphi_2^3)] \\ &= [(\psi, g\varphi_1^3 + f\varphi_2^3)] \\ &= [(\psi, \varphi_1 + \varphi_2)] \\ &= [(\psi, \varphi_1) + (\psi, \varphi_2)] \end{aligned}$$

and so

$$[(\tilde{\psi}, \tilde{\varphi}_1)] +_{12} [(\tilde{\psi}, \tilde{\varphi}_2)] = [(\psi, \varphi_1)] +_{12} [(\psi, \varphi_2)].$$

□

*Remark 14.7.* We can do exactly the same thing but for a map that has the first element different and the second element the same.

**Definition.** Let  $\mathcal{H}_1$  and  $\mathcal{H}_2$  be complex Hilbert spaces with sesqui-linear inner products  $\langle \cdot | \cdot \rangle_{\mathcal{H}_1}$  and  $\langle \cdot | \cdot \rangle_{\mathcal{H}_2}$ , respectively. Then the *composite Hilbert space* is the Hilbert space with set

$$\mathcal{H}_1 \otimes \mathcal{H}_2 := \overline{F(\mathcal{H}_1 \times \mathcal{H}_2)} / \sim,$$

where the overline indicates the topological closure, and with sesqui-linear inner product: for  $\psi_1, \psi_2 \in \mathcal{H}_1$  and  $\varphi_1, \varphi_2 \in \mathcal{H}_2$ ,

$$\langle [(\psi_1, \varphi_1)] | [(\psi_2, \varphi_2)] \rangle_{\mathcal{H}_1 \otimes \mathcal{H}_2} := \langle \psi_1 | \psi_2 \rangle_{\mathcal{H}_1} \cdot \langle \varphi_1 | \varphi_2 \rangle_{\mathcal{H}_2},$$

extended by linearity, with respect to which the closure is taken (i.e. the topology is derived from here).  $\diamond$

*Remark 14.8.* Note, we need to take the topological closure as the free vector space only considers *finite* linear combinations, but our Hilbert spaces could be infinite dimensional.

*Proof.* (that we have a sesqui-linear inner product).

(i) Conjugate symmetry.

$$\begin{aligned} \overline{\langle [(\psi_1, \varphi_1)] | [(\psi_2, \varphi_2)] \rangle_{\mathcal{H}_1 \otimes \mathcal{H}_2}} &= \overline{\langle \psi_1 | \psi_2 \rangle_{\mathcal{H}_1} \cdot \langle \varphi_1 | \varphi_2 \rangle_{\mathcal{H}_2}} \\ &= \overline{\langle \psi_2 | \psi_1 \rangle_{\mathcal{H}_1} \cdot \langle \varphi_2 | \varphi_1 \rangle_{\mathcal{H}_2}} \\ &=: \langle [(\psi_2, \varphi_2)] | [(\psi_1, \varphi_1)] \rangle_{\mathcal{H}_1 \otimes \mathcal{H}_2} \end{aligned}$$

(ii) Linearity in second argument. The extension by linearity means

$$\begin{aligned} \left\langle [(\psi_1, \varphi_1)] \left| \sum_i z_i [(\psi_i, \varphi_i)] \right. \right\rangle_{\mathcal{H}_1 \otimes \mathcal{H}_2} &:= \sum_i z_i \langle \psi_1 | \psi_i \rangle_{\mathcal{H}_1} \cdot \langle \varphi_1 | \varphi_i \rangle_{\mathcal{H}_2} \\ &= \sum_i z_i \langle [(\psi_1, \varphi_1)] | [(\psi_i, \varphi_i)] \rangle_{\mathcal{H}_1 \otimes \mathcal{H}_2}, \end{aligned}$$

for  $z_i \in \mathbb{C}$ .

(iii) Positive-definiteness. As  $\langle \cdot | \cdot \rangle_{\mathcal{H}_1}, \langle \cdot | \cdot \rangle_{\mathcal{H}_2} \geq 0$  it follows that<sup>28</sup>  $\langle - | - \rangle_{\mathcal{H}_1 \otimes \mathcal{H}_2} \geq 0$ . Then from

$$(0_{\mathcal{H}_1}, \varphi) = (0 \cdot \psi, \varphi) \sim 0(\psi, \varphi) \sim (\psi, 0 \cdot \varphi) = (\psi, 0_{\mathcal{H}_2}),$$

we have

$$[(0_{\mathcal{H}_1}, \varphi)] = [(\psi, 0_{\mathcal{H}_2})] =: 0_{\mathcal{H}_1 \otimes \mathcal{H}_2}.$$

Finally, from

$$\begin{aligned} 0 &= \langle [(\psi, \varphi)] | [(\psi, \varphi)] \rangle_{\mathcal{H}_1 \otimes \mathcal{H}_2} \\ &:= \langle \psi | \psi \rangle_{\mathcal{H}_1} \cdot \langle \varphi | \varphi \rangle_{\mathcal{H}_2}, \end{aligned}$$

which implies either  $\psi = 0_{\mathcal{H}_1}$  and/or  $\varphi = 0_{\mathcal{H}_2}$ , and so  $[\psi, \varphi] = 0_{\mathcal{H}_1 \otimes \mathcal{H}_2}$ .

---

<sup>28</sup>We shall use ‘-’ for empty slots on the composite space.

□

We also need to check that the sesqui-linear inner product is well defined.

*Proof.* The proof follows a similar method to the proof of Proposition 14.6. We shall just show the first two results here in order to save space.

(i) Assume  $(\widetilde{\psi_1}, \widetilde{\varphi_1}) = (\psi_1, \varphi_1)$ .

(a) If  $(\widetilde{\psi_2}, \widetilde{\varphi_2}) = (\psi_2, \varphi_2)$ . The inner product result follows trivially.

(b)  $(\widetilde{\psi_2}, \widetilde{\varphi_2}) = (\psi_2, \varphi_2^3) + (\psi_2, \varphi_2^4)$ , where  $\varphi_2 = \varphi_3 + \varphi_4$ ,

$$\begin{aligned}
\left\langle [(\widetilde{\psi_1}, \widetilde{\varphi_1})] \middle| [(\widetilde{\psi_2}, \widetilde{\varphi_2})] \right\rangle_{12} &= \left\langle [(\psi_1, \varphi_1)] \middle| [(\psi_2, \varphi_2^3) + (\psi_2, \varphi_2^4)] \right\rangle_{12} \\
&= \left\langle [(\psi_1, \varphi_1)] \middle| [(\psi_2, \varphi_2^3)] \right\rangle_{12} + \left\langle [(\psi_1, \varphi_1)] \middle| [(\psi_2, \varphi_2^4)] \right\rangle_{12} \\
&= \left\langle [(\psi_1, \varphi_1)] \middle| [(\psi_2, \varphi_2^3)] \right\rangle_{12} + \left\langle [(\psi_1, \varphi_1)] \middle| [(\psi_2, \varphi_2^4)] \right\rangle_{12} \\
&:= \langle \psi_1 | \psi_2 \rangle_1 \langle \varphi_1 | \varphi_2^3 \rangle_2 + \langle \psi_1 | \psi_2 \rangle_1 \langle \varphi_1 | \varphi_2^4 \rangle_2 \\
&= \langle \psi_1 | \psi_2 \rangle_1 \cdot \left( \langle \varphi_1 | \varphi_2^3 \rangle_2 + \langle \varphi_1 | \varphi_2^4 \rangle_2 \right) \\
&= \langle \psi_1 | \psi_2 \rangle_1 \langle \varphi_1 | \varphi_2^3 + \varphi_2^4 \rangle_2 \\
&= \langle \psi_1 | \psi_2 \rangle_1 \langle \varphi_1 | \varphi_2 \rangle_2 \\
&=: \left\langle [(\psi_1, \varphi_1)] \middle| [(\psi_2, \varphi_2)] \right\rangle_{12}.
\end{aligned}$$

□

We introduce the new notation

$$\psi \boxtimes \varphi := [(\psi, \varphi)].$$

Here we have used a  $\boxtimes$  for the tensor product of two vectors. We have done this in order to highlight the fact that it is *not* the same thing as  $\otimes$ , which is the tensor product between vector spaces. We will, however, end up using  $\otimes$  for *all* tensor products later, as this is the common notation. It is important to remember that they are distinctly different objects, and, if in doubt, we should go back to the definitions to clarify the circumstance.

In this new notation we can rewrite the definition for the sesqui-linear inner product simply as

$$\left\langle \psi \boxtimes \varphi \middle| \widetilde{\psi} \boxtimes \widetilde{\varphi} \right\rangle_{\mathcal{H}_1 \times \mathcal{H}_2} := \left\langle \psi \middle| \widetilde{\psi} \right\rangle_{\mathcal{H}_1} \langle \varphi | \widetilde{\varphi} \rangle_{\mathcal{H}_2},$$

extended by linearity.

*Example 14.9.* This example acts as a further warning that its important that we consider the space  $F(\mathcal{H}_1 \times \mathcal{H}_2)$  and not just  $\mathcal{H}_1 \times \mathcal{H}_2$ .

Let  $\mathcal{H}_1 = \mathcal{H}_2 = \mathbb{C}^2$ . Then we can express the elements at 2x1 matrices, in which case we can consider  $\boxtimes$  to be the outer product. Note then that

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} \boxtimes \begin{pmatrix} 0 \\ 1 \end{pmatrix} - \begin{pmatrix} 0 \\ 1 \end{pmatrix} \boxtimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

is in  $\mathcal{H}_1 \otimes \mathcal{H}_2$ , but it *cannot* be written as  $\psi \boxtimes \varphi$  for some  $\psi \in \mathcal{H}_1$  and  $\varphi \in \mathcal{H}_2$ .



**Theorem 14.10.** Let  $\{e_i\}_{i=1,\dots,\dim(\mathcal{H}_1)}$  and  $\{f_i\}_{i=1,\dots,\dim(\mathcal{H}_2)}$  be a Schauder (ON)-bases for  $\mathcal{H}_1$  and  $\mathcal{H}_2$  respectively. Then we can construct a Schauder (ON)-basis for  $\mathcal{H}_1 \otimes \mathcal{H}_2$  as

$$\{e_i \boxtimes f_j\}_{\substack{i=1,\dots,\dim(\mathcal{H}_1) \\ j=1,\dots,\dim(\mathcal{H}_2)}}$$

**Corollary 14.11.** We can rewrite

$$\mathcal{H}_1 \otimes \mathcal{H}_2 := \left\{ \sum_{i=1}^{\dim(\mathcal{H}_1)} \sum_{j=1}^{\dim(\mathcal{H}_2)} a_{ij} e_i \boxtimes f_j \mid a_{ij} \in \mathbb{C}, \sum_{i,j} |a_{ij}|^2 < \infty \right\},$$

from which it also follows that

$$\dim(\mathcal{H}_1 \otimes \mathcal{H}_2) = \dim(\mathcal{H}_1) \cdot \dim(\mathcal{H}_2).$$

## 14.2 Practical Rules for Tensor Products of Vectors

This short section just highlights a couple rules obeyed by  $\boxtimes$ .

(i) Let  $\psi_1, \psi_2 \in \mathcal{H}_1$ ,  $\varphi_1, \varphi_2 \in \mathcal{H}_2$  and  $\alpha, \beta \in \mathbb{C}$ . Then the following holds

$$(\psi_1 + \alpha\psi_2) \boxtimes (\varphi_1 + \beta\varphi_2) = \psi_1 \boxtimes \varphi_1 + \alpha\psi_2 \boxtimes \varphi_1 + \beta\psi_1 \boxtimes \varphi_2 + \alpha\beta\psi_2 \boxtimes \varphi_2.$$

(ii) Given that  $\{e_i \boxtimes f_j\}$  is a basis, we have:

$$\forall \Psi \in \mathcal{H}_1 \otimes \mathcal{H}_2 \quad \exists a_{ij} \in \mathbb{C} : \Psi := \sum_{i,j} a_{ij} e_i \boxtimes f_j.$$

*Remark 14.12.* Note, obviously, that the order matters when taking a tensor product. In other words, in general

$$\psi \boxtimes \varphi \neq \varphi \boxtimes \psi.$$

Note, its not even a case of ‘choosing the right  $\psi$  and  $\varphi$ ’, as the LHS is an element of  $\mathcal{H}_1 \otimes \mathcal{H}_2$  whereas the RHS is an element of  $\mathcal{H}_2 \otimes \mathcal{H}_1$ . So, unless the two spaces are the same, they could never be equal.

## 14.3 Tensor Product Between Operators

**Definition.** Let  $A : \mathcal{H}_1 \rightarrow \mathcal{H}_1$  and  $B : \mathcal{H}_2 \rightarrow \mathcal{H}_2$  be linear maps. Then we define their tensor product as

$$\begin{aligned} A \hat{\otimes} B : \mathcal{H}_1 \otimes \mathcal{H}_1 &\rightarrow \mathcal{H}_1 \otimes \mathcal{H}_2 \\ \psi \boxtimes \varphi &\mapsto (A \hat{\otimes} B)(\psi \boxtimes \varphi) := (A\psi) \boxtimes (B\varphi). \end{aligned}$$

◇

**Theorem 14.13.** If  $A : \mathcal{H}_1 \rightarrow \mathcal{H}_1$  and  $B : \mathcal{H}_2 \rightarrow \mathcal{H}_2$  are self adjoint then their tensor product  $A \hat{\otimes} B$  is also self adjoint on  $\mathcal{H}_1 \otimes \mathcal{H}_2$ .

*Proof.* We have  $A = A^*$  and  $B = B^*$ , i.e. that their domains coincide and  $A\psi = A^*\psi$  for all  $\psi \in \mathcal{D}_A$  and similarly for  $B$  and  $B^*$ . Then we have

$$\begin{aligned} A \hat{\otimes} B : \mathcal{D}_A \otimes \mathcal{D}_B &\rightarrow \mathcal{H}_1 \otimes \mathcal{H}_2 \\ \psi \boxtimes \varphi &\mapsto (A\psi) \boxtimes (B\varphi), \end{aligned}$$

and

$$\begin{aligned} A^* \hat{\otimes} B^* : \mathcal{D}_A \otimes \mathcal{D}_B &\rightarrow \mathcal{H}_1 \otimes \mathcal{H}_2 \\ \psi \boxtimes \varphi &\mapsto (A^*\psi) \boxtimes (B^*\varphi) = (A\psi) \boxtimes (B\varphi), \end{aligned}$$

and so the domain coincides and they have the same result for all  $\psi \boxtimes \varphi \in \mathcal{D}_A \otimes \mathcal{D}_B$ . So it is self adjoint.  $\square$

**Theorem 14.14.** *If  $A : \mathcal{H}_1 \rightarrow \mathcal{H}_1$  and  $B : \mathcal{H}_2 \rightarrow \mathcal{H}_2$  are self adjoint then*

- (i)  $\sigma(A \hat{\otimes} B) = \overline{\sigma(A) \cdot \sigma(B)}$ , where the overline is the topological closure and the  $\cdot$  indicates all possible products of elements in the sets.
- (ii)  $\sigma(A \hat{\otimes} \text{id}_{\mathcal{H}_2} + \text{id}_{\mathcal{H}_1} \hat{\otimes} B) = \overline{\sigma(A) + \sigma(B)}$ , where again the overline is the topological closure.

An application of the second of these results finds use when you know how to measure the observable  $A$  on system 1 and  $B$  on system 2, then you can measure them on the composite system.

#### 14.4 Symmetric and Antisymmetric Tensor Products

Recalling Remark 14.12, if we do have  $\mathcal{H}_1 = \mathcal{H}_2$  it is possible to define a symmetric and a antisymmetric tensor product. These definitions are important in quantum mechanics as they allow us to categorise particles according to their so called *exchange statistics*. The symmetric composite system concerns a system of two (and by induction, any number) of *bosons*, whereas the antisymmetric one corresponds to *fermions*. These are both examples of what are known as *indistinguishable particles*, meaning that two fermions of the same type (two electrons, say) cannot be distinguished from each other. A good analogy is to consider two identical looking balls. Imagine being in a room with the two balls on the floor. Someone asks you to leave the room and then calls you back in. They then ask you whether the two balls, still in the same places on the floor, have switched places or not? Of course there is no way for you to know, as they look identical, and you weren't present when they potentially could have switched.

The version in QM is related to whether they live on the same Hilbert space. Recalling Remark 13.28, we see that this means that, not only are they allowed to move within the same physical space, they also have the same angular momentum (or *spin*). For the two particles to be indistinguishable, their composite Hilbert spaces must be the same. For example, if a divider was put between the balls, and you knew the balls could only move along the floor, you would know that they couldn't possibly have changed places — this could correspond to one electron having  $L^2(U, \lambda)$  and the other having  $L^2(V, \lambda)$  where  $U, V \subset \mathbb{R}^3$  with  $U \cap V = \emptyset$ .

**Definition.** Let  $\psi, \varphi \in \mathcal{H}$ , then we define their *symmetric* tensor product as

$$\psi \sqcup \varphi := \frac{1}{2}(\psi \boxtimes \varphi + \varphi \boxtimes \psi),$$

which is an element of the *symmetric composite Hilbert space*, defined as

$$\mathcal{H} \odot \mathcal{H} := \left\{ \sum_{i,j=1}^{\dim(\mathcal{H})} a_{ij} e_i \sqcup e_j \mid a_{ij} \in \mathbb{C}, \sum_{i,j} |a_{ij}|^2 < \infty \right\},$$

where  $\{e_i\}$  is a basis of  $\mathcal{H}$ . ◇

Remark 14.15. Note it follows from the definition that  $a_{ij} = a_{ji}$  for a symmetric composite Hilbert space.

**Definition.** Let  $\psi, \varphi \in \mathcal{H}$ , then we define their *antisymmetric* tensor product as

$$\psi \boxminus \varphi := \frac{1}{2}(\psi \boxtimes \varphi - \varphi \boxtimes \psi),$$

which is an element of the *antisymmetric composite Hilbert space*, defined as

$$\mathcal{H} \oslash \mathcal{H} := \left\{ \sum_{i,j=1}^{\dim(\mathcal{H})} a_{ij} e_i \boxminus e_j \mid a_{ij} \in \mathbb{C}, \sum_{i,j} |a_{ij}|^2 < \infty \right\}$$

◇

Remark 14.16. Note it follows from the definition that  $a_{ij} = -a_{ji}$  for a antisymmetric composite Hilbert space.

Remark 14.17. For  $\psi \in \mathcal{H}$  we have  $\psi \boxminus \psi = 0$ , which is known as the *Pauli exclusion principle for Fermions*.

Remark 14.18. For  $\psi, \varphi \in \mathcal{H}$  where  $\psi$  and  $\varphi$  are linearly independent, then

$$\psi \boxminus \varphi = \frac{1}{2}(\psi \boxtimes \varphi - \varphi \boxtimes \psi) \neq \tilde{\psi} \boxtimes \tilde{\varphi},$$

for some  $\tilde{\psi}, \tilde{\varphi} \in \mathcal{H}$ . Which again emphasises that its important we consider the space of all linear combinations.

**Definition.** Let  $A, B : \mathcal{H} \rightarrow \mathcal{H}$  be linear operators. Then we can define the *symmetric tensor product of linear maps* as

$$\begin{aligned} A \hat{\odot} B : \mathcal{H} \odot \mathcal{H} &\rightarrow \mathcal{H} \odot \mathcal{H} \\ \psi \sqcup \varphi &\mapsto (A \hat{\odot} B)(\psi \sqcup \varphi) := (A\psi) \sqcup (B\varphi). \end{aligned}$$

◇

**Definition.** Let  $A, B : \mathcal{H} \rightarrow \mathcal{H}$  be linear operators. Then we can define the *antisymmetric tensor product of linear maps* as

$$\begin{aligned} A \hat{\oslash} B : \mathcal{H} \oslash \mathcal{H} &\rightarrow \mathcal{H} \oslash \mathcal{H} \\ \psi \boxminus \varphi &\mapsto (A \hat{\oslash} B)(\psi \boxminus \varphi) := (A\psi) \boxminus (B\varphi). \end{aligned}$$

◇

## 14.5 Collapse of Notation

As mentioned before, we shall now change our notation to that of the standard literature. That is

$$\begin{aligned}\otimes, \boxtimes, \widehat{\otimes} &\rightarrow \otimes, \\ \odot, \square, \widehat{\odot} &\rightarrow \odot, \\ \oslash, \boxminus, \widehat{\oslash} &\rightarrow \wedge.\end{aligned}$$

## 14.6 Entanglement

As has been stressed many times, recall

$$\{\psi \otimes \varphi \mid \psi \in \mathcal{H}_1, \varphi \in \mathcal{H}_2\} \subsetneq \mathcal{H}_1 \otimes \mathcal{H}_2.$$

**Definition.** We call an element  $\Psi \in \mathcal{H}_1 \otimes \mathcal{H}_2$  *simple* if there exists a  $\psi \in \mathcal{H}_1$  and a  $\varphi \in \mathcal{H}_2$  such that

$$\Psi = \psi \otimes \varphi.$$

If it is not of this form (i.e. you need linear combinations) then it is called *non-simple*.  $\diamond$

Recall: A state  $\rho : \mathcal{H} \rightarrow \mathcal{H}$  is called *pure* if

$$\exists \psi \in \mathcal{H} : \forall \alpha \in \mathcal{H} : \rho_\psi(\alpha) = \frac{\langle \psi | \alpha \rangle}{\langle \psi | \psi \rangle} \psi,$$

or, equivalently, we can think of

$$\rho_\psi(\cdot) := \frac{\langle \psi | \cdot \rangle}{\langle \psi | \psi \rangle} \psi.$$

**Definition.** Let  $\Psi \in \mathcal{H}_1 \otimes \mathcal{H}_2$ . Then a pure state  $\rho_\Psi$  on the composite system is called *non-entangled* if there exists  $\rho_\psi$  and  $\rho_\varphi$  for  $\psi \in \mathcal{H}_1$  and  $\varphi \in \mathcal{H}_2$  such that<sup>29</sup>

$$\rho_\Psi = \rho_\psi \otimes \rho_\varphi.$$

Otherwise, the state is called *entangled*.  $\diamond$

**Lemma 14.19.** A state  $\rho_\Psi$  is non-entangled if and only if  $\Psi$  is simple.

*Proof.* Assume  $\Psi$  is simple. Then

$$\begin{aligned}\rho_\Psi(\cdot) &= \frac{\langle \Psi | \cdot \rangle_{12}}{\langle \Psi | \Psi \rangle_{12}} \Psi \\ &= \frac{\langle \psi \otimes \varphi | \cdot \rangle_{12}}{\langle \psi \otimes \varphi | \psi \otimes \varphi \rangle_{12}} \psi \otimes \varphi \\ &= \frac{\langle \psi | \cdot \rangle_1 \langle \varphi | \cdot \rangle_2}{\langle \psi | \psi \rangle_1 \langle \varphi | \varphi \rangle_2} \psi \otimes \varphi \\ &= \left( \frac{\langle \psi | \cdot \rangle_1}{\langle \psi | \psi \rangle_1} \psi \right) \otimes \left( \frac{\langle \varphi | \cdot \rangle_2}{\langle \varphi | \varphi \rangle_2} \varphi \right) \\ &= (\rho_\psi(\cdot)) \otimes (\rho_\varphi(\cdot)),\end{aligned}$$

<sup>29</sup>Note the tensor product here is that between linear operators.

where in the last two lines the  $\otimes$  is the tensor product between linear maps.

The reverse part of the proof (starting from  $\rho_\Psi$  non-entangled) follows from working backwards through the above.  $\square$

**Lemma 14.20.** *A state  $\rho_\Psi$  is entangled if and only if  $\Psi$  is non-simple.*

*Proof.* This proof is trivial given the previous one, as if  $\Psi$  is non-simple then  $\rho_\Psi$  cannot be non-entangled, and vice versa.  $\square$

## 15 Total Spin of Composite System

The lecture aims to answer the following question: "What is the total angular momentum (or spin) of a bi-partite system if we know the spin of each constituent system?"

More precisely, in the context of quantum mechanics, consider a spin- $j_A$  system with Hilbert space  $\mathcal{H}_A$  and angular momentum operators  $A_1, A_2, A_3$  and a spin- $j_B$  system with Hilbert space  $\mathcal{H}_B$  and angular momentum operators  $B_1, B_2, B_3$ . Then what is the spin of the composite system with Hilbert space  $\mathcal{H}_A \otimes \mathcal{H}_B$  and how do we construct the angular momentum operators for this composite system?

**Proposition 15.1.** *The operators  $A_i \otimes \text{id}_{\mathcal{H}_B}$  for  $i = 1, 2, 3$ , satisfy the spin commutation relations. Similarly for  $\text{id}_{\mathcal{H}_A} \otimes B_i$ .*

*Proof.* We shall use the general expression involving the Levi-Civita symbol. Consider the action on a general element  $\alpha \otimes \beta \in \mathcal{H}_A \otimes \mathcal{H}_B$ ,

$$\begin{aligned} [A_i \otimes \text{id}_{\mathcal{H}_B}, A_j \otimes \text{id}_{\mathcal{H}_B}](\alpha \otimes \beta) &:= (A_i \otimes \text{id}_{\mathcal{H}_B})((A_j \alpha) \otimes \beta) - (A_j \otimes \text{id}_{\mathcal{H}_B})((A_i \alpha) \otimes \beta) \\ &= A_i(A_j \alpha) \otimes \beta - A_j(A_i \alpha) \otimes \beta \\ &= ((A_i A_j - A_j A_i) \alpha) \otimes \beta \\ &= ([A_i, A_j] \alpha) \otimes \beta \\ &= i\epsilon_{ijk}(A_k \alpha) \otimes \beta \\ &= i\epsilon_{ijk}(A_k \otimes \text{id}_{\mathcal{H}_B})(\alpha \otimes \beta), \end{aligned}$$

which because  $\alpha \otimes \beta$  was arbitrary (or equivalently by the linearity of the operators) this holds for *any* element of  $\mathcal{H}_A \otimes \mathcal{H}_B$ .

The method is identical for the  $\text{id}_{\mathcal{H}_A} \otimes B_i$  case.  $\square$

Now before moving on recall (page 137) that we have an ON-eigenbasis<sup>30</sup> for each constituent system. That is if  $A^2$  is the Casimir operator for the spin- $j_A$  system then we have the ON-eigenbasis

$$\{\alpha_{j_A}^{m_A}\}_{m_A=-j_A, \dots, j_A}$$

with

$$A^2 \alpha_{j_A}^{m_A} = j_A(j_A + 1) \alpha_{j_A}^{m_A}.$$

Similarly we have  $B^2$  and  $\{\beta_{j_B}^{m_B}\}$ ,  $m_B = -j_B, \dots, j_B$ .

### 15.1 Total Spin

*Remark 15.2.* From now on we shall simply write  $\mathbb{1}$  instead of  $\text{id}_{\mathcal{H}_A}$  and  $\text{id}_{\mathcal{H}_B}$ , and the placement relative to the tensor product will indicate which is meant.

In everything that follows it is important to note that  $j_A$  and  $j_B$  are fixed. This condition shall come in use later.

---

<sup>30</sup>ON here stands for orthonormal.

**Definition.** We define the self adjoint angular momentum operators  $J_1, J_2, J_3$  on the composite space  $\mathcal{H}_A \otimes \mathcal{H}_B$  as

$$J_i := A_i \otimes \mathbb{1} + \mathbb{1} \otimes B_i,$$

and we call them the *total spin operators*.  $\diamond$

*Proof.* (that they obey the spin commutation relations)

Consider

$$\begin{aligned} [A_i \otimes \mathbb{1}, \mathbb{1} \otimes B_j](\alpha \otimes \beta) &:= (A_i \otimes \mathbb{1})(\alpha \otimes B_j \beta) - (\mathbb{1} \otimes B_j)(A_i \alpha \otimes \beta) \\ &= (A_i \alpha) \otimes (B_j \beta) - (A_i \alpha) \otimes (B_j \beta) \\ &= 0, \end{aligned}$$

which from the fact that the commutator bracket is antisymmetric in its entries, along with Proposition 15.1 gives the result.  $\square$

**Definition.** We define the Casimir operator for the composite system as always,

$$J^2 := \sum_{i=1}^3 J_i \circ J_i.$$

$\diamond$

**Definition.** We define the *total ladder operators* as

$$J_{\pm} := A_{\pm} \otimes \mathbb{1} + \mathbb{1} \otimes B_{\pm}.$$

$\diamond$

We now want to find the common eigenvalues of  $J^2$  and one of the total spin operators,  $J_3$  say. We will show the following results:

$$\begin{aligned} \sigma(J^2) &= \{|j_A - j_B|, \dots, j_A + j_B\} \\ \sigma(J_3) &= \{-(j_A + j_B), \dots, j_A + j_B\}. \end{aligned}$$

Remark 15.3. Note from Theorem 14.14, we can already obtain the second of these two results. That is

$$\begin{aligned} \sigma(J_3) &:= \sigma(A_3 \otimes \mathbb{1} + \mathbb{1} \otimes B_3) \\ &= \overline{\sigma(A_3) + \sigma(B_3)} \\ &= \overline{\{-j_A, \dots, j_A\} + \{-j_B, \dots, j_B\}} \\ &= \{-(j_A + j_B), \dots, j_A + j_B\}. \end{aligned}$$

### 15.2 Eigenbasis For The Composite System in Terms of Simultaneous Eigenvectors of $A^2 \otimes \mathbb{1}$ , $\mathbb{1} \otimes B^2$ , $A_3 \otimes \mathbb{1}$ and $\mathbb{1} \otimes B^3$

We already know that  $\{\alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B}\}$  for  $m_A = -j_A, \dots, j_A$  and  $m_B = -j_B, \dots, j_B$ , are common eigenvectors of all four operators with eigenvalues

$$\begin{aligned} (A^2 \otimes \mathbb{1})(\alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B}) &= j_A(j_A + 1)\alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B} \\ (\mathbb{1} \otimes B^2)(\alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B}) &= j_B(j_B + 1)\alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B} \\ (A_3 \otimes \mathbb{1})(\alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B}) &= m_A \alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B} \\ (\mathbb{1} \otimes B_3)(\alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B}) &= m_B \alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B}. \end{aligned}$$

It also follows from the definition of the composite inner product that it is an ON-eigenbasis. That is,

$$\begin{aligned} \left\langle \alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B} \middle| \alpha_{j'_A}^{m'_A} \otimes \beta_{j'_B}^{m'_B} \right\rangle_{12} &:= \left\langle \alpha_{j_A}^{m_A} \middle| \alpha_{j'_A}^{m'_A} \right\rangle_1 \left\langle \beta_{j_B}^{m_B} \middle| \beta_{j'_B}^{m'_B} \right\rangle_2 \\ &= \delta_{j_A, j'_A} \delta_{m_A, m'_A} \delta_{j_B, j'_B} \delta_{m_B, m'_B}. \end{aligned}$$

So we have an ON-eigenbasis for the eigenvectors of these operators. As we shall see, this basis shall be crucial to finding the eigenvectors of  $J^2$  and  $J_3$ , and so their spectra.

### 15.3 Conversion to Eigenbasis in Terms of $J^2$ , $J_3$ , $A^2 \otimes \mathbb{1}$ , $\mathbb{1} \otimes B^2$

The first thing we note here is that not only is  $J^2$  a Casimir operator of  $J_1, J_2, J_3$  but so are  $A^2 \otimes \mathbb{1}$  and  $\mathbb{1} \otimes B^2$ . This is seen straight from the linearity of the commutator bracket,

$$\begin{aligned} [A^2 \otimes \mathbb{1}, J_i] &:= [A^2 \otimes \mathbb{1}, A_i \otimes \mathbb{1} + \mathbb{1} \otimes B_i] \\ &= [A^2 \otimes \mathbb{1}, A_i \otimes \mathbb{1}] + [A^2 \otimes \mathbb{1}, \mathbb{1} \otimes B_i] \\ &= 0, \end{aligned}$$

as each bracket vanishes. Similarly for  $\mathbb{1} \otimes B^2$ . We also have, using *Corollary 11.11* that

$$[J^2, A^2 \otimes \mathbb{1}] = 0 = [J^2, \mathbb{1} \otimes B^2].$$

We can therefore consider eigenvectors of  $J_3$  that are not only common to  $J^2$  but also to  $A^2 \otimes \mathbb{1}$  and  $\mathbb{1} \otimes B^2$ , and so we have a simultaneous eigenbasis,  $\{\xi_{j, j_A, j_B}^m\}$  which satisfies

$$\begin{aligned} J^2 \xi_{j, j_A, j_B}^m &= j(j+1) \xi_{j, j_A, j_B}^m \\ J_3 \xi_{j, j_A, j_B}^m &= m \xi_{j, j_A, j_B}^m \\ (A^2 \otimes \mathbb{1}) \xi_{j, j_A, j_B}^m &= j_A(j_A + 1) \xi_{j, j_A, j_B}^m \\ (\mathbb{1} \otimes B^2) \xi_{j, j_A, j_B}^m &= j_B(j_B + 1) \xi_{j, j_A, j_B}^m. \end{aligned}$$

Now since we already have the ON-eigenbasis  $\{\alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B}\}$  for  $A^2 \otimes \mathbb{1}$  and  $\mathbb{1} \otimes B^2$  it follows (by the definition of a basis) that this new basis can be expanded as<sup>31</sup>

$$\xi_{j, j_A, j_B}^m = \sum_{m_A = -j_A}^{j_A} \sum_{m_B = -j_B}^{j_B} \left\langle \alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B} \middle| \xi_{j, j_A, j_B}^m \right\rangle \alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B}.$$

<sup>31</sup>We shall drop the subscript on the inner product here to lighten notation, but obviously it is the one for the composite Hilbert space.



**Definition.** We define the *Clebsch-Gordan coefficients* (CGc) as

$$C_{j,j_A,j_B}^{m,m_A,m_B} := \left\langle \alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B} \left| \xi_{j,j_A,j_B}^m \right. \right\rangle,$$

and so we can rewrite the previous expression as


$$\xi_{j,j_A,j_B}^m = \sum_{m_A=-j_A}^{j_A} \sum_{m_B=-j_B}^{j_B} C_{j,j_A,j_B}^{m,m_A,m_B} \alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B}.$$

◇

Remark 15.4. The Clebsch-Gordan coefficients are just complex numbers. Although they might be rather difficult to calculate in practice, the method should now be clear. All we need to do is calculate the CGc and then we instantly have our new eigenbasis, and so we get the spectra for  $J^2$  and  $J_3$ .

As just noted, they are pretty hard to calculate, stemming from the fact that  $\xi_{j,j_A,j_B}^m$  appears both on the LHS and within the inner product, however we can do it indirectly. This forms the remainder of this lecture.

The strategy is as follows: start from some convenient eigenvector  $\xi_{j,j_A,j_B}^j$  and its associated CGCs, then use the ladder operators to obtain the eigenvector  $\xi_{j,j_A,j_B}^{j\pm 1}$  and the resulting CGCs. We will then change the value of  $j$  itself and repeat the process. In this manner we will build up a table of CGCs.



#### 15.4 Value of $m$

Consider the action of  $J_3$  on both bases,

$$\begin{aligned} J_3 \xi_{j,j_A,j_B}^m &= m \xi_{j,j_A,j_B}^m \\ J_3 (\alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B}) &= (m_A + m_B) (\alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B}). \end{aligned}$$

Then, from the fact that the CGCs are simply complex numbers and the fact that  $J_3$  is linear, it follows from the expansion equation that we require

$$m = m_A + m_B.$$

In other words, whenever  $m \neq m_A + m_B$ , we require that the CGc vanishes. We can, therefore, place this as a constraint on our summands giving us

$$\xi_{j,j_A,j_B}^m = \sum_{\substack{m_A, m_B \\ m_A + m_B = m}} C_{j,j_A,j_B}^{m, m_A, m_B} (\alpha_{j_A}^{m_A} \otimes \beta_{j_B}^{m_B}),$$

where we have left the ranges of  $m_A/m_B$  out, but they are of course just  $-j_A, \dots, j_A$  and  $-j_B, \dots, j_B$ .

### 15.5 Clebsch-Gordan Coefficients for Maximal $j$

We are now in a position to choose our convenient initial eigenvector. It follows from the ranges of  $m_A$  and  $m_B$  along with the condition  $m = m_A + m_B$  and  $m = -j, \dots, j$  that the maximum value  $j$  can take is  $j_A + j_B$ . It follows then that

$$\xi_{j_A+j_B, j_A, j_B}^{j_A+j_B} = C_{j_A+j_B, j_A, j_B}^{j_A+j_B, j_A, j_B} (\alpha_{j_A}^{j_A} \otimes \beta_{j_B}^{j_B}),$$

and all other CGCs at the level  $C_{j_A+j_B, j_A, j_B}^{j_A+j_B, -, -}$  vanish. Then from the fact that both eigenbases are normalised we know that

$$|C_{j_A+j_B, j_A, j_B}^{j_A+j_B, j_A, j_B}|^2 = 1,$$

and so the two eigenvectors vary only by a complex phase. However, seeing as we are only interested in eigenvalues here, and an *overall* phase plays no effect on the eigenvalue, we are free to set this phase however we like. We choose it such that

$$C_{j_A+j_B, j_A, j_B}^{j_A+j_B, j_A, j_B} = 1.$$

We can now start applying the ladder operators to lower the value of  $m = j_A + j_B$ . Using Proposition 13.24 we have

$$\begin{aligned} J_- \xi_{j_A+j_B, j_A, j_B}^{j_A+j_B} &= \sqrt{(j_A + j_B)(j_A + j_B + 1) - (j_A + j_B)(j_A + j_B - 1)} \xi_{j_A+j_B, j_A, j_B}^{j_A+j_B-1} \\ &= \sqrt{2(j_A + j_B)} \xi_{j_A+j_B, j_A, j_B}^{j_A+j_B-1} \end{aligned}$$

However we equally have

$$\begin{aligned} J_- (\alpha_{j_A}^{j_A} \otimes \beta_{j_B}^{j_B}) &= (A_- \otimes 1 + 1 \otimes B_-) (\alpha_{j_A}^{j_A} \otimes \beta_{j_B}^{j_B}) \\ &= \sqrt{2j_A} (\alpha_{j_A}^{j_A-1} \otimes \beta_{j_B}^{j_B}) + \sqrt{2j_B} (\alpha_{j_A}^{j_A} \otimes \beta_{j_B}^{j_B-1}). \end{aligned}$$

Then, equating these two, we obtain

$$\begin{aligned} C_{j_A+j_B, j_A, j_B}^{j_A+j_B-1, j_A-1, j_B} &= \sqrt{\frac{j_A}{j_A + j_B}} \\ C_{j_A+j_B, j_A, j_B}^{j_A+j_B-1, j_A, j_B-1} &= \sqrt{\frac{j_B}{j_A + j_B}}, \end{aligned}$$

with all other CGCs at this level vanishing.

We can repeat this process to obtain the CGCs at the level  $C_{j_A+j_B, j_A, j_B}^{j_A+j_B-2, -, -}$ , and iterate until we reach  $m = -(j_A + j_B)$ , which is where it must terminate.

## 15.6 Clebsch-Gordan Coefficients For Lower Than Max $j$

We now wish to reduce  $j$  itself to produce the second column of our table. We first need to ask what the next highest allowed  $j$  value is. Recalling that  $j \in \mathbb{N}_0/2$  we might try  $j_A + j_B - 1/2$ , however this is not allowed. The answer to why follows simply from the fact that  $j_A$  and  $j_B$  themselves are fixed, so all we can change is  $m_A$  and  $m_B$ , which must change in integer steps. Combining this with the  $m = m_A + m_B$ , which holds generally, we would not be able to get  $m = -j, \dots, j$ . That is, the next CGCs are of level  $C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, -, -}$ . Then using the fact that there are only two ways to obtain this (either  $m_A \rightarrow m_A - 1$  or  $m_B \rightarrow m_B - 1$ ) we have

$$\xi_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1} = C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A-1, j_B}(\alpha_{j_A}^{j_A-1} \otimes \beta_{j_B}^{j_B}) + C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A, j_B-1}(\alpha_{j_A}^{j_A} \otimes \beta_{j_B}^{j_B-1}).$$

We then use the fact that the eigenvectors in this equation are all orthonormal to obtain

$$1 = |C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A-1, j_B}|^2 + |C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A, j_B-1}|^2,$$

and we also use the fact that the RHS eigenvectors are the same here as with the  $J_-$  case above however the LHS eigenvectors are necessarily orthogonal to give

$$\begin{aligned} \overline{C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A-1, j_B}} \cdot C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A-1, j_B} &= -\overline{C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A, j_B-1}} \cdot C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A, j_B-1} \\ \sqrt{\frac{j_A}{j_A + j_B}} \cdot C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A-1, j_B} &= -\sqrt{\frac{j_B}{j_A + j_B}} \cdot C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A, j_B-1} \\ C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A-1, j_B} &= -\sqrt{\frac{j_B}{j_A}} \cdot C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A, j_B-1}. \end{aligned}$$

Solving simultaneously,

$$\begin{aligned} \left(\frac{j_B}{j_A} + 1\right) |C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A, j_B-1}|^2 &= 1 \\ |C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A, j_B-1}| &= \sqrt{\frac{j_A}{j_A + j_B}} \\ \Rightarrow |C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A-1, j_B}| &= -\sqrt{\frac{j_B}{j_A + j_B}} \end{aligned}$$

Finally we just fix the phases as we want to give

$$C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A, j_B-1} = \sqrt{\frac{j_A}{j_A + j_B}}, \quad C_{j_A+j_B-1, j_A, j_B}^{j_A+j_B-1, j_A-1, j_B} = -\sqrt{\frac{j_B}{j_A + j_B}}.$$

We can then apply the  $J_-$  operator as before to move down this column. We can repeat this process of lowering  $j$  again to obtain the third column, and iterate until we reach  $j = |j_A - j_B|$ , where it must terminate. We see that this is the termination point quickly from  $m = -j, \dots, j$  along with  $m = m_A + m_B$ . On the next page I have included a table (from David J. Griffiths' QM book) for some calculated CGCs. As we can see... they're not pretty things.

## 15.7 Total Spin of Composite System

We conclude, then, that in quantum mechanics when we want to compose a spin- $j_A$  system with a spin- $j_B$  system we do *not* just get a spin- $j_A + j_B$  system, but instead we get the direct sum

$$(\text{spin-}_{j_A+j_B}) \oplus (\text{spin-}_{j_A+j_B-1}) \oplus \dots \oplus (\text{spin-}_{|j_A-j_B|}).$$



## 16 Quantum Harmonic Oscillator

As has been remarked, the world and everything in it are quantum by nature. There is no ‘classical’ ball which we make quantum, there is a quantum ball that we approximate classically. Equally there isn’t a ‘classical’ harmonic oscillator which we use to construct the quantum one. We will thus *not* entertain any type of so called ‘*quantisation*’ idea — that of starting from the classical system and somehow transforming it into the quantum counter part. We shall demonstrate explicitly in the first section why this is not a good idea, but a quick argument explains it.

Imagine you have some general theory. Of course you can obtain any special theory related to it by taking approximations/constraints, however you have no real hope of doing the opposite — you should not expect to be able to obtain the general theory by ‘unapproximating’ the special one. Quantum mechanics is the general theory, with classical mechanics is the special one. It is therefore a ridiculous idea to try and obtain quantum theory this way.

### 16.1 What People Say

Despite the clear message above, people still choose to do such a thing; they take the equations governing a system classically and ‘replace them’ with the quantum versions. To be fair, it is not that the physics community are unaware of the above fact, it is simply that they argue ‘we only do it in special cases where we know no problems arise.’

However, even in said special circumstances, we argue, it is still a terribly misleading and potentially devastating (theoretically speaking!) idea. We shall quickly highlight why this is.

The procedure is as follows. Take the function representing your classical observable  $f(p, q)$ , where  $p$  is the position and  $q$  the momentum, and simply rewrite the function but replacing  $p$  with the quantum mechanical operator  $P$  and  $q$  with  $Q$ .

$$f(p, q) \rightsquigarrow f(P, Q)$$

For example the energy observable for the harmonic oscillator

$$h(p, q) = \frac{1}{2m}p^2 + \frac{m\omega^2}{2}q^2 \rightsquigarrow H := h(P, Q) = \frac{1}{2m}P \circ P + \frac{m\omega^2}{2}Q \circ Q$$

It follows from

$$P, Q : S(\mathbb{R}) \rightarrow S(\mathbb{R}),$$

with

$$(P\psi)(x) := -i\hbar\psi'(x), \quad (Q\psi)(x) := x\psi(x),$$

that

$$H : S(\mathbb{R}) \rightarrow S(\mathbb{R}),$$

with

$$(H\psi)(x) := -\frac{\hbar^2}{2m}\psi''(x) + \frac{m\omega^2}{2}x^2\psi(x).$$

This often appears as

$$H := -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + \frac{m\omega^2}{2} x^2,$$

or for a more general case (i.e. not a harmonic oscillator) as

$$H := -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V(x),$$

where  $V(x)$  is the potential associated to the system.

This all looks very nice, and indeed it is correct, however there is a serious problem here. Classically we could add

$$(pq - qp)g(p, q),$$

for some other observable of the system  $g(p, q)$  without changing anything, as the bracket vanishes. That is

$$f(p, q) = f(p, q) + (pq - qp)g(p, q).$$

However if we then applied the ‘ $\rightsquigarrow$ ’ approach to this we would get

$$f(P, Q) = f(P, Q) + [P, Q]g(P, Q) = f(P, Q) + i\hbar g(P, Q),$$

which is obviously not true for general  $g(P, Q)$ . So it appears that even in these simple cases where ‘there is no danger’, there is a serious theoretical problem. For this reason we shall just not do this at all, and instead simply define what we mean by the energy observable (or *Hamiltonian*) of our system and proceed from there.

## 16.2 The Quantum Harmonic Oscillator

In keeping with Axiom 1, we need an underlying Hilbert space; we use  $\mathcal{H} = L^2(\mathbb{R})$ . We also have (in agreement with Axiom 4) an energy observable, known as the Hamiltonian of the system,

$$H := \frac{1}{2m} P \circ P + \frac{1}{2} m\omega^2 Q \circ Q.$$

However, a note must be made. If  $H$  is to be an observable, it must be self adjoint. But in the above expression we have used the essentially self adjoint  $Q, P : S(\mathbb{R}) \rightarrow S(\mathbb{R})$ . This is not a large worry as we can simply take their unique self adjoint extensions. We still have a problem though. Although  $P \circ P$  and  $Q \circ Q$  (as the self adjoint extensions) will be self adjoint, their *sum* need not be, as the adjoint does not necessarily distribute across the addition.

What we shall do is consider the essentially self adjoint operators throughout, and then at the end we shall present Theorem that allows us to conclude that  $H$  (constructed from the essentially self adjoint operators) is essentially self adjoint, and so a unique self adjoint extension exists.

As above, we shall not employ a different notation for the self adjoint and essentially self adjoint operators, but instead infer which we are dealing with by considering the domains.

### 16.3 The Energy Spectrum

Recall that the spectrum of an operator is given by

$$\sigma(H) = \sigma_p(H) \cup \sigma_c(H).$$

The aim of this lecture is to calculate  $\sigma_p(H)$  and show that  $\sigma_c(H) = \emptyset$ .

**Definition.** Consider the operators  $Q, P : S(\mathbb{R}) \rightarrow S(\mathbb{R})$ . Then define  $a_{\pm} : S(\mathbb{R}) \rightarrow S(\mathbb{R})$  via

$$a_{\pm} := \sqrt{\frac{m\omega}{2\hbar}}Q \mp \frac{i}{\sqrt{2\hbar m\omega}}P.$$

◇

**Corollary 16.1.** *We can re-express the Hamiltonian as*

$$H = \hbar\omega \left( a_+ a_- + \frac{1}{2} \text{id}_{S(\mathbb{R})} \right).$$

*Proof.* The proof follows from direct substitution. Let

$$\alpha := \sqrt{\frac{m\omega}{2\hbar}}, \quad \beta := \frac{1}{\sqrt{2\hbar m\omega}}$$

$$\begin{aligned} H &= \hbar\omega \left( (\alpha Q - i\beta P)(\alpha Q + i\beta P) + \frac{1}{2} \text{id}_{S(\mathbb{R})} \right) \\ &= \hbar\omega \left( \alpha^2 Q \circ Q + \beta^2 P \circ P + i\alpha\beta(QP - PQ) + \frac{1}{2} \text{id}_{S(\mathbb{R})} \right) \\ &= \hbar\omega \left( \alpha^2 Q \circ Q + \beta^2 P \circ P + i\alpha\beta[Q, P] + \frac{1}{2} \text{id}_{S(\mathbb{R})} \right) \\ &= \hbar\omega \left( \frac{m\omega}{2\hbar} Q \circ Q + \frac{1}{2\hbar m\omega} P \circ P + i\frac{1}{2\hbar}(\hbar) \text{id}_{S(\mathbb{R})} + \frac{1}{2} \text{id}_{S(\mathbb{R})} \right) \\ &= \frac{1}{2m} P \circ P + \frac{m\omega^2}{2} Q \circ Q, \end{aligned}$$

where we have used  $[Q, P] = i\hbar \text{id}_{S(\mathbb{R})}$ . □

**Proposition 16.2.** *The following commutation relations hold:*

$$(i) \quad [a_-, a_+] = \text{id}_{S(\mathbb{R})},$$

$$(ii) \quad [H, a_+] = \hbar\omega a_+,$$

$$(iii) \quad [H, a_-] = -\hbar\omega a_-.$$

*Proof.* They all follow from direct substitution, using  $H$  as written in the previous Corollary.



(i)

$$\begin{aligned}
[a_-, a_+] &= [\alpha Q + i\beta P, \alpha Q - i\beta P] \\
&= \alpha^2[Q, Q] - i\alpha\beta[Q, P] + i\beta\alpha[P, Q] + \beta^2[P, P] \\
&= -i2\alpha\beta[Q, P] \\
&= -i2\alpha\beta(i\hbar) \text{id}_{S(\mathbb{R})} \\
&= \frac{2\hbar}{2\hbar} \text{id}_{S(\mathbb{R})} \\
&= \text{id}_{S(\mathbb{R})},
\end{aligned}$$

where we have made use of the linearity of the commutator bracket.

(ii)

$$\begin{aligned}
[H, a_+] &= \left[ \hbar\omega a_+ a_- + \frac{1}{2} \text{id}_{S(\mathbb{R})}, a_+ \right] \\
&= \hbar\omega [a_+ a_-, a_+] + \frac{1}{2} [\text{id}_{S(\mathbb{R})}, a_+] \\
&= \hbar\omega (a_+ [a_-, a_+] + [a_+, a_+] a_-) \\
&= \hbar\omega a_+ \text{id}_{S(\mathbb{R})} \\
&= \hbar\omega a_+
\end{aligned}$$

(iii) This follows exactly analogously to (ii).

□

Remark 16.3. Strictly speaking in the previous proof we should have considered the action of the commutator on an element of  $S(\mathbb{R})$  and showed that the expressions hold for an arbitrary element. Doing it this way will return the same results, however this will not always be true, and so care must be taken in future.

There are four more basic facts that allow us to obtain the spectrum in its entirety. We claim that, for the  $H$ -eigenvalue  $\psi$ , the following hold:

- (i)  $H(a_+\psi) = (E + \hbar\omega)(a_+\psi)$ ,
- (ii)  $\|a_+\psi\| \geq \|\psi\| > 0$ ,
- (iii)  $H(a_-\psi) = (E - \hbar\omega)(a_-\psi)$ ,
- (iv)  $E \geq \frac{1}{2}\hbar\omega$ .

*Proof.* (i) From the previous result we have

$$\begin{aligned}
H(a_+\psi) &= a_+(H\psi) + [H, a_+]\psi \\
&= Ea_+\psi + \hbar\omega a_+\psi \\
&= (E + \hbar\omega)(a_+\psi)
\end{aligned}$$

(ii) Given that  $(a_+)^* = a_-$  and vice versa,<sup>32</sup>

$$\begin{aligned}
\|a_+\psi\|^2 &= \langle a_+\psi | a_+\psi \rangle \\
&= \langle \psi | (a_+)^* a_+ \psi \rangle \\
&= \langle \psi | a_- a_+ \psi \rangle \\
&= \langle \psi | a_+ a_- \psi \rangle + \langle \psi | [a_-, a_+] \psi \rangle \\
&= \langle a_- \psi | a_- \psi \rangle + \langle \psi | \text{id}_{S(\mathbb{R})} \psi \rangle \\
&\geq \langle \psi | \psi \rangle \\
&= \|\psi\|^2,
\end{aligned}$$

where we used the fact that the inner product is non-negative definite in the second to last step (i.e. the first term is non-negative). The result follows from taking the square root and imposing the condition that the norm is non-negative definite.

(iii) This is done exactly analogously to (i).

(iv) Consider

$$\begin{aligned}
E \langle \psi | \psi \rangle &= \langle \psi | E \psi \rangle \\
&= \langle \psi | H \psi \rangle \\
&= \hbar\omega \left( \langle \psi | a_+ a_- \psi \rangle + \frac{1}{2} \langle \psi | \text{id}_{S(\mathbb{R})} \psi \rangle \right) \\
&= \hbar\omega \left( \langle a_- \psi | a_- \psi \rangle + \frac{1}{2} \langle \psi | \text{id}_{S(\mathbb{R})} \psi \rangle \right) \\
&\geq \frac{\hbar\omega}{2} \langle \psi | \psi \rangle.
\end{aligned}$$

Then from the fact that  $\psi$  is an eigenvector (and so cannot be the zero vector), the inner product is non-vanishing and we can divide through by it, giving the result.  $\square$

We can, thus, draw some conclusions. For any  $H$ -eigenvector,  $\psi$ , with eigenvalue  $E$  we have:

1. From (i) and (ii) it follows that  $a_+\psi$  is a eigenvector, as (i) tells us it obeys the eigenvalue equation and (ii) tells us its not the zero vector. Thus we know that the sequence

$$\{(a_+)^n \psi\}_{n \in \mathbb{N}_0}$$

where the power indicates  $n$ -th order composition of operators, is a sequence of eigenvectors with corresponding eigenvalues

$$\{E + n\hbar\omega\}_{n \in \mathbb{N}_0}.$$

---

<sup>32</sup>To show this you need to consider the definition of the adjoint and work from there, as you don't know that it will distribute across the addition in the definitions.

2. (iii) and (iv) tell us that the sequence of eigenvectors

$$\{(a_-)^n \psi\}_{n \in \mathbb{N}_0}$$

*must* terminate for some  $n = N \in \mathbb{N}$ . That is, we can not continue to keep lowering the eigenvalue  $E$  forever, as (iv) says it bounded from below. Note this tells us that  $a_- \psi$  is *not* strictly a  $H$ -eigenvalue (just as  $J_\pm$  weren't for  $\Omega$  and  $J_3$ ).

In other words there is a non-vanishing  $\psi_0 \in S(\mathbb{R})$  defined as

$$\psi_0 := (a_-)^N \psi$$

such that  $a_- \psi_0 = 0_{S(\mathbb{R})}$ . It follows, then, from the definition of the Hamiltonian that

$$\begin{aligned} H\psi_0 &= \hbar\omega a_+ a_- \psi_0 + \frac{\hbar\omega}{2} \psi_0 \\ &= \frac{\hbar\omega}{2} \psi_0, \end{aligned}$$

and so it has the lowest possible eigenvalue, by (iv).

3. The entire sequence (as defined above) of eigenvalues is

$$\left\{ \hbar\omega \left( n + \frac{1}{2} \right) \right\}_{n \in \mathbb{N}_0}.$$

Equivalently, we say the  $n$ -th eigenvector

$$\psi_n := (a_+)^n \psi_0$$

has the corresponding eigenvalue

$$E_n := \hbar\omega \left( n + \frac{1}{2} \right)$$

4. Considering again  $a_- \psi_0 = 0_{S(\mathbb{R})}$  along with the definition of  $a_-$  we have

$$\left( \sqrt{\frac{m\omega}{2\hbar}} x + i \frac{1}{\sqrt{2\hbar m\omega}} (-i\hbar) \frac{d}{dx} \right) \psi_0(x) = 0,$$

which is just a ODE. We can solve this using separation of variables; rearranging, we have

$$\psi_0'(x) = -\frac{m\omega}{\hbar} x \psi_0(x),$$

which using standard separation of variables technique gives

$$\begin{aligned} \ln |\psi_0(x)| &= -\frac{m\omega}{2\hbar} x^2 + C \\ \psi_0(x) &= \pm e^C e^{-\frac{m\omega}{2\hbar} x^2} \\ \psi_0(x) &= A e^{-\frac{m\omega}{2\hbar} x^2}, \end{aligned}$$

for complex constants  $C$  and  $A := \pm e^C$ .

Imposing a normalisation condition, we can then write the  $n$ -th eigenvector in terms of the  $n$ -th *Hermit polynomial*,  $H_n$ , as

$$\psi_n \propto H_n \left( \sqrt{\frac{m\omega}{\hbar}} x \right) e^{-\frac{m\omega}{2\hbar} x^2}.$$

**Corollary 16.4.** *From 4. we note that (up to the usual ambiguity of a complex multiple) there is only one eigenvector to each eigenvalue. That is we have the 1-dimensional eigenspace*

$$\text{Eig}_H(E_n) = \text{span}_{\mathbb{C}}(\psi_n),$$

*which tells us not only that  $\psi_0$  exists in the first place, but that it is unique.*

Remark 16.5. At the end of the last corollary we said that we confirmed the existence of  $\psi_0$  in the first place. This might seem like a strange comment given the whole calculation, however it is actually rather important. To illustrate why Dr. Schuller mentions a doctoral proposal he once saw in which the student had derived some truly impressive formulae, only to have someone point out that towards the start of his calculation he had 0, and so everything that followed could have just been a repercussion of that (i.e.  $0 \cdot n = 0$  for any  $n$  in your space). It is therefore to check that the things you are using actually exist, in this case  $\psi_0$  doesn't vanish and so is an eigenvector.

The Hermit polynomial expression is equally an important result as it tells us that  $\psi_n \in S(\mathbb{R})$  (as all polynomials are in  $S(\mathbb{R})$ ), which it needs to be if we are to act on it with our operators. Moreover, one can show that the set

$$\{\psi_n \mid n \in \mathbb{N}_0\}$$

is an ON-eigenbasis for  $L^2(\mathbb{R})$ , which leads us to the theorem promised at the start of the lecture.

**Theorem 16.6.** *If a symmetric operator has as its eigenvectors an ON-basis, the operator is guaranteed to be essentially self adjoint.*

This theorem tells us that  $H$  is essentially self adjoint, and the fact that we have an ON-eigenbasis for  $L^2(\mathbb{R})$  tells us that the continuous spectrum is empty.

## 17 Measurements of Observables

So far we have discussed the spectrum of an observable, which tells us all the *possible* measurement outcomes, but tells us nothing about the actual act of taking a measurement itself. This comes through axioms 3 and 5. In order to illustrate these two axioms we will repeatedly use the quantum harmonic oscillator as an example, but it is important to note the methods are not specific to this case. Any restrictions required for the methods to hold will be clearly stated.

This lecture can be read in two ways. One could read sections 4 and 5 first (on how you prepare a given state) and then return to read sections 1-3 (on how you take measurements of this state); or one could simply read it as presented (i.e. 1-5). Both reading orders have their advantages, but we present it here in the order taught by Dr. Schuller. Also in correspondence with the lecture given, we shall also translate some of the notation into the commonly used bra-ket notation (see lecture 4), even though we do not use it in this course. All these expressions shall appear in blue.

### 17.1 Spectral Decomposition of $H$ For The Quantum Harmonic Oscillator

Recall: We found an ON-basis of  $H$ -eigenvalues which we labelled  $\psi_n$  obeying

$$H\psi_n = E_n\psi_n,$$

with

$$E_n = \hbar\omega\left(n + \frac{1}{2}\right).$$

More precisely we derived

$$\psi_0(x) = \sqrt{\frac{m\omega}{2\hbar}} \exp\left(-\frac{m\omega}{2\hbar}x^2\right),$$

and

$$\psi_n(x) := A_n(a_+)^n\psi_0(x) \propto H_n\left(\sqrt{\frac{m\omega}{\hbar}}x\right) \exp\left(-\frac{m\omega}{2\hbar}x^2\right).$$

The only thing we will actually use in this lecture is the fact that the  $\{\psi_n\}$  is an ON-basis,

$$\langle\psi_n|\psi_m\rangle = \delta_{nm},$$

and the fact that the spectrum is given by

$$\sigma(H) = \left\{ \hbar\omega\left(n + \frac{1}{2}\right) \mid n \in \mathbb{N}_0 \right\}.$$

The key to understanding measurement theory in quantum mechanics is that you know the spectral decomposition of the observable(s) you want to measure. In order to obtain the spectral decomposition of  $H$  we consider the projectors

$$P_n(\cdot) := \langle\psi_n|\cdot\rangle \psi_n = |\psi_n\rangle \langle\psi_n|.$$

Note that this operator is bounded as

$$\sup_{\varphi \in \mathcal{H}} \frac{\|P_n(\varphi)\|_{\mathcal{H}}}{\|\varphi\|_{\mathcal{H}}^2} = \sup_{\varphi \in \mathcal{H}} \frac{|\langle \psi_n | \varphi \rangle|^2 \|\psi_n\|^2}{\|\varphi\|^2} = \sup_{\varphi \in \mathcal{H}} \frac{|\langle \psi_n | \varphi \rangle|^2}{\|\varphi\|^2} < \infty.$$

We can, therefore, employ the operator norm on  $\mathcal{L}(\mathcal{H})$  to decide convergence of the following sum with the result

$$\sum_{n=0}^{\infty} P_n = \text{id}_{\mathcal{H}}.$$

**Definition.** For every Borel set  $\Omega \in \mathbb{R}$ , define

$$P_H(\Omega) := \sum_{\substack{n \\ E_n \in \Omega}} P_n,$$

i.e. the sum over the projectors such that the energy eigenvalue corresponding to the state<sup>33</sup> corresponding to  $\psi_n$  is within your Borel set.  $\diamond$

Remark 17.1. From now on we shall drop the  $E_n \in \Omega$  on the sum, to lighten the notation, but it is important to remember that it belongs there whenever we use  $P_H$ . It will prove highly instrumental to the results that follow.

Example 17.2. Let  $\Omega = \{E_m\}$ , i.e. just the set containing the single eigenvalue  $E_m$ . Clearly then

$$P_H(\Omega) = P_m.$$

Example 17.3. Let  $\Omega = \{E_m, E_k\}$ . Then we have

$$P_H(\Omega) = P_m + P_k.$$

**Proposition 17.4.** *The map  $P_H : \sigma(\mathbb{R}) \rightarrow \mathcal{L}(\mathcal{H})$  is a projection valued measure, and in fact corresponds to the projection valued measure that appears in the spectral theorem for  $H$ . That is*

$$\begin{aligned} H &= \int_{\mathbb{R}} \lambda P_H(d\lambda) \\ &= \sum_{n=0}^{\infty} E_n \cdot P_n \\ &= \sum_{n=0}^{\infty} \hbar\omega \left( n + \frac{1}{2} \right) |\psi_n\rangle \langle \psi_n|. \end{aligned}$$

Remark 17.5. The above proposition makes sense. The Hamiltonian (the energy operator) is given by the energy eigenvalues multiplied by a projector that projects the state into a state whose energy eigenvalue was the prefactor. This is clearly just the eigenvector equation.

---

<sup>33</sup>Again recall  $\psi_n$  are not the states themselves,  $\rho_{\psi_n}$  are

Remark 17.6. Note there was nothing specifically special about the fact that we were considering the Hamiltonian above. Indeed the same method holds for *any* observable you wish to measure. First find an ON-basis of eigenvectors for your operator,  $A$  say, and then define the PVM associated to the observable

$$P_A : \sigma(\mathbb{R}) \rightarrow \mathcal{L}(\mathcal{H})$$


in the same way and then plug it into the spectral theorem.

## 17.2 Probability to Measure a Certain Value

As stated in the opening of this lecture, the method is the same for all observables and their measurements, but we shall use the energy of the quantum harmonic oscillator as our working example.

Recall, the spectrum of  $H$  is the set of principally possible measurement outcome results. We want to show this pictorially, as it will help with understanding what's to follow.

As we measuring anything in the real world, we need some kind of measuring device. You feed in the thing you want to take a measurement of and the meter on the device tells you the measurement value. The one slightly different, but highly important, difference to note with quantum mechanics is that the device potentially alters the thing you were measuring. We shall draw this as follows.



The  $H$  tells us that it is the device associated to the observable  $H$ , the scale markings tell us the spectrum<sup>34</sup>, the arrow tells us the *actual* measurement made,  $\rho_b$  is the state before the measurement and  $\rho_a$  is the state after the measurement.

Remark 17.7. Note that the pointer here will not move continuously between the notches; it moves between the values by jumping between them. In other words, it can no point at in between two notches, as this would not be part of the spectrum.

Recall: a *state* of a quantum system is a self adjoint operator that is:


<sup>34</sup>Here we are considering the spectrum of the harmonic oscillator, and so the notches are evenly spaced. Clearly this will not always be true. In general we have notches of varying separation as well as ‘blocks’ for the continuous parts of the spectrum.

- (i) Trace-class:  $\text{Tr}(\rho) = \sum_{e_n} \langle e_n | \rho(e_n) \rangle < \infty$ , where  $\{e_n\}$  is *any* ON-basis.
- (ii) Unit trace:  $\text{Tr}(\rho) = 1$ .
- (iii) Positive:  $\forall \varphi \in \mathcal{H}, \langle \varphi | \rho(\varphi) \rangle \geq 0$ .<sup>35</sup>

Also recall: Axiom 3 asserts that the probability to obtain a measurement outcome (a ‘pointer position’) within a Borel set  $\Omega$  when measuring an observable  $H$  on a system in state  $\rho$  is given by

$$\text{Tr}(P_H(\Omega) \circ \rho),$$

where  $P_H$  is the unique PVM from the spectral decomposition of  $H$ . In terms of our picture it asks the question ‘What is the probability that the pointer points within the range  $\Omega$  on the scale?’



*Remark 17.8.* We wish to emphasise this point again here. The spectrum of an observable only tells you the *possible* measurements and the results of last lecture give you information on the *probability* of each possible measurement. This is where the probabilistic nature enters into quantum mechanics. When a measurement is made, the result is concrete. You get exactly that result. This in turn effects the state of your system, giving a (potentially) new state. This is where the indeterminate nature of quantum mechanics enters.

That is, prior to the measurement you can only say with what probability you get one of the possible final states, but once the measurement is made, it is exactly that one, and which final state you get depends on which measurement result you get. This is the *quantum* behaviour of the system.

As we shall see in section 5 there is another form of probability concerned with quantum mechanics, but this probability does not stem from the quantum nature of the system itself. It stems from the ‘ignorance’ of the experimenter/the equipment in order to be able to distinguish which measurement was made. This results in what are known as *mixed states*.

### 17.3 Measurements of Pure States

One can think of pure states as the most precise information one can obtain about a quantum system. Recall that any pure state can be written as

$$\rho_\psi = \frac{|\psi\rangle\langle\psi|}{\langle\psi|\psi\rangle} = \frac{|\psi\rangle\langle\psi|}{\langle\psi|\psi\rangle},$$

<sup>35</sup>Note that this should really be called ‘non-negative’, however this is just how it is named.



for *some*  $\psi \in \mathcal{H}$ .

*Remark 17.9.* Again we emphasise that people often refer to  $\psi$  as being the pure state itself. This might still seem forgivable, but as mentioned previously this is in fact uncountably infinitely incorrect, as we have a complex scaling ambiguity: for any  $\lambda \in \mathbb{C} \setminus \{0\}$ ,

$$\rho_{\lambda\psi} = \rho_\psi.$$

One might then say ‘Ok, just take the normalised  $\psi$  elements,’ but again this is still incorrect as multiplying by  $e^{i\alpha}$  for  $\alpha \in \mathbb{C}$  would still give the same result. One could say, then, ‘a state of the system is given by an element of the Hilbert space, up to arbitrary rescaling.’

Now let's employ  $\text{Tr}(P_H(\Omega) \circ \rho_\varphi)$  to calculate the probability to measure a certain energy of the harmonic oscillator for the state  $\rho_\varphi$ . As we are dealing with the harmonic oscillator, which has a purely discrete spectrum, we can simply make our Borel sets such that they contain only one measurement (one notch on the scale). We have then, for some ON-basis  $\{e_n\}$

$$\text{Tr}(P_H(\{E_k\}) \circ \rho_\varphi) = \sum_n \langle e_n | (P_H(\{E_k\}) \circ \rho_\varphi) e_n \rangle.$$

Now seeing as  $e_n$  need only be *some* ON-basis, we are free to use our ON-eigenbasis  $\{\psi_n\}$ , giving<sup>36</sup>

$$\begin{aligned} \text{Tr}(P_H(\{E_k\}) \circ \rho_\varphi) &= \sum_n \langle \psi_n | (P_H(\{E_k\}) \circ \rho_\varphi) \psi_n \rangle \\ &= \sum_n \left\langle \psi_n \left| \sum_m \langle \psi_m | \rho_\varphi(\psi_n) \rangle \psi_m \right. \right\rangle \\ &= \sum_n \langle \psi_n | \langle \psi_k | \rho_\varphi(\psi_n) \rangle \psi_k \rangle \\ &= \sum_n \left\langle \psi_n \left| \left\langle \psi_k \left| \frac{\langle \varphi | \psi_n \rangle}{\langle \varphi | \varphi \rangle} \varphi \right. \right\rangle \psi_k \right\rangle \\ &= \sum_n \frac{\langle \varphi | \psi_n \rangle}{\langle \varphi | \varphi \rangle} \langle \psi_k | \varphi \rangle \langle \psi_n | \psi_k \rangle \\ &= \frac{\langle \varphi | \psi_n \rangle}{\langle \varphi | \varphi \rangle} \langle \psi_n | \varphi \rangle \\ &= \frac{|\langle \varphi | \psi_n \rangle|^2}{\|\varphi\|^2} \\ &= \frac{\|P_k \varphi\|^2}{\|\varphi\|^2}, \end{aligned}$$

where we have used the fact that  $\|\psi_n\| = 1$  to get to the last line.

We now note that although we do not require  $\varphi$  to be an eigenvector of  $H$ , we can always express it as a linear combination of the ON-eigenbasis  $\{\psi_n\}$ . The following two

---

<sup>36</sup>Recall that in the definition of  $P_H$  the sum is taken such that the energy eigenvalue with that index is within the Borel set.

examples shall highlight this point and demonstrate how one could almost instantly determine the probabilities of obtaining a given energy measurement given the expression for  $\varphi$  corresponding to a pure state.

Example 17.10. First imagine that  $\varphi$  is an eigenvector of  $H$ , then we clearly have

$$\varphi = A\psi_\ell$$

for  $A \in \mathbb{C}$  and some *fixed*  $\ell$ . Plugging this into the formula we obtain

$$\begin{aligned} \text{Tr}(P_H(\{E_k\}) \circ \rho_\varphi) &= \frac{\|P_k(A\psi_\ell)\|^2}{\|A\psi_\ell\|^2} \\ &= \frac{\|\langle \psi_k | A\psi_\ell \rangle \psi_k\|^2}{|A|^2 \|\psi_\ell\|^2} \\ &= \frac{|A \langle \psi_k | \psi_\ell \rangle|^2 \|\psi_k\|^2}{|A|^2} \\ &= \delta_{k\ell}, \end{aligned}$$

so the probability of obtaining energy measurement  $E_k$  vanishes unless  $\psi_\ell = \psi_k$ , in which case we are certain to get that measurement.

Remark 17.11. In the above we could call  $\varphi$  an energy-eigenstate. In this case the use of the word ‘state’ is truly forgivable as all the other eigenvectors with the same eigenvalue would produce the same state (as in both cases we have the same scaling ambiguity). However, if we wish to avoid confusing ourselves, we need not say this.

Example 17.12. Now let’s assume  $\varphi$  isn’t a  $H$ -eigenvector, but is a linear combination, say

$$\varphi = A\psi_p + B\psi_q,$$

for  $A, B \in \mathbb{C}$  and  $p \neq q$ . Then we have

$$\begin{aligned} \text{Tr}(P_H(\{E_k\}) \circ \rho_\varphi) &= \frac{\|P_k(A\psi_p + B\psi_q)\|^2}{\|A\psi_p + B\psi_q\|^2} \\ &= \frac{\|\langle \psi_k | A\psi_p + B\psi_q \rangle \psi_k\|^2}{\langle A\psi_p + B\psi_q | A\psi_p + B\psi_q \rangle} \\ &= \frac{\|(A\delta_{kp} + B\delta_{kq})\psi_k\|^2}{|A|^2 + |B|^2} \\ &= \frac{|A\delta_{kp} + B\delta_{kq}|^2}{|A|^2 + |B|^2} \\ &= \frac{|A|^2 \delta_{kp}^2}{|A|^2 + |B|^2} + \frac{|B|^2 \delta_{kq}^2}{|A|^2 + |B|^2}, \end{aligned}$$

where we have used the fact that  $\langle \psi_p | \psi_q \rangle = 0$  in the denominator and then in the last step used the fact that  $\delta_{kp}\delta_{kq} = 0$  as  $p \neq q$ .

So we see that the coefficients  $A$  and  $B$  tell us the measurement probabilities. The extension to linear combinations with more elements follows trivially to give: if

$$\varphi = \sum_i c_i \psi_i = \sum_i c_i |\psi_i\rangle,$$

for  $c_i \in \mathbb{C}$  then the probability to measure energy  $E_k$  is

$$\text{Tr}(P_H(\{E_k\}) \circ \rho_\varphi) = \frac{\sum_i |c_i|^2 \delta_{ki}}{\sum_i |c_i|^2} = \frac{|c_k|^2}{\sum_i |c_i|^2}.$$

## 17.4 Preparation of Pure States

Axiom 5 asserts that upon measurement of  $H$  the state  $\rho_b$  is projected to the state  $\rho_a$  given by<sup>37</sup>


$$\rho_a := \frac{P_H(\Omega)\rho_b P_H(\Omega)}{\text{Tr}(P_H(\Omega)\rho_b P_H(\Omega))},$$

if  $\Omega$  is the Borell set in which the *actual, really observed, really having happened*, measurement (pointer reading) *came* to lie — that is it depends on the measurement result (past tense!).

This fact, however, can be used to our advantage in order to prepare a state of our choosing. We shall consider the following two cases separately:

- (i) The observable  $H$  with a discrete, non-degenerate spectrum,
- (ii) Allowing for degeneracy of the spectrum.

For (i) we can prepare a pure state  $\rho_{\psi_k}$  where  $\psi_k$  is an eigenvalue of  $H$  by the following device



We feed in a general pure state of our system into the  $H$  device, which measures the energy of that pure state. It then sends this measurement output into the filter device. The state post measurement is then fed into the filter device, which is designed to only let something pass through it if the measurement was  $E_k$ . In this way the *only* pure state that can leave is  $\rho_{\psi_k}$ .

Note, although the final output is guaranteed to be  $\rho_{\psi_k}$ , that does not mean that the output from the  $H$  device is always  $\rho_{\psi_k}$  — it could be *any* of the possible output states. It


<sup>37</sup>This is known as ‘wave-function collapse’ in the literature. Dr. Schuller does not like using the wave analogy and so, if anything, he called this ‘collapse of the state’.

is also important that  $H$  is non-degenerate, otherwise the filter would let multiple different states through, all of which gave  $E_k$  as their measurement output.

For (ii) we allow for degeneracy of  $H$ . We overcome this by considering a *maximal set of mutually commuting observables*,  $\{A_1, \dots, A_f\}$ , for which there are common eigenvectors  $\psi_{a_1, \dots, a_f}$  with

$$A_i \psi_{a_1, \dots, a_f} = a_i \psi_{a_1, \dots, a_f}.$$


The maximal set means that these states are uniquely determined using these operators; that is we have a subset of eigenvectors which we differentiate using this maximal set of commuting operators. The device looks like



### 17.5 Mixed States

Mixed states encode the ‘ignorance’ (in the sense of lack of knowledge) on behalf of the experimenter/equipment. For example, the experimenter’s inability to see exactly where the pointer is pointing, and so takes a guess at the reading. This introduces further uncertainty into which final state we obtain. It is important to note, though, that this is *not* an inherently quantum mechanical property.

The typical set up for preparing a mixed state is as follows:



The ‘probability’ generator here is some method of choosing which input (left) to output (right), where there is a probability  $p$  to use the top input ( $\psi_k$ ). For example it could be a

person rolling a dice that says "If I roll a '1' then I shall use the top input, otherwise I'll use the bottom one," in which case  $p = 1/6$ . Normalisation is taken care of by requiring the other possible outcome to have probability  $(1 - p)$ .

Remark 17.13. It is very important to realise the the output for a mixed state is the sum of two *states*; it is not the state made from the sum of two eigenvectors, as was the case with Example 17.12. That is

$$p\rho_{\psi_k} + (1 - p)\rho_{(1-p)\psi_\ell} \neq \rho_{p\psi_k + (1-p)\psi_\ell}.$$

We highlight this point here as it demonstrates one of the misleading aspects of using bracket notation. People often talk about a pure state as one that can be written as a linear superposition of the eigenstates (as with Example 17.12), writing

$$|\Psi\rangle = a|\psi_k\rangle + b|\psi_\ell\rangle,$$

for  $a, b \in \mathbb{C}$  and  $k \neq \ell$ , where the normalisation condition requires  $|a|^2 + |b|^2 = 1$ . But if we were to think of  $|\Psi\rangle$  as the *state* then this would look like a mixed state — it is the superposition of two pure *states*.

In order to differentiate a pure state from a mixed state they introduce the density matrices, which are the  $\rho$ s we've been using, and say that the density matrix of a mixed state is of the form

$$\rho_{\text{mixed}} = \sum_i p_i |\psi_i\rangle \langle \psi_i|,$$

where  $p_i$  is the probability of being in the corresponding state, but then going back to the start of section 17.3, we see this is just the same as what we wrote for a mixed state, without any of the potential confusion.

## 18 The Fourier Operator

This lecture will begin our systematic approach to the study of the so-called *Schödinger operator*:

$$H : \mathcal{D}_H \rightarrow L^2(\mathbb{R}^d),$$

with

$$(H\psi)(x) := -\frac{\hbar^2}{2m}(\Delta\psi)(x) + V(x)\psi(x),$$

where  $\Delta := \partial_1^2 + \dots + \partial_d^2$  is the *Laplacian operator* and  $V(x)$  is the potential. The *Fourier operator* is an indispensable tool in conducting the study.

We will start by expanding on Lemma 12.19. We will then use the fact that the Schwartz space is densely defined on  $L^2(\mathbb{R}^d)$  along with the BLT theorem to provide a prescription for taking the Fourier transform on  $L^2(\mathbb{R}^d)$ .

### 18.1 The Fourier Operator on Schwartz Space

Recalling the definition of the Schwartz space, it is clear that the following facts hold: If  $f \in S(\mathbb{R}^d)$  then

- (i)  $Q^k f \in S(\mathbb{R}^d)$  for all  $k = 1, \dots, d$ , where  $Q^k$  is the  $k$ -th position operator.
- (ii)  $P_k f \in S(\mathbb{R}^d)$  for all  $k = 1, \dots, d$ , where  $P_k$  is the  $k$ -th momentum operator.

We shall use these facts in the following calculations.

**Definition.** The *Fourier operator* on Schwartz space is the linear map  $\mathfrak{F} : S(\mathbb{R}^d) \rightarrow S(\mathbb{R}^d)$  with

$$(\mathfrak{F}f)(x) := \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d y e^{-ixy} f(y),$$

where  $xy := x_1 y_1 + \dots + x_d y_d$ . ◇

*Remark 18.1.* We are using the  $1/(2\pi)^{d/2}$  convention in the definition above. As we shall see, all that is required is a ‘total of’  $1/(2\pi)$  between the Fourier operator and its inverse. This convention is often used as it makes comparing to the inverse easier. Other conventions (for example having  $1/(2\pi)$  appear in the inverse and just have unit coefficient in the above) find use in certain cases (for example if you were only concerned with taking  $\mathfrak{F}$ ).

*Remark 18.2.* We shall also call the action of the Fourier operator on a function  $f \in S(\mathbb{R}^d)$  the Fourier *transform* of the function.

We now wish to make some remarks on notation.

- (i) Particularly in physics, it is often intuitive to think of  $f$  as a function on ‘position space’ (in the sense that its argument is a position space), while thinking of the Fourier transform  $\mathfrak{F}f$  as a function on momentum space. Thus, one often relabels the variables as follows

$$(\mathfrak{F}f)(p) := \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d y e^{-ipx} f(x).$$

While this does have its advantages at times (the famous example being the motivation behind the derivation of Heisenberg's uncertainty relation, a truly vital relation in quantum mechanics), it can also lead to misconceptions. For example, when thought of this way, one may think that you can not take the double Fourier transform  $\mathfrak{F}(\mathfrak{F}f)$ , as the first one gives a momentum space, which the second does not 'act on'. However from the definition given, this is clearly nonsense — of course you can take it twice as  $\mathfrak{F} : S(\mathbb{R}^d) \rightarrow S(\mathbb{R}^d)$ .

We shall, however, stick to this notation, but we should not be fooled by what we can take the Fourier transform of because of it.

- (ii) Recall that  $(Q^k f)(x) = x^k f(x)$  which is just a real number. It is therefore totally meaningless to write something of the form

$$\mathfrak{F}(x^k f(x)).$$

However having to define the operators each time and then taking the Fourier transform of their action on a function could end up quite lengthy, so instead we introduce the following notations

$$\widehat{x^k f(x)} := \mathfrak{F}(Q^k f),$$

and

$$\mathfrak{F}(x \mapsto x^k f(x)),$$

and similarly for other operators.

The former of these two is how one usually sees the Fourier transform written, and it often just written as

$$\hat{g} := \mathfrak{F}g,$$

where  $g$  is the result of the action of the operator on  $f$  (so here  $g := Q^k f$ ).

**Proposition 18.3.** *Let  $f \in S(\mathbb{R}^d)$  and  $\gamma \in \mathbb{N}_0^d$ . Then*

$$(i) \quad \mathfrak{F}((-i)^{|\gamma|} \partial_\gamma f)(p) = p_\gamma \cdot (\mathfrak{F}f)(p).$$

$$(ii) \quad \mathfrak{F}(x \mapsto x^\gamma f)(p) = i^{|\gamma|} [\partial_\gamma (\mathfrak{F}f)](p).$$

*Proof.* We shall prove these both by induction.

- (i) Let  $\gamma = k$ , and so  $|\gamma| = 1$ . Then, using integration by parts, we have

$$\begin{aligned} \mathfrak{F}(-i \partial_k f)(p) &:= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d x e^{-ipx} (-i) (\partial_k f)(x) \\ &= -\frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d x (-ip_k) e^{-ipx} (-i) f(x) + \left[ \frac{-i}{(2\pi)^{d/2}} e^{-ipx} f(x) \right]_{\partial \mathbb{R}^d} \\ &= p_k \cdot \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d x e^{-ipx} f(x) \\ &= p_k \cdot (\mathfrak{F}f)(p), \end{aligned}$$

where we have used the fact that the elements of the Schwartz space are rapidly decaying to remove the boundary term.

Now assume it is true for  $|\gamma| = n$ . Then, if  $\gamma'$  is the next step, from the fact that  $\partial_k f \in S(\mathbb{R}^d)$  we have

$$\begin{aligned}\mathfrak{F}((-i)^{n+1}\partial_{\gamma_1}\partial_{\gamma_2}\dots\partial_{\gamma_{n+1}}f)(p) &= \mathfrak{F}((-i)^n\partial_{\gamma_1}\partial_{\gamma_2}\dots\partial_{\gamma_n}(-i\partial_{\gamma_{n+1}}f))(p) \\ &= p_{\gamma_1}p_{\gamma_2}\dots p_{\gamma_n} \cdot \mathfrak{F}(-i\partial_{\gamma_{n+1}}f)(p) \\ &= p_{\gamma_1}\dots p_{\gamma_n}p_{\gamma_{n+1}} \cdot (\mathfrak{F}f)(p) \\ &=: p_{\gamma'} \cdot (\mathfrak{F}f)(p)\end{aligned}$$

(ii) Again let  $\gamma = k$ , then

$$\begin{aligned}\mathfrak{F}(x \mapsto x^k f(x))(p) &:= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d x e^{ipx} x^k f(x) \\ &= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d x \frac{\partial}{\partial p^k} (e^{-ipx}) f(x) \\ &= i \frac{\partial}{\partial p^k} \left[ \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d x e^{-ipx} f(x) \right] \\ &=: i \left( \frac{\partial}{\partial p^k} \mathfrak{F}f \right)(p).\end{aligned}$$

Then using the notation  $\partial_k$  gives the result here. In fact, we should have really written  $\partial_k(p \mapsto e^{-ipx})$  on the second line — i.e. you take the derivative before you evaluate at  $x$ . Now assume its true for  $|\gamma| = n$ . Then, if  $\gamma'$  is the next step, we have

$$\begin{aligned}\mathfrak{F}(x \mapsto x^{\gamma_1}\dots x^{\gamma_{n+1}} f(x))(p) &= \mathfrak{F}\left(x \mapsto x^{\gamma_1}\dots x^{\gamma_n} (y \mapsto y^{\gamma_{n+1}} f(y))\right)(p) \\ &= i^n [\partial_{\gamma_1}\dots\partial_{\gamma_n} \mathfrak{F}(y \mapsto y^{\gamma_{n+1}} f(y))](p) \\ &= i^{n+1} [\partial_{\gamma'} (\mathfrak{F}f)](p)\end{aligned}$$

□

**Proposition 18.4.** *Let  $f \in S(\mathbb{R}^d)$ .*

(i) *Let  $a \in \mathbb{R}^d$ , then*

$$\widehat{f(x-a)}(p) = e^{iap} \cdot \widehat{f(x)}(p).$$

(ii) *Let  $\lambda \in \mathbb{C}$ , then*

$$\widehat{f(\lambda x)}(p) = \frac{1}{\lambda^d} \widehat{f(x)}\left(\frac{p}{\lambda}\right).$$

*Proof.* (i) Using the change of variables  $y = x - a$ , we have

$$\begin{aligned}\widehat{f(x-a)}(p) &:= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d x e^{-ipx} f(x-a) \\ &= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d y e^{-ip(y+a)} f(y) \\ &= e^{-ipa} \cdot \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d x e^{-ipx} f(x) \\ &=: e^{-ipa} \widehat{f(x)}(p),\end{aligned}$$



where we relabelled  $y \rightarrow x$  again.

(ii) Using the change of variables  $y = \lambda x$

$$\begin{aligned}\widehat{f(\lambda x)}(p) &= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d x e^{-ipx} f(\lambda x) \\ &= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} \frac{1}{\lambda^d} d^d y e^{-ip \frac{y}{\lambda}} f(y) \\ &= \frac{1}{\lambda^d} \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d x e^{-i \frac{p}{\lambda} x} f(x) \\ &= \frac{1}{\lambda^d} \widehat{f(x)}\left(\frac{p}{\lambda}\right),\end{aligned}$$

where again we relabelled  $y \rightarrow x$ .

□

## 18.2 Inverse of Fourier Operator

*Remark 18.5.* This is often also called ‘the inverse Fourier transform’.

**Lemma 18.6.** *Let  $x \in \mathbb{R}^d$  and  $z \in \mathbb{C}$  with  $\text{Re}(z) > 0$ . Then the following is true.*

$$\widehat{\exp\left(-\frac{z}{2}x^2\right)}(p) = \frac{1}{z^{d/2}} \exp\left(-\frac{1}{2z}p^2\right).$$

*Proof.* We shall prove this for the case  $d = 1$ . Let

$$G_z(x) := \exp\left(-\frac{z}{2}x^2\right).$$

Then we have

$$\begin{aligned}(\partial G_z)(x) &= -zxG_z(x) \\ ip \cdot (\mathfrak{F}G_z)(p) &= -iz[\partial(\mathfrak{F}G_z)](p),\end{aligned}$$

which is an ODE for  $\mathfrak{F}G_z$ . Solving by separation (as done when considering the quantum harmonic oscillator) we arrive at

$$(\mathfrak{F}G_z)(p) = A \exp\left(-\frac{p^2}{2z}\right).$$

Plugging in  $p = 0$  and the definitions for the LHS gives

$$\frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} dx 1 \cdot e^{-\frac{z}{2}x^2} = A.$$

Then employing the fact that the integral above is holomorphic<sup>38</sup> we extend the standard integral result

$$\int_{\mathbb{R}} dx e^{-i\sigma x^2} = \sqrt{\frac{\pi}{\sigma}},$$

---

<sup>38</sup>See ‘Fourier Series, Fourier Transform and Their Application to Mathematical Physics’ by V. Serov Capter 16

for  $\sigma \in \mathbb{R}$  to the case we are considering, giving

$$A = \frac{1}{\sqrt{z}}.$$

□

**Theorem 18.7.** *The Fourier operator  $\mathfrak{F} : S(\mathbb{R}^d) \rightarrow S(\mathbb{R}^d)$  is invertable with inverse*

$$(\mathfrak{F}^{-1}g)(x) = \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d p e^{+ipx} g(p).$$

*Proof.* Need to show that  $[\mathfrak{F}^{-1}(\mathfrak{F}f)](x) = f(x)$ . In order to do so, we shall have to introduce a regulator

$$\lim_{\varepsilon \rightarrow 0} e^{-\frac{\varepsilon}{2}p^2} = 1$$

into the integral. We shall then use the fact that  $(\mathfrak{F}f)(p)$  will be dominant and the fact that we are using Lebesgue integrals to pull out the limit. We shall also use Fubini's theorem to move the order of the integrals.

$$\begin{aligned} [\mathfrak{F}^{-1}(\mathfrak{F}f)](x) &:= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d p e^{ipx} (\mathfrak{F}f)(p) \\ &= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d p \lim_{\varepsilon \rightarrow 0} e^{-\frac{\varepsilon}{2}p^2} e^{ipx} (\mathfrak{F}f)(p) \\ &= \lim_{\varepsilon \rightarrow 0} \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d p e^{-\frac{\varepsilon}{2}p^2} e^{ipx} (\mathfrak{F}f)(p) \\ &:= \lim_{\varepsilon \rightarrow 0} \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d p e^{-\frac{\varepsilon}{2}p^2} e^{ipx} \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d y e^{-ipy} f(y) \\ &= \lim_{\varepsilon \rightarrow 0} \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d y \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d p e^{-\frac{\varepsilon}{2}p^2} e^{ipx} e^{-ipy} f(y) \\ &= \lim_{\varepsilon \rightarrow 0} \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d y \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d p e^{-\frac{\varepsilon}{2}p^2} e^{-ip(y-x)} f(y) \\ &= \lim_{\varepsilon \rightarrow 0} \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d y \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d p e^{-\frac{\varepsilon}{2}p^2} e^{ipx} e^{-ipy} f(y) \\ &= \lim_{\varepsilon \rightarrow 0} \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d z \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d p e^{-\frac{\varepsilon}{2}p^2} e^{-ipz} f(z+x) \\ &= \lim_{\varepsilon \rightarrow 0} \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d z \left[ \exp\left(-\frac{\varepsilon}{2}p^2\right)(z) \right] f(z+x) \\ &= \lim_{\varepsilon \rightarrow 0} \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d z \frac{1}{\varepsilon^{d/2}} \exp\left(-\frac{1}{2\varepsilon}z^2\right) f(z+x) \\ &= \lim_{\varepsilon \rightarrow 0} \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d z' \varepsilon^{d/2} \frac{1}{\varepsilon^{d/2}} \exp\left(-\frac{1}{2\varepsilon}(\varepsilon z')^2\right) f(\sqrt{\varepsilon}z' + x) \\ &= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d z' \exp\left(-\frac{1}{2}(z')^2\right) \lim_{\varepsilon \rightarrow 0} f(\sqrt{\varepsilon}z' + x) \\ &= \frac{1}{(2\pi)^{d/2}} (2\pi)^{d/2} f(x) \\ &= f(x), \end{aligned}$$

where we have used the substitutions  $z = y + x$  and then  $z = \sqrt{\varepsilon}z'$  along with the standard integral result used in the previous lemma. □

### 18.3 Extension of $\mathfrak{F}$ to $L^2(\mathbb{R}^d)$

We already know that  $\mathfrak{F}$  is densely defined on  $L^2(\mathbb{R}^2)$ , so if we can show it is bounded the BLT theorem will tell us there is a unique, bounded extension of  $\mathfrak{F}$  on  $L^2(\mathbb{R}^d)$ .

**Theorem 18.8** (Parseval's Theorem). *Let  $f \in S(\mathbb{R}^d)$ , then*

$$\int_{\mathbb{R}^d} d^d p |\mathfrak{F}f(p)|^2 = \int_{\mathbb{R}^d} d^d x |f(x)|^2.$$

*Proof.* The proof follows by direct calculation.

$$\begin{aligned} \int_{\mathbb{R}^d} d^d p |\mathfrak{F}f(p)|^2 &= \int_{\mathbb{R}^d} d^d p \left| \int_{\mathbb{R}^d} d^d x e^{-ipx} f(x) \right|^2 \\ &= \int_{\mathbb{R}^d} d^d p \int_{\mathbb{R}^d} d^d x \left| e^{-ipx} f(x) \right|^2 \\ &= \int_{\mathbb{R}^d} d^d p \int_{\mathbb{R}^d} d^d x |f(x)|^2 \\ &= \int_{\mathbb{R}^d} d^d x |f(x)|^2, \end{aligned}$$

where we used the fact that the integral is over a real domain. □

It follows from Parseval's theorem, then, that

$$\begin{aligned} \|\mathfrak{F}\| &:= \sup_{f \in S(\mathbb{R}^d)} \frac{\|\mathfrak{F}f\|_{S(\mathbb{R}^d)}^2}{\|f\|_{S(\mathbb{R}^d)}^2} \\ &= \sup_{f \in S(\mathbb{R}^d)} \frac{\sqrt{\int_{\mathbb{R}^d} d^d p |\mathfrak{F}f(p)|^2}}{\sqrt{\int_{\mathbb{R}^d} d^d x |f(x)|^2}} \\ &= \sup_{f \in S(\mathbb{R}^d)} \sqrt{\frac{\int_{\mathbb{R}^d} d^d p |\mathfrak{F}f(p)|^2}{\int_{\mathbb{R}^d} d^d x |f(x)|^2}} \\ &= 1, \end{aligned}$$

so we have a unique extension

$$\mathfrak{F} : L^2(\mathbb{R}^2) \rightarrow L^2(\mathbb{R}^2).$$

In practice if we wanted to take the Fourier transform of a function  $f \in L^2(\mathbb{R}^d) \setminus S(\mathbb{R}^d)$  then we do it via the following prescription: Let  $\{f_n\}_{n \in \mathbb{N}} \subset S(\mathbb{R}^d)$  with  $\lim_{n \rightarrow \infty} f_n = f$ , then

$$\mathfrak{F}f = \mathfrak{F}\left(\lim_{n \rightarrow \infty} f_n\right) = \lim_{n \rightarrow \infty} (\mathfrak{F}f_n),$$

where we have used the fact that  $\mathfrak{F}$  is bounded to remove the limit.

## 18.4 Convolutions

**Definition.** The *convolution* of two functions  $f, g \in L^1(\mathbb{R}^d)$ , written  $f * g$ , is the  $L^1(\mathbb{R}^d)$  function defined pointwise by

$$(f * g)(x) := \int_{\mathbb{R}^d} d^d y f(x - y) g(y).$$

◇

**Lemma 18.9.** *The convolution of two functions is symmetric, i.e.*

$$f * g = g * f.$$

*Proof.* The result comes from simple change of variables along with the commutativity of the complex multiplication,

$$\begin{aligned} (f * g)(x) &:= \int_{\mathbb{R}^d} d^d y f(x - y) g(y) \\ &= (-1)^{2d} \int_{\mathbb{R}^d} d^d z f(z) g(x - z) \\ &= \int_{\mathbb{R}^d} d^d z g(x - z) f(z) \\ &= \int_{\mathbb{R}^d} d^d y g(x - y) f(y) \\ &=: (g * f)(x), \end{aligned}$$

where the  $(-1)^{2d}$  term comes from the fact that  $dy = -dz$  along with the fact that the integral limits swap. Since  $x \in \mathbb{R}^d$  was arbitrary, we have the result.  $\square$

**Lemma 18.10.** *The convolution is associative, i.e.*

$$(f * g) * h = f * (g * h).$$

*Proof.* By direct calculation using Fubini's Theorem, and the previous lemma:

$$\begin{aligned}
((f * g) * h)(x) &= (h * (f * g))(x) \\
&:= \int_{\mathbb{R}^d} d^d y h(x - y)(f * g)(x) \\
&= \int_{\mathbb{R}^d} d^d y h(x - y)(g * f)(x) \\
&:= \int_{\mathbb{R}^d} d^d y h(x - y) \int_{\mathbb{R}^d} d^d z g(z - x)f(z) \\
&= \int_{\mathbb{R}^d} d^d u \int_{\mathbb{R}^d} d^d z h(u)g(z - u - y)f(z) \\
&= \int_{\mathbb{R}^d} d^d z \left( \int_{\mathbb{R}^d} d^d u g(z - y - u)h(u) \right) f(z) \\
&= \int_{\mathbb{R}^d} d^d z (g * h)(z - y)f(z) \\
&= \int_{\mathbb{R}^d} d^d x (g * h)(x - y)f(x) \\
&= ((g * h) * f)(x) \\
&= (f * (g * h))(x),
\end{aligned}$$

which holds for all  $x \in \mathbb{R}^d$ , giving the result.  $\square$

**Lemma 18.11.** *The convolution is distributive across addition, i.e.*

$$f * (g + h) = f * g + f * h,$$

where the addition is defined pointwise.

*Proof.* This follows from the linearity of the Lebesgue integral:

$$\begin{aligned}
(f * (g + h))(x) &:= \int_{\mathbb{R}^d} d^d y f(x - y)(g + h)(x) \\
&= \int_{\mathbb{R}^d} d^d y f(x - y)g(x) + \int_{\mathbb{R}^d} d^d y f(x - y)h(x) \\
&= (f * g)(x) + (f * h)(x),
\end{aligned}$$

which holds for all  $x \in \mathbb{R}^d$ , giving the result.  $\square$

**Theorem 18.12.** *The Fourier transform of the convolution of two functions is proportional to the product of their Fourier transforms, explicitly*

$$\mathfrak{F}(f * g) = (2\pi)^{d/2} \mathfrak{F}(f) \cdot \mathfrak{F}(g).$$

*Proof.* By direct calculation,

$$\begin{aligned}
\mathfrak{F}(f * g)(p) &:= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d x e^{-ipx} (f * g)(x) \\
&:= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d x \left( e^{-ipx} \int_{\mathbb{R}^d} d^d y f(x-y) g(y) \right) \\
&= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d z \left( e^{-ip(z+y)} \int_{\mathbb{R}^d} d^d y f(z) g(y) \right) \\
&= \frac{1}{(2\pi)^{d/2}} \int_{\mathbb{R}^d} d^d z \int_{\mathbb{R}^d} d^d y e^{-ipz} f(z) e^{-ipy} g(y) \\
&= (2\pi)^{d/2} \left( \frac{1}{(2\pi)^{2/pi}} \int_{\mathbb{R}^d} d^d z e^{ipz} f(z) \right) \cdot \left( \frac{1}{(2\pi)^{2/pi}} \int_{\mathbb{R}^d} d^d y e^{ipy} g(y) \right) \\
&= (2\pi)^{2/d} ((\mathfrak{F}f)(p)) \cdot ((\mathfrak{F}g)(p)),
\end{aligned}$$

where we have used the fact that we can consider the convolution integration variable (the  $y$ ) as a constant when relabelling the Fourier transform variable, and used the fact that the Fourier transform of a function is finite to make the integral of an integral into a product of integrals. Finally since this is true for all  $p \in \mathbb{R}^d$  the result follows.  $\square$

## 19 The Schrodinger Operator

The Schrödinger operator for a vanishing potential is given by

$$H_{\text{free}} := -\frac{\hbar^2}{2m} \Delta$$

and it corresponds to the energy observable for a free particle<sup>39</sup> of mass  $m$ .

This lecture aims to derive the spectrum of  $H_{\text{free}}$  and use it to study the time evolution of pure states. We shall consider  $d = 3$  throughout this lecture, and shall make use of the results of last lecture heavily. We shall also use units such that  $m = \frac{1}{2}\hbar^2$  in order to lighten notation; i.e.  $H_{\text{free}} = -\Delta$ .

### 19.1 Domain of Self Adjointness of $H_{\text{free}}$

If we want to talk about  $H_{\text{free}}$  being an observable, we need to show that it is self adjoint on some domain. From (i) in Proposition 18.3 it follows that

$$\mathfrak{F}(-\Delta\psi)(p) = |p|^2(\mathfrak{F}\psi)(p) =: (P^2\hat{\psi})(p),$$

where  $|p|^2 := p_1^2 + p_2^2 + p_3^2$ . In other words,

$$\mathfrak{F}(-\Delta\psi) =: P^2\hat{\psi}.$$

*Remark 19.1.* The physicist says "in momentum space the Laplacian acts simply by multiplication of the norm of the momentum,  $|p|^2$ ".

We can now rewrite the above by inserting  $\text{id}_{L^2(\mathbb{R}^3)} = \mathfrak{F}^{-1}\mathfrak{F}$  to give

$$\mathfrak{F} \circ H_{\text{free}} \circ \mathfrak{F}^{-1} \circ \mathfrak{F}\psi = P^2 \circ \mathfrak{F}\psi,$$

from which it follows

$$\mathfrak{F}H_{\text{free}}\mathfrak{F}^{-1} = P^2,$$

whose maximal domain is

$$\mathcal{D}_{P^2} = \{\hat{\psi} \in L^2(\mathbb{R}^3) \mid P^2\hat{\psi} \in L^2(\mathbb{R}^3)\}.$$

**Theorem 19.2.** *A maximally defined real multiplication operator is self adjoint on its maximal domain.*

*Proof.* Let

$$\begin{aligned} A : \mathcal{D}_A &\rightarrow \mathcal{H} \\ \psi &\mapsto a\psi \end{aligned}$$

---

<sup>39</sup>Free particle here means what we think of classically as a free particle, it experiences no potential.

where  $a \in \mathbb{R}$  and  $\mathcal{D}$  is the maximal domain of  $A$  (i.e. there are no elements outside this domain such that  $a\psi \in \mathcal{H}$ ). This operator is clearly symmetric as, for  $\psi, \varphi \in \mathcal{D}$

$$\begin{aligned}\langle \psi | A \varphi \rangle &= \langle \psi | a \varphi \rangle \\ &= a \langle \psi | \varphi \rangle \\ &= \langle \bar{a} \psi | \varphi \rangle \\ &= \langle a \psi | \varphi \rangle \\ &= \langle A \psi | \varphi \rangle.\end{aligned}$$

We have therefore  $A \subseteq A^*$ , which means  $\mathcal{D}_A \subseteq \mathcal{D}_{A^*}$  with  $A^*\psi = A\psi$  for all  $\psi \in \mathcal{D}_A$ , but  $\mathcal{D}_A$  is maximal so there is no  $\psi \notin \mathcal{D}_A$  such that  $A^*\psi = a\psi \in \mathcal{H}$  and so  $\mathcal{D}_{A^*} \subseteq \mathcal{D}_A$ . Therefore  $A = A^*$  on this maximal domain.  $\square$

From this theorem, then, we have that  $\mathfrak{F}H_{\text{free}}\mathfrak{F}^{-1}$  is self adjoint on the domain  $\mathcal{D}_{P^2}$ .

## 19.2 Spectrum of $H_{\text{free}}$

One can quickly find the spectrum of  $H_{\text{free}}$  using the resolvent map,

$$R_{P^2}(z) = \frac{1}{|p|^2 - z},$$

from which it clearly follows that the resolvent set is simply

$$\rho(P^2) = \{z \in \mathbb{C} \mid z \neq |p|^2\}.$$

Then using the fact that  $|p|^2 \in \mathbb{R}$  with  $|p|^2 > 0$ <sup>40</sup> we have

$$\rho(P^2) = \mathbb{C} \setminus \mathbb{R}_0^+.$$

Then finally using the definition of the spectrum as the complement of the resolvent set we get

$$\sigma(H_{\text{free}}) = \sigma(P^2) = \mathbb{R}_0^+.$$

*Remark 19.3.* This is not the method used in the lectures, which I<sup>41</sup> find more confusing. As I don't fully understand the latter parts of the proof provided (I think the main idea is introducing a form of the spectral theorem in which the integral is performed over the spectrum of the operator and then use the characteristic function Dr. Schuller introduced) I shall not type it up here to avoid potential confusion to the readers. If you do follow the complete method please feel free to contact me and I can add it and give you credit.

**Proposition 19.4.** *For every self adjoint operator there is always some transformation which transforms the operator into a mere multiplication operator.*

*Remark 19.5.* Once you know the transformation the self adjoint operator of interest, the spectrum always follows by the same method.

<sup>40</sup>We use a strict equality as if  $|p|^2 = 0$  then there is no momentum and so the operator  $P^2$  just maps it to 0.

<sup>41</sup>I being Richie



### 19.3 Time Evolution

Recall that Axiom 4 tells us the evolution of a state is given by<sup>42</sup>

$$\rho_{\psi_{t_2}} = e^{-i(t_2-t_1)H} \rho_{\psi_{t_1}} e^{i(t_2-t_1)H},$$

which for a pure state becomes

$$\frac{\langle \psi_{t_2} | \cdot \rangle}{\langle \psi_{t_2} | \psi_{t_2} \rangle} \psi_{t_2} = e^{-i(t_2-t_1)H} \frac{\langle \psi_{t_1} | \cdot \rangle}{\langle \psi_{t_1} | \psi_{t_1} \rangle} \psi_{t_1} e^{i(t_2-t_1)H}.$$

If we now choose to view the RHS as the following composition

$$\frac{1}{\langle \psi_{t_1} | \psi_{t_2} \rangle} \left( e^{-i(t_2-t_1)H} \langle \psi_{t_1} | \cdot \rangle \right) \circ \left( \psi_{t_1} e^{i(t_2-t_1)H} \right) = \frac{1}{\langle \psi_{t_1} | \psi_{t_2} \rangle} \left( e^{-i(t_2-t_1)H} | \psi_{t_1} \rangle \right) \circ \left( \langle \psi_{t_1} | e^{i(t_2-t_1)H} \right),$$

it follows that we can represent the time evolution of a pure *state* via the evolution of the *Hilbert space element* as

$$\psi_{t_2} = e^{-i(t_2-t_1)H} \psi_{t_1}.$$

*Remark 19.6.* We are assuming here that  $H$  is time-independent. If it wasn't you would simply use an integral in the exponential.

*Remark 19.7.* We should note that in the above time is viewed as a parameter, not a coordinate — i.e. this is *not* a spacetime picture. The elements  $\psi_{t_1}$  and  $\psi_{t_2}$  are simply elements of the Hilbert space, each of which is associated to a different time. This can be compared to saying classically that the position of a particle is an element of  $\mathbb{R}^3$ , and at a later time its position is still an element of  $\mathbb{R}^3$ , although potentially a different one.

Now for the free particle we have

$$\mathfrak{F}^{-1} P^2 \mathfrak{F} = H_{\text{free}},$$

which, along with the fact that  $P^2$  is a multiplicative operator, gives us

$$e^{-itH_{\text{free}}} \mathfrak{F}^{-1} = \mathfrak{F}^{-1} e^{-itP^2}.$$

Acting both sides on  $\widehat{\psi} := \mathfrak{F}\psi$  gives

$$e^{-itH_{\text{free}}} \psi = \mathfrak{F}^{-1} \left( p \mapsto e^{-it|p|^2} \cdot \widehat{\psi}(p) \right),$$

which is a convolution of functions. It is then tempting to use

$$f * g = (2\pi)^{d/2} \mathfrak{F}^{-1} ((\mathfrak{F}f) \cdot (\mathfrak{F}g))$$

to give us<sup>43</sup>

$$e^{-itH_{\text{free}}} \psi(x) = \frac{1}{(2\pi)^{3/2}} \left( \mathfrak{F}^{-1} (p \mapsto e^{-it|p|^2}) * \psi \right)(x),$$

<sup>42</sup>We're using units such that  $\hbar = 1$ .

<sup>43</sup>Remember  $d = 3$  here.

and then use Lemma 18.6 to give

$$\mathfrak{F}^{-1}(p \mapsto e^{-it|p|^2})(x) = \frac{1}{(2t)^{3/2}} e^{-\frac{1}{4it}|x|^2},$$

however there is a problem with both of these steps.

Firstly for us to use the convolution theorem we require both functions to be  $L^1(\mathbb{R}^3)$ . For  $\psi$  we can simply take the intersection  $\psi \in L^2(\mathbb{R}^3) \cap L^1(\mathbb{R}^3)$ , however the exponential term is unavoidably not in  $L^1(\mathbb{R}^3)$  (if you take the absolute value you get 1, and then integrating over all of  $\mathbb{R}^3$  gives a divergent result). On top of that, in order to use Lemma 18.6 we require the real part of the coefficient to be strictly positive (in order to avoid the branch cut), but  $\text{Re}(it) = 0$ .

Luckily we can fix both of these problems with the same step, regularisation. We regularise both by introducing a positive, real factor into the exponential and then taking the limit,

$$e^{-it|p|^2} = \lim_{\varepsilon \rightarrow 0} e^{-(it+\varepsilon)|p|^2}.$$

The addition of  $\varepsilon$  stops the integral diverging (because of the minus sign) and we also have  $\text{Re}(it + \varepsilon) = \varepsilon > 0$  and so we can use Lemma 18.6.

So, using the continuity of the product and the inverse Fourier transform we have

$$\begin{aligned} e^{-itH_{\text{free}}}\psi(x) &= \left[ \lim_{\varepsilon \rightarrow 0} \left( x \mapsto \frac{1}{(2\pi)^{3/2}} \frac{1}{(2(it+\varepsilon))^{3/2}} \exp\left(-\frac{|x|^2}{4(it+\varepsilon)}\right) \right) * \psi \right](x) \\ &:= \lim_{\varepsilon \rightarrow 0} \frac{1}{(4\pi(it+\varepsilon))^{3/2}} \int_{\mathbb{R}^3} d^3y \exp\left(-\frac{|x-y|^2}{4(it+\varepsilon)}\right) \psi(y). \end{aligned}$$

Finally using dominated convergence to take the limit inside the integral, we have<sup>44</sup>

$$\psi_{t_2}(x) = \frac{1}{(i4\pi t)^{3/2}} \int_{\mathbb{R}^3} d^3y \exp\left(-\frac{|x-y|^2}{i4t}\right) \psi_{t_1}(y).$$

## 19.4 Physical Interpretation (for the Patient Observer)

After massaging the above result a bit we arrive at the famous ‘*spreading of the wavefunction*’ result.

First start by expanding

$$|x-y|^2 = |x|^2 + |y|^2 - 2xy$$

to give

$$\psi_{t_2}(x) = \frac{\exp\left(-\frac{|x|^2}{i4t}\right)}{(i4\pi t)^{3/2}} \int_{\mathbb{R}^3} d^3y \exp\left(-i\frac{x}{2t}y\right) \exp\left(-\frac{|y|^2}{i4t}\right) \psi_{t_1}(y),$$

which is a Fourier transform with result

$$\psi_{t_2}(x) = \frac{\exp\left(-\frac{|x|^2}{i4t}\right)}{(i2t)^{3/2}} \cdot \overline{\exp\left(-\frac{|y|^2}{i4t}\right) \psi_{t_1}(y)}\left(\frac{x}{2t}\right).$$


---

<sup>44</sup>Note  $t = t_2 - t_1$  here.

We now use the fact that we have a patient observer (i.e. one who watches for a long time) to take the asymptotic behaviour<sup>45</sup> to give

$$\psi_{t_2}(x) \sim \frac{\exp\left(-\frac{|x|^2}{i4t}\right)}{(i2t)^{3/2}} \hat{\psi}_{t_1}\left(\frac{x}{2t}\right),$$

which, if the  $\psi$ s were viewed as ‘waves’ (i.e. plots on a graph) would indicate that the ‘wave spreads out over time’. In other words, simplifying to  $\mathbb{R}$  instead of  $\mathbb{R}^3$  we’d have something along the following diagram.



So the function appears to spread out (keeping the area under it constant) over time.

---

<sup>45</sup>That is take the limit  $t \rightarrow \infty$  at places where it won't cause problems.

## 20 Periodic Potentials

This lecture aims to look at periodic potentials and find the most general information we can about their energy spectrum. In order to do this we will use so-called *Rigged Hilbert Spaces*.<sup>46</sup>


**Definition.** The Hamiltonian for a periodic potential is of the form

$$H = -\frac{\hbar^2}{2m} \Delta + V(x),$$


with

- (i) Periodicity in  $V(x)$ , i.e.  $V(x + a) = V(x)$  for all  $x \in \mathbb{R}^3$  where  $a$  is the periodicity of the system
- (ii)  $V$  is pointwise continuous and bounded.

◇




As we shall see, by making no assumptions apart from the above, we will be able to extract a remarkable generic conclusion about the energy spectrum of particle moving in a generic periodic potential. This is truly a amazing result as the potential can even be discontinuous (countably) infinite times! A huge application of this formalism is in the study of solid state physics, where the periodic potential comes from that generated by a regular lattice of so-called *lattice constant*  $a$ ,



<sup>46</sup>I am currently reading up on these, and will add an additional section to the end of these notes once I have a better idea on them.

As we shall see the general result is that the energy spectrum comes in continuous, open intervals in  $\mathbb{R}$ , known as *bands*.



## 20.1 Basics of Rigged Hilbert Space

As mentioned at the start, we wish to make use of rigged Hilbert spaces<sup>47</sup> in order to find the spectrum of the energy observable. The basic reason behind this is that rigged Hilbert spaces essentially extend what we usually think of as the eigenvalue equation

$$H\psi = E\psi,$$

where  $E$  is a discrete value in  $\mathbb{R}$ , to the case where  $E$  can be continuous. This is known as the *generalised eigenvalue equation*.

We do this because ultimately we know that the spectrum will be continuous intervals, however even if it was purely discrete, or a combination of both, the theory of rigged Hilbert spaces would still account for this.

The basic idea behind rigged Hilbert spaces is to consider elements  $\Psi$  that satisfy the generalised eigenvalue equation, but do not lie in  $L^2(\mathbb{R}^d)$ . It turns out that they lie in the adjoint of a densely defined subspace, which for us is the Schwartz space. In this way we construct our so-called *Gelfand Triple*:

$$S(\mathbb{R}^d) \subset L^2(\mathbb{R}^d) \subset S^*(\mathbb{R}^d).$$

The easiest way to see that we need such a construction here is that, as the Hamiltonian is constructed using derivative operators, its eigenvalues are likely to be of the form

$$\Psi \propto e^{iE},$$

which is clearly not square integrable (it's modulus is 1).

*Remark 20.1.* It is important to note here that a rigged Hilbert space is not some extension of the physics or of quantum mechanics, but indeed it is the most natural mathematical structure required in order to study quantum mechanics. In fact it is the rigged Hilbert space structure which provides the full mathematical foundation in order to understand Dirac's bra-ket notation, and it introduces the well known *Dirac delta function*. This gives the first insight into what a rigged Hilbert space is — it is the equipping (i.e. the 'rigging') of a Hilbert space with a theory of distributions.

**Proposition 20.2.** *Any  $H$ -eigenvector  $\Psi \in S^*(\mathbb{R}^d) \setminus L^2(\mathbb{R}^d)$  has a purely continuous energy spectrum.*

---

<sup>47</sup>Again, coming soon!

## 20.2 Fundamental Solutions and Fundamental Matrix

**Definition.** A set of solutions  $\{\psi_1, \dots, \psi_n\}$  for a system of linear, homogeneous ordinary differential operators is known as a *fundamental set of solutions* if

- (i) They are linearly independent
- (ii) Any other solution to the ODEs can be expanded using the set  $\{\psi_1, \dots, \psi_n\}$ .

In other words, they form a basis for the solution space.  $\diamond$

**Proposition 20.3.** *The cardinality of the fundamental set of solutions of a system of  $n$ -th order ODEs is  $n$ .*

Example 20.4. Let the system of ODEs just be the single equation

$$\ddot{\psi}(x) + \omega^2 \psi(x) = 0,$$

for some non-vanishing  $\omega \in \mathbb{R}$ . The the fundamental set of solutions is

$$\psi_1 = \cos(\omega x) \quad \text{and} \quad \psi_2 = \sin(\omega x).$$

It is easy enough to see that these two solutions do indeed form a basis for the solution space.

**Lemma 20.5.** *Let  $\{\psi_1, \dots, \psi_n\}$  be a set of fundamental solutions for some system of ODEs. Then the set  $\{c_1\psi_1, \dots, c_n\psi_n\}$  for  $c_i \in \mathbb{F}$  (the underlying field) is also a set of fundamental solutions.*

As our Hamiltonian is a second order ODE there are 2 fundamental solutions. These fundamental solutions depend on the value of  $E$  and so we label them as  $\{\psi_1^E, \psi_2^E\}$ . We remove the ambiguity in the coefficients by requiring

$$\begin{aligned} \psi_1^E(0) &= 1, & (\psi_1^E)'(0) &= 0 \\ \psi_2^E(0) &= 0, & (\psi_2^E)'(0) &= 1. \end{aligned}$$

**Definition.** An *entire function* is a complex valued function that is holomorphic ( $\mathbb{C}$ -differentiable in the neighbourhood of a point) at all finite points in the complex domain.  $\diamond$

**Theorem 20.6.** *The fundamental solutions  $\psi_1^E$  and  $\psi_2^E$  are entire functions on  $E$ .*

Remark 20.7. Note in order to make the above theorem true, we require that our eigenvalues are complex,  $E \in \mathbb{C}$ . This is clearly unphysical, however do this here in order to exploit the strong results of complex analysis, and then we shall restrict ourselves to  $E \in \mathbb{R}$  at the end.

**Definition.** The *fundamental matrix* of a system of linear, homogeneous ODEs, with the fundamental set of solutions  $\{\psi_1, \dots, \psi_n\}$  is the matrix

$$M(x) = \begin{pmatrix} \psi_1(x) & \dots & \psi_n(x) \\ \psi_1'(x) & \dots & \psi_n'(x) \\ \vdots & \dots & \vdots \\ \psi_1^{(n)}(x) & \dots & \psi_n^{(n)}(x) \end{pmatrix}$$

◇

**Lemma 20.8.** *The determinant fundamental matrix for our system is constant<sup>48</sup>*

$$[\det M^E]'(x) = 0.$$

*Proof.* By direct calculation, and using the generalised eigenvalue equation,

$$\begin{aligned} [\psi_1^E(\psi_2^E)' - (\psi_1^E)'\psi_2^E]'(x) &= [(\psi_1^E)'(\psi_2^E)' + \psi_1^E(\psi_2^E)'' - (\psi_2^E)'(\psi_1^E)' - (\psi_1^E)''\psi_2^E](x) \\ &= \left[ \psi_1^E \left( -\frac{2m}{\hbar^2}(E - V)\psi_2^E \right) - \left( -\frac{2m}{\hbar^2}(E - V)\psi_1^E \right) \psi_1^E \right](x) \\ &= 0. \end{aligned}$$

□

**Corollary 20.9.** *It follows trivially from the conditions we placed on  $\psi_1^E$  and  $\psi_2^E$  that  $\det M^E(x) = 1$ .*

### 20.3 Translation Operator

**Definition.** The *translation operator* is the linear operator such that its action on an element in its domain is given by

$$(T\psi)(x) := \tilde{\psi}(x) := \psi(x + a),$$

for some  $a \in \mathbb{C}$ .

◇

**Proposition 20.10.** *Let  $T$  be the translation operator with  $a$  being the periodicity of our system. Then the  $T$  commutes with the Hamiltonian,*

$$[T, H] = 0.$$

*Proof.* Consider the action on an element  $\psi$  in the codomain of both operators,

$$\begin{aligned} [T, H]\psi(x) &= T(H\psi)(x) - H(T\psi)(x) \\ &= T\left(-\frac{\hbar^2}{2m}\Delta\psi + V\psi\right)(x) - H(T\psi)(x) \\ &= \left(-\frac{\hbar^2}{2m}\Delta T\psi + VT\psi\right)(x) - H(\tilde{\psi})(x) \\ &= H(T\psi)(x) - H(T\psi)(x) \\ &= 0, \end{aligned}$$

where we have used the fact that the translation by a constant doesn't effect the result of differentiating a function and the fact that  $T$  is linear. □

**Lemma 20.11.** *Let  $\psi^E$  be a  $H$ -eigenvector of our system. Then  $\tilde{\psi}^E := T\psi^E$  is also a  $H$ -eigenvector.*

---

<sup>48</sup>Note we also label  $M$  with a superscript  $E$ .

*Proof.* From the commutation result,

$$\begin{aligned} H(T\psi^E)(x) &= T(H\psi^E)(x) \\ H\widetilde{\psi}^E(x) &= T(E\psi^E)(x) \\ &= E(T\psi^E)(x) \\ &= E\widetilde{\psi}^E(x). \end{aligned}$$

□

So we have that the translated solution is also a solution. We now use the fact that  $\{\psi_1^E, \psi_2^E\}$  is a fundamental set of solutions to expand the translated fundamental solution,

$$\psi_j^E(x+a) = \sum_{i=1}^2 a_j^i \psi_i^E(x),$$

for  $j = 1, 2$ .

Now consider the case for  $x = 0$ , then we have

$$\begin{aligned} \psi_j^E(a) &= a_j^1 \psi_1^E(0) + a_j^2 \psi_2^E(0) \\ &= a_j^1, \end{aligned}$$

and similarly

$$(\psi_j^E)'(a) = a_j^2,$$

from which it follows that

$$a_j^i = (M^E(a))^i_j.$$

So we have that a general solution is of the form

$$\psi^E(x+a) = \sum_{\ell=1}^2 \sum_{i=1}^2 A^\ell(M^E(a))^i_j \psi_i^E(x),$$

for  $A^1, A^2 \in \mathbb{C}$ .

Remark 20.12. As we showed previously the translation operator and the Hamiltonian commute, and so they share common eigenvectors. We shall label these eigenvectors as follows

$$\begin{aligned} H\psi^{E,\lambda} &= E\psi^{E,\lambda}, \\ T\psi^{E,\lambda} &= \lambda\psi^{E,\lambda}. \end{aligned}$$

We can reformulate the second eigenvalue equation as follows: using  $T\psi^{E,\lambda}(x) = \lambda\psi^{E,\lambda}(x+a)$  we have

$$\begin{aligned} \psi^{E,\lambda}(x+a) &= \lambda\psi^{E,\lambda}(x) \\ \sum_{\ell=1}^2 \sum_{i=1}^2 A_\lambda^\ell(M^E(a))^i_\ell \psi_i^E(x) &= \lambda \sum_{i=1}^2 A_\lambda^i \psi_i^E(x), \end{aligned}$$



where we have introduced a  $\lambda$  label (not an index!) on  $A$  to indicate that it corresponds to a specific  $\lambda$ . Now using the linear independence of the fundamental solutions, we can equate coefficients, then noting that the LHS is just matrix multiplication we see that we can write  $A_\lambda$  as a column matrix with two entries

$$A_\lambda = \begin{pmatrix} A_\lambda^1 \\ A_\lambda^2 \end{pmatrix},$$

which is an eigenvalue of the fundamental matrix at  $a$  with eigenvalue  $\lambda$ .

If we let  $\lambda_1$  and  $\lambda_2$  be the two eigenvalues of  $M^E(a)$ , then there is some basis such that

$$M^E(a) = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix},$$

from which we have<sup>49</sup>

$$\begin{aligned} \det M^E(a) &= \lambda_1 \cdot \lambda_2 \\ \text{Tr } M^E(a) &= \lambda_1 + \lambda_2. \end{aligned}$$

Then using the result  $\det M^E(a) = 1$ , we have the condition

$$\lambda_1 = \frac{1}{\lambda_2},$$

and we just want some way to find the trace to work out what the second condition is.

## 20.4 Application to Our Quantum Problem

**Theorem 20.13** (Floquet<sup>50</sup>). *Let  $V(x)$  be a complex, piecewise continuous, periodic function with minimum period  $\pi$ . Then the solutions to the ODE*

$$y'' + V(x)y = 0$$

*are linearly independent and can be written as*

$$\begin{aligned} f_1(x) &= e^{i\theta x} p_1(x) \\ f_2(x) &= e^{-i\theta x} p_2(x), \end{aligned}$$

*for  $\theta \in [-\pi, \pi)$  and where  $p_1(x)$  and  $p_2(x)$  are periodic with the same period as  $V(x)$ .*

Remark 20.14. Floquet's theorem also tells us that the eigenvalues are simply

$$\lambda_1 = e^{i\theta} \quad \text{and} \quad \lambda_2 = e^{-i\theta}.$$

We have, then, that

$$\psi_1^E(a) + \psi_2^E(a) = \text{Tr } M^E(a) = e^{i\theta} + e^{-i\theta} = 2 \cos \theta.$$

We now define the function

$$\gamma_V(E) := \frac{1}{2}(\psi_1^E(a) + \psi_2^E(a)) = \cos \theta,$$

where the subscript indicates that its for the type of potentials we're considering.

<sup>49</sup>Note these results are basis independent for any transformation given by an endomorphism.

<sup>50</sup>Based on a combination of the one given in the lectures and the one given on [Wolfram](#).

## 20.5 Energy Bands

Recall that  $\psi_1^E$  and  $\psi_2^E$  are entire functions of  $E$ , and so their sum is also an entire function. From this it follows that the restriction of  $\gamma_V$  to the reals,

$$\gamma_V|_{\mathbb{R}} : \mathbb{R} \rightarrow \mathbb{C},$$

is at least smooth. Thus we know that


- (i) If  $(E_0, \theta_0)$  solves the equation  $\gamma_V(E_0) = \cos \theta_0$ , then any  $E$  in a sufficiently small neighbourhood of  $E_0$  solves the equation  $\gamma_V(E) = \cos \theta$  for some  $\theta$ .
- (ii) Equivalently, if  $E_1$  does *not* solve  $\gamma_V(E_1) = \cos(\theta_1)$  for *any*  $\theta_1 \in ([\pi, \pi])$ , then no  $E$  is any sufficiently small neighbourhood will.

From these conditions we can draw the remarkable conclusion: For any periodic, piecewise continuous and bounded potential, the energy spectrum is a countable union of open intervals, known as energy bands.


*Remark 20.15.* Note the fact that we only have continuous parts to our spectrum is consistent with our rigged Hilbert space ideas; the functions  $\psi^{E,\lambda}$  contain a phase factor and then a periodic function, and so are not square integrable, but they are bounded and so are elements of  $S^*(\mathbb{R}^d) \setminus L^2(\mathbb{R}^d)$ .

## 20.6 Quantitative Calculation (Outline)

To obtain the precise intervals (bands) one needs to find (perhaps numerically) the fundamental solutions. One obtains, for instance, for a potential of the form



results in energy bands of the form



## 21 Relativistic Quantum Mechanics

As we shall see the transition into relativistic quantum mechanics is highly non-trivial; in the sense that we don't simply add a new term onto our expressions that accounts for the relativistic effects. This lecture is meant as a very brief overview/introduction to quantum field theory, and so does not claim to be self contained in any sense. The main idea we want to highlight is how the ideas change once we start accounting for relativistic effects, and what the repercussions of those changes are.

### 21.1 Heuristic Derivation of the Schrödinger Equation

Schrödinger recognised that he could obtain the Schrödinger equation from the *classical* energy-momentum relation

$$E = \frac{p^2}{2m} + V,$$

using the substitutions

$$E \rightsquigarrow i\hbar\partial_t, \quad P_a \rightsquigarrow -i\hbar\partial_a$$

giving

$$i\hbar\partial_t\psi = -\frac{\hbar^2}{2m}\partial_a\partial^a\psi + V\psi,$$

for  $\psi : \mathbb{R}^3 \rightarrow \mathbb{C}$ .

If we want to get the probabilistic interpretation that quantum mechanics is built on, we need to introduce an object that

- (i) Is non-negative definite,
- (ii) Integrates to unity,
- (iii) This integral doesn't change in time.

As we have been using all along the such needed object is

$$\rho := |\psi|^2.$$

We might ask ourselves 'How does one come up with the idea to such an object?', the answer for which comes from the following.

Firstly its clear that  $\rho \geq 0$ , by definition of the inner product. We can also always arrange for the integral over all space to be unity by normalisation. Now consider the Schrödinger equation and its complex conjugate

$$\begin{aligned} i\hbar\partial_t\psi &= -\frac{\hbar^2}{2m}\partial_a\partial^a\psi + V\psi \\ -i\hbar\partial_t\bar{\psi} &= -\frac{\hbar^2}{2m}\partial_a\partial^a\bar{\psi} + V\bar{\psi}. \end{aligned}$$

If we multiply the former from the right by  $\bar{\psi}$  and the latter from the left by  $\psi$ , then subtract the two results we arrive, after rearranging a bit, at

$$\begin{aligned} [(\partial_t \psi) \bar{\psi} + \psi (\partial_t \bar{\psi})] &= \frac{i\hbar}{2m} [\psi (\partial_a \partial^a \bar{\psi}) - (\partial_a \partial^a \psi) \bar{\psi}] \\ \partial_t (\psi \bar{\psi}) &= \frac{i\hbar}{2m} \partial_a [\psi (\partial^a \bar{\psi}) - (\partial^a \psi) \bar{\psi}]. \end{aligned}$$

Then defining

$$\rho := \psi \bar{\psi}, \quad j^a := -\frac{i\hbar}{2m} [\psi (\partial^a \bar{\psi}) - (\partial^a \psi) \bar{\psi}],$$

we arrive at the continuity equation

$$\partial_t \rho + \partial_a j^a = 0,$$

from which it follows that

$$\begin{aligned} \partial_t \int_{\mathbb{R}^3} d^3x \rho(x) &= \int_{\mathbb{R}^3} d^3x (\partial_t \rho)(x) \\ &= \int_{\mathbb{R}^3} d^3x (\partial_a j^a)(x) \\ &= 0, \end{aligned}$$

where we have used the fact that we are integrating over all of  $\mathbb{R}^3$  (which has no boundary/surface) with the fact that  $\partial_a j^a$  is a purely surface term.

## 21.2 Heuristic Derivation of the Relativistic Schrödinger Equation (Klein-Gordon Equation)

A note is made on the structural difference between non-relativistic spacetime and the spacetime of general relativity. For a much deeper discussion of this the reader is directed to Dr. Schuller's International Winter School on Gravity and Light<sup>51</sup>

Schroödinger, quite courageously, tried to apply the method of the previous section to the relativistic energy-momentum relation ( $c = 1$  here),

$$E^2 = p^2 + m^2,$$

which gives

$$-\hbar^2 \partial_t^2 = -\hbar^2 \partial_a \partial^a + m^2,$$

which, after rearranging, gives the so-called *Klein-Gordon equation*<sup>52</sup>

$$(\square + m^2)\phi = 0,$$

where we have introduced the d'Alembert operator

$$\square := \partial_t^2 - \partial_a \partial^a.$$

<sup>51</sup>Available via [YouTube](#).

<sup>52</sup>It is named such as Oskar Klein and Walter Gordan also arrived at this result after Schroödinger, and proceeded to try and interpret it as the description of relativistic electrons, which we shall see shortly is not the case.

The question we now have to ask is ‘can we still obtain some probability interpretation using  $\phi$ ?’ The answer is no, as we shall now show.

In correlation to the non-relativistic case, in order to ensure the integral is a constant in time, we wish to find a  $J^\mu$  such that

$$\partial_\mu J^\mu = \partial_0 J^0 + \partial_a J^a = 0.$$

Again similarly to before, by considering the Klein-Gordon equation and its complex conjugate we arrive at

$$J^\mu := (\partial^\mu \phi) \bar{\psi} - \psi (\partial^\mu \bar{\psi}),$$

and Gauss’ theorem tells us that the *only* candidate for the probability amplitude  $\rho$  is

$$\rho := J^0 := (\partial^0 \phi) \bar{\psi} - \psi (\partial^0 \bar{\psi}).$$

This all looks fine, in fact it looks exactly like the non-relativistic case. However there is one subtle, yet highly important, difference. In the Schrödinger equation we had only first order time derivatives, whereas the Klein-Gordon equation is second order in time derivatives. This means that for the latter we can prescribe as initial conditions not only  $\phi(t)$  but its derivative  $(\partial_t \phi)(t)$  at some time. We can thus choose these initial conditions such that  $\rho < 0$  at some time, which violates the interpretation of  $\rho$  as a probability density. This is a problem that cannot be removed at this level, the reason for which we shall soon see.

*Remark 21.1.* Historically, Schrödinger actually arrived at the relativistic equation first (as he knew this was ultimately where he wanted to go), however when running into the problem highlighted above he decided instead to consider the non-relativistic case and ended up with the Schrödinger equation.

*Remark 21.2.* Note some people often refer to the time and spatial derivatives being on an *equal footing* in the Klein-Gordon equation. This is a highly misleading choice of words. They are not on an equal footing, as the temporal derivatives come with a positive sign, whereas the spatial ones come with a negative sign. This is not just some little difference to be brushed over. Indeed, without this minus sign stems from Maxwell’s equations, and without it the Klein-Gordon equation would be physically useless; the gist being that if time and space were on an equal footing then we would not be able to predict the future, which is the main driving force of physics. What people should say is that they are on a *similar* footing.

### 21.3 Dirac Equation

So, as we have seen, it is this second derivative that causes us the problems, so the natural question is ‘How do we avoid it?’ The immediate answer is to try considering

$$E = \sqrt{p^2 + m^2},$$

and then using the substitutions as above. However this is no better as now the RHS is the square root of a differential operator, and so in the expansion about  $m^2$  we will end up with a theory with infinite spatial derivative order. Not good at all!

Dirac then asked the question ‘What if I use a different substitution prescription such that the whole of the RHS becomes a single order derivative?’ Following this thought, after several calculations, he arrives at the so-called *Dirac equation*

$$(i\gamma^\mu \partial_\mu - m\mathbb{1}_4)\Psi = 0,$$

where the  $\gamma^\mu$  are  $4 \times 4$  matrices satisfying the anticommutation relation

$$\{\gamma^\mu, \gamma^\nu\} := \gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2\eta^{\mu\nu},$$

known as the *Dirac algebra*, where  $\eta^{\mu\nu}$  is the Minkowski metric given by<sup>53</sup>


$$\eta^{\mu\nu} = \text{diag}(-1, 1, 1, 1).$$

The vector  $\Psi$  here is a 4 component object known as a spinor, where (after some work) we can think of two of the components being a particle and the other two being the associated antiparticle.

So the Dirac equation introduces antimatter into the mix, however it turns out that it still doesn’t fix the probability problem addressed by the Klein-Gordon equation!

#### 21.4 The Deep Route of the Problem

It turns out the problem stems from perhaps the most well-known equation in the world...  $E = mc^2$ , which tells us that we need not conserve particle number. For example the interaction between two particles could result in any number of resulting particles (provided the energy is in sufficiently large)



If this information is contained within  $E = mc^2$ , it is also contained in the relativistic energy-momentum equation  $E^2 = p^2 + m^2c^4$ , and so we were unjustified to look for an equation in the relativistic context that describes *one* (or any fixed number) of particles. In other words, our theory needs to account for this non-conservation of particle number, including have no particles (i.e. the vacuum). Mathematically the idea is to construct a direct sum of Hilbert spaces, each of which describes a different number of particles. This is known as the *Fock space*,

$$\mathcal{F} := \mathbb{C} \oplus \mathcal{H} \oplus (\mathcal{H} \otimes \mathcal{H}) \oplus (\mathcal{H} \otimes \mathcal{H} \otimes \mathcal{H}) \oplus \dots$$

---

<sup>53</sup>Using the  $(-, +, +, +)$  signature.

We can then construct the inner product on  $\mathcal{F}$  from the inner products on the Hilbert spaces, all of which are obtained from the inner product on  $\mathcal{H}$ . That is if  $\psi, \varphi \in \mathcal{F}$  given by

$$\begin{aligned}\psi &= a_0 \oplus a_1 \psi_1 \oplus \sum_{ij} a_2^{ij} (\psi_{2i} \otimes \psi_{2j}) \oplus \dots \\ \varphi &= b_0 \oplus b_1 \varphi_1 \oplus \sum_{ij} b_2^{ij} (\varphi_{2i} \otimes \varphi_{2j}) \oplus \dots,\end{aligned}$$

then the inner product is given by

$$\langle \psi | \varphi \rangle = \overline{a_0} b_0 + \overline{a_1} b_1 \langle \psi_1 | \varphi_1 \rangle_{\mathcal{H}} + \sum_{ijkl} \overline{a_2^{ij}} b_2^{kl} \langle \psi_{2i} \otimes \psi_{2j} | \varphi_{2k} \otimes \varphi_{2l} \rangle_{\mathcal{H} \otimes \mathcal{H}} + \dots$$

*Remark 21.3.* For the Klein-Gordon equation it is actually the *symmetrised* tensor products we need, giving this symmetric Fock space

$$\odot \mathcal{F} := \mathbb{C} \oplus (\mathcal{H} \odot \mathcal{H}) \oplus (\mathcal{H} \odot \mathcal{H} \odot \mathcal{H}) \oplus \dots,$$

and it describes *bosonic* systems.

Similarly for the Dirac equation we require the *anti-symmetrised* tensor product, giving

$$\wedge \mathcal{F} := \mathbb{C} \oplus (\mathcal{H} \wedge \mathcal{H}) \oplus (\mathcal{H} \wedge \mathcal{H} \wedge \mathcal{H}) \oplus \dots,$$


which describes *fermionic* systems.

We claim that once you lift the Klein-Gordon and Dirac equations onto their respective Fock spaces, that the problem of negative probability density vanishes.

## 21.5 Feynman Diagrams

Richard Feynman developed a pictorial representation of the highly complicated interaction of particles. The basic idea is to take a perturbation expansion of the interaction and represent each order by a set of diagrams. The order of the diagram is associated with the number of so-called *vertices* present.

For example the first few terms for the process of electron-electron scattering would be drawn as:



The first diagram (which has no vertices) is the zeroth order diagram, the second one (with two vertices) is the second order diagram and the last two (which both have 4 vertices) are the fourth order diagrams. The furthest most left and right arrows (i.e. the ones that have a non-vertexed end) are known as *external lines*, and the other ones are known as *internal lines*. Particles represented by internal lines are often referred to *virtual* particles.

Remark 21.4. One should be careful when it comes to drawing the arrows on the internal lines, however, as (unless the rest of the diagram indicates otherwise) we could have a particle or an antiparticle (whose arrow points the opposite way). It is for this reason that the so-called *loop* in the third diagram does not have arrows. On the final diagram we do draw the arrows, as conservation of electric charge forces us to ensure our virtual particles are electrons (not positrons, the anti-electron).

Note also on loop internal lines we have simply written  $e$  and not  $e^-$  or  $e^+$  (the positron), this further indicates that we do not know which is which, only that one must be an electron and the other a positron.

Mathematically Feynman diagrams correspond to integral equations, and there is a set of rules (cleverly named *Feynman Rules*) which tell you how to convert the diagrams into these integrals. They are a indispensable tool when it comes to studying relativistic particle physics, as not only are they quicker to draw then writing out integrals, they have some incredibly elegant properties (such as so-called *crossing symmetry*) which make the calculations significantly easier. However, we shall not go into any more detail here; the unfamiliar reader is directed to the massive resource of information in textbooks/on the internet.



## FURTHER READINGS

### Mathematical quantum mechanics

- Ballentine, *Quantum Mechanics: A Modern Development* (Second edition), World Scientific 2014
- Faddeev, Yakubovskii, *Lectures on Quantum Mechanics for Mathematics Students*, American Mathematical Society 2009
- Folland, *Quantum Field Theory: A Tourist Guide for Mathematicians*, American Mathematical Society 2008
- Gieres, *Mathematical surprises and Dirac's formalism in quantum mechanics*  
<https://arxiv.org/abs/quant-ph/9907069>
- Hall, *Quantum Theory for Mathematicians*, Springer 2013
- Mackey, *Mathematical Foundations of Quantum Mechanics*, Dover Publications 2004
- Moretti, *Spectral Theory and Quantum Mechanics: With an Introduction to the Algebraic Formulation*, Springer 2013
- Parthasarathy, *Mathematical Foundations of Quantum Mechanics*, Hindustan Book Agency 2005
- Strocchi, *An Introduction to the Mathematical Structure of Quantum Mechanics: A Short Course for Mathematicians*, World Scientific 2008
- Takhtajan, *Quantum Mechanics for Mathematicians*, American Mathematical Society 2008

### Standard quantum mechanics textbooks

- Griffiths, *Introduction to Quantum Mechanics* (Second edition), Cambridge University Press 2016
- Nazarov & Danon, *Advanced Quantum Mechanics*, Cambridge University Press, 2013

### Linear Algebra

- Friedberg, Insel, Spence, *Linear Algebra* (4th Edition), Pearson 2002
- Jänich, *Linear algebra*, Springer 1994
- Lang, *Linear Algebra* (Third edition), Springer 1987
- Shakarchi, *Solutions Manual for Lang's Linear Algebra*, Springer 1996

## Topology

- Adamson, *A General Topology Workbook*, Birkhäuser 1995
- Kalajdziewski, *An Illustrated Introduction to Topology and Homotopy*, CRC Press 2015
- Munkres, *Topology* (Second edition), Pearson 2014

## Functional analysis

- Aliprantis, Burkinshaw, *Principles of Real Analysis (Third Edition)*, Academic Press 1998
- Aliprantis, Burkinshaw, *Problems in Real Analysis: A Workbook with Solutions*, Academic Press 1998
- Day, *Normed Linear Spaces*, Springer 1973
- Halmos, *A Hilbert Space Problem Book*, Springer 1982
- Hunter, Nachtergaele, *Applied Analysis*, World Scientific, 2001
- Kadison, Ringrose, *Fundamentals of the Theory of Operator Algebras. Volumes I-II*, American Mathematical Society 1997
- Leoni, *A first Course in Sobolev Spaces*, American Mathematical Society 2009
- Rynne, Youngson, *Linear Functional Analysis (Second Edition)*, Springer 2008
- Serov, *Fourier Series, Fourier Transform and Their Applications to Mathematical Physics* Springer 2017

## Measure theory and Integration

- Bartle, *A Modern Theory of Integration*, American Mathematical Society 2001
- Bartle, *Solutions Manual to A Modern Theory of Integration*, American Mathematical Society 2001
- Halmos, *Measure Theory*, Springer 1982
- Nelson, *A User-friendly Introduction to Lebesgue Measure and Integration*, American Mathematical Society 2015
- Rana, *An Introduction to Measure and Integration*, American Mathematical Society 2002