A - MELANOME :

I- Maladie localisée :

PEMBROLIZUMAB J1=J21 [1-4]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Immunothérapie	Pembrolizumab	Intraveineuse	200.0 mg	Dans 200 cc de SS 0,9%

*[1] Luke JJ, et al. *Pembrolizumab versus Placebo as Adjuvant Therapy in Resected Stage IIB or IIC Melanoma (KEYNOTE-716): A Randomised, Double-Blind, Phase 3 Trial*. Lancet. 2021;398(10311):1192-1203.*

[2] Eggermont AMM, et al. Adjuvant Pembrolizumab versus Placebo in Resected Stage III Melanoma (KEYNOTE-054). N Engl J Med. 2018;378(19):1789-1801.

[3] Patel SP, et al. Neoadjuvant-Adjuvant or Adjuvant-Only Pembrolizumab in Advanced Melanoma. N Engl J Med. 2023;388(9):813-823.

[4] Study of mRNA-4157 (V940) in Combination With Pembrolizumab (MK-3475) in Participants With Melanoma (V940-001). ClinicalTrials.gov. 2023.*

Nivolumab/ipilimumab J1=J21 [5-10]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Immunothérapie	Ipilimumab	Intraveineuse	1 mg/kg	Dans 50 cc de SS 0,9% en 30 minutes
1	Immunothérapie	Nivolumab	Intraveineuse	3 mg/kg	Dans 100 cc de SS 0.9% en 30 minutes

[5] Reijers ILM, Menzies AM, van Akkooi ACJ, et al. Personalized response-directed surgery and adjuvant therapy after neoadjuvant ipilimumab and nivolumab in high-risk stage III melanoma: the PRADO trial. Nat Med 2022;28:1178-1188. 3 Blank CU, Lucas MW,

[6] Scolyer RA, et al. Neoadjuvant Nivolumab and Ipilimumab in Resectable Stage III Melanoma N Engl J Med 2024;391:1696-1708.

[7] Versluis JM, Menzies AM, Sikorska K, et al. Survival update of neoadjuvant ipilimumab plus nivolumab in macroscopic stage III melanoma in the OpACIN and OpACIN-neo trials. Ann Oncol 2023;34:420-430.

[8] Blank CU, Reijers L, Saw RPM, et al. Survival data of PRADO: a phase 2 study of personalized response-driven surgery and adjuvant therapy after neoadjuvant ipilimumab (IPI) and nivolumab (NIVO) in resectable stage III melanoma [abstract]. J Clin Oncol 2022;40(Suppl): Abstract 9501.

[9] Versluis JM, Menzies AM, Sikorska K, et al. Survival update of neoadjuvant ipilimumab plus nivolumab in macroscopic stage III melanoma in the OpACIN and OpACIN-neo trials. Ann Oncol 2023;34:420-430.

[10] Reijers ILM, Menzies AM, Versluis JM, et al. The impact of response-directed surgery and adjuvant therapy on long-term survival after neoadjuvant ipilimumab plus nivolumab in stage III melanoma: Three-year data of PRADO and OpACIN-neo [abstract]. J Clin Oncol 2023;41(Suppl): Abstract 101.

II- Maladie avancée/ métastatique :

Dacarbazine J1=J21 [11]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Pré-médication	Chlorure De Sodium	Intraveineuse	250.0 ml	0.9% 250 cc SS en 30 min
2	Pré-médication	Methylprednisolone	Intraveineuse	120.0 mg	IVL 30min avant chimiothérapie
3	Pré-médication	Ondansetron	Intraveineuse	8.0 mg	IVL 30min avant chimiothérapie
4	Pré-médication	Metoclopramide	Intraveineuse	10.0 mg	IVL
5	Chimiothérapie	Dacarbazine	Intraveineuse	850.0 mg/M2	dans 250 cc SS ou SG en 30 min puis rinçage
6	Rinçage	Chlorure De Sodium	Intraveineuse	250.0 ml	0.9% 250 cc SS en 30 min

[11] Huncharek M, Caubert JF, McGarry R. Single agent DTIC versus combination chemotherapy with or without immunotherapy in metastatic melanoma: a meta-analysis of 3273 patients from 20 randomised trials. Melanoma Research 2001; 11 (1): 75-81.

CARBOPLATINE-PACLITAXEL J1=J21 [2,3]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Pré-médication	Omeprazole	Orale	20.0 mg	20mg 1 cp le jour de la chimio
1	Pré-médication	Anti Histaminiques H1	Orale	10.0 mg	1cp 4h avant Paclitaxel
2	Pré-médication	Methylprednisolone	Intraveineuse	120.0 mg	IVL 30min avant chimiothérapie
3	Pré-médication	Ondansetron	Intraveineuse	8.0 mg	IVL 30min avant chimiothérapie
4	Chimiothérapie	Paclitaxel	Intraveineuse	175.0 mg/M2	Dans 500 ml SG 5% ou SS 0.9% en 3h
5	Chimiothérapie	Carboplatine	Intraveineuse	AUC 5.0	AUC 5 100 ml SG 5% puis rinçage

[12] ECOG E2603: Flaherty KT, Lee SJ, Zhao F, Schuchter LM, Flaherty L, Kefford R, Atkins MB, Leming P, Kirkwood JM. Phase III trial of carboplatin and paclitaxel with or without sorafenib in metastatic melanoma. J Clin Oncol. 2013 Jan 20;31(3):373-9. Epub 2012 Dec 17.

[13] BCH-MM-131101: Yan X, Sheng X, Chi Z, Si L, Cui C, Kong Y, Tang B, Mao L, Wang X, Lian B, Li S, Bai X, Zhou L, Dai J, Yao H, Guo J. Randomized Phase II Study of Bevacizumab in Combination With Carboplatin Plus Paclitaxel in Patients With Previously Untreated Advanced Mucosal Melanoma. J Clin Oncol. 2021 Mar 10;39(8):881-889. Epub 2021 Jan 14.

PEMBROLIZUMAB J1=J21 [14-17]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Immunothérapie	Pembrolizumab	Intraveineuse	200.0 mg	Dans 200 cc de SS 0,9%

[14] KEYNOTE-001melanoma: Hamid O, Robert C, Daud A, Hodi FS, Hwu WJ, Kefford R, Wolchok JD, Hersey P, Joseph RW, Weber JS, Dronca R, Gangadhar TC, Patnaik A, Zarour H, Joshua AM, Gergich K, Elassaiss-Schaap J, Algazi A, Mateus C, Boasberg P, Tumeh PC, Chmielowski B, Ebbinghaus SW, Li XN, Kang SP, Ribas A. Safety and tumor responses with lambrolizumab (anti-PD-1) in melanoma. N Engl J Med. 2013 Jul 11;369(2):134-44. Epub 2013 Jun 2.

A. Update: Robert C, Ribas A, Wolchok JD, Hodi FS, Hamid O, Kefford R, Weber JS, Joshua AM, Hwu WJ, Gangadhar TC, Patnaik A, Dronca R, Zarour H, Joseph RW, Boasberg P, Chmielowski B, Mateus C, Postow MA, Gergich K, Elassaiss-Schaap J, Li XN, Iannone R, Ebbinghaus SW, Kang SP, Daud A. Anti-programmed-death-receptor-1 treatment with pembrolizumab in ipilimumab-refractory advanced melanoma: a randomised dose-comparison cohort of a phase 1 trial. Lancet. 2014 Sep 20;384(9948):1109-17. Epub 2014 Jul 14.

B. Update: Ribas A, Hamid O, Daud A, Hodi FS, Wolchok JD, Kefford R, Joshua AM, Patnaik A, Hwu WJ, Weber JS, Gangadhar TC, Hersey P, Dronca R, Joseph RW, Zarour H, Chmielowski B, Lawrence DP, Algazi A, Rizvi NA, Hoffner B, Mateus C, Gergich K, Lindia JA, Giannotti M, Li XN, Ebbinghaus S, Kang SP, Robert C. Association of pembrolizumab with tumor response and survival among patients with advanced melanoma. JAMA. 2016 Apr 19;315(15):1600-9. Erratum in: JAMA. 2016 Jun 14;315(22):2472.

C. Update: Hamid O, Robert C, Daud A, Hodi FS, Hwu WJ, Kefford R, Wolchok JD, Hersey P, Joseph R, Weber JS, Dronca R, Mitchell TC, Patnaik A, Zarour HM, Joshua AM, Zhao Q, Jensen E, Ahsan S, Ibrahim N, Ribas A. Five-year survival outcomes for patients with advanced melanoma treated with pembrolizumab in KEYNOTE-001. Ann Oncol. 2019 Apr 1;30(4):582-588.

[15]KEYNOTE-002: Ribas A, Puzanov I, Dummer R, Schadendorf D, Hamid O, Robert C, Hodi FS, Schachter J, Pavlick AC, Lewis KD, Cranmer LD, Blank CU, O'Day SJ, Ascierto PA, Salama AK, Margolin KA, Loquai C, Eigentler TK, Gangadhar TC, Carlino MS, Agarwala SS, Moschos SJ, Sosman JA, Goldinger SM, Shapira-Frommer R, Gonzalez R, Kirkwood JM, Wolchok JD, Eggermont A, Li XN, Zhou W, Zernhelt AM, Lis J, Ebbinghaus S, Kang SP, Daud A. Pembrolizumab versus investigator-choice chemotherapy for ipilimumab-refractory melanoma (KEYNOTE-002): a randomised, controlled, phase 2 trial. Lancet Oncol. 2015 Aug;16(8):908-18. Epub 2015 Jun 23.

a. HRQoL analysis: Schadendorf D, Dummer R, Hauschild A, Robert C, Hamid O, Daud A, van den Eertwegh A, Cranmer L, O'Day S, Puzanov I, Schachter J, Blank C, Salama A, Loquai C, Mehnert JM, Hille D, Ebbinghaus S, Kang SP, Zhou W, Ribas A. Health-related quality of life in the randomised KEYNOTE-002 study of pembrolizumab versus chemotherapy in patients with ipilimumab-refractory melanoma. Eur J Cancer. 2016 Nov;67:46-54. Epub 2016 Sep 2.

b. Update: Hamid O, Puzanov I, Dummer R, Schachter J, Daud A, Schadendorf D, Blank C, Cranmer LD, Robert C, Pavlick AC, Gonzalez R, Hodi FS, Ascierto PA, Salama AKS, Margolin KA, Gangadhar TC, Wei Z, Ebbinghaus S, Ibrahim N, Ribas A. Final analysis of a randomised trial comparing pembrolizumab versus investigator-choice chemotherapy for ipilimumab-refractory advanced melanoma. Eur J Cancer. 2017 Nov;86:37-45.

[16]KEYNOTE-006: Robert C, Schachter J, Long GV, Arance A, Grob JJ, Mortier L, Daud A, Carlino MS, McNeil C, Lotem M, Larkin J, Lorigan P, Neyns B, Blank CU, Hamid O, Mateus C, Shapira-Frommer R, Kosh M, Zhou H, Ibrahim N, Ebbinghaus S, Ribas A; KEYNOTE-006 investigators. Pembrolizumab versus ipilimumab in advanced melanoma. N Engl J Med. 2015 Jun 25;372(26):2521-32. Epub 2015 Apr 19.

a. Update: Schachter J, Ribas A, Long GV, Arance A, Grob JJ, Mortier L, Daud A, Carlino MS, McNeil C, Lotem M, Larkin J, Lorigan P, Neyns B, Blank C, Petrella TM, Hamid O, Zhou H, Ebbinghaus S, Ibrahim N, Robert C. Pembrolizumab versus ipilimumab for advanced melanoma: final overall survival results of a multicentre, randomised, open-label phase 3 study (KEYNOTE-006). Lancet. 2017 Oct 21;390(10105):1853-1862. Epub 2017 Aug 16.

b. PRO analysis: Petrella TM, Robert C, Richtig E, Miller WH Jr, Masucci GV, Walpole E, Lebbe C, Steven N, Middleton MR, Hille D, Zhou W, Ibrahim N, Cebon J. Patient-reported outcomes in KEYNOTE-006, a randomised study of pembrolizumab versus ipilimumab in patients with advanced melanoma. Eur J Cancer. 2017 Nov;86:115-124. Epub 2017 Oct 4.

c. Update: Robert C, Ribas A, Schachter J, Arance A, Grob JJ, Mortier L, Daud A, Carlino MS, McNeil CM, Lotem M, Larkin JMG, Lorigan P, Neyns B, Blank CU, Petrella TM, Hamid O, Su SC, Krepler C, Ibrahim N, Long GV. Pembrolizumab versus ipilimumab in advanced melanoma (KEYNOTE-006): post-hoc 5-year results from an open-label, multicentre, randomised, controlled, phase 3 study. Lancet Oncol. 2019 Sep;20(9):1239-1251. Epub 2019 Jul 22.

d. Update: Robert C, Carlino MS, McNeil C, Ribas A, Grob JJ, Schachter J, Nyakas M, Kee D, Petrella TM, Blaustein A, Lotem M, Arance A, Daud AI, Hamid O, Larkin J, Anderson J, Krepler C, Grebennik D, Long GV. Seven-Year Follow-Up of the Phase III KEYNOTE-006 Study: Pembrolizumab Versus Ipilimumab in Advanced Melanoma. J Clin Oncol. 2023 Aug 20;41(24):3998-4003. Epub 2023 Jun 22.

[17] ECHO-301/KEYNOTE-252: Long GV, Dummer R, Hamid O, Gajewski TF, Caglevic C, Dalle S, Arance A, Carlino MS, Grob JJ, Kim TM, Demidov L, Robert C, Larkin J, Anderson JR, Maleski J, Jones M, Diede SJ, Mitchell TC. Epacadostat plus pembrolizumab versus placebo plus pembrolizumab in patients with unresectable or metastatic melanoma (ECHO-301/KEYNOTE-252): a phase 3, randomised, double-blind study. Lancet Oncol. 2019 Aug;20(8):1083-1097. Epub 2019 Jun 17.

IPILIMUMAB J1=J21 [18 - 23]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Immunothérapie	Ipilimumab	Intraveineuse	3 mg/kg	Dans 200 cc de SS 0,9% ou SG 5% en 90 minutes

[18] Margolin K, Ernstoff MS, Hamid O, et al. Ipilimumab in patients with melanoma and brain metastases: an open-label, phase 2 trial. Lancet Oncol 2012;13:459- 465.

[19] Weber JS, Kahler KC, Hauschild A. Management of immune-related adverse events and kinetics of response with ipilimumab. J Clin Oncol 2012;30:2691- 2697.

[20] Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with metastatic melanoma. N Eng J Med 2010;363:711-723.

[21] Robert C, Thomas L, Bondarenko I, et al. Ipilimumab plus dacarbazine for previously untreated metastatic melanoma. N
Engl J Med 2011;364:2517-2526.

[22] Maio M, Grob JJ, Aamdal S, et al. Five-year survival rates for treatment-naive patients with advanced melanoma who received ipilimumab plus dacarbazine in a phase III trial. J Clin Oncol 2015;33:1191-1196.

[23] Ascierto PA, Del Vecchio M, Robert C, et al. Ipilimumab 10 mg/kg versus ipilimumab 3 mg/kg in patients with unresectable or metastatic melanoma: a randomised, double-blind, multicentre, phase 3 trial. Lancet Oncol 2017;18:611- 622.

Nivolumab J1=J28 [24 - 26]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Immunothérapie	Nivolumab	Intraveineuse	480 mg	Dans 250 cc de SS 0,9% en 60 minutes

[24] Weber JS, D'Angelo SP, Minor D, et al. Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): a randomised, controlled, open-label, phase 3 trial. Lancet Oncol 2015;16:375-384.

[25] Larkin J, Minor D, D'Angelo S, et al. Overall survival in patients with advanced melanoma who received nivolumab versus investigator's choice chemotherapy in CheckMate 037: A randomized, controlled, open-label phase III trial. J Clin Oncol 2018;36:383-390.

[26] Robert C, Long GV, Brady B, et al. Nivolumab in previously untreated melanoma without BRAF mutation. N Engl J Med 2015;372:320-330.

Nivolumab J1=J14 [24 - 26]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Immunothérapie	Nivolumab	Intraveineuse	240 mg	Dans 100 cc de SS 0,9% en 30 minutes

[24] Weber JS, D'Angelo SP, Minor D, et al. Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): a randomised, controlled, open-label, phase 3 trial. Lancet Oncol 2015;16:375-384.

[25] Larkin J, Minor D, D'Angelo S, et al. Overall survival in patients with advanced melanoma who received nivolumab versus investigator's choice chemotherapy in CheckMate 037: A randomized, controlled, open-label phase III trial. J Clin Oncol 2018;36:383-390.

[26] Robert C, Long GV, Brady B, et al. Nivolumab in previously untreated melanoma without BRAF mutation. N Engl J Med 2015;372:320-330.

Nivolumab/ipilimumab J1=J21 [18 - 26]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Immunothérapie	Ipilimumab	Intraveineuse	1 mg/kg	Dans 50 cc de SS 0,9% en 30 minutes
1	Immunothérapie	Nivolumab	Intraveineuse	3 mg/kg	Dans 100 cc de SS 0.9% en 30 minutes

[18] Margolin K, Ernstoff MS, Hamid O, et al. Ipilimumab in patients with melanoma and brain metastases: an open-label, phase 2 trial. Lancet Oncol 2012;13:459- 465.

[19] Weber JS, Kahler KC, Hauschild A. Management of immune-related adverse events and kinetics of response with ipilimumab. J Clin Oncol 2012;30:2691- 2697.

[20] Hodi FS, O'Day SJ, McDermott DF, et al. Improved survival with ipilimumab in patients with metastatic melanoma. N Eng J Med 2010;363:711-723.

[21] Robert C, Thomas L, Bondarenko I, et al. Ipilimumab plus dacarbazine for previously untreated metastatic melanoma. N
Engl J Med 2011;364:2517-2526.

[22] Maio M, Grob JJ, Aamdal S, et al. Five-year survival rates for treatment-naive patients with advanced melanoma who received ipilimumab plus dacarbazine in a phase III trial. J Clin Oncol 2015;33:1191-1196.

[23] Ascierto PA, Del Vecchio M, Robert C, et al. Ipilimumab 10 mg/kg versus ipilimumab 3 mg/kg in patients with unresectable or metastatic melanoma: a randomised, double-blind, multicentre, phase 3 trial. Lancet Oncol 2017;18:611- 622.

[24] Weber JS, D'Angelo SP, Minor D, et al. Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): a randomised, controlled, open-label, phase 3 trial. Lancet Oncol 2015;16:375-384.

[25] Larkin J, Minor D, D'Angelo S, et al. Overall survival in patients with advanced melanoma who received nivolumab versus investigator's choice chemotherapy in CheckMate 037: A randomized, controlled, open-label phase III trial. J Clin Oncol 2018;36:383-390.

[26] Robert C, Long GV, Brady B, et al. Nivolumab in previously untreated melanoma without BRAF mutation. N Engl J Med 2015;372:320-330.

Dabrafenib/ Trametinib J1=J28 [27 - 33]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Thérapie ciblée	Dabrafenib	Orale	300 mg	Si présentation 50mg : 3cp x 2/jour à prendre à distance des repas (1 heure avant ou 2 heures après un repas). Si présentation 75mg : 2cp x 2/j à prendre à distance des repas (1 heure avant ou 2 heures après un repas).
1	Thérapie ciblée	Trametinib	orale	2mg	1 cp/jour à prendre à distance des repas (1 heure avant ou 2 heures après un repas)

[27] Long GV, Stroyakovskiy D, Gogas H, et al. Dabrafenib and trametinib versus dabrafenib and placebo for Val600 BRAF-mutant melanoma: a multicentre, double-blind, phase 3 randomised controlled trial. Lancet 2015;386:444-451.

[28] Long GV, Flaherty KT, Stroyakovskiy D, et al. Dabrafenib plus trametinib versus dabrafenib monotherapy in patients with metastatic BRAF V600E/K-mutant melanoma: long-term survival and safety analysis of a phase 3 study. Ann Oncol 2017;28:1631-1639.

[29] Robert C, Karaszewska B, Schachter J, et al. Improved overall survival in melanoma with combined dabrafenib and trametinib. N Engl J Med 2015;372:30-39.

[30] Davies MA, Saiag P, Robert C, et al. Dabrafenib plus trametinib in patients with BRAF(V600)-mutant melanoma brain metastases (COMBI-MB): a multicentre, multicohort, open-label, phase 2 trial. Lancet Oncol 2017;18:863-873.

[31] Long GV, Weber JS, Infante JR, et al. Overall survival and durable responses in patients with BRAF V600-mutant metastatic melanoma receiving dabrafenib combined with trametinib. J Clin Oncol 2016;34:871-878. Erratum in: J Clin Oncol 2019;37:355.

[32] Long GV, Eroglu Z, Infante J, et al. Long-term outcomes in patients with BRAF V600-mutant metastatic melanoma who received dabrafenib combined with trametinib. J Clin Oncol 2018;36:667-673.

[33] Hauschild A, Grob JJ, Demidov LV, et al. Dabrafenib in BRAF-mutated metastatic melanoma: a multicentre, open-label, phase 3 randomised controlled trial. Lancet 2012;380:358-365.

Dabrafenib J1=J28 [33]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Thérapie ciblée	Dabrafenib	Orale	300 mg	Si présentation 50mg : 3cp x 2/j à prendre à distance des repas (1 heure avant ou 2 heures après un repas). Si présentation 75mg : 2cp x 2/j à prendre à distance des repas (1 heure avant ou 2 heures après un repas).

[33] Hauschild A, Grob JJ, Demidov LV, et al. Dabrafenib in BRAF-mutated metastatic melanoma: a multi-centre, open-label, phase 3 randomised controlled trial. Lancet 2012;380:358-365.

Vemurafenib J1=J28 [34 - 37]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Thérapie Ciblée	Vemurafenib	Orale	1920 Mg	4Cp X 2/Jour Avant Ou Après Repas

324] Sosman JA, Kim KB, Schuchter L, et al. Survival in BRAF V600-mutant advanced melanoma treated with vemurafenib. N Engl J Med 2012;366:707-714.

[35] McArthur GA, Chapman PB, Robert C, et al. Safety and efficacy of vemurafenib in BRAF(V600E) and BRAF(V600K) mutation-positive melanoma (BRIM-3): extended follow-up of a phase 3, randomised, open-label study. Lancet Oncol 2014;15:323-332.

[36] Chapman PB, Robert C, Larkin J, et al. Vemurafenib in patients with BRAFV600 mutation-positive metastatic melanoma: final overall survival results of the randomized BRIM-3 study. Ann Oncol 2017;28:2581-2587.

[37] McArthur GA, Maio M, Arance A, et al. Vemurafenib in metastatic melanoma patients with brain metastases: an open-label, single-arm, phase 2, multicentre study. Ann Oncol 2017;28:634-641.

Vemurafenib/ Cobimetinib J1=J28 [38-40]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Thérapie ciblée	Vemurafenib	Orale	1920 mg	4cp x 2/jour avant ou après repas
1	Thérapie ciblée	Cobimetinib	Orale	60 mg	3 cp/jour en une prise, avant ou après repas, pendant 21 jours, puis repos de 07 jours

[38] Larkin J, Ascierto PA, Dreno B, et al. Combined vemurafenib and cobimetinib in BRAF-mutated melanoma. N Engl J Med 2014;371:1867-1876.

[39] Ascierto PA, McArthur GA, Dreno B, et al. Cobimetinib combined with vemurafenib in advanced BRAF(V600)-mutant melanoma (coBRIM): updated efficacy results from a randomised, double-blind, phase 3 trial. Lancet Oncol 2016;17:1248-1260.

[40] Ribas A, Gonzalez R, Pavlick A, et al. Combination of vemurafenib and cobimetinib in patients with advanced BRAF(V600)-mutated melanoma: a phase 1b study. Lancet Oncol 2014;15:954-965.

Encorafenib/ Binimetinib J1=J28 [41-42]

Ordre	Type	Produit / Protocole	Voie	Dose	Observation
J1					
1	Thérapie ciblée	Encorafenib	Orale	450.0 mg	6 cp de 75mg/j en une prise, à distance des repas (1h avant ou 2h après un repas)
1	Thérapie ciblée	Binimetinib	Orale	90.0 mg	3 cp de 15mg x2/jour avant ou après repas.

[41] Dummer R, Ascierto PA, Gogas HJ, et al. Encorafenib plus binimetinib versus vemurafenib or encorafenib in patients with BRAF-mutant melanoma (COLUMBUS): a multicentre, open-label, randomised phase 3 trial. Lancet Oncol 2018;19:603-615.

[42] Dummer R, Ascierto PA, Gogas HJ, et al. Overall survival in patients with BRAFmutant melanoma receiving encorafenib plus binimetinib versus vemurafenib or encorafenib (COLUMBUS): a multicentre, open-label, randomised, phase 3 trial. Lancet Oncol 2018;19:1315-1327.